


Solutions

1. a) $-2xe^{-x^2} \tan x + e^{-x^2} \sec^2 x$
 b) $f'(u) = 2 \cos(u^2 + 2u) \cdot -\sin(u^2 + 2u) \cdot (2u + 2)$; $f'(1) = -2 \cos 3 \cdot \sin 3 \cdot 4 = -4 \sin 6$
 c) $\frac{(x^2 + 1) \cdot 2(2x + 1) \cdot 2 - (2x + 1)^2 \cdot 2x}{(x^2 + 1)^2} \Big|_{x=1} = \frac{24 - 18}{4} = \frac{3}{2}$

2. Differentiating implicitly, $y^3 + 3xy^2y' - 4xy - 2x^2y' + 5 = 0$
 $\Rightarrow 1 + 6y' - 8 - 8y' + 5 = 0$ at $(2, 1) \Rightarrow y'(2) = -1$;
 tangent line: $y - 1 = -(x - 2)$, or $y = -x + 3$.




3. $y' = 6x^2 - 6x$; $= 0$ if $x = 0, 1$; $y(0) = 1$, max; $y(1) = 0$, min
 $y'' = 12x - 6$; $= 0$ if $x = \frac{1}{2}$; $y(\frac{1}{2}) = \frac{1}{2}$, point of inflection

4. Maximize $A = xy$, where $2x + y = 48$.

$$A = xy = x(48 - 2x) = 48x - 2x^2 \Rightarrow A' = 48 - 4x;$$

$$A' = 0 \text{ if } x = 12; \text{ then } y = 24, \text{ and } A = 288 \text{ sq. feet.}$$

Since $x = 0$ or $y = 0$ gives $A = 0$, the above must be a maximum.



5. $\lim_{h \rightarrow 0} \frac{(10 + h)^7 - 10^7}{h} = \frac{d}{dx} x^7 \Big|_{x=10} = 7 \cdot 10^6$

6. a) cont., but not diff. b) neither, since undefined at $x = 0$
 c) both, since the two functions have the same value 1 and the same derivative 0 when $x = 0$.

7. Let $V =$ volume of oil, $r =$ disc radius, $h =$ disc thickness; then $V = \pi r^2 h$.

Using mks units: data: when $r = 1$, $\frac{dh}{dt} = -.004$; to find: $\frac{dh}{dt}$, when $r = 2$.

$$h = \frac{V}{\pi r^2} \Rightarrow \frac{dh}{dt} = -\frac{2V}{\pi r^3} \cdot \frac{dr}{dt}. \text{ Substituting in the data: } -.004 = -\frac{2V}{\pi} \cdot .1 \Rightarrow \frac{2V}{\pi} = .04.$$

Therefore when $r = 2$, $\frac{dh}{dt} = -\frac{2V}{\pi \cdot 8} \cdot .1 = -.0005$; so thickness decreases at .5 mm/sec.

8. Separating variables, the equation becomes $e^y dy = e^{2x} dx$.

Integrating both sides, $e^y = \frac{1}{2}e^{2x} + c$; substituting, $y(0) = 0 \Rightarrow c = \frac{1}{2}$.

Solving for y , $y = \ln \frac{1}{2}(e^{2x} + 1) = \ln(e^{2x} + 1) - \ln 2$.

9. $\int_0^1 x^3(x^4 + 3)^{1/2} dx = \frac{1}{6}(x^4 + 3)^{3/2} \Big|_0^1 = \frac{1}{6}(8 - 3\sqrt{3})$. (Put $u = x^4 + 3$, so $du = 4x^3$.)

10. Graph is above x -axis for $x \geq 1$. Area $= \int_1^2 \frac{\ln^3 x}{x} dx = \frac{1}{4} \ln^4 x \Big|_1^2 = \frac{1}{4}(\ln 2)^4$.

11. Volume $= \int_0^a \pi \cdot \frac{1}{(2x + 1)^2} dx = \frac{-\pi/2}{2x + 1} \Big|_0^a = \frac{\pi}{2} \left(1 - \frac{1}{2a + 1}\right)$, which $\rightarrow \frac{\pi}{2}$, as $a \rightarrow \infty$.

12. $\int_0^{\pi/2} \sin x dx \approx \frac{\pi}{4} \left(\frac{1}{2} \sin 0 + \sin \frac{\pi}{4} + \frac{1}{2} \sin \frac{\pi}{2}\right) = \frac{\pi}{4} \left(\frac{\sqrt{2}}{2} + \frac{1}{2}\right) \approx 3.1 \cdot \frac{2.4}{8} \approx .9$