

18.156 Differential Analysis II

Lecture 21

Instructor: Larry Guth
Trans.: Kevin Sackel

Lecture 21: 1 April 2015

21.1 Finishing the Sierpinski Theorem

Recall from our wishful thinking that we wished in part to bound $|\widehat{Pf}(n)|$ for $n \in \mathbb{Z}^2$. We also had $\widehat{Pf}(n) = \widehat{f}(n)$, so the following estimate is useful.

Proposition 1. *If $f = \chi_{B(R)}$ in \mathbb{R}^2 , then*

$$|\widehat{f}(w)| \lesssim CR^{1/2}|w|^{-3/2}.$$

Proof Sketch. We have that by rotational invariance,

$$\begin{aligned} |\widehat{f}(w)| &= \left| \int_{B(R)} e^{iw \cdot x} dx \right| \\ &= \left| \int_{B(R)} e^{i|w|x_1} dx_1 dx_2 \right| \\ &= 2 \left| \int_{-R}^R (R^2 - x^2)^{1/2} e^{i|w|x} dx \right| \end{aligned}$$

We first integrate by parts with $u = (R^2 - x^2)^{1/2}$ and $dv = e^{iwx} dx$ to obtain

$$|\widehat{f}(w)| = \frac{1}{|w|} \left| \int_{-R}^R (R^2 - x^2)^{1/2} x e^{iwx} dx \right|.$$

If we apply the triangle inequality immediately, we only obtain a bound of about Rw^{-1} . Instead, we would like something better. The idea is to break up the domain $[-R, R]$ into two regions, one an interval of the form $[-S, S]$ and the other the set with $S < |x| < R$. On the former region, we can integrate by parts again, and on the outer region we can use the triangle inequality, and then optimize the result by changing S . It turns out that taking $S = 1/|w|$ is the right choice, but either way, we get the estimate in the proposition. \square

Remark 2. Also, of course, $|\widehat{f}(w)| \leq \pi R^2$ by the triangle inequality directly, and this is a better estimate for small R .

Now, we have $N(R) = Pf(0)$, and our wishful thinking would have us hope that this is $\pi R^2 + \sum_{n \in \mathbb{Z}^2 \setminus \{0\}} \widehat{f}(n)$. But this sum isn't even absolutely summable given our estimate $|\widehat{f}(n)| \lesssim R^{1/2}|n|^{-3/2}$. Instead, we need to do something slightly different. Our trick is to approximate f by smooth functions so that we get better estimates and convergent sums. In order to do this, consider η a smooth bump function with support in $B(1)$ and $\int \eta = 1$. Set $f_{R,\epsilon} := \chi_{B(R)} * \eta_\epsilon$, where $\eta_\epsilon(x) = \epsilon^{-2}\eta(x/\epsilon)$ still has integral 1, but the support becomes more and more localized to 0 as $\epsilon \rightarrow 0$. For now, we will fix R and simply write $f_\epsilon := f_{R,\epsilon}$, and we will go back after the next proposition. The function f_ϵ is now smooth and compactly supported, and so we get the convergence properties needed in the sums we are looking at, and in particular, we can prove the following:

Proposition 3. $|Pf_\epsilon(0) - \pi R^2| \leq CR^{1/2}w^{-1/2}$.

Proof. Because f_ϵ is smooth, any sum we write in this proof will converge, as the reader can verify. Recall that the Fourier transform commutes with convolution, so we have

$$Pf_\epsilon(0) = \sum_{n \in \mathbb{Z}^2} \widehat{Pf_\epsilon}(n) = \sum_{n \in \mathbb{Z}^2} \widehat{f_\epsilon}(n) = \sum_{n \in \mathbb{Z}^2} \widehat{f}(n)\widehat{\eta_\epsilon}(n).$$

The term with $n = 0$ is just πR^2 , and so the left hand side of the inequality of our proposition is just

$$\left| \sum_{n \in \mathbb{Z}^2 - \{0\}} \widehat{f}(n)\widehat{\eta_\epsilon}(n) \right|.$$

Now, since η is compactly supported, we get $|\widehat{\eta_\epsilon}(n)| \lesssim (1 + |n|\epsilon)^{-10000}$, and so applying the triangle inequality to the sum at hand and using our estimates for $|\widehat{f}(n)|$ and $|\widehat{\eta_\epsilon}(n)|$, we find that

$$|Pf_\epsilon(0) - \pi R^2| \lesssim R^{1/2} \sum_{n \neq 0} |n|^{-3/2} (1 + |n|\epsilon)^{-10000} \sim R^{1/2} \sum_{0 < |n| < \epsilon^{-1}} |n|^{-3/2} \sim R^{1/2}w^{-1/2}.$$

□

Proof of Sierpinski Theorem. Note that $f_\epsilon \geq 1$ on $B(R - \epsilon)$, and so we have $N(R) \leq Pf_{R+\epsilon,\epsilon}(0)$, and so we find

$$\begin{aligned} N(R) &\leq \pi(R + \epsilon)^2 + CR^{1/2}\epsilon^{-1/2} \\ &= \pi R^2 + CR\epsilon + CR^{1/2}\epsilon^{-1/2} \\ &\leq \pi R^2 + CR^{2/3} \end{aligned}$$

where the last inequality comes from choosing $\epsilon = R^{-1/3}$ (which is the best possible choice by AM-GM). There is a similar lower bound, and so we find $|E(R)| \lesssim R^{2/3}$ as desired. □