

FINITE LATTICES AND JORDAN-HÖLDER SETS¹⁾

RICHARD P. STANLEY

1. Introduction

In this paper we extend some aspects of the theory of 'supersolvable lattices' [3] to a more general class of finite lattices which includes the upper-semimodular lattices. In particular, all conjectures made in [3] concerning upper-semimodular lattices will be proved. For instance, we will prove that if L is finite upper-semimodular and if L' denotes L with any set of 'levels' removed, then the Möbius function of L' alternates in sign. Familiarity with [3] will be helpful but not essential for the understanding of the results of this paper. However, many of the proofs are identical to the proofs in [3] (once the machinery has been suitably generalized) and will be omitted.

2. Admissible labelings

Let L be a finite lattice with bottom $\hat{0}$ and top $\hat{1}$, such that every maximal chain of L has the same length n . Hence L has a rank function ϱ satisfying $\varrho(\hat{0})=0$, $\varrho(\hat{1})=n$, and $\varrho(y)=1+\varrho(x)$ whenever y covers x in L . We call L a *graded* lattice.

Let I denote the set of join-irreducible elements of L . A *labeling* ω of L is any map $\omega: I \rightarrow \mathbf{P}$, where \mathbf{P} denotes the positive integers. A labeling ω is said to be *natural* if $z, z' \in I$ and $z \leq z'$ implies $\omega(z) \leq \omega(z')$. If $x < y$ in L and ω is a fixed labeling of L , define

$$\gamma(x, y) = \min \{ \omega(z) \mid z \in I, x < x \vee z \leq y \}.$$

Thus, $\gamma(x, y)$ is the least label of a join-irreducible which is less than or equal to y but not less than or equal to x . Note that $\gamma(x, y)$ is always defined since y is a join of join-irreducibles. We are now able to make the key definition of this paper. A labeling ω is said to be *admissible* if whenever $x < y$ in L , there is a *unique* unrefinable chain $x = x_0 < x_1 < \dots < x_m = y$ between x and y (so $m = \varrho(y) - \varrho(x)$) such that

$$\gamma(x_0, x_1) \leq \gamma(x_1, x_2) \leq \dots \leq \gamma(x_{m-1}, x_m). \quad (1)$$

We then call the pair (L, ω) an *admissible* lattice. Our motivation for this definition is that admissibility seems to be the weakest condition for which Theorem 3.1 holds.

¹⁾ The research was supported by a Miller Research Fellowship at the University of California at Berkeley.

The idea for this definition came from [3, Cor. 1.3] and its relation to [3, Thm. 2.1]. Our present Theorem 3.1 is a generalization of [3, Thm. 2.1].

We first note a simple property of admissible labelings.

2.1. PROPOSITION. *Let ω be an admissible labeling of a finite graded lattice L . Then ω is natural.*

Proof. Suppose $z, z' \in I$ with $z < z'$ and $\omega(z) > \omega(z')$. Since z is join-irreducible, it covers a unique element x . Similarly z' covers a unique element y . Hence any unrefinable chain between x and z' has the form $x < z = y_0 < y_1 < \dots < y_m = y < z'$ (possibly $m=0$ so $z=y$). Since z is join-irreducible, it follows from the definition of γ that $\gamma(x, z) = \omega(z)$. Similarly $\gamma(y, z') = \omega(z')$. Since $\omega(z) > \omega(z')$, ω cannot be admissible. \square

We know of two main classes of admissible lattices. The first class is given by the next proposition.

First recall that a lattice L of finite length is said to be *upper-semimodular* if it is a graded lattice whose rank function ϱ satisfies $\varrho(x) + \varrho(y) \geq \varrho(x \vee y) + \varrho(x \wedge y)$ for all $x, y \in L$. Equivalently, L is upper-semimodular if whenever x covers y , then $x \vee z$ covers or equals $y \vee z$, for all $x, y, z \in L$.

2.2. PROPOSITION. *Let L be a finite upper-semimodular lattice and ω a natural labeling of L such that whenever z and z' are incomparable join-irreducibles then $\omega(z) \neq \omega(z')$. (Such a labeling of L is clearly possible; in fact, an injective natural labeling can always be found.) Then ω is admissible.*

To prove this result, we first need a lemma.

2.3. LEMMA. *Let (L, ω) satisfy the hypotheses of Proposition 2.2, and let $x < y$ in L . Let z be a minimal element of the set J of all join-irreducibles z' of L satisfying $\omega(z') = \gamma(x, y)$ and $x < x \vee z' \leq y$. (J is not empty by definition of $\gamma(x, y)$.) Then $x \vee z$ covers x .*

Proof. Let I denote as before the set of join-irreducibles of L . Let $I' \subseteq I$ be the set of all $z' \in I$ satisfying $z' < z$. Let $z' \in I'$. Since $z' < z$, $x \leq x \vee z' \leq y$. Since ω is natural, $\omega(z') \leq \omega(z)$. If $\omega(z') < \omega(z)$, then by definition of $\gamma(x, y)$ we cannot have $x < x \vee z' \leq y$, so $x = x \vee z'$. On the other hand, if $\omega(z') = \omega(z)$, then by hypothesis we cannot have $x < x \vee z' \leq y$, so once again $x = x \vee z'$. Thus $x = x \vee z'$ for all $z' \in I'$. Let $w = \bigvee_{z' \in I'} z'$. Since z is join-irreducible, $w < z$. Since $x = x \vee z'$ for all $z' \in I'$, we have $x \vee w = x$.

Now if z doesn't cover w , then $w < w' < z$ for some $w' \in L$. But then there is a new join-irreducible $v < z$ such that $w < w \vee v \leq w'$, contradicting the definition of w . Hence z covers w . But by upper-semimodularity, if z covers w , then $x \vee z$ covers or equals $x \vee w = x$. By assumption, $x < x \vee z$, so $x \vee z$ covers x . \square

Proof of Proposition 2.2. Let $x < y$ in L , and let $m = \varrho(y) - \varrho(x)$. We first show the existence of an unrefinable chain $x = x_0 < x_1 < \dots < x_m = y$ between x and y satis-

fyng (1). Let z_1 be a minimal element of the set J_1 of join-irreducibles z satisfying $\omega(z) = \gamma(x_0, y)$ and $x < x \vee z \leq y$. Let $x_1 = x_0 \vee z_1$. By Lemma 2.3, x_1 covers x_0 , while by definition $x_1 \leq y$.

If $m = 1$, we are done, so assume $m \geq 2$. Let z_2 be a minimal element of the set J_2 of join-irreducibles z satisfying $\omega(z) = \gamma(x_1, y)$ and $x_1 < x_1 \vee z \leq y$. Let $x_2 = x_1 \vee z_2$. Once again by Lemma 2.3 x_2 covers x_1 , while again by definition $x_2 \leq y$. Now by definition of $\gamma(x_0, y)$ we have $\omega(z_1) = \gamma(x_0, y) \leq \omega(z_2) = \gamma(x_1, y)$. Continuing in this way, after m steps we get an unrefinable chain $x = x_0 < x_1 < \dots < x_m = y$ satisfying $\gamma(x_0, y) \leq \gamma(x_1, y) \leq \dots \leq \gamma(x_{m-1}, y)$. But clearly by definition of γ and the x_i 's, $\gamma(x_i, y) = \gamma(x_i, x_{i+1})$. Hence we have constructed a chain C satisfying (1).

It remains to show the uniqueness of C . We shall prove the following two results:

- (i) If $x' \in L$ is such that x' covers x , $x' \leq y$, and $\gamma(x, x') = \gamma(x, y)$, then $x' = x_1$;
- (ii) If $x = x'_0 < x'_1 < \dots < x'_m = y$ is any unrefinable chain satisfying (1), then $\gamma(x'_1, x) = \gamma(x, y)$.

Thus (i) and (ii) imply that x'_1 is uniquely determined, viz., $x'_1 = x_1$ (where $x_1 = x_0 \vee z_1$ as defined above). Hence the proof of the proposition follows by induction on m .

Proof of (i). Suppose $x'' \neq x'$ also is such that x'' covers x , $x'' \leq y$, and $\gamma(x, x'') = \gamma(x, y)$. Thus there exist $z', z'' \in I$ such that $\omega(z') = \omega(z'') = \gamma(x, y)$, $x \vee z' = x'$, $x \vee z'' = x''$. Since x' and x'' both cover x , they are incomparable. Hence z' and z'' are incomparable. Thus by hypothesis $\omega(z') \neq \omega(z'')$, a contradiction. Hence x'' cannot exist.

Proof of (ii). Let $x = x'_0 < x'_1 < \dots < x'_m = y$ be an unrefinable chain satisfying (1). Hence $\gamma(x, x'_1) \geq \gamma(x, y)$. Suppose $\gamma(x, x'_1) > \gamma(x, y)$. Let $z \in I$ satisfy $\omega(z) = \gamma(x, y)$ and $x < x \vee z \leq y$. Let i be the least positive integer for which $x \vee z \leq x'_i$. (Clearly i exists since $x \vee z \leq x'_m$.) Then $x'_{i-1} \vee z = x'_i$, so $\gamma(x'_{i-1}, x'_i) = \gamma(x, y) < \gamma(x, x'_1)$. Thus (1) cannot hold. \square

The second main class of admissible lattices are the *supersolvable lattices* [3]. If L is a finite lattice and Δ a maximal chain of L , we call the pair (L, Δ) a supersolvable lattice (or SS-lattice) if the sublattice of L generated by Δ and any chain in L is distributive. It is easily seen that if (L, Δ) is an SS-lattice, then L is graded (cf. [3, §1]).

2.4. PROPOSITION. *Let (L, Δ) be an SS-lattice with Δ given by $\hat{0} = x_0 < x_1 < \dots < x_n = \hat{1}$. Define a labeling $\omega: I \rightarrow \mathbf{P}$ by letting $\omega(z)$ be the least positive integer t for which $z \leq x_t$. Then ω is admissible*

Proof. Recall that an interval $[u, v]$ of a lattice is *prime* if it contains exactly two elements, i.e., if v covers u . In a distributive lattice D , two prime intervals $[x, y]$ and $[u, v]$ are said to be *projective* if there is a unique join-irreducible z such that $y = x \vee z$ and $v = u \vee z$. This is easily seen to be equivalent to the usual definition of projectivity (e.g., [1, p. 14]) if one thinks of D as being coordinatized by a ring of sets.

If y covers x in L , then it is easily seen [3, p. 198] that there is a unique positive integer t , which we denote by $\gamma'(x, y)$, for which the prime intervals $[x, y]$ and $[x_{t-1}, x_t]$ are projective in the distributive lattice D_{xy} generated by Δ and $\{x, y\}$. In [3, Cor. 1.3], it was shown that for any $x' < y'$ in L , there is a unique unrefinable chain $x' = x'_0 < x'_1 < \dots < x'_m = y'$ between x' and y' such that

$$\gamma'(x'_0, x'_1) < \gamma'(x'_1, x'_2) < \dots < \gamma'(x'_{m-1}, x'_m).$$

Hence it suffices to prove that $\gamma(x, y) = \gamma'(x, y)$ whenever y covers x .

We shall need the following elementary facts concerning projectivity in a finite distributive lattice D . The proofs are immediate from the above definition of projectivity.

- (a) The prime intervals $[x, y]$ and $[u, v]$ are projective in D if and only if $x = (x \vee u) \wedge y$ and $y = (x \vee v) \wedge y$.
- (b) If $[x, y]$ and $[u, v]$ are projective prime intervals in D , then $y \not\leq u$.
- (c) Suppose $[w, z]$ is a prime interval in D and z is join-irreducible. If y covers x in D and $z \leq y$, $z \not\leq x$, then $[w, z]$ and $[x, y]$ are projective.

We proceed to prove that if y covers x in L , then $\gamma(x, y) = \gamma'(x, y)$. By definition of $\gamma(x, y)$, there is a join-irreducible z satisfying $x \vee z = y$ and $\omega(z) = \gamma(x, y)$. Let w be the unique element of L covered by z , and set $s = \omega(z)$. By (c), $[w, z]$ and $[x_{s-1}, x_s]$ are projective in the distributive lattice D_{wz} generated by Δ and $\{w, z\}$, so $\gamma'(w, z) = s$. If z' is a join-irreducible of L such that $z' < z$, then it follows from (b) (taking D to be generated by Δ and $\{z, z'\}$) that $\omega(z') \neq \omega(z)$. Since $\omega(z') \leq \omega(z)$, thus $\omega(z') < \omega(z)$.

We claim that $w \leq x$. It suffices to prove $z' \leq x$ for all join-irreducibles $z' \leq w$. If z' is such a join-irreducible, then by the above $\omega(z') < \omega(z)$. Hence by the definition of z , $x \vee z' = y$. But $x \vee z' \leq y$ since $z \leq y$. Since y covers x , we must have $z' \leq x$. Hence $w \leq x$.

We need to show $\omega(z) = t$, i.e., $s = t$. By (a) and (c) this is equivalent to $w = (w \vee x_{t-1}) \wedge z$ and $z = (w \vee x_t) \wedge z$. Since $\gamma'(x, y) = t$, we know by (a) that

$$x = (x \vee x_{t-1}) \wedge y \tag{2}$$

$$y = (x \vee x_t) \wedge y. \tag{3}$$

Since $w \leq x$, $z \not\leq x$, $z \leq y$, and z covers w , from (2) we get $w = x \wedge z = (x \vee x_{t-1}) \wedge z$. Thus since $w \leq x$ and $w \leq z$, $w \leq (w \vee x_{t-1}) \wedge z \leq (x \vee x_{t-1}) \wedge z = w$ so $w = (w \vee x_{t-1}) \wedge z$ as desired. To prove the other equality $z = (w \vee x_t) \wedge z$, we need to show $w \vee x_t \geq z$. Since w is the only element which z covers, this is equivalent to $x_t \not\leq w$. But if $x_t \leq w$, then $x_t \leq x$ since $w \leq x$. From (3) this would imply $y = x \wedge y = x$, a contradiction. \square

It follows from Proposition 2.4 that the theory of SS -lattices, as developed in [3], is a special case of the theory of admissible lattices. A large class of examples of SS -lattices, some of which are not semimodular, is given in [3, §2].

3. Jordan-Hölder sequences

Let (L, ω) be an admissible finite graded lattice. Let $x \leq y$ in L , and suppose K is an unrefinable chain in L between x and y given by $x = x_0 < x_1 < \dots < x_m = y$. Define the *Jordan-Hölder sequence* (or *J-H sequence*) associated with K to be the sequence a_1, a_2, \dots, a_m of positive integers given by $a_i = \gamma(x_{i-1}, x_i)$. We shall denote this sequence by π_K and shall write

$$\pi_K = (a_1, a_2, \dots, a_m).$$

In [3] π_K was called a ‘*J-H permutation*’ but here repetitions among the a_i are possible.

Now define the *Jordan-Hölder set* (or *J-H set*) $\mathcal{J}_{xy}(L, \omega)$ of $(L, \omega; x, y)$ (denoted \mathcal{J}_{xy} for short) to be the set of all *J-H sequences* π_K , including repetitions, as K ranges over all unrefinable chains between x and y . It follows from the definition of an admissible labeling that there is a unique element $\pi_K = (a_1, \dots, a_m)$ of \mathcal{J}_{xy} satisfying $a_1 \leq a_2 \leq \dots \leq a_m$. If $x = \hat{0}$ and $y = \hat{1}$, we denote $\mathcal{J}_{xy}(L, \omega)$ simply by $\mathcal{J}(L, \omega)$ or just \mathcal{J} , and call it the *J-H set of* (L, ω) .

If $k \in \mathbf{P}$, let \mathbf{k} denote the set $\{1, 2, \dots, k\}$. We also write $S = \{m_1, m_2, \dots, m_s\}_<$ to signify that $S = \{m_1, m_2, \dots, m_s\}$ and $m_1 < m_2 < \dots < m_s$. Suppose L is a finite graded lattice and $[x, y]$ is an interval of L of length (rank) m , i.e., $\varrho(y) - \varrho(x) = m$. If $\{m_1, \dots, m_s\}_< = S \subseteq \mathbf{m} - \mathbf{1}$, define $\alpha_{xy}(S)$ to be the number of chains

$$x < y_1 < \dots < y_s < y$$

in L satisfying $\varrho(y_i) - \varrho(x) = m_i, i = 1, 2, \dots, s$. Thus if $S = \{k\}$, then $\alpha_{xy}(S)$ is the number of elements z of $[x, y]$ of rank k in $[x, y]$ (i.e., $\varrho(z) - \varrho(x) = k$). Moreover, $\alpha_{xy}(\phi) = 1$ and $\alpha_{xy}(\mathbf{m} - \mathbf{1})$ is the total number of unrefinable chains in L between x and y . Now define for $S \subseteq \mathbf{m} - \mathbf{1}$,

$$\beta_{xy}(S) = \sum_{T \subseteq S} (-1)^{|S-T|} \alpha_{xy}(T),$$

so by the Principle of Inclusion-Exclusion [2],

$$\alpha_{xy}(S) = \sum_{T \subseteq S} \beta_{xy}(T).$$

As mentioned in [3, p. 198], if $L_{xy}(S)$ denotes the partially ordered set of all $z \in L$ satisfying either (a) $z = x$; (b) $z = y$; or (c) $x < z < y$ and $\varrho(z) - \varrho(x) \in S$, then

$$\mu_S(x, y) = (-1)^{s+1} \beta_{xy}(S), \tag{4}$$

where μ_S is the Möbius function of $L_{xy}(S)$ and $|S| = s$. For this reason we call the function $\beta_{xy}(\cdot)$ the *rank-selected Möbius invariant* of the interval $[x, y]$.

If $\pi = (a_1, a_2, \dots, a_m)$ is a finite sequence of integers, then a pair $a_j > a_{j+1}$ is called a *descent* of π , and the set

$$D(\pi) = \{j : a_j > a_{j+1}\}$$

is called the *descent set* of π . We can now state the fundamental combinatorial property of *J-H* sets. This result is a direct generalization of [3, Thm. 1.2]. The proof is identical to the proof of [3, Thm. 1.2], except that here the definition of an admissible lattice plays the role of Lemma 3.1 of [3]. Thus no condition is needed about distributive sublattices of L .

3.1. THEOREM. *Let (L, ω) be an admissible lattice, and let $[x, y]$ be an interval of L of length m . If $S \subseteq \mathbf{m} - \mathbf{1}$, then the number of sequences π in the *J-H* set $\mathcal{J}_{xy}(L, \omega)$ with descent set $D(\pi) = S$ is equal to $\beta_{xy}(S)$. (The reader is reminded that $\mathcal{J}_{xy}(L, \omega)$ contains one sequence π for each maximal chain of $[x, y]$, so that repeated sequences are taken into account.)*

3.2. COROLLARY. *Let (L, ω) be an admissible lattice. If $[x, y]$ is an interval of L of length m and if $S \subseteq \mathbf{m} - \mathbf{1}$, then $\beta_{xy}(S) \geq 0$. \square*

In view of (4), Corollary 3.2 may be restated as follows:

3.2'. COROLLARY. *Let (L, ω) be an admissible lattice of length n , and let $S \subseteq \mathbf{n} - \mathbf{1}$. Then the Möbius function μ_S of the rank-selected partially ordered set $L(S)$ alternates in sign; i.e., if $[x, y]$ is an interval in $L(S)$ of length k , then*

$$(-1)^k \mu_S(x, y) \geq 0. \quad \square$$

Since by Proposition 2.2 every finite upper-semimodular lattice has an admissible labeling, Corollary 3.2' applies to all such lattices, and in particular, to finite geometric lattices.

3.3. COROLLARY. *Let (L, ω) be an admissible lattice and $[x, y]$ an interval of L of length m . Let μ denote the Möbius function of L . Then $(-1)^m \mu(x, y)$ is equal to the number of unrefinable chains $x = x_0 < x_1 < \dots < x_m = y$ between x and y satisfying*

$$\gamma(x_0, x_1) > \gamma(x_1, x_2) > \dots > \gamma(x_{m-1}, x_m).$$

Proof. Let $S = \mathbf{m} - \mathbf{1}$ in Theorem 3.1, and use (4). \square

4. Applications

We shall state those results in [3] proved for *SS*-lattices which remain true for admissible lattices. The proofs are exactly the same as in the *SS*-case once suitable

analogues are given for two concepts in [3]. First, the role of the ‘induced M -chain Δ_{xy} between x and y ’ is replaced by the unique unrefinable chain $x = x_0 < x_1 < \dots < x_m = y$ between x and y satisfying $\gamma(x_0, x_1) \leq \gamma(x_1, x_2) \leq \dots \leq \gamma(x_{m-1}, x_m)$. Secondly, we need a replacement for statement (A) in the proof of Theorem 5.2 of [3]. Although a direct analogue of (A) can be given, it is simpler to use the following fact:

4.1. LEMMA. *If $[x, y]$ is an interval of an upper-semimodular admissible lattice (L, ω) such that y is the join of atoms of $[x, y]$, then there is an unrefinable chain $x = x_0 < x_1 < \dots < x_m = y$ between x and y such that $\gamma(x_0, x_1) > \gamma(x_1, x_2) > \dots > \gamma(x_{m-1}, x_m)$.*

Proof. Recall that a *geometric lattice* is an upper-semimodular lattice whose join-irreducibles are its atoms. If L' denotes the partially ordered set of all elements of $[x, y]$ which are a join of atoms of $[x, y]$ (including x as the void join), then L' has the structure of a geometric lattice (though L' is not necessarily a sublattice of L). If μ denotes the Möbius function of L and μ' that of L' , then from [2, Cor. on p. 349] we conclude $\mu(x, y) = \mu'(x, y)$. Hence by [2, §7, Thm. 4], $\mu(x, y) \neq 0$. The desired result now follows from Corollary 3.3. \square

The reader can now verify that the proofs of the following results are the same as the analogous results for SS -lattices given in [3].

4.2. PROPOSITION. (Generalizes [3, Prop. 3.3]). *Let (L, ω) be an admissible lattice, and let $[x, y]$ be an interval of L length m . Let $S \subseteq \mathbf{m} - 1$. If $\beta_{xy}(S) > 0$ and $T \subseteq S$, then $\beta_{xy}(T) > 0$. \square*

Suppose L is a finite geometric lattice. Then L is upper-semimodular, so by Proposition 2.2 L possesses an admissible labeling. Moreover, every interval of L is a geometric lattice, and the Möbius function of L is never 0. It follows from (4) and Proposition 4.2 that Corollary 3.2' can be strengthened in the case of geometric lattices as follows:

4.3. COROLLARY. *Let L be a finite geometric lattice of rank n , and let $S \subseteq \mathbf{n} - 1$. Then the Möbius function μ_S of the rank-selected partially ordered set $L(S)$ strictly alternates in sign; i.e., if $[x, y]$ is an interval in $L(S)$ of length k , then*

$$(-1)^k \mu_S(x, y) > 0. \quad \square$$

For some related properties of geometric lattices, see the next section.

Recall [3, §5] that a *Loewy chain* between x and y in a lattice L of finite length is a chain $x = x_0 < x_1 < \dots < x_r = y$ such that each $x_i, i \in \mathbf{r}$, is the join of the atoms of the interval $[x_{i-1}, x_i]$.

4.4. PROPOSITION. (Generalizes [3, Lemma 5.1]). *Let (L, ω) be an admissible*

lattice with $[x, y]$ an interval of length m . Let K be an unrefinable chain in L between x and y :

$$K: x = y_0 < y_1 < \dots < y_m = y.$$

Let $0 < m_1 < m_2 < \dots < m_r = m$. Then the subchain

$$x = y_0 < y_{m_1} < y_{m_2} < \dots < y_{m_r} = y$$

of K is a Loewy chain between x and y if

$$m - 1 - D(\pi_K) \subseteq \{m_1, m_2, \dots, m_{r-1}\}. \quad \square$$

4.5. THEOREM. (Generalizes [3, Thm. 5.2]). Let L be a finite upper-semimodular lattice with $[x, y]$ an interval of L of length m . Let $S = \{m_1, m_2, \dots, m_s\} \subset \leq m - 1$. There exists a chain C ,

$$C: x = y_0 < y_1 < \dots < y_s < y_{s+1} = y$$

satisfying the two conditions

- (i) $q(y_i) - q(x) = m_i, 1 \leq i \leq s$ (where q as usual is the rank function of L);
- (ii) C is a Loewy chain between x and y ,

if and only if $\beta_{xy}((m - 1) - S) > 0$.

Now recall [3, §6] that if q is a fixed positive integer, then a q -lattice is a lattice L of finite length with the property that every interval $[x, y]$ of L for which y is the join of atoms of $[x, y]$ is isomorphic to the lattice of subspaces of a projective geometry of degree q (or to a Boolean algebra if $q = 1$). Such a lattice is necessarily upper-semimodular [3, pp. 213–214] and hence possesses an admissible labeling. A q -lattice, however, need not be supersolvable, so the next proposition is strictly stronger than the corresponding Lemma 6.4 of [3]. For instance, let L' be the lattice of subgroups of a finite abelian p -group of type (3,3). Let L be L' truncated above rank 3, i.e., identify all elements of L' of rank at least 4. Then L is a p -lattice but is not supersolvable.

4.6. PROPOSITION. (Replaces [3, Lemma 6.4]). Let (L, ω) be an admissible q -lattice of rank n . Let $S \subseteq n - 1$, with $(n - 1) - S = \{j_1, j_2, \dots, j_{t-1}\} \subset$. Also let $j_0 = 0, j_t = n$. Define $N(S)$ to be the number of maximal chains K of L satisfying $D(\pi_K) \supseteq S$, where $D(\pi_K)$ is the descent set of the J - H sequence π_K . Then $N(S) = q^k M$, where

$$k = \sum_{n=1}^t \binom{j_r - j_{r-1}}{2} \tag{5}$$

and where M is the number of Loewy chains

$$\hat{0} = y_0 < y_1 < \dots < y_t = \hat{1} \tag{6}$$

such that $q(y_i) = j_i, 0 \leq i \leq t$.

Since Proposition 4.6 is not a strict analogue of [3, Lemma 6.4], we shall give a proof.

Proof. If K is a maximal chain of L such that $D(\pi_K) \ni S$, then by Proposition 4.4 the subchain C of K consisting of all $x \in K$ such that $\varrho(x) = j_i (0 \leq i \leq t)$ is a Loewy chain. Hence it suffices to prove that if we have a Loewy chain (6) with $\varrho(y_i) = j_i$, then the number of refinements of C to a maximal chain K satisfying $D(\pi_K) \ni S$ is equal to q^k , where k is given by (5).

Assume we have such a Loewy chain C . Since L is a q -lattice, each interval $[y_{r-1}, y_r]$ ($1 \leq r \leq t$) is a projective geometry of degree q (or a Boolean algebra if $q = 1$). Hence $\mu(y_{r-1}, y_r) = (-1)^b q^{k_r}$, where $b = j_r - j_{r-1}$ and $k_r = \binom{j_r - j_{r-1}}{2}$. Now by Corollary 3.3 the number of maximal chains $y_{r-1} = z_0 < z_1 < \dots < z_b = y_r$ of the interval $[y_{r-1}, y_r]$ such that

$$\gamma(z_0, z_1) > \gamma(z_1, z_2) > \dots > \gamma(z_{b-1}, z_b)$$

is just $(-1)^b \mu(y_{r-1}, y_r) = q^{k_r}$. Hence the total number of refinements of C to a maximal chain K satisfying $D(\pi_K) \ni S$ is equal to $q^{k_1} q^{k_2} \dots q^{k_t} = q^k$, and the proof follows. \square

4.7. COROLLARY. (Generalizes [3, Corollary 6.5]). *Let L be a q -lattice of rank n , and let $S \subseteq \mathbf{n-1}$, with $(\mathbf{n-1}) - S = \{j_1, j_2, \dots, j_{t-1}\}_<$ and $j_0 = 0, j_t = n$. Then $\beta(S)$ is divisible by q^k , where k is given by (6).*

The derivation of Corollary 4.7 from Proposition 4.6 is not quite as trivial as the derivation of [3, Corollary 6.5] from [3, Lemma 6.4], so we shall give a proof.

Proof. Fix an admissible labeling ω of L . By Theorem 3.1, $\beta(S)$ is equal to the number of maximal chains K of L satisfying $D(\pi_K) = S$. Hence if $N(S)$ is defined as in Proposition 4.6, we have

$$N(S) = \sum_{T \ni S} \beta(T),$$

so

$$\beta(S) = \sum_{T \ni S} (-1)^{|T-S|} N(T). \tag{7}$$

Suppose we have $\mathbf{n-1} \ni T \ni S$ where $(\mathbf{n-1}) - T = \{i_1, i_2, \dots, i_{s-1}\}_<$ and $(\mathbf{n-1}) - S = \{j_1, j_2, \dots, j_{t-1}\}_<$, and $i_0 = j_0 = 0, i_s = j_t = n$. An easy computation shows that

$$\sum_{r=1}^s \binom{i_r - i_{r-1}}{2} \geq \sum_{r=1}^t \binom{j_r - j_{r-1}}{2}.$$

It follows from Proposition 4.6 that each term $N(T)$ appearing in (7) is divisible by q^k , so the proof follows. \square

4.8. THEOREM. (Generalizes [3, Theorem 6.6]). *Let L be a q -lattice of rank n , and let $S \subseteq \mathbf{n}-1$ with $|S|=s$. Then $\beta(S)$ is divisible by $q^{Q(n,s)}$, where*

$$Q(n,s) = \frac{1}{2} \left[\frac{n}{n-s} \right] \left(n+s - (n-s) \left[\frac{n}{n-s} \right] \right)$$

(brackets denote the integer part). *This result is best possible in the sense that given n and $0 \leq s \leq n-1$, there exists a q -lattice (which can even be chosen to be modular) of rank n and a set $S \subseteq \mathbf{n}-1$ of cardinality s such that $\beta(S) = q^{Q(n,s)}$ (see [3, p. 216]). \square*

5. The broken circuit theorem

In this section we shall point out the connection between our work and the so-called ‘broken circuit theorem’ of G.-C. Rota [2, Prop. 1, p. 358], which generalizes to arbitrary finite geometric lattices a result of Whitney on graphs. The reader should be warned that [2, Prop. 1, p. 358] is *false* when $k > 1$. However, the proof is valid when $k = 1$, and this is the case which will concern us here.

We proceed to describe the broken circuit theorem. Let L be a finite geometric lattice of rank n , and let a_1, a_2, \dots, a_t be an ordering of the atoms A of L . A subset C of A is called a *circuit* if the rank of the join of the elements of C is $|C|-1$, while the rank of the join of the elements of any proper subset C' of C is $|C'|$. A subset $B = \{a_{i_1}, a_{i_2}, \dots, a_{i_j}\}$ of A is called a *broken circuit* if there exists an atom a_m such that $m > i_r$, for $r = 1, 2, \dots, j$, and such that $B \cup \{a_m\}$ is a circuit. Note that the notion of a circuit depends only on L , while that of a broken circuit also depends on the ordering chosen for the elements of A .

BROKEN CIRCUIT THEOREM (G.-C. Rota). *Let L be a finite geometric lattice of rank n with an ordering a_1, a_2, \dots, a_t of the atoms of L . Let μ be the Möbius function of L . Then $(-1)^n \mu(\hat{0}, \hat{1})$ is equal to the number of sets of n atoms of L not containing any broken circuit. \square*

Given an ordering a_1, a_2, \dots, a_t of the atoms of a finite geometric lattice L of rank n , define a labeling ω of L by $\omega(a_i) = t - i + 1$, so $i < j$ implies $\omega(a_i) > \omega$. By Proposition 2.2, (L, ω) is an admissible lattice. Let $\hat{0} = x_0 < x_1 < \dots < x_n = \hat{1}$ be a maximal chain K in L satisfying

$$\gamma(x_0, x_1) > \gamma(x_1, x_2) > \dots > \gamma(x_{n-1}, x_n). \tag{8}$$

We know by Corollary 3.3 that the number of such maximal chains K is $(-1)^n \mu(\hat{0}, \hat{1})$. We would like to relate this fact to the Broken Circuit Theorem by constructing an explicit one-to-one correspondence λ between maximal chains K satisfying (8) and sets of n atoms of L containing no broken circuit.

This correspondence λ is defined as follows. Given a maximal chain K satisfying (8), let $\lambda(K)$ be the set $\{b_1, b_2, \dots, b_n\}$ of those n atoms defined by $\omega(b_j) = \gamma(x_{j-1}, x_j)$.

5.1. PROPOSITION. *The function λ defines a one-to-one correspondence between maximal chains K of L satisfying (8), and sets of n atoms of L containing no broken circuit.*

Proof. We first prove that $\lambda(K)$ contains no broken circuits. Suppose $B = \{b_{i_1}, b_{i_2}, \dots, b_{i_s}\}$ is a broken circuit contained in $\lambda(K)$ with $i_1 < i_2 < \dots < i_s$, i.e., $\omega(b_{i_1}) > \omega(b_{i_2}) > \dots > \omega(b_{i_s})$. By definition of broken circuit, there exists an atom a of L such that $\omega(b_{i_r}) > \omega(a)$ for $r = 1, 2, \dots, s$ and $B \cup \{a\}$ is a circuit. By definition of the b_i 's and γ , $x_{i_{s-1}} \vee b_{i_s} = x_{i_s}$ and $b_{i_r} \leq x_{i_{s-1}}$ for $r = 1, 2, \dots, s-1$. Hence since $B \cup \{a\}$ is a circuit, $x_{i_{s-1}} \vee a = x_{i_s}$. By definition of γ , this means $\omega(b_{i_s}) < \omega(a)$, a contradiction. Hence $\lambda(K)$ contains no broken circuit.

Now let $B = \{b_1, b_2, \dots, b_n\}$ be a set of n atoms containing no broken circuit, with $\omega(b_1) > \omega(b_2) > \dots > \omega(b_n)$. Recall that a *basis* of L is a set of n atoms c_1, c_2, \dots, c_n of L such that $\varrho(c_1 \vee c_2 \vee \dots \vee c_n) = n$. Equivalently, a basis is a set of n atoms containing no circuit. Now note that B is a basis, since if it contained a circuit it would contain a broken circuit. If $\lambda(K) = B$, then K must be given by $x_j = b_1 \vee b_2 \vee \dots \vee b_j$, so λ is injective. It remains to prove that these x_j 's satisfy $\gamma(x_{j-1}, x_j) = \omega(b_j)$, which shows λ is surjective. By definition of the x_i 's, $x_{j-1} \vee b_j = x_j$. Suppose a is an atom such that $x_{j-1} \vee a = x_j$ and $\omega(a) < \omega(b_j)$. Thus the set $\{b_1, b_2, \dots, b_j, a\}$ contains a circuit C . Moreover, $a \in C$ since the b_i 's are independent. Since $\omega(b_1) > \omega(b_2) > \dots > \omega(b_j)$ and $\omega(b_j) > \omega(a)$, $\omega(a) < \omega(b_i)$ for $1 \leq i \leq j$. Hence $C - \{a\}$ is a broken circuit, a contradiction. This completes the proof. \square

REFERENCES

- [1] Garrett Birkhoff, *Lattice Theory*, 3rd Ed., American Mathematical Society, Providence, Rhode Island, 1967.
- [2] G.-C. Rota, *On the foundations of combinatorial theory: I: Theory of Möbius functions*, *Z. Wahrscheinlichkeitstheorie*, 2 (1964), 340-368.
- [3] R. Stanley, *Supersolvable lattices*, *Algebra Universalis*, 2 (1972), 197-217.

*University of California at Berkeley
Berkeley, California
U.S.A.*