

Let

$$F(x) = x + c_2x^2 + c_3x^3 + \dots \in \mathbb{C}[[x]].$$

**Easy:**  $F(F(x)) = x \Rightarrow F(x) = x$ .

What about  $F(-F(-x)) = x$ ?

**Fact #1.**  $c_2, c_4, \dots$  can be arbitrary and uniquely determine  $c_3, c_5, \dots$

**Example.**  $c_3 = c_2^2$

$$c_5 = 3c_4c_2 - 2c_2^4$$

$$c_7 = 13c_2^6 - 18c_4c_2^3 + 2c_4^2 + 4c_2c_6$$

**1.** What are the coefficients?

**Fact #2** (EC1, Exercise 1.41). *Given  $F(-F(-x)) = x$ , there is a unique*

$$G(x) = x + b_2x^2 + b_4x^4 + b_6x^6 + \dots$$

*such that  $F(x) = G^{\langle -1 \rangle}(-G(-x))$ .*

**Example.**  $b_2 = -\frac{1}{2}c_2$

$$b_4 = \frac{1}{8}(5c_2^3 - 4c_4)$$

$$b_6 = -\frac{1}{16}(49c_2^5 - 56c_2^2c_4 + 8c_6)$$

**2.** What are the coefficients?

**Example.**

$$F(x) = \frac{x}{1+2x} \Rightarrow b_{2n} = (-1)^{n-1} E_{2n-1}$$

**Fact #3** (Aguiar). For any  $A(x) = x + a_2x^2 + a_3x^3 + \dots$  there are unique

$$G(x) = x + d_3x^3 + d_5x^5 + \dots$$

$$F(x) = x + c_2x^2 + c_3x^3 + \dots$$

such that

$$F(-F(-x)) = x \text{ and } A(x) = G(F(x)).$$

**Example.**  $c_2 = a_2, \quad d_3 = a_3 - a_2^2$

$$c_4 = a_4 - 3a_3a_2 + 3a_2^3$$

$$d_5 = a_5 + 3a_2^2a_3 - 3a_2a_4 - a_2^4$$

**3.** What are the coefficients?