

# The $X$ -Descent Set of a Permutation

Richard P. Stanley  
M.I.T. and U. Miami

July 4, 2022

# The Descent Set of a Permutation

$$w = a_1 a_2 \cdots a_n \in \mathfrak{S}_n$$

**descent set** of  $w$ :  $\mathbf{Des}(w) = \{1 \leq i \leq n - 1 : a_i > a_{i+1}\}$

# The Descent Set of a Permutation

$$w = a_1 a_2 \cdots a_n \in \mathfrak{S}_n$$

**descent set** of  $w$ :  $\text{Des}(w) = \{1 \leq i \leq n-1 : a_i > a_{i+1}\}$

Fix  $n$ . For  $S \subseteq [n-1]$ , define

$$F_S = \sum_{\substack{1 \leq i_1 \leq i_2 \leq \cdots \leq i_n \\ i_j < i_{j+1} \text{ if } j \in S}} x_{i_1} x_{i_2} \cdots x_{i_n},$$

known as **(Gessel's) fundamental quasisymmetric function**.

# The Descent Set of a Permutation

$$w = a_1 a_2 \cdots a_n \in \mathfrak{S}_n$$

**descent set** of  $w$ :  $\text{Des}(w) = \{1 \leq i \leq n-1 : a_i > a_{i+1}\}$

Fix  $n$ . For  $S \subseteq [n-1]$ , define

$$F_S = \sum_{\substack{1 \leq i_1 \leq i_2 \leq \cdots \leq i_n \\ i_j < i_{j+1} \text{ if } j \in S}} x_{i_1} x_{i_2} \cdots x_{i_n},$$

known as **(Gessel's) fundamental quasisymmetric function**.

**Theorem.**  $\sum_{w \in \mathfrak{S}_n} F_{\text{Des}(w)} = (x_1 + x_2 + \cdots)^n$

## The case $n = 3$

$w$	$F_{\text{Des}(w)}$
123	$\sum_{1 \leq a \leq b \leq c} x_a x_b x_c$
132	$\sum_{1 \leq a \leq b < c} x_a x_b x_c$
213	$\sum_{1 \leq a < b \leq c} x_a x_b x_c$
231	$\sum_{1 \leq a \leq b < c} x_a x_b x_c$
312	$\sum_{1 \leq a < b \leq c} x_a x_b x_c$
321	$\sum_{1 \leq a < b < c} x_a x_b x_c$
	$(x_1 + x_2 + \dots)^3$

## $X$ -descent sets

$$X \subseteq \mathcal{E}_n := \{(i, j) : 1 \leq i \leq n, 1 \leq j \leq n, i \neq j\}$$

## $X$ -descent sets

$$X \subseteq \mathcal{E}_n := \{(i, j) : 1 \leq i \leq n, 1 \leq j \leq n, i \neq j\}$$

**$X$ -descent** of  $w = a_1 \cdots a_n \in \mathfrak{S}_n$ : an index  $1 \leq i \leq n - 1$  for which  $(a_i, a_{i+1}) \in X$

## $X$ -descent sets

$$X \subseteq \mathcal{E}_n := \{(i, j) : 1 \leq i \leq n, 1 \leq j \leq n, i \neq j\}$$

**$X$ -descent** of  $w = a_1 \cdots a_n \in \mathfrak{S}_n$ : an index  $1 \leq i \leq n - 1$  for which  $(a_i, a_{i+1}) \in X$

**$X$ -descent set  $X\text{Des}(w)$** : set of  $X$ -descents



## $X$ -descent sets

$$X \subseteq \mathcal{E}_n := \{(i, j) : 1 \leq i \leq n, 1 \leq j \leq n, i \neq j\}$$

**$X$ -descent** of  $w = a_1 \cdots a_n \in \mathfrak{S}_n$ : an index  $1 \leq i \leq n - 1$  for which  $(a_i, a_{i+1}) \in X$

**$X$ -descent set  $X\text{Des}(w)$** : set of  $X$ -descents

**Example.** (a)  $X = \{(i, j) : n - 1 \geq i > j \geq 1\}$ :  $X\text{Des} = \text{Des}$  (the ordinary descent set)

(b)  $X = \{(i, j) \in [n] \times [n] : i \neq j\}$ :  $X\text{Des}(w) = [n - 1]$ , where  $[n - 1] = \{1, 2, \dots, n - 1\}$

# Symmetric functions

**Symmetric function:**  $f = f(x_1, x_2, \dots)$ , a power series of bounded degree with rational coefficients, invariant under any permutation of the  $x_i$ 's.

**partition** of  $n$ :  $\lambda = (\lambda_1, \lambda_2, \dots)$ ,  $\lambda_1 \geq \lambda_2 \geq \dots \geq 0$ ,  $\sum \lambda_i = n$ , denoted  $\lambda \vdash n$

# Symmetric functions

**Symmetric function:**  $f = f(x_1, x_2, \dots)$ , a power series of bounded degree with rational coefficients, invariant under any permutation of the  $x_i$ 's.

**partition** of  $n$ :  $\lambda = (\lambda_1, \lambda_2, \dots)$ ,  $\lambda_1 \geq \lambda_2 \geq \dots \geq 0$ ,  $\sum \lambda_i = n$ , denoted  $\lambda \vdash n$

**Example. Power sums:**  $p_k = \sum_i x_i^k$  (with  $p_0 = 1$ ),

$$p_\lambda = p_{\lambda_1} p_{\lambda_2} \cdots,$$

a  $\mathbb{Q}$ -basis for the space of symmetric functions

# Symmetric functions

**Symmetric function:**  $f = f(x_1, x_2, \dots)$ , a power series of bounded degree with rational coefficients, invariant under any permutation of the  $x_i$ 's.

**partition** of  $n$ :  $\lambda = (\lambda_1, \lambda_2, \dots)$ ,  $\lambda_1 \geq \lambda_2 \geq \dots \geq 0$ ,  $\sum \lambda_i = n$ , denoted  $\lambda \vdash n$

**Example. Power sums:**  $p_k = \sum_i x_i^k$  (with  $p_0 = 1$ ),

$$p_\lambda = p_{\lambda_1} p_{\lambda_2} \cdots,$$

a  $\mathbb{Q}$ -basis for the space of symmetric functions

**Schur functions**  $s_\lambda$ : another  $\mathbb{Q}$ -basis, not defined here

## A generating function for the XDescent set

Define  $U_X = \sum_{w \in \mathfrak{S}_n} F_{\text{XDes}(w)}$ .

## A generating function for the XDescent set

Define  $U_X = \sum_{w \in \mathfrak{S}_n} F_{\text{XDes}(w)}$ .

**Example.**  $n = 3$ ,  $X = \{(1, 3), (2, 1), (3, 1), (3, 2)\}$

$w$	$\text{XDes}(w)$
123	$\emptyset$
132	$\{1, 2\}$
213	$\{1, 2\}$
231	$\{2\}$
312	$\{1\}$
321	$\{1, 2\}$

$$U_X = F_\emptyset + F_1 + F_2 + 3F_{1,2} = p_1^3 - p_2 p_1 + p_3 = s_3 + s_{21} + 2s_{111}$$

## First easy theorem

**Theorem.** (a)  $U_X$  is a  $p$ -integral symmetric function, i.e.,  
 $U_X = \sum_{\lambda} c_{\lambda} p_{\lambda}$ , where  $c_{\lambda} \in \mathbb{Z}$ .

## First easy theorem

**Theorem.** (a)  $U_X$  is a  $p$ -integral symmetric function, i.e.,  
 $U_X = \sum_{\lambda} c_{\lambda} p_{\lambda}$ , where  $c_{\lambda} \in \mathbb{Z}$ .

**Proof.** Consider the coefficient of a monomial, say  $\mathbf{m} = x_1^2 x_2^3 x_4^2$   
(where  $n = 7$ ). Recall

$$U_X = \sum_{w \in \mathfrak{S}_n} F_{\text{XDes}(w)}$$

$$F_S = \sum_{\substack{1 \leq i_1 \leq i_2 \leq \dots \leq i_n \\ i_j < i_{j+1} \text{ if } j \in S}} x_{i_1} x_{i_2} \cdots x_{i_n}.$$



## First easy theorem

**Theorem.** (a)  $U_X$  is a  $p$ -integral symmetric function, i.e.,  
 $U_X = \sum_{\lambda} c_{\lambda} p_{\lambda}$ , where  $c_{\lambda} \in \mathbb{Z}$ .

**Proof.** Consider the coefficient of a monomial, say  $\mathbf{m} = x_1^2 x_2^3 x_4^2$   
(where  $n = 7$ ). Recall

$$U_X = \sum_{w \in \mathfrak{S}_n} F_{\text{XDes}(w)}$$

$$F_S = \sum_{\substack{1 \leq i_1 \leq i_2 \leq \dots \leq i_n \\ i_j < i_{j+1} \text{ if } j \in S}} x_{i_1} x_{i_2} \cdots x_{i_n}.$$

Let  $w = a_1 a_2 \cdots a_7$ . Thus  $\mathbf{m}$  appears in  $F_{\text{XDes}(w)}$  if and only if  
 $(a_1, a_2), (a_3, a_4), (a_4, a_5), (a_6, a_7) \notin X$ .

## Proof continued

Let  $w = a_1 a_2 \cdots a_7$ . Thus  $m$  appears in  $F_{\text{XDes}(w)}$  if and only if  $(a_1, a_2), (a_3, a_4), (a_4, a_5), (a_6, a_7) \notin X$ .

## Proof continued

Let  $w = a_1 a_2 \cdots a_7$ . Thus  $m$  appears in  $F_{\text{XDes}(w)}$  if and only if  $(a_1, a_2), (a_3, a_4), (a_4, a_5), (a_6, a_7) \notin X$ .

Write  $w = a_1 a_2 \cdot a_3 a_4 a_5 \cdot a_6 a_7 = u_1 u_2 u_3$  (juxtaposition of words).  
Then  $x_1^3 x_2^2 x_4^2$  appears in  $F_{\text{XDes}(w')}$ , where  $w' = u_2 u_1 u_3$ .  
Generalizing shows that  $U_X$  is a symmetric function.

## Proof continued

Let  $w = a_1 a_2 \cdots a_7$ . Thus  $\mathfrak{m}$  appears in  $F_{\text{XDes}(w)}$  if and only if  $(a_1, a_2), (a_3, a_4), (a_4, a_5), (a_6, a_7) \notin X$ .

Write  $w = a_1 a_2 \cdot a_3 a_4 a_5 \cdot a_6 a_7 = u_1 u_2 u_3$  (juxtaposition of words). Then  $x_1^3 x_2^2 x_4^2$  appears in  $F_{\text{XDes}(w')}$ , where  $w' = u_2 u_1 u_3$ . Generalizing shows that  $U_X$  is a symmetric function.

Also  $x_1^2 x_2^3 x_4^2 = \mathfrak{m}$  appears in  $F_{\text{XDes}(w'')}$ , where  $w'' = u_3 u_2 u_1$ . Generalizing shows that the coefficient of  $x_1^{\alpha_1} x_2^{\alpha_2} \cdots$  in  $U_X$  is an integer multiple of  $\alpha_1! \alpha_2! \cdots$ .

## Proof continued

Let  $w = a_1 a_2 \cdots a_7$ . Thus  $\mathfrak{m}$  appears in  $F_{\text{XDes}(w)}$  if and only if  $(a_1, a_2), (a_3, a_4), (a_4, a_5), (a_6, a_7) \notin X$ .

Write  $w = a_1 a_2 \cdot a_3 a_4 a_5 \cdot a_6 a_7 = u_1 u_2 u_3$  (juxtaposition of words). Then  $x_1^3 x_2^2 x_4^2$  appears in  $F_{\text{XDes}(w')}$ , where  $w' = u_2 u_1 u_3$ . Generalizing shows that  $U_X$  is a symmetric function.

Also  $x_1^2 x_2^3 x_4^2 = \mathfrak{m}$  appears in  $F_{\text{XDes}(w'')}$ , where  $w'' = u_3 u_2 u_1$ . Generalizing shows that the coefficient of  $x_1^{\alpha_1} x_2^{\alpha_2} \cdots$  in  $U_X$  is an integer multiple of  $\alpha_1! \alpha_2! \cdots$ .

Well-known and easy that this implies  $U_X$  is  $p$ -integral (given that  $U_X$  is a symmetric function).  $\square$

## Second easy theorem

$\omega$ : linear transformation on symmetric functions given by  
 $\omega(p_\lambda) = (-1)^{n-\ell(\lambda)} p_\lambda$  for  $\lambda \vdash n$ , where  $\ell(\lambda) = \#\{i : \lambda_i > 0\}$ .

## Second easy theorem

$\omega$ : linear transformation on symmetric functions given by  $\omega(p_\lambda) = (-1)^{n-\ell(\lambda)} p_\lambda$  for  $\lambda \vdash n$ , where  $\ell(\lambda) = \#\{i : \lambda_i > 0\}$ .

**Note.**  $\omega^2 = 1$

## Second easy theorem

$\omega$ : linear transformation on symmetric functions given by  $\omega(p_\lambda) = (-1)^{n-\ell(\lambda)} p_\lambda$  for  $\lambda \vdash n$ , where  $\ell(\lambda) = \#\{i : \lambda_i > 0\}$ .

**Note.**  $\omega^2 = 1$

**Theorem.** Let  $\overline{X} = \mathcal{E}_n - X$ . Then  $\omega U_X = U_{\overline{X}}$ .



## Second easy theorem

$\omega$ : linear transformation on symmetric functions given by  $\omega(p_\lambda) = (-1)^{n-\ell(\lambda)} p_\lambda$  for  $\lambda \vdash n$ , where  $\ell(\lambda) = \#\{i : \lambda_i > 0\}$ .

**Note.**  $\omega^2 = 1$

**Theorem.** Let  $\overline{X} = \mathcal{E}_n - X$ . Then  $\omega U_X = U_{\overline{X}}$ .

**Proof.**

## Second easy theorem

$\omega$ : linear transformation on symmetric functions given by  $\omega(p_\lambda) = (-1)^{n-\ell(\lambda)} p_\lambda$  for  $\lambda \vdash n$ , where  $\ell(\lambda) = \#\{i : \lambda_i > 0\}$ .

**Note.**  $\omega^2 = 1$

**Theorem.** Let  $\overline{X} = \mathcal{E}_n - X$ . Then  $\omega U_X = U_{\overline{X}}$ .

**Proof.** Exercise.

## Special case

**record set**  $\text{rec}(w)$  for  $w = a_1 \cdots a_n \in \mathfrak{S}_n$ :

$\text{rec}(w) = \{0 \leq i \leq n-1 : a_i > a_j \text{ for all } j < i\}$ . Thus always  $0 \in \text{rec}(w)$ .

**record partition**  $\text{rp}(w)$ : if  $\text{rec}(w) = \{r_0, \dots, r_j\}_<$ , then  $\text{rp}(w)$  is the numbers  $r_1 - r_0, r_2 - r_1, \dots, n - r_j$  arranged in decreasing order.

## Special case

**record set**  $\text{rec}(w)$  for  $w = a_1 \cdots a_n \in \mathfrak{S}_n$ :

$\text{rec}(w) = \{0 \leq i \leq n-1 : a_i > a_j \text{ for all } j < i\}$ . Thus always  $0 \in \text{rec}(w)$ .

**record partition**  $\text{rp}(w)$ : if  $\text{rec}(w) = \{r_0, \dots, r_j\}_<$ , then  $\text{rp}(w)$  is the numbers  $r_1 - r_0, r_2 - r_1, \dots, n - r_j$  arranged in decreasing order.

**Theorem** (conjectured by **RS**, proved by **I. Gessel**). *Let  $X$  have the property that if  $(i, j) \in X$  then  $i > j$ . Then*

$$U_X = \sum_{\substack{w \in \mathfrak{S}_n \\ X\text{Des}(w) = \emptyset}} p_{\text{rp}(w)}.$$

*In particular,  $U_X$  is  $p$ -positive.*

## An example

$$n = 4, X = \{(2, 1), (3, 2), (4, 3)\}$$

$w$	$\text{rec}(w)$
1234	1111
1342	211
1423	31
2314	211
2341	211
2413	31
3124	31
3142	22
3412	31
4123	4
4231	4

## An example

$$n = 4, X = \{(2, 1), (3, 2), (4, 3)\}$$

$w$	$\text{rec}(w)$
1234	1111
1342	211
1423	31
2314	211
2341	211
2413	31
3124	31
3142	22
3412	31
4123	4
4231	4

$$\Rightarrow U_X = p_1^4 + 3p_2p_1^2 + 4p_3p_1 + p_2^2 + 2p_4$$

## A generalization

**Theorem (D. Grinberg)** *Suppose that  $(i, j) \in X \Rightarrow (j, i) \notin X$ .  
Then  $U_X$  is  $p$ -positive.*

## A generalization

**Theorem (D. Grinberg)** *Suppose that  $(i, j) \in X \Rightarrow (j, i) \notin X$ .  
Then  $U_X$  is  $p$ -positive.*

In fact, Grinberg has a combinatorial interpretation of the coefficients (not given here).



## Connection with chromatic symmetric functions

$P$ : partial ordering of  $[n]$

$$Y_P = \{(i, j) : i >_P j\}$$

$\text{inc}(P)$ : incomparability graph of  $P$ , i.e., vertex set  $[n]$ , edges  $ij$  if  $i \parallel j$  in  $P$

$X_G$ : chromatic symmetric function of the graph  $G$  (generalizes the chromatic polynomial)

## Connection with chromatic symmetric functions

$P$ : partial ordering of  $[n]$

$$Y_P = \{(i, j) : i >_P j\}$$

$\text{inc}(P)$ : incomparability graph of  $P$ , i.e., vertex set  $[n]$ , edges  $ij$  if  $i \parallel j$  in  $P$

$X_G$ : chromatic symmetric function of the graph  $G$  (generalizes the chromatic polynomial)

**Theorem.**  $U_{Y_P} = X_{\text{inc}(P)}$

## Reverse succession-free permutations

Let  $X = \{(2, 1), (3, 2), \dots, (n, n - 1)\}$ .

$f_n = \#\{w \in \mathfrak{S}_n : \text{XDes}(w) = \emptyset\}$  (**rs-free** permutations)

## Reverse succession-free permutations

Let  $X = \{(2, 1), (3, 2), \dots, (n, n - 1)\}$ .

$f_n = \#\{w \in \mathfrak{S}_n : \text{XDes}(w) = \emptyset\}$  (**rs-free** permutations)

**Known result.** 
$$\sum_{n \geq 0} f_n \frac{x^n}{n!} = \frac{e^{-x}}{(1-x)^2}$$

## Reverse succession-free permutations

Let  $X = \{(2, 1), (3, 2), \dots, (n, n - 1)\}$ .

$f_n = \#\{w \in \mathfrak{S}_n : \text{XDes}(w) = \emptyset\}$  (**rs-free** permutations)

**Known result.** 
$$\sum_{n \geq 0} f_n \frac{x^n}{n!} = \frac{e^{-x}}{(1-x)^2}$$

**Theorem.** 
$$U_X = \sum_{i=1}^n f_i s_{i, 1^{n-i}}$$

(generating function for  $w \in \mathfrak{S}_n$  by positions of reverse successions)

## Reverse succession-free permutations

Let  $X = \{(2, 1), (3, 2), \dots, (n, n - 1)\}$ .

$f_n = \#\{w \in \mathfrak{S}_n : \text{XDes}(w) = \emptyset\}$  (**rs-free** permutations)

**Known result.** 
$$\sum_{n \geq 0} f_n \frac{x^n}{n!} = \frac{e^{-x}}{(1-x)^2}$$

**Theorem.** 
$$U_X = \sum_{i=1}^n f_i s_{i, 1^{n-i}}$$

(generating function for  $w \in \mathfrak{S}_n$  by positions of reverse successions)

**Example.**  $n = 4$ :  $U_X = 11s_4 + 3s_{31} + s_{211} + s_{1111}$

## Sketch of proof

**Theorem.**  $U_X = \sum_{i=1}^n f_i s_{i,1}^{n-i}$

## Sketch of proof

**Theorem.** 
$$U_X = \sum_{i=1}^n f_i s_{i,1^{n-i}}$$

**Proof.** For  $S \subseteq [n-1]$ , take coefficient of  $F_S$  on both sides.



## Sketch of proof

**Theorem.** 
$$U_X = \sum_{i=1}^n f_i s_{i,1^{n-i}}$$

**Proof.** For  $S \subseteq [n-1]$ , take coefficient of  $F_S$  on both sides.

**Left-hand side:**  $\#\{w \in \mathfrak{S}_n : \text{XDes}(w) = S\}$

## Sketch of proof

**Theorem.**  $U_X = \sum_{i=1}^n f_i s_{i,1^{n-i}}$

**Proof.** For  $S \subseteq [n-1]$ , take coefficient of  $F_S$  on both sides.

**Left-hand side:**  $\#\{w \in \mathfrak{S}_n : \text{XDes}(w) = S\}$

**Right-hand side:** Use

$$s_{i,1^{n-i}} = \sum_{S \in \binom{[n-1]}{n-i}} F_S.$$

To show:  $f_i = \#\{w \in \mathfrak{S}_n : \text{XDes}(w) = S\}$  if  $\#S = n - i$ .

## Conclusion of proof

To show:  $f_i = \#\{w \in \mathfrak{S}_n : \text{XDes}(w) = S\}$  if  $\#S = n - i$ .

## Conclusion of proof

To show:  $f_i = \#\{w \in \mathfrak{S}_n : \text{XDes}(w) = S\}$  if  $\#S = n - i$ .

Will define a bijection

$$\{w \in \mathfrak{S}_n : \text{XDes}(w) = S\} \rightarrow \{u \in \mathfrak{S}_i : \text{XDes}(u) = \emptyset\}.$$

## Conclusion of proof

To show:  $f_i = \#\{w \in \mathfrak{S}_n : \text{XDes}(w) = S\}$  if  $\#S = n - i$ .

Will define a bijection

$$\{w \in \mathfrak{S}_n : \text{XDes}(w) = S\} \rightarrow \{u \in \mathfrak{S}_i : \text{XDes}(u) = \emptyset\}.$$

**Example.**  $w = 3247651$ , so  $S = \{1, 4, 5\}$ ,  $n = 7$ ,  $i = 4$ . Factor  $w$ :

$$w = 32 \cdot 4 \cdot 765 \cdot 1.$$

Let  $1 \rightarrow 1$ ,  $32 \rightarrow 2$ ,  $4 \rightarrow 3$ ,  $765 \rightarrow 4$ : get

$$w \rightarrow 2341 = u. \quad \square$$

## A $q$ -analogue for $X = \{(2, 1), (3, 2), \dots, (n, n - 1)\}$

Let  $U_X(q) = \sum_{w \in \mathfrak{S}_n} q^{\text{des}(w^{-1})} F_{X\text{Des}(w)}$ , where  $\text{des}$  denotes the number of (ordinary) descents.

$U_X(q)$  is the generating function for  $w \in \mathfrak{S}_n$  by positions of reverse successions and by  $\text{des}(w^{-1})$ .

$$f_n(q) = \sum_{\substack{w \in \mathfrak{S}_n \\ X\text{Des}(w) = \emptyset}} q^{\text{des}(w^{-1})}$$

## A $q$ -analogue for $X = \{(2, 1), (3, 2), \dots, (n, n - 1)\}$

Let  $U_X(q) = \sum_{w \in \mathfrak{S}_n} q^{\text{des}(w^{-1})} F_{X\text{Des}(w)}$ , where  $\text{des}$  denotes the number of (ordinary) descents.

$U_X(q)$  is the generating function for  $w \in \mathfrak{S}_n$  by positions of reverse successions and by  $\text{des}(w^{-1})$ .

$$f_n(q) = \sum_{\substack{w \in \mathfrak{S}_n \\ X\text{Des}(w) = \emptyset}} q^{\text{des}(w^{-1})}$$

**Theorem.**  $U_X(q) = \sum_{i=1}^n q^{n-i} f_i(q) s_{i, 1^{n-i}}$

## Digraph interpretation

We can also regard  $X$  as a **digraph**, with edges  $i \rightarrow j$  if  $(i, j) \in X$ .

A **Hamiltonian path** in  $X$  is a permutation  $a_1 a_2 \cdots a_n \in \mathfrak{S}_n$  such that  $(a_i, a_{i+1}) \in X$  for  $1 \leq i \leq n - 1$ . Define

$$\mathbf{ham}(X) = \# \text{ Hamiltonian paths in } X$$



# Digraph interpretation

We can also regard  $X$  as a **digraph**, with edges  $i \rightarrow j$  if  $(i, j) \in X$ .

A **Hamiltonian path** in  $X$  is a permutation  $a_1 a_2 \cdots a_n \in \mathfrak{S}_n$  such that  $(a_i, a_{i+1}) \in X$  for  $1 \leq i \leq n - 1$ . Define

$$\mathbf{ham}(X) = \# \text{ Hamiltonian paths in } X$$

NOTE.

- ▶  $w \in \mathfrak{S}_n$  is a Hamiltonian path in  $X$  if and only if  $X\text{Des}(w) = [n - 1]$ .

## Digraph interpretation

We can also regard  $X$  as a **digraph**, with edges  $i \rightarrow j$  if  $(i, j) \in X$ .

A **Hamiltonian path** in  $X$  is a permutation  $a_1 a_2 \cdots a_n \in \mathfrak{S}_n$  such that  $(a_i, a_{i+1}) \in X$  for  $1 \leq i \leq n - 1$ . Define

$$\mathbf{ham}(X) = \# \text{ Hamiltonian paths in } X$$

NOTE.

- ▶  $w \in \mathfrak{S}_n$  is a Hamiltonian path in  $X$  if and only if  $X\text{Des}(w) = [n - 1]$ .
- ▶  $w$  is a Hamiltonian path in  $\overline{X}$  if and only if  $X\text{Des}(w) = \emptyset$ .

## Connection with $U_X$

**Theorem.** Let  $U_X = \sum_{\lambda} c_{\lambda} p_{\lambda}$ . Then  $\text{ham}(\overline{X}) = \sum_{\lambda} c_{\lambda}$ .

## Connection with $U_X$

**Theorem.** Let  $U_X = \sum_{\lambda} c_{\lambda} p_{\lambda}$ . Then  $\text{ham}(\overline{X}) = \sum_{\lambda} c_{\lambda}$ .

**Proof.** Recall  $U_X = \sum_{w \in \mathfrak{S}_n} F_{\text{XDes}(w)}$ . Since  $w \in \mathfrak{S}_n$  is a Hamiltonian path in  $\overline{X}$  if and only if  $\text{XDes}(w) = \emptyset$ ,

$$\text{ham}(\overline{X}) = \#\{w \in \mathfrak{S}_n : \text{XDes}(w) = \emptyset\}.$$

Note

$$[x_1^n] F_S = \begin{cases} 1, & S = \emptyset \\ 0, & \text{otherwise.} \end{cases}$$

Also for  $\lambda \vdash n$ ,  $[x_1^n] p_{\lambda} = 1$ .

## Connection with $U_X$

**Theorem.** Let  $U_X = \sum_{\lambda} c_{\lambda} p_{\lambda}$ . Then  $\text{ham}(\bar{X}) = \sum_{\lambda} c_{\lambda}$ .

**Proof.** Recall  $U_X = \sum_{w \in \mathfrak{S}_n} F_{\text{XDes}(w)}$ . Since  $w \in \mathfrak{S}_n$  is a Hamiltonian path in  $\bar{X}$  if and only if  $\text{XDes}(w) = \emptyset$ ,

$$\text{ham}(\bar{X}) = \#\{w \in \mathfrak{S}_n : \text{XDes}(w) = \emptyset\}.$$

Note

$$[x_1^n] F_S = \begin{cases} 1, & S = \emptyset \\ 0, & \text{otherwise.} \end{cases}$$

Also for  $\lambda \vdash n$ ,  $[x_1^n] p_{\lambda} = 1$ .

Take coefficient of  $x_1^n$  on both sides of

$$U_X = \sum_{w \in \mathfrak{S}_n} F_{\text{XDes}(w)} = \sum_{\lambda} c_{\lambda} p_{\lambda}. \quad \square$$

## Simple corollary

**Corollary.** Let  $U_X = \sum_{\lambda} c_{\lambda} p_{\lambda}$  as before. Then

$$\text{ham}(X) = \sum_{\lambda} (-1)^{n-\ell(\lambda)} c_{\lambda}.$$

## Simple corollary

**Corollary.** Let  $U_X = \sum_{\lambda} c_{\lambda} p_{\lambda}$  as before. Then

$$\text{ham}(X) = \sum_{\lambda} (-1)^{n-\ell(\lambda)} c_{\lambda}.$$

Recall  $\omega p_{\lambda} = (-1)^{n-\ell(\lambda)} p_{\lambda}$  and  $\omega U_X = U_{\bar{X}}$ . Now apply  $\omega$  to  $U_X = \sum_{\lambda} c_{\lambda} p_{\lambda}$  and use previous theorem:

$$\text{ham}(\bar{X}) = \sum_{\lambda} c_{\lambda}. \quad \square$$

# Berge's theorem

**Theorem (C. Berge).**  $\text{ham}(X) \equiv \text{ham}(\bar{X}) \pmod{2}$

**Proof (D. Grinberg).** Let  $U_X = \sum_{\lambda} c_{\lambda} p_{\lambda}$ . To prove:

$$\sum (-1)^{n-\ell(\lambda)} c_{\lambda} \equiv \sum c_{\lambda} \pmod{2}.$$



# Berge's theorem

**Theorem (C. Berge).**  $\text{ham}(X) \equiv \text{ham}(\bar{X}) \pmod{2}$

**Proof (D. Grinberg).** Let  $U_X = \sum_{\lambda} c_{\lambda} p_{\lambda}$ . To prove:

$$\sum (-1)^{n-\ell(\lambda)} c_{\lambda} \equiv \sum c_{\lambda} \pmod{2}.$$

Obvious since  $(-1)^{n-\ell(\lambda)} = \pm 1$ .  $\square$

# Tournaments

**tournament:** a digraph  $X$  with vertex set  $[n]$  (say), such that for all  $1 \leq i < j \leq n$ , exactly one of  $(i, j) \in X$  or  $(j, i) \in X$ .

# Tournaments

**tournament**: a digraph  $X$  with vertex set  $[n]$  (say), such that for all  $1 \leq i < j \leq n$ , exactly one of  $(i, j) \in X$  or  $(j, i) \in X$ .

**Theorem (D. Grinberg)**. *Let  $X$  be a tournament. Then*

$$U_X = \sum_w 2^{\text{nsc}(w)} p_{\rho(w)},$$

where  $w$  ranges over all permutations in  $\mathfrak{S}_n$  of odd order such that every nonsingleton cycle of  $w$  is a (directed) cycle of  $X$ , and where  $\text{nsc}(w)$  denotes the number of nonsingleton cycles of  $w$ .

# Tournaments

**tournament**: a digraph  $X$  with vertex set  $[n]$  (say), such that for all  $1 \leq i < j \leq n$ , exactly one of  $(i, j) \in X$  or  $(j, i) \in X$ .

**Theorem (D. Grinberg)**. *Let  $X$  be a tournament. Then*

$$U_X = \sum_w 2^{\text{nsc}(w)} p_{\rho(w)},$$

where  $w$  ranges over all permutations in  $\mathfrak{S}_n$  of odd order such that every nonsingleton cycle of  $w$  is a (directed) cycle of  $X$ , and where  $\text{nsc}(w)$  denotes the number of nonsingleton cycles of  $w$ .

Special case of a result for **any**  $X$ .

## A corollary

**Theorem** (repeated). *Let  $X$  be a tournament. Then*

$$U_X = \sum_w 2^{\text{nsc}(w)} p_{\rho(w)},$$

*where  $w$  ranges over all permutations in  $\mathfrak{S}_n$  of odd order such that every nonsingleton cycle of  $w$  is a (directed) cycle of  $X$ , and where  $\text{nsc}(w)$  denotes the number of nonsingleton cycles of  $w$ .*

**Corollary.** *If  $X$  is a tournament, then*

$$U_X \in \mathbb{Z}[p_1, 2p_3, 2p_5, 2p_7, \dots].$$

## A corollary

**Theorem** (repeated). *Let  $X$  be a tournament. Then*

$$U_X = \sum_w 2^{\text{nsc}(w)} p_{\rho(w)},$$


*where  $w$  ranges over all permutations in  $\mathfrak{S}_n$  of odd order such that every nonsingleton cycle of  $w$  is a (directed) cycle of  $X$ , and where  $\text{nsc}(w)$  denotes the number of nonsingleton cycles of  $w$ .*

**Corollary.** *If  $X$  is a tournament, then*

$$U_X \in \mathbb{Z}[p_1, 2p_3, 2p_5, 2p_7, \dots].$$


**Note.** Thus  $U_X$  can be written uniquely as a linear combination of Schur's "shifted Schur functions"  $P_\lambda$ , where  $\lambda$  has distinct parts. Can anything worthwhile be said about the coefficients?

## An example



$w$	$2^{\text{nsc}(w)} p_{\rho(w)}$
(1)(2)(3)(4)	$p_1^4$
(1, 2, 4)(3)	$2p_3p_1$
(1, 3, 4)(2)	$2p_3p_1$

## An example



$w$	$2^{\text{nsc}(w)} p_{\rho(w)}$
(1)(2)(3)(4)	$p_1^4$
(1, 2, 4)(3)	$2p_3p_1$
(1, 3, 4)(2)	$2p_3p_1$

$$\Rightarrow U_X = p_1^4 + 4p_3p_1 = 5P_4 - 2P_{3,1}$$



## An application to Hamiltonian paths

**Observation** (repeated). Let  $U_x = \sum_{\lambda} c_{\lambda} p_{\lambda}$ . Then

$$\text{ham}(X) = \sum_{\lambda} (-1)^{n-\ell(\lambda)} c_{\lambda}.$$

**Theorem** (repeated). Let  $X$  be a tournament. Then

$$U_X = \sum_w 2^{\text{nsc}(w)} p_{\rho(w)},$$

where  $w$  ranges over all permutations in  $\mathfrak{S}_n$  of odd order such that every nonsingleton cycle of  $w$  is a (directed) cycle of  $X$ , and where  $\text{nsc}(w)$  denotes the number of nonsingleton cycles of  $w$ .

## An application to Hamiltonian paths

**Observation** (repeated). Let  $U_x = \sum_{\lambda} c_{\lambda} p_{\lambda}$ . Then

$$\text{ham}(X) = \sum_{\lambda} (-1)^{n-\ell(\lambda)} c_{\lambda}.$$

**Theorem** (repeated). Let  $X$  be a tournament. Then

$$U_X = \sum_w 2^{\text{nsc}(w)} p_{\rho(w)},$$

where  $w$  ranges over all permutations in  $\mathfrak{S}_n$  of odd order such that every nonsingleton cycle of  $w$  is a (directed) cycle of  $X$ , and where  $\text{nsc}(w)$  denotes the number of nonsingleton cycles of  $w$ .

**Corollary.** Let  $X$  be a tournament. Then

$$\text{ham}(X) = \sum_w 2^{\text{nsc}(w)}.$$

# Rédei's theorem

**Corollary** (repeated). *Let  $X$  be a tournament. Then*

$$\text{ham}(X) = \sum_w 2^{\text{ns}(w)}.$$

# Rédei's theorem

**Corollary** (repeated). *Let  $X$  be a tournament. Then*

$$\text{ham}(X) = \sum_w 2^{\text{ns}(w)}.$$

Since  $c_{1^n} = 1$  for **all**  $X$  (immediate from  $U_X = \sum_{w \in \mathfrak{S}_n} F_{X \text{Des}(w)}$ ), we conclude:

## Rédei's theorem

**Corollary** (repeated). *Let  $X$  be a tournament. Then*

$$\text{ham}(X) = \sum_w 2^{\text{nsc}(w)}.$$

Since  $c_1^n = 1$  for **all**  $X$  (immediate from  $U_X = \sum_{w \in \mathfrak{S}_n} F_{X \text{Des}(w)}$ ), we conclude:

**Theorem** (L. Rédei, 1934) *Every tournament has an odd number of Hamiltonian paths.*

## The final slide

