

MIT OpenCourseWare  
<http://ocw.mit.edu>

18.085 Computational Science and Engineering I  
Fall 2008

For information about citing these materials or our Terms of Use, visit: <http://ocw.mit.edu/terms>.

Name \_\_\_\_\_

November 4, 2002

Grading 1

2

3

-----

**Problem 1 (33 points)**

This question is about a fixed-free hanging bar (made of 2 materials) with a point load at  $x = \frac{3}{4}$ :

$$-\frac{d}{dx} \left( c(x) \frac{du}{dx} \right) = \delta \left( x - \frac{3}{4} \right)$$

$$u(0) = 0$$

$$w(1) = 0$$

Suppose that

$$c(x) = \begin{cases} 1, & x < \frac{1}{2} \\ 4, & x > \frac{1}{2} \end{cases}$$

a) Which of  $u$ ,  $\frac{du}{dx}$ , and  $w = c \frac{du}{dx}$  have jumps at (i)  $x = \frac{1}{2}$  and (ii)  $x = \frac{3}{4}$ ?

b) Solve for  $w(x)$  and draw its graph from  $x = 0$  to  $x = 1$ .

c) Solve for  $u(x)$  and draw its graph from  $x = 0$  to  $x = 1$ .

## Problem 2 (34 points)

a)

(i) Find the real part  $u(x, y)$  and the imaginary part  $s(x, y)$  of

$$f(z) = \frac{1}{z} = \frac{1}{x + iy}$$

(ii) Also find  $u(r, \theta)$  and  $s(r, \theta)$  for the same function expressed in polar coordinates:

$$f(z) = \frac{1}{z} = \frac{1}{re^{i\theta}}$$

b) Draw the equipotential curve  $u(x, y) = \frac{1}{2}$  and the streamline  $s(x, y) = \frac{1}{2}$ . (I suggest to use  $x$ - $y$  coordinates and "clear out" denominators.) What shapes are these two curves?

c) What can you say about  $u(x, y)$  (what condition does it satisfy) along the line  $s = \frac{1}{2}$ ?

### Problem 3 (33 points)

a). Suppose that the Laplacian of  $F(x,y)$  is zero:

$$\frac{\partial^2 F}{\partial x^2} + \frac{\partial^2 F}{\partial y^2} = 0.$$

Show that  $u = \frac{\partial F}{\partial y}$  and  $s = \frac{\partial F}{\partial x}$  satisfy the Cauchy-Riemann equations.

b). Which of these vector fields are gradients of some function  $u(x,y)$  and what is that function? Does  $u(x,y)$  solve Laplace's equation  $\text{div}(\text{grad } u) = 0$ ?

(i)  $v(x,y) = (x^2, y^2)$

(ii)  $v(x,y) = (y^2, x^2)$

(iii)  $v(x,y) = (x+y, x-y)$

c) (i) Find the solution to Laplace's equation inside the unit circle  $r^2 = x^2 + y^2 = 1$  if the boundary condition on the circle is  $u = u_0(\theta) = \frac{1}{2} + \cos \theta + \cos 2\theta$ . (OK to use polar coordinates.)  
(ii) Find the numerical value of the solution  $u$  at the center and at the point  $x = \frac{1}{2}, y = 0$ .