

Practice Mid Term Exam 1

OCTOBER 25, 2011

Problem 1:

Let X_1, X_2, X_3, \dots be a Markov chain on a finite state space $S = \{1, \dots, N\}$ with transition matrix P . Among the following statements, say which implies which.

- (a) There exists a probability distribution $\bar{\pi}$ such that $\lim_{n \rightarrow \infty} \pi P^n = \bar{\pi}$ for every probability distribution π .
- (b) $\lim_{n \rightarrow \infty} P^n = \begin{bmatrix} \bar{\pi} \\ \vdots \\ \bar{\pi} \end{bmatrix}$, for some probability distribution $\bar{\pi}$.
- (c) There exists a probability distribution $\bar{\pi}$ such that $\bar{\pi}P = \bar{\pi}$.
- (d) 1 is an eigenvalue of P with multiplicity 1, and all other eigenvalues λ have $|\lambda| < 1$.
- (e) $P_{ij} > 0$ for all $i, j \in S$.
- (f) There exists $n > 0$ such that $P_{ij}^n > 0$ for all $i, j \in S$.
- (g) $P_{ij}^n > 0$ for all $i, j \in S$ and $n > 0$.
- (h) For all $i, j \in S$ there exists $n > 0$ such that $P_{ij}^n > 0$.
- (i) X_1, X_2, X_3, \dots is an irreducible Markov chain.
- (j) X_1, X_2, X_3, \dots is an irreducible aperiodic Markov chain.

Problem 2:

Let X_1, X_2, X_3, \dots be a Markov chain on \mathbb{Z} such that $X_0 = 0$ and, conditioned on $X_n = i$, we have

$$X_{n+1} = \begin{cases} i - 1 & \text{with prob. } \alpha \\ i & \text{with prob. } \beta \\ i + 1 & \text{with prob. } \gamma \end{cases},$$

where $\alpha, \beta, \gamma \geq 0$ are such that $\alpha + \beta + \gamma = 1$. Let $T_k = \inf\{n \geq 0 \mid X_n = k\}$ be the time of first passage through k , and let $u_k(s) = \mathbb{E}[s^{T_k}]$, for $|s| < 1$.

- (a) Show that, for every $k \geq 1$, we have $u_k(s) = (u_1(s))^k$.
- (b) Compute $u_1(s)$.

Problem 3:

Let X_1, X_2, X_3, \dots be a Markov chain on $S = \{1, 2, \dots, 11\}$ with transition matrix

$$P = \begin{bmatrix} 1/2 & 1/2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 1/2 & 1/2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1/2 & 0 & 1/2 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 1/2 & 0 & 0 & 0 & 1/2 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1/2 & 0 & 1/2 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 1/2 & 0 & 1/2 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 0 & 1 \\ 0 & 0 & 0 & 0 & 0 & 1 & 0 & 0 & 0 & 0 & 0 \end{bmatrix}.$$

- (a) List all communicating classes, specifying if they are transient or recurrent classes.
- (b) For each recurrent communicating class, decide whether it is aperiodic or periodic, and in the latter case, find its period.

Problem 4:

Let X_1, X_2, X_3, \dots be a sequence of independent, identically distributed random variables, with values in $S = \{1, 2, 3\}$ and probability distribution $\mathbb{P}[X = 1] = 1/2$, $\mathbb{P}[X = 2] = 1/3$, $\mathbb{P}[X = 3] = 1/6$. Explain why this defines a Markov chain, and compute the transition matrix P . Compute the equilibrium distribution $\bar{\pi}$.

Problem 5: .

Customers arrive at a certain facility according to a Poisson process of rate λ . It is known that exactly 5 customers arrive in the first hour. Each customer, independently from the other customers, spends a time T in the store that is an exponential random variable of rate α , and then leave the store. Compute the probability that the store is empty at the end of this first hour.

Problem 6: .

Let $Y_n, n = 0, 1, 2, \dots$ be a Markov chain with transition matrix P , and let N_t be a Poisson process of rate λ . Consider the continuous time process $X_t = Y_{N_t}$. Argue that it is a continuous time Markov chain, and find its infinitesimal generating matrix A .