

**HOMEWORK 1 FOR 18.706, FALL 2012**  
**DUE WEDNESDAY, FEBRUARY 22.**

- (1) Show that the converse to Schur Lemma is false by constructing a three dimensional algebra  $A$  over  $\mathbb{C}$  and a two dimensional module  $M$  over  $A$ , such that  $M$  is reducible but  $\text{End}_A(M) \cong \mathbb{C}$ .

Can a module whose endomorphisms form a division ring be decomposable?
- (2) Let  $R$  be the subring in  $\text{Mat}_2(\mathbb{R})$  given by  $R = \{(a_{ij}) \mid a_{21} = 0, a_{22} \in \mathbb{Q}\}$ . Show that  $R$  is left Artinian and Noetherian but it is neither right Artinian nor right Noetherian.
- (3) This problem illustrates that a finite dimensional algebra may have indecomposable modules of arbitrarily large dimension.

Let  $k$  be a field and  $I \subset k[x, y]$  be the ideal generated by  $x, y$ . Let  $A = k[x, y]/I^2$ . Show that  $M_n = I^n/I^{n+2}$  is an indecomposable  $A$ -module. Construct an example of an infinite dimensional indecomposable  $A$ -module.
- (4) Describe the socle and co-socle filtration of the free rank one module for the following rings.
  - (a)  $R = \mathbb{Z}/72$ .
  - (b)  $R = k[D_4]$ , where  $k$  is a field of characteristic two, and  $D_4$  denotes the dihedral group of order 8 (the group of symmetries of the square).
- (5) Let  $Q$  be a quiver, i.e. a finite oriented graph. Let  $A(Q)$  be the path algebra of  $Q$  over a field  $k$ , i.e. the algebra whose basis is formed by paths in  $Q$  (compatible with orientations, and including paths of length 0 from a vertex to itself), and multiplication is concatenation of paths (if the paths cannot be concatenated, the product is zero).
  - (a) Represent the algebra of upper triangular matrices as  $A(Q)$ .
  - (b) Show that  $A(Q)$  is finite dimensional iff  $Q$  is acyclic, i.e. has no oriented cycles.
  - (c) For any acyclic  $Q$ , decompose  $A(Q)$  (as a left module) in a direct sum of indecomposable modules, and classify the simple  $A(Q)$ -modules.
  - (d) Find a condition on  $Q$  under which  $A(Q)$  is isomorphic to  $A(Q)^{op}$ , the algebra  $A(Q)$  with opposite multiplication. Use this to give an example of an algebra  $A$  that is not isomorphic to  $A^{op}$ .
- (6) This problem provides examples showing that the conclusion of the Krull-Schmidt Theorem does not hold without the finiteness assumption on the module.

- (a) Let  $R$  be a Dedekind domain<sup>1</sup> which is not a principal ideal domain (e.g.  $R = \mathbb{Z}[\sqrt{-5}]$  or  $R = \mathbb{C}[x, y]/(y^2 - x^3 - 1)$ ). Show that the conclusion of Krull-Schmidt Theorem does not hold for finitely generated projective  $R$  modules.  
(Hint: you can use that  $I \oplus J \cong R \oplus IJ$  for nonzero ideals  $I, J \subset R$ ).
- (b) Let  $A$  be the algebra of smooth real functions on the real line, such that  $a(x+1) = a(x)$ . Let  $M$  be the  $A$ -module of smooth functions on the line such that  $b(x+1) = -b(x)$ .  
Show that  $M$  is indecomposable and not isomorphic to  $A$ , and that  $M \oplus M \cong A \oplus A$  as a left  $A$ -module. Thus the conclusion of Krull-Schmidt theorem does not hold in this case.

---

<sup>1</sup>Recall that this means that  $R$  is a Noetherian (commutative) domain where every ideal is a product of prime ideals. Rings of integers in number fields provide important examples of Dedekind domains. Another class of examples comes from coordinates rings of smooth affine algebraic curves over a field.