


Brownian motion

18.S995 - L02

Typical length scales





<http://www2.estrellamountain.edu/faculty/farabee/BIOBK/biobookcell2.html>

Brownian motion



“Brownian” motion


Jan Ingen-Housz (1730-1799)



1784/1785:

über betrügen könnte, darf man nur in den Brennpunct eines Mikroskops einen Tropfen Weingeist sammt etwas gestoßener Kohle setzen; man wird diese Körperchen in einer verwirrten beständigen und heftigen Bewegung erblicken, als wenn es Thierchen wären, die sich reißend unter einander fortbewegen.


Robert Brown (1773-1858)



Linnean society, London

1827: irregular motion of pollen in fluid

<http://www.brianjford.com/wbbrownc.htm>



W. Sutherland (1858-1911)




Source: www.theage.com.au

$$D = \frac{RT}{6\pi\eta aC}$$

Phil. Mag. **9**, 781 (1905)

A. Einstein (1879-1955)



Source: wikipedia.org

$$\langle x^2(t) \rangle = 2Dt$$

$$D = \frac{RT}{N} \frac{1}{6\pi kP}$$

Ann. Phys. **17**, 549 (1905)

M. Smoluchowski
(1872-1917)





Source: wikipedia.org

$$D = \frac{32}{243} \frac{mc^2}{\pi\mu R}$$

Ann. Phys. **21**, 756 (1906)

Jean Baptiste Perrin (1870-1942, Nobel prize 1926)



- ▶ colloidal particles of radius $0.53\mu\text{m}$
- ▶ successive positions every 30 seconds joined by straight line segments
- ▶ mesh size is $3.2\mu\text{m}$

Mouvement brownien et réalité moléculaire, Annales de chimie et de physique VIII 18, 5-114 (1909)


Les Atomes, Paris, Alcan (1913)

experimental evidence for
atomistic structure of matter

Norbert Wiener

(1894-1964)


MIT




Relevance in biology

- **intra**cellular transport
- **inter**cellular transport
- microorganisms must beat BM to achieve directed locomotion
- tracer diffusion = important experimental “tool”
- generalized BMs (polymers, membranes, etc.)

Polymers & filaments ($D=1$)




Drosophila oocyte




Physical parameters
(e.g. bending rigidity)
from fluctuation
analysis


Brownian tracer particles in a bacterial suspension



Bacillus subtilis



Tracer colloids



Basic idea

Split dynamics into


- deterministic part (drift)
- random part (diffusion)

$$\dot{x} = f(t, x(t)) + \text{noise}$$

Typical problems

Determine


- noise 'structure'
- transport coefficients
- first passage (escape) times



$$\dot{x} = f(t, x(t)) + \text{noise}$$

Probability space $(\Omega, \mathcal{F}, \mathbb{P})$

$$\mathcal{F} = \{\emptyset, A, B, A \cap B, A \cup B, \dots, \Omega\}$$



$$\mathbb{P}[A \cup B] = \mathbb{P}[A] + \mathbb{P}[B] - \mathbb{P}[A \cap B]$$


Expectation values of random variables

$$X : \Omega \rightarrow \mathbb{R}^n$$

$$\mathbb{E}[f(X)] = \int d\mathbb{P} f(x) = \int dx p(x) f(x)$$

$$p(x) \geq 0, \quad \int dx p(x) = 1$$

$$\mathbb{E}[\alpha f(X) + \beta g(X)] = \alpha \mathbb{E}[f(X)] + \beta \mathbb{E}[g(X)]$$



1.1 Random walks

1.1.1 Unbiased random walk (RW)

Consider the one-dimensional unbiased RW (fixed initial position $X_0 = x_0$, N steps of length ℓ)

$$X_N = x_0 + \ell \sum_{i=1}^N S_i \quad (1.1)$$

where $S_i \in \{\pm 1\}$ are iid. random variables (RVs) with $\mathbb{P}[S_i = \pm 1] = 1/2$. Noting that ¹

$$\mathbb{E}[S_i] = -1 \cdot \frac{1}{2} + 1 \cdot \frac{1}{2} = 0, \quad (1.2)$$

$$\mathbb{E}[S_i S_j] = \delta_{ij} \mathbb{E}[S_i^2] = \delta_{ij} \left[(-1)^2 \cdot \frac{1}{2} + (1)^2 \cdot \frac{1}{2} \right] = \delta_{ij}, \quad (1.3)$$

we find for the first moment of the RW

$$\mathbb{E}[X_N] = x_0 + \ell \sum_{i=1}^N \mathbb{E}[S_i] = x_0 \quad (1.4)$$

Second moment (uncentered)

$$\begin{aligned}\mathbb{E}[X_N^2] &= \mathbb{E}\left[\left(x_0 + \ell \sum_{i=1}^N S_i\right)^2\right] \\ &= \mathbb{E}\left[x_0^2 + 2x_0\ell \sum_{i=1}^N S_i + \ell^2 \sum_{i=1}^N \sum_{j=1}^N S_i S_j\right] \\ &= x_0^2 + 2x_0 \cdot 0 + \ell^2 \sum_{i=1}^N \sum_{j=1}^N \mathbb{E}[S_i S_j] \\ &= x_0^2 + 2x_0 \cdot 0 + \ell^2 \sum_{i=1}^N \sum_{j=1}^N \delta_{ij} \\ &= x_0^2 + \ell^2 N.\end{aligned}\tag{1.5}$$

Variance


The variance (second centered moment)

$$\begin{aligned}\mathbb{E} [(X_N - \mathbb{E}[X_N])^2] &= \mathbb{E}[X_N^2 - 2X_N\mathbb{E}[X_N] + \mathbb{E}[X_N]^2] \\ &= \mathbb{E}[X_N^2] - 2\mathbb{E}[X_N]\mathbb{E}[X_N] + \mathbb{E}[X_N]^2 \\ &= \mathbb{E}[X_N^2] - \mathbb{E}[X_N]^2\end{aligned}\tag{1.6}$$

therefore grows linearly with the number of steps:

$$\mathbb{E} [(X_N - \mathbb{E}[X_N])^2] = \ell^2 N.\tag{1.7}$$

Continuum limit




$$X_N = x_0 + \ell \sum_{i=1}^N S_i$$

Let

$$x_0 = 0, \quad N = t/\tau$$

$$P(N, K) := \mathbb{P}[X_N/\ell = K]$$

Continuum limit



$$\begin{aligned}
 P(N, K) &= \left(\frac{1}{2}\right)^N \binom{N}{\frac{N-K}{2}} \\
 &= \left(\frac{1}{2}\right)^N \frac{N!}{((N+K)/2)! ((N-K)/2)!}.
 \end{aligned} \tag{1.8}$$


The associated probability density function (PDF) can be found by defining

$$p(t, x) := \frac{P(N, K)}{2\ell} = \frac{P(t/\tau, x/\ell)}{2\ell} \tag{1.9}$$

and considering limit $\tau, \ell \rightarrow 0$ such that

$$D := \frac{\ell^2}{2\tau} = \text{const}, \tag{1.10}$$

Continuum limit



(pset I)

yielding the Gaussian

$$p(t, x) \simeq \sqrt{\frac{1}{4\pi Dt}} \exp\left(-\frac{x^2}{4Dt}\right) \quad (1.11)$$

Eq. (1.11) is the fundamental solution to the diffusion equation,

$$\partial_t p_t = D \partial_{xx} p, \quad (1.12)$$

where $\partial_t, \partial_x, \partial_{xx}, \dots$ denote partial derivatives. The mean square displacement of the continuous process described by Eq. (1.11) is


$$\mathbb{E}[X(t)^2] = \int dx x^2 p(t, x) = 2Dt, \quad (1.13)$$

in agreement with Eq. (1.7).

Remark One often classifies diffusion processes by the (asymptotic) power-law growth of the mean square displacement,

$$\mathbb{E}[(X(t) - X(0))^2] \sim t^\mu. \quad (1.14)$$

- $\mu = 0$: Static process with no movement.
- $0 < \mu < 1$: Sub-diffusion, arises typically when waiting times between subsequent jumps can be long and/or in the presence of a sufficiently large number of obstacles (e.g. slow diffusion of molecules in crowded cells).
- $\mu = 1$: Normal diffusion, corresponds to the regime governed by the standard Central Limit Theorem (CLT).
- $1 < \mu < 2$: Super-diffusion, occurs when step-lengths are drawn from distributions with infinite variance (Lévy walks; considered as models of bird or insect movements).
- $\mu = 2$: Ballistic propagation (deterministic wave-like process).



1.1.2 Biased random walk (BRW)


Consider a one-dimensional hopping process on a discrete lattice (spacing ℓ), defined such that during a time-step τ a particle at position $X(t) = \ell j \in \ell\mathbb{Z}$ can either

- (i) jump a fixed distance ℓ to the left with probability λ , or
- (ii) jump a fixed distance ℓ to the right with probability ρ , or
- (iii) remain at its position x with probability $(1 - \lambda - \rho)$.

Assuming that the process is Markovian (does not depend on the past), the evolution of the associated probability vector $P(t) = (P(t, x)) = (P_j(t))$, where $x = \ell j$, is governed by the master equation

$$P(t + \tau, x) = (1 - \lambda - \rho) P(t, x) + \rho P(t, x - \ell) + \lambda P(t, x + \ell). \quad (1.15)$$

Master equations



$$P(t + \tau, x) = (1 - \lambda - \rho) P(t, x) + \rho P(t, x - \ell) + \lambda P(t, x + \ell). \quad (1.15)$$

Technically, ρ , λ and $(1 - \lambda - \rho)$ are the non-zero-elements of the corresponding transition matrix $W = (W_{ij})$ with $W_{ij} > 0$ that governs the evolution of the column probability vector $P(t) = (P_j(t)) = (P(t, y))$ by

$$P_i(t + \tau) = W_{ij} P_j(t) \quad (1.16a)$$

or, more generally, for n steps

$$P(t + n\tau) = W^n P(t). \quad (1.16b)$$

The stationary solutions are the eigenvectors of W with eigenvalue 1. To preserve normalization, one requires $\sum_i W_{ij} = 1$.

Continuum limit Define the density $p(t, x) = P(t, x)/\ell$. Assume τ, ℓ are small, so that we can Taylor-expand

$$p(t + \tau, x) \simeq p(t, x) + \tau \partial_t p(t, x) \tag{1.17a}$$

$$p(t, x \pm \ell) \simeq p(t, x) \pm \ell \partial_x p(t, x) + \frac{\ell^2}{2} \partial_{xx} p(t, x) \tag{1.17b}$$

Continuum limit Define the density $p(t, x) = P(t, x)/\ell$. Assume τ, ℓ are small, so that we can Taylor-expand

$$p(t + \tau, x) \simeq p(t, x) + \tau \partial_t p(t, x) \quad (1.17a)$$

$$p(t, x \pm \ell) \simeq p(t, x) \pm \ell \partial_x p(t, x) + \frac{\ell^2}{2} \partial_{xx} p(t, x) \quad (1.17b)$$

Neglecting the higher-order terms, it follows from Eq. (1.15) that

$$\begin{aligned} p(t, x) + \tau \partial_t p(t, x) &\simeq (1 - \lambda - \rho) p(t, x) + \\ &\rho [p(t, x) - \ell \partial_x p(t, x) + \frac{\ell^2}{2} \partial_{xx} p(t, x)] + \\ &\lambda [p(t, x) + \ell \partial_x p(t, x) + \frac{\ell^2}{2} \partial_{xx} p(t, x)]. \end{aligned} \quad (1.18)$$

Continuum limit Define the density $p(t, x) = P(t, x)/\ell$. Assume τ, ℓ are small, so that we can Taylor-expand

$$p(t + \tau, x) \simeq p(t, x) + \tau \partial_t p(t, x) \quad (1.17a)$$

$$p(t, x \pm \ell) \simeq p(t, x) \pm \ell \partial_x p(t, x) + \frac{\ell^2}{2} \partial_{xx} p(t, x) \quad (1.17b)$$

Neglecting the higher-order terms, it follows from Eq. (1.15) that

$$\begin{aligned} p(t, x) + \tau \partial_t p(t, x) &\simeq (1 - \lambda - \rho) p(t, x) + \\ &\rho [p(t, x) - \ell \partial_x p(t, x) + \frac{\ell^2}{2} \partial_{xx} p(t, x)] + \\ &\lambda [p(t, x) + \ell \partial_x p(t, x) + \frac{\ell^2}{2} \partial_{xx} p(t, x)]. \end{aligned} \quad (1.18)$$

Dividing by τ , one obtains the advection-diffusion equation

$$\partial_t p = -u \partial_x p + D \partial_{xx} p \quad (1.19a)$$

with drift velocity u and diffusion constant D given by²

$$u := (\rho - \lambda) \frac{\ell}{\tau}, \quad D := (\rho + \lambda) \frac{\ell^2}{2\tau}. \quad (1.19b)$$

Time-dependent solution

Dividing by τ , one obtains the advection-diffusion equation

$$\partial_t p = -u \partial_x p + D \partial_{xx} p \quad (1.19a)$$

with drift velocity u and diffusion constant D given by²

$$u := (\rho - \lambda) \frac{\ell}{\tau}, \quad D := (\rho + \lambda) \frac{\ell^2}{2\tau}. \quad (1.19b)$$

We recover the classical diffusion equation (1.12) from Eq. (1.19a) for $\rho = \lambda = 0.5$. The time-dependent fundamental solution of Eq. (1.19a) reads

$$p(t, x) = \sqrt{\frac{1}{4\pi Dt}} \exp\left(-\frac{(x - ut)^2}{4Dt}\right) \quad (1.20)$$

Remarks Note that Eqs. (1.12) and Eq. (1.19a) can both be written in the current-form

$$\partial_t p + \partial_x j_x = 0 \quad (1.21)$$

with

$$j_x = up - D\partial_x p, \quad (1.22)$$

reflecting conservation of probability. Another commonly-used representation is

$$\partial_t p = \mathcal{L}p, \quad (1.23)$$

where \mathcal{L} is a linear differential operator; in the above example (1.19b)

$$\mathcal{L} := -u \partial_x + D \partial_{xx}. \quad (1.24)$$

Stationary solutions, if they exist, are eigenfunctions of \mathcal{L} with eigenvalue 0.

(useful later when discussing Brownian motors)