

INTERSECTION PATTERNS OF CURVES

JACOB FOX, JÁNOS PACH, AND CSABA D. TÓTH

ABSTRACT

We prove that for every $k \in \mathbb{N}$ there is a constant $c_k > 0$ with the following property. Every set of $n > 1$ continuous curves in the plane, any pair of which intersect in at most k points, has two disjoint subsets A and B , each of size at least $c_k n$, such that either every curve in A intersects all curves in B , or no curve in A intersects any curve in B . This statement does not remain true if we drop the condition on the number of intersection points per pair.

1. Introduction

A classical result of Erdős and Szekeres [10] in Ramsey theory states that every graph on n vertices contains a clique or an independent set of size[†] at least $\frac{1}{2} \log n$. This bound is tight up to a constant factor: Erdős [7] showed that there exists a graph on n vertices, for every integer $n > 1$, with no clique or independent set with more than $2 \log n$ vertices. Erdős and Hajnal [8] proved that certain graphs contain much larger cliques or independent sets: For every hereditary family \mathcal{F} of graphs other than the family of all graphs, there is a constant $c(\mathcal{F}) > 0$ such that every graph in \mathcal{F} on n vertices contains a clique or an independent set of size at least $e^{c(\mathcal{F})\sqrt{\log n}}$. (A family of graphs is *hereditary* if it is closed under taking induced subgraphs.) They also asked whether this bound can be improved to $n^{c(\mathcal{F})}$. This motivates the following definitions.

DEFINITION 1. A family \mathcal{F} of finite graphs has the *Erdős-Hajnal property* if there is a constant $c(\mathcal{F}) > 0$ such that every graph in \mathcal{F} on n vertices contains a clique or an independent set of size at least $n^{c(\mathcal{F})}$.

A complete bipartite graph is said to be *balanced* if its vertex classes differ in size by at most one. A balanced complete bipartite graph with n vertices is called a *bi-clique of size n* .

DEFINITION 2. A family \mathcal{F} of finite graphs has the *strong Erdős-Hajnal property* if there is a constant $b(\mathcal{F}) > 0$ such that for every graph G in \mathcal{F} on $n > 1$ vertices, G or its complement \overline{G} contains a bi-clique of size at least $b(\mathcal{F})n$.

Alon *et al.* [2] proved that if a hereditary family of graphs has the strong Erdős-Hajnal property, then it also has the Erdős-Hajnal property. For partial results on the Erdős-Hajnal problem, see [3], [4], [5], and [9].

Given a family \mathcal{S} of subsets of \mathbb{R}^d , its *intersection graph* $G_{\mathcal{S}}$ is defined as the graph on the vertex set \mathcal{S} , in which two vertices $S, T \in \mathcal{S}$ are connected by an edge if and only if S and T

2000 *Mathematics Subject Classification* 05D10, 05C62 (primary), 05C35, 06A07 (secondary).

The first author is supported by an NSF Graduate Research Fellowship and a Princeton Centennial Fellowship. The second author is supported by NSF Grant CCF-05-14079, and by grants from NSA, PSC-CUNY, the Hungarian Research Foundation OTKA, and BSF.

[†]All logarithms in this paper are of base two.

have nonempty intersection. As noted by Ehrlich, Even, and Tarjan [6], not every graph can be realized as an intersection graph of connected sets in the plane. For instance, the bipartite graph on 15 vertices formed by replacing each edge of K_5 by a path of length 2 has no such realization. This implies, using the above result of Erdős and Hajnal, that the intersection graph of any n connected sets in the plane contains a clique or an independent set of size $e^{c\sqrt{\log n}}$, for some absolute constant $c > 0$. This general bound has been improved for intersection graphs of certain geometric objects. Pach and Solymosi [?] proved that the family of intersection graphs of line segments in the plane has the strong Erdős-Hajnal property. Later, Alon *et al.* [2] generalized this result and showed that the intersection graphs of semialgebraic sets of constant description complexity in \mathbb{R}^d also have the strong Erdős-Hajnal property. Recently, Basu [4] has further extended this result to a broader class of algebraically defined sets.

A (simple) *curve* in the plane is the range of a continuous (injective) function $f : I \rightarrow \mathbb{R}^2$ whose domain is a closed interval $I \subset \mathbb{R}$. A collection of curves is *k-intersecting* if any two of them intersect in at most k points. A 1-intersecting collection of curves is also called a family of *pseudo-segments*. A collection of connected portions of algebraic curves of maximum degree d in general position is d^2 -intersecting (by Bézout's theorem). Clearly, the family of intersection graphs of k -intersecting collections of curves is hereditary. The main result of this paper is the following.

THEOREM 1.1. *Let C be a k -intersecting collection of $n > 1$ curves in the plane such that the number of intersecting pairs of curves is at least εn^2 , for some $\varepsilon > 0$. Then the intersection graph of C contains a bi-clique of size at least $c_k \varepsilon^c n$, where c is an absolute constant and $c_k > 0$ depends only on k .*

The constant c in Theorem 1.1 can be taken to be 64, but probably this value is far from optimal.

A *separator* for a graph $G = (V, E)$ is a subset $V_0 \subset V$ such that there is a partition $V = V_0 \cup V_1 \cup V_2$ with $|V_1|, |V_2| \leq \frac{2}{3}|V|$ and no vertex in V_1 is adjacent to any vertex in V_2 . According to the Lipton-Tarjan separator theorem [18], every planar graph with n vertices has a separator of size $O(\sqrt{n})$. Fox and Pach [14] proved the following generalization of this result: The intersection graph of any finite collection of curves in the plane with a total of m intersection points (*crossings*) among them has a separator of size $O(\sqrt{m})$.

Suppose now that C is a k -intersecting collection of $n > 1$ curves in the plane that does not satisfy the condition in Theorem 1.1, for some $\varepsilon > 0$. That is, C has fewer than εn^2 intersecting pairs of curves. If $\varepsilon = \varepsilon_k$ is sufficiently small, then we can apply the above mentioned separator theorem from [14] with $m < k\varepsilon n^2$ to conclude that the complement of the intersection graph of C contains a bi-clique of size at least constant times n . Therefore, Theorem 1.1 implies

COROLLARY 1.2. *For every $k \in \mathbb{N}$, the family of intersection graphs of k -intersecting collections of curves in the plane has the strong Erdős-Hajnal property. That is, for every $k \in \mathbb{N}$, there is a constant $c_k > 0$ such that the intersection graph of every k -intersecting collection of n curves or its complement contains a bi-clique of size $c_k n$.*

The last statement settles a problem raised in [20]. Note that Corollary 1.2 is false if we drop the condition that the number of intersection points between two curves is bounded by a constant. Indeed, combining a result of Fox [11] with the fact that the incomparability graph of any partially ordered set can be obtained as the intersection graph of a collection of curves [20, 21], we can conclude that for every $\varepsilon > 0$, there is a collection of n curves in the plane whose intersection graph G contains no bi-clique of size $\frac{14n}{\varepsilon \log n}$, and \overline{G} contains no bi-clique of size n^ε .

A *Jordan region* is a subset of the plane that is homeomorphic to a closed disk. We say that a Jordan region R *contains* another Jordan region S if S lies in the interior of R . Clearly, if two

Jordan regions intersect, then either their boundary curves intersect or one contains the other in its interior. A *crossing* between a pair of Jordan regions is defined as an intersection point of their boundaries or a containment between them. (Two Jordan regions may have arbitrarily many crossings.) A family of Jordan regions is *k-intersecting* if the family of the boundary curves are *k-intersecting*.

Define an *r-region* to be a subset of the plane that is the union of at most r Jordan regions. Call these (at most r) Jordan regions of an *r-region* the *components* of the *r-region*. A family of *r-regions* is *k-intersecting* if the family of all their components are *k-intersecting*. We are assuming here that no two *r-regions* in a *k-intersecting* family of *r-regions* have a common component, although this assumption is not necessary.

We prove the following generalization of Corollary 1.2.

THEOREM 1.3. *For $r, k \in \mathbb{N}$, the family of intersection graphs of k -intersecting collections of r -regions has the strong Erdős-Hajnal property.*

Alon *et al.* [2] proved that the intersection graphs of collections of semi-algebraic sets of constant description complexity in \mathbb{R}^d have the strong Erdős-Hajnal property. Theorem 1.3 generalizes the planar case of this result. Indeed, the boundary of a semi-algebraic set of bounded description complexity in the plane is the union of a bounded number of algebraic curves of bounded degree, any two of which either intersect in a bounded number of points or overlap. By slightly perturbing the semi-algebraic sets, while maintaining their intersection pattern and their description complexity, we can assume that their boundaries do not overlap. By slightly “fattening” these sets, if necessary, we can also assume that each of them is the union of a constant number of Jordan regions.

The paper is organized as follows. Sections 2–5 add up to the proof of Theorem 1.1. Given a *k-intersecting* collection of curves, it is enough to find a large bi-clique in the intersection graph of some portions of the curves. If we truncate the curves at their intersection points with one or two special curves (called *grounds*), we obtain a set of grounded or double grounded curves (defined in Section 3). The arrangement of curves that connect points on two grounds which are disjoint except for one endpoint of one of the grounds is incident to the other ground is homeomorphic to an arrangement of curves connecting points on two parallel lines, $x = 0$ and $x = 1$, which we call *[0, 1]-curves*. Intersection graphs of *[0, 1]-curves* can, in turn, be realized as incomparability graphs.

In Section 2, we prove some auxiliary results on the structure of $K_{t,t}$ -free comparability and incomparability graphs. In Section 3, we introduce grounded and double-grounded collections of curves and prove a lemma on the intersection pattern of double grounded curves with other curves (Lemma 3.2). Section 4 is the heart of the proof of Theorem 1.1: we prove Theorem 1.1 for *[0, 1]-curves* (Theorem 4.1). In Section 5, we put the pieces of the puzzle together and prove Theorem 1.1. We do not make any serious attempt to optimize absolute constants in our statements and proofs.

Theorem 1.3 will be established in Section 6. The last section contains some applications of our results as well as a few open problems.

There are two different types of crossings of curves. An intersection point p of a pair of curves is a *proper crossing* for the pair of curves if any local deformation still preserves the existence of an intersection point. Otherwise, p is a *touching point* for the pair of curves. A collection C of curves in the plane is in *general position* if no three curves in C contain the same point, there are no touching points, and no point is an endpoint of two curves in C . Note that we can modify a pair of curves in the neighborhood of a touching point p so that the resulting curves have two proper crossings in the neighborhood of p . Therefore, if C is a *k-intersecting* collection of curves in the plane, then by slightly moving the curves in a small neighborhood of each intersection point, we obtain a $2k$ -intersecting collection C' of curves in the plane in

general position with the same intersection graph as C . Consequently, we will assume without loss of generality throughout the remainder of the paper that every collection of curves in the plane we consider will be in general position.

2. The Structure of Comparability and Incomparability Graphs

A *comparability graph* of a poset $(P, <)$ is a graph on the vertex set P , where two elements of P are adjacent if and only if they are comparable. The *incomparability graph* of a poset $(P, <)$ is defined analogously, except that now two elements of P are connected by an edge if and only if they are incomparable.

In this section, we prove a few auxiliary results on comparability graphs and incomparability graphs of partially ordered sets (*posets*). The central role that partial orders play in the analysis of intersection patterns of curves can be explained by the next lemma (Lemma 2.1). It shows that in the important special case when the curves connect two parallel lines in the strip bounded by them, their intersection pattern can be fully described by a partial order.

We use the following terminology. A $[0, 1]$ -*curve* is a curve in the plane that has one endpoint on the line $L_1 : x = 0$, the other endpoint on the line $L_2 : x = 1$, and the rest of the curve is contained in the open strip between L_1 and L_2 . A curve in the plane is called *x-monotone* if it intersects any vertical line in at most one point. Define a partial order $<$ on $[0, 1]$ -curves as follows: let $\alpha < \beta$ if α and β are disjoint and the y -coordinate of $\alpha \cap L_1$ is less than the y -coordinate of $\beta \cap L_1$ (that is, if α lies below β). For a collection C of $[0, 1]$ -curves with distinct endpoints, the intersection graph is the same as the incomparability graph of the poset $(C, <)$. It is easy to show that every incomparability graph can be represented as the intersection graph of *x-monotone* $[0, 1]$ -curves.

LEMMA 2.1. [16, 21, 20] *The following three statements are equivalent:*

- (i) G is an incomparability graph.
- (ii) G is the intersection graph of a collection of $[0, 1]$ -curves.
- (iii) G is the intersection graph of a collection of *x-monotone* $[0, 1]$ -curves. □

A graph is said to be *H-free* if it does not contain H as a (not necessarily induced) subgraph. A graph is *r-degenerate* if each of its subgraphs contains a vertex of degree at most r . It is easy to see that every r -degenerate graph with $n > r$ vertices has at most $rn - \binom{r+1}{2}$ edges.

Every poset is the intersection of its linear extensions. The *dimension* of a poset is the minimum number of linear extensions whose intersection is that poset. We use the notation $[n] := \{1, \dots, n\}$. For a permutation π of $[n]$, let $P_\pi = ([n], <_\pi)$ be the 2-dimensional partial order on $[n]$ given by $i <_\pi j$ if and only if $i < j$ and $\pi(i) < \pi(j)$.

The following lemma was established in [12]; we include its very short proof for completeness.

LEMMA 2.2. [12] *If G is a $K_{t,t}$ -free incomparability graph of a 2-dimensional poset with n elements, then G is $(2t - 2)$ -degenerate. In particular, G has at most $(2t - 2)n - \binom{2t-1}{2}$ edges.*

Proof. Since the family of incomparability graphs of 2-dimensional posets is hereditary, it is enough to find a vertex of degree at most $2t - 2$. Every 2-dimensional poset on n vertices can be realized as P_π , for a suitable permutation π . Suppose for contradiction that the degree of every vertex of the incomparability graph of P_π is at least $2t - 1$. Notice that every $i \in [n]$ is incomparable with at most $i - 1 + \pi(i) - 1$ other elements of $[n]$. Since each element $i \in [n]$ is incomparable with at least $2t - 1$ other elements of $[n]$, we have $\pi(i) \geq t + 1$ for $i \in [t]$ and $i \geq t + 1$ for $\pi(i) \in [t]$. In particular, every $i \in [t]$ is incomparable with every element j with $\pi(j) \in [t]$. Hence, the incomparability graph contains $K_{t,t}$, which is a contradiction. □

A partially ordered set (P, \prec) is called an *interval order* if one can assign an interval (x_L, x_R) to each $x \in P$ such that $x_R < y_L$ if and only if $x \prec y$.

LEMMA 2.3. *If G is a $K_{t,t}$ -free comparability graph of an interval order (P, \prec) with n elements, then G has at most $2(t-1)n$ edges.*

Proof. Let $P = \{I_1, \dots, I_n\}$ be a set of n intervals representing the interval order (P, \prec) . If there is an integer $i \in [n]$ such that at least t intervals are to the left of I_i (that is, there are at least t intervals I_j with $I_j \prec I_i$) and at least t intervals to the right of I_i , then the comparability graph of the interval order (P, \prec) contains $K_{t,t}$. So, for every $i \in [n]$, there are fewer than t intervals to the left of I_i or there are fewer than t intervals to the right of I_i . Partition P into two subsets $P = L \cup R$ such that $I_i \in L$ if and only if there are at least t intervals to the left of I_i . If L has at least t elements, then there are at least t intervals to the left of the left-most endpoint p of all intervals in L , and all intervals of L are to the right of p , which contradicts our assumption that G is $K_{t,t}$ -free. Hence, we have $|L| < t$. Since each interval in R has at most $t-1$ intervals on its left, the number of edges of G is at most $|L|n + (t-1)|R| \leq 2(t-1)n$. \square

The *edge density* of a graph with n vertices and m edges is the fraction $m/\binom{n}{2}$ of vertex pairs connected by an edge. Next, we show that if an incomparability graph is fairly dense but does not contain a large bi-clique, then it has a large induced subgraph whose edge density is close to 1.

LEMMA 2.4. *If G is a $K_{t,t}$ -free incomparability graph with n vertices and $m \geq \sqrt{5tn^3}$ edges, then G contains an induced subgraph with at least $\frac{m^2}{5n^3}$ vertices and edge density at least $1 - 10t\frac{n^3}{m^2}$.*

Proof. Label the elements of the poset with $[n]$ such that the usual order of integers is a linear extension of the partial order, that is, assume that $i < j$ whenever $i \prec j$ in the poset $([n], \prec)$. Denote by G the incomparability graph of $([n], \prec)$. Let $S \subset [n]$ be the set of integers such that $i \in S$ if and only if i is incomparable with at least $\frac{m}{n}$ other elements of $[n]$. So we have $\frac{m}{n}n + n|S| \geq 2m$, or equivalently, $|S| \geq \frac{m}{n}$.

For every $i \in S$, let $u(i)$ denote its *up-degree*, that is, the number of integers $j \in [n]$ such that $i < j$ and j is incomparable with i ; and similarly let $d(i)$ denote its *down-degree*, which is the number of integers $j \in [n]$ such that $j < i$ and j is incomparable with i . Partition S into two subsets $S = U \cup D$ such that $i \in U$ if and only if $u(i) \geq d(i)$. Notice that each element of U has up-degree at least $\frac{m}{2n}$ and each element of D has down-degree at least $\frac{m}{2n}$. Since $|S| \geq \frac{m}{n}$, we have $|U| \geq \frac{m}{2n}$ or $|D| \geq \frac{m}{2n}$. Without loss of generality, assume $|U| \geq \frac{m}{2n}$.

Partition $[n]$ into at most $\frac{5}{2}\frac{n^2}{m}$ intervals, each of length at most $\frac{m}{2n}$. Pick one of these intervals, I , that contains at least $|U|/(\frac{5}{2}\frac{n^2}{m}) \geq \frac{m^2}{5n^3}$ elements of U . We next show that the edge density of the incomparability graph of $(I \cap U, \prec)$ is close to 1. Denote by i the largest element of I . For every $j \in I \cap U$, let F_j be the set of integers f such that $j < f \leq i$ and $j \prec f$; and let H_j be the set of integers h such that $i < h \leq n$ and h is incomparable with j . Notice that every element $f \in F_j$ is incomparable with every element $h \in H_j$, otherwise $j \prec f \prec h$ would contradict that every element of H_j is incomparable with j . Notice, further, that $|F_j| \leq |H_j|$ since $\frac{m}{2n} \leq u(j) = |H_j| + (i - j - |F_j|)$ and $i - j \leq |I| \leq \frac{m}{2n}$. It follows that $|F_j| < t$, otherwise already the incomparability graph of $F_j \cup H_j$ would contain $K_{t,t}$. Therefore, each element $j \in I \cap U$ is incomparable with all but at most $t-1$ elements $k \in I$ with $j < k$. Hence, the

edge density of the subgraph induced by $I \cap U$ is at least

$$1 - \frac{(t-1)|I \cap U|}{\binom{|I \cap U|}{2}} \geq 1 - \frac{10tn^3}{m^2}.$$

□

3. Double-Grounded Curves

A collection C of curves in the plane is *grounded* if there is a closed Jordan curve γ (which is a *ground* for C) such that each curve in C has one endpoint on γ and the rest of the curve is in the exterior of γ . A collection C of curves in the plane is *double-grounded* if there are disjoint closed Jordan curves γ_1 and γ_2 such that each curve $\alpha \in C$ has one endpoint on γ_1 and the other endpoint on γ_2 , and the rest of α is disjoint from both γ_1 and γ_2 .

A collection C of n grounded curves can naturally be labeled cyclically along a ground γ . Start by assigning the label 0 to any curve in C and then proceed to label the remaining curves clockwise, breaking ties arbitrarily, so that the $(i+1)$ -th curve along γ has label i . The labels are elements of \mathbb{Z}_n . We can define the distance between a pair of grounded curves in C as the cyclic distance between the labels, that is, the distance $d(i, j)$ between the curve labeled i and the one labeled j is $\min(|i-j|, n-|i-j|)$. We let $[i, j]$ denote the cyclic interval of elements $\{i, i+1, \dots, j\}$.

LEMMA 3.1. *Suppose k be a positive integer, A is a collection of double-grounded curves with grounds γ_1 and γ_2 , and B is a collection of curves such that each curve in B intersects γ_1 and γ_2 in at most k points. If, for some $d \geq 6$,*

- (i) *each curve in A intersects at most d other curves in A ; and*
- (ii) *each curve of B intersects at least $15kd$ curves in A ,*

then there are subsets $A' \subset A$ and $B' \subset B$ of size $|A'| \geq d/3$ and $|B'| \geq \frac{d|B|}{3|A|}$ such that every curve in A' intersects every curve in B' .

Proof. We may assume without loss of generality that the Jordan regions bounded by γ_1 and γ_2 are disjoint. Define the cyclic labeling for A based on their endpoints along γ_1 . Consider two intersecting curves $\alpha_i, \alpha_j \in A$ labeled i and j , respectively. The endpoints of α_i and α_j partition γ_1 into two simple curves $\gamma_1[ij]$ and $\gamma_1[ji]$. One of $\gamma_1[ij]$ and $\gamma_1[ji]$ together with portions of α_i and α_j forms a closed Jordan curve $\hat{\gamma}_1$ that separates γ_2 from the other part of γ_1 . Every curve in A whose endpoint at γ_1 is separated from γ_2 by $\hat{\gamma}_1$ must intersect α_i or α_j . It follows that if the cyclic distance of two intersecting curves in A is at least $2d$, then one of them must intersect at least d curves of A . Since we assume that no curve in A intersects more than d other curves of A , the cyclic distance between any two intersecting curves in A is less than $2d$.

Split every curve $\beta \in B$ at its intersection point(s) with the grounds γ_1 and γ_2 into at most $2k+1$ pieces such that the relative interior of each piece is disjoint from the grounds (refer to Fig. 4). For every $\beta \in B$, let β' be a piece of β that intersects the most curves in A . So β' intersects at least $\frac{15kd}{2k+1} \geq 5d$ curves in A . Let α_i and α_j be curves that intersect β' and have maximum cyclic distance. Since β' intersects at least $5d$ curves in A , and $|A| \geq 15kd \geq 15d$, there are two curves $\alpha_i, \alpha_j \in A$ that intersect β' and whose cyclic distance is at least $5d-1$. There are cyclic intervals $\Phi \subset [i, j]$ and $\Psi \subset [j, i]$, each of cardinality at least d , such that every curve with label in Φ or Ψ has cyclic distance at least $2d$ from both α_i and α_j . Indeed, there are at least $(5d-2) - 2(2d-1) = d$ curves in each of the cyclic intervals $[i, j]$ and $[j, i]$ at distance at least $2d$ from both α_i and α_j . Every curve with label in Φ intersects β' or every curve with label in Ψ intersects β' (see Fig. 1).

Partition $\mathbb{Z}_{|A|} = I_1 \cup \dots \cup I_s$ into cyclic intervals of size at least $d/3$ and at most $d/2$. The number s of cyclic intervals is at most $3|A|/d$. For every $j \in [s]$, let A_j denote the set of curves




FIGURE 1. A collection A of double-grounded curves with grounds γ_1 and γ_2 . Label the curves cyclically by their endpoints along γ_1 . If the cyclic distance of two curves $\alpha_i, \alpha_j \in A$ is at least $5d - 1$, then every curve $\beta \in B$ that intersects both α_i and α_j must intersect all curves with label in Φ or all curves with label in Ψ .

in A whose labels are in I_j . Since every $\beta \in B$ intersects a set of curves in A whose labels belong to a cyclic interval of size d , for each $\beta \in B$, there is an index $j \in [s]$ such that β intersects every curve in A_j . By the pigeonhole principle, there is an index $j \in [s]$ and a subset $B' \subset B$ with $|B'| \geq \frac{|B|}{s} \geq \frac{d|B|}{3|A|}$ such that every curve in B' intersects every curve in A_j . Letting $A' = A_j$, our proof is complete. \square

The next lemma, which follows easily from the previous lemma, is used multiple times in the proof of Theorem 1.1.

LEMMA 3.2. *Let $k \geq 1$ and $m \geq 144$ be integers, let A be a collection of double-grounded curves with grounds γ_1 and γ_2 , and let B be a collection of curves such that every curve in B intersects γ_1 and γ_2 in at most k points. If*

- (i) *there are at most m intersecting pairs in A , and*
- (ii) *there are at least $20k\sqrt{m}|B|$ intersecting pairs in $A \times B$,*

then there are subsets $A' \subset A$ and $B' \subset B$ of size $|A'| \geq \sqrt{m}/7$ and $|B'| > km|B|/|A|^2$ such that every curve in A' intersects every curve in B' .

Proof. Let $A_1 \subseteq A$ be the set of all curves in A that intersect at most $d = \lfloor \sqrt{m}/2 \rfloor$ other curves in A . There are at most $2m/d \leq 5\sqrt{m}$ curves in $A \setminus A_1$, and so there are at most $5\sqrt{m}|B|$ intersecting pairs in $(A \setminus A_1) \times B$. There are still at least $15k\sqrt{m}|B|$ intersecting pairs in $A_1 \times B$. Let $B_1 \subseteq B$ be the set of curves in B that intersect at least $8k\sqrt{m}$ curves in A_1 . There are at least $7k\sqrt{m}|B|/|A_1| \geq 7k\sqrt{m}|B|/|A|$ elements in B_1 . Applying Lemma 3.1 to A_1 and B_1 , there are subsets $A' \subset A_1$ and $B' \subset B_1$ of size $|A'| \geq d/3 \geq \sqrt{m}/7$ and $|B'| \geq d|B_1|/(3|A_1|) > km|B|/|A|^2$ such that every curve in A' intersects every curve in B' . \square

4. Proof of Theorem 1.1 for $[0, 1]$ -Curves

Any collection of $[0, 1]$ -curves between the lines $L_1 : x = 0$ and $L_2 : x = 1$ can also be regarded as a collection of *double-grounded curves*, where L_1 and L_2 act as the grounds. In this section, we establish Theorem 1.1 for $[0, 1]$ -curves. This special case is the heart of the proof.

THEOREM 4.1. *Given a k -intersecting collection C of n $[0, 1]$ -curves with at least εn^2 intersecting pairs, its intersection graph contains a bi-clique of size at least $c_k \varepsilon^2 n$, where $c_k > 0$ depends only on k .*

As discussed at the end of the introduction, by slightly moving curves if necessary we may assume without loss of generality that the collection of $[0, 1]$ -curves are in general position.

Roughly speaking, our strategy is to try to prove Theorem 4.1 by induction on k . The base case $k = 1$ is resolved in Lemma 4.2 below. Ideally, for $k \geq 2$, we would like to find a curve $\Gamma \in C$ such that the portions of the $[0, 1]$ -curves that intersect Γ and stretch between L_1 and Γ form a $(k - 1)$ -intersecting family C' of at least $\Omega_k(\varepsilon^2 n)$ curves with $\Omega_k(|C'|^2)$ intersecting pairs. If this were the case, we could use induction on k , because the Jordan-Schönflies theorem (see, e.g., [19]) implies C' is homeomorphic to a collection of $[0, 1]$ -curves where Γ acts as the second line L_2 . Unfortunately, we will not be able to exclude that a small number of pairs of curves in C' intersect in more than $k - 1$ points. This motivates the following definition.

For $\varepsilon, \delta \in (0, 1]$ and $k, n \in \mathbb{N}$ with $k \geq 1$, $n \geq 2$, let $f(\varepsilon, \delta, k, n)$ denote the largest integer $t \in \mathbb{N}$ with the following property: For any collection C of n $[0, 1]$ -curves in the plane in general position with at least εn^2 intersecting pairs, at most δn^2 pairs of which intersect in more than k points, the intersection graph of C contains $K_{t,t}$.

LEMMA 4.2. *For every $\varepsilon \in (0, 1]$ and for any integer $n > 1$, we have*

$$f\left(\varepsilon, \frac{\varepsilon}{2}, 1, n\right) \geq \frac{\varepsilon}{4}n.$$

Proof. Let C be a collection of n $[0, 1]$ -curves in the plane in general position with at least εn^2 intersecting pairs, at most $\frac{\varepsilon}{2}n^2$ of which intersect in more than 1 point. So there are at least $\frac{\varepsilon}{2}n^2$ pairs of curves in C that intersect in exactly one point.

Define a two-dimensional partial order \prec on C so that $\alpha \prec \beta$ if the endpoint of α lies below the endpoint of β on L_i , $i = 1, 2$. Since there are no touching intersection points, then each pair of curves in C that intersect in exactly one point are incomparable by \prec . Therefore, the incomparability graph of (C, \prec) , which is a subgraph of the intersection graph of C , has at least $\frac{\varepsilon}{2}n^2$ edges. By Lemma 2.2, the incomparability graph of (C, \prec) and hence also the intersection graph of C contains $K_{t,t}$ with $t \geq \frac{\varepsilon}{4}n$. \square

Our next theorem is the main tool in the proof of Theorem 4.1.

THEOREM 4.3. *For all $\varepsilon > 0$, $\delta > 0$ with $4000\delta \leq \varepsilon^4$, $k \geq 2$, $n > 1$, and $t \leq \frac{\varepsilon^2 n}{10^7 k}$, we have*

$$f(\varepsilon, \delta, k, n) \geq \min(t, f(\varepsilon', \delta', k - 1, n')),$$

where $\varepsilon' = \frac{1}{10^6 k^2}$, $\delta' = 10^6 \left(\frac{\delta}{\varepsilon^4} + \frac{t}{\varepsilon^2 n}\right)$, and $n' \geq \frac{\varepsilon^2 n}{250}$.

Before proving Theorem 4.3, we first use it to establish Theorem 4.1. Applying Theorem 4.3 successively $k - 1$ times with $t = (10k)^{-30k} \varepsilon^2 n$, and then Lemma 4.2, we obtain that

$$f(\varepsilon, 0, k, n) \geq (10k)^{-30k} \varepsilon^2 n,$$

which proves Theorem 4.1. Indeed, we obtain

$$\begin{aligned} f(\varepsilon, 0, k, n) &\geq \min(t, f(\varepsilon', \delta_1, k-1, n_1)) \geq \dots \geq \min(t, f(\varepsilon', \delta_i, k-i, n_i)) \geq \dots \\ &\geq \min(t, f(\varepsilon', \delta_{k-1}, 1, n_{k-1})) \geq \min(t, f(\varepsilon', \varepsilon'/2, 1, n_{k-1})) \geq \min(t, \varepsilon' n_{k-1}/4) = t, \end{aligned}$$

where $\varepsilon' = \frac{1}{10^6 k^2}$, $n_1 \geq \frac{\varepsilon^2 n}{250}$,

$$n_i \geq \frac{\varepsilon'^2}{250} n_{i-1} \geq \frac{\varepsilon^2}{250} \left(\frac{\varepsilon'^2}{250} \right)^{i-1} n \geq 10^{-3} (10k)^{-15(i-1)} \varepsilon^2 n \quad \text{for } 2 \leq i \leq k-1 \text{ by induction on } i,$$

$$\delta_1 = \frac{10^6 t}{\varepsilon^2 n} = 10^6 (10k)^{-30k}, \text{ and}$$

$$\delta_i = 10^6 \left(\frac{\delta_{i-1}}{\varepsilon'^4} + \frac{t}{\varepsilon'^2 n_{i-1}} \right) = 10^{30} k^8 \delta_{i-1} + 10^{18} k^4 (10k)^{-30k} \frac{\varepsilon^2 n}{n_{i-1}} \leq 10^6 (10k)^{30(i-1-k)}$$

for $2 \leq i \leq k-1$ by induction on i .




FIGURE 2. Three $[0, 1]$ -curves, α , β , and γ , with the property that the endpoint of γ along L_i lies above the corresponding endpoints of α and β ($i = 1, 2$); and γ intersects both α and β . On the left: If the intersection points $\alpha \cap \gamma$ and $\beta \cap \gamma$ span disjoint subcurves of γ , then α and β must intersect. In the middle: If the intersection points $\alpha \cap \gamma$ and $\beta \cap \gamma$ span overlapping subcurves of γ , then there is a point $p \in \beta \cap \gamma$ contained in both subcurves of γ . On the right: If α and β intersect, then the portions of $\beta_\alpha \subset \beta$ and $\gamma_\alpha \subset \gamma$ between the line L_1 and their first intersections with α do not intersect at p .

The following simple observation is an important tool for the proof of Theorem 4.3.

OBSERVATION 1. (See Fig. 2. left side.) Suppose that α , β , and γ are $[0, 1]$ -curves such that

- (i) for $i = 1, 2$, the endpoints of α and β on line L_i lie below the endpoint of γ ;
- (ii) γ intersects both α and β ; and
- (iii) the intersection points of $\alpha \cap \gamma$ and $\beta \cap \gamma$ lie in disjoint subcurves of γ .

Then α and β must intersect. \square

Proof of Theorem 4.3: Let C be a collection of n $[0, 1]$ -curves with $m \geq \varepsilon n^2$ intersecting pairs, at most δn^2 of which intersect in more than k points. Suppose further that G , the intersection of graph of C , is $K_{t,t}$ -free. Since G is $K_{t,t}$ -free and has at least one edge, then $1 < t \leq \frac{\varepsilon^2 n}{10^7 k}$, so $n > 10^7 k \varepsilon^{-2}$. By Lemma 2.4, G has an induced subgraph G_1 with $n_1 \geq \frac{m^2}{5n^3} \geq \frac{\varepsilon^2 n}{5} > 10^6 k$ vertices and edge density at least $1 - 10t \frac{n^3}{m^2} \geq 1 - \frac{10t}{\varepsilon^2 n} > .99$. So the number m_1 of edges of G_1 is at least $.99 \binom{n_1}{2} \geq .49 n_1^2$. Let $C_1 \subseteq C$ denote the collection of curves that correspond to the vertices of G_1 .

Define a linear order \prec_1 on C_1 such that $\alpha \prec_1 \beta$ if the endpoint of α on the line $L_1 : x = 0$ lies below the corresponding endpoint of β . Also define the 2-dimensional partial order \prec_2 on C_1 where $\alpha \prec_2 \beta$ if and only if $\alpha \prec_1 \beta$ and the endpoint of α on the line $L_2 : x = 1$ lies below the corresponding endpoint of β . Note that if $\alpha, \beta \in C_1$ are incomparable by \prec_2 , then they must intersect, so the incomparability graph of the poset (C_1, \prec_2) is a subgraph of the intersection graph G_1 of C_1 .

For every curve $\gamma \in C_1$, let $C_1(\gamma)$ be the set of all curves $\alpha \in C_1$ such that $\alpha \prec_1 \gamma$ and $\alpha \cap \gamma \neq \emptyset$; and let $d(\gamma) = |C_1(\gamma)|$. Note that $\sum_{\gamma \in C_1} d(\gamma)$ is the number $e(G_1)$ of edges of G_1 . (Refer to Fig. 2.) For two intersecting curves $\alpha, \beta \in C_1$, let α_β denote the portion of α between its endpoint on L_1 and its first intersection point with β . Let T be the set of triples

$$T = \{(\alpha, \beta, \gamma) \in C_1 \times C_1 \times C_1 : \alpha, \beta \in C_1(\gamma), \alpha \neq \beta, \text{ and } \gamma_\alpha \cap \beta \neq \emptyset\}.$$

We now give a lower bound on the cardinality of T . For any two curves $\alpha, \beta \in C_1(\gamma)$, if we follow γ from its endpoint at L_1 , we either meet α before β or vice versa: We have either $\gamma_\alpha \cap \beta \neq \emptyset$ or $\gamma_\beta \cap \alpha \neq \emptyset$. Therefore, for each γ , there are at least $\binom{d(\gamma)}{2}$ triples in T with last coordinate γ . So

$$\begin{aligned} |T| &\geq \sum_{\gamma \in C_1} \binom{d(\gamma)}{2} = \frac{1}{2} \sum_{\gamma \in C_1} (d(\gamma)^2 - d(\gamma)) = -\frac{m_1}{2} + \frac{1}{2} \sum_{\gamma \in C_1} d(\gamma)^2 \\ &\geq -\frac{m_1}{2} + \frac{1}{2} n_1 \left(\frac{\sum_{\gamma \in C_1} d(\gamma)}{n_1} \right)^2 = -\frac{m_1}{2} + \frac{m_1^2}{2n_1} \geq \frac{n_1^3}{10}, \end{aligned}$$

where we use $\sum_{\gamma \in C_1} d(\gamma) = m_1$, Jensen's inequality for the convex function $f(x) = x^2$, $m_1 \geq .49n_1^2$, and $n_1 > 10^6$.

A triple $(\alpha, \beta, \gamma) \in T$ is called *bad* if $\alpha \cap \beta \neq \emptyset$ and $\beta \cap \gamma = \beta_\alpha \cap \gamma_\alpha$, that is, if all the intersection points of β and γ are between their portions stretching between L_1 and α (Fig. 2, left). All other triples of T are *good* (Fig. 2, middle). We next give an upper bound on the number of bad triples. Distinguish two kinds of bad triples: A bad triple of the *first* kind contains at least one pair of curves incomparable under \prec_2 ; in every remaining bad triple (α, β, γ) , of the *second* kind, all pairs are pairwise comparable under \prec_2 but the intersection points $\alpha \cap \gamma$ and $\beta \cap \gamma$ span disjoint subcurves along γ .

Since the incomparability graph of the 2-dimensional poset (C_1, \prec_2) is a subgraph of the intersection graph G_1 , applying Lemma 2.2, it follows that (C_1, \prec_2) contains fewer than $2tn_1$ incomparable pairs. Every incomparable pair can participate in up to $2n_1$ ordered triples $(\alpha, \beta, \gamma) \in T$. So there are at most $4tn_1^2$ bad triples of the first kind.

Next, we consider bad triples of the second kind. For every curve $\gamma \in C_1$, let $C_2(\gamma)$ be the set of all curves $\alpha \in C_1$ with $\alpha \prec_2 \gamma$ and $\alpha \cap \gamma \neq \emptyset$. Every curve $\alpha \in C_2(\gamma)$ determines a subcurve $\gamma[\alpha]$ of γ spanned by the intersection points $\alpha \cap \gamma$. By Observation 1, if two curves $\alpha, \beta \in C_2(\gamma)$ determine disjoint subcurves $\gamma[\alpha], \gamma[\beta] \subset \gamma$, then $\alpha \cap \beta \neq \emptyset$. Therefore, the intersection graph G_1 contains the comparability graph of the interval order on $C_2(\gamma)$, where α precedes β in the interval order if $\gamma[\alpha]$ and $\gamma[\beta]$ are disjoint and $\gamma[\alpha]$ lies closer to L_1 than $\gamma[\beta]$ along γ . Since G_1 is $K_{t,t}$ -free, for each $\gamma \in C_1$, the number of pairs $\alpha, \beta \in C_2(\gamma)$ where $\gamma[\alpha]$ is disjoint from $\gamma[\beta]$ is at most $2(t-1)|C_2(\gamma)| < 2tn_1$ by Lemma 2.3. Therefore, there are at most $2tn_1^2$ triples of the second kind. The total number of bad triples is at most $6tn_1^2$.

There are at least $n_1^3/10$ triples in T , but at most $6tn_1^2$ bad triples. Let $D \subset C_1$ be the set where $\alpha \in D$ if there are at least $60tn_1$ bad triples with first coordinate α . The size of D is at most $n_1/10$, so there are at most $\frac{n_1}{10} \binom{n_1}{2} < n_1^3/20$ triples in T with first coordinate in D . Therefore, there are at least $n_1^3/20$ triples in T with first coordinate in $C_1 \setminus D$. Furthermore, there are at most $4\delta n^2 n_1 \leq (100\delta/\varepsilon^4)n_1^3 \leq n_1^3/40$ triples in T where the first coordinate intersects the second or third coordinate in more than k points. Hence, there is a curve $\hat{\alpha} \in C_1 \setminus D$ such that at least $n_1^2/40$ triples in T have first coordinate $\hat{\alpha}$, and $\hat{\alpha}$ intersects the second and third

coordinates in at most k points; but at most $60tn_1$ bad triples have first coordinate $\hat{\alpha}$. Fix such an $\hat{\alpha}$, and let

$$A = \{\gamma_{\hat{\alpha}} : \hat{\alpha} \in C_1(\gamma) \text{ and } |\hat{\alpha} \cap \gamma| \leq k\} \quad \text{and}$$

$$B = \{\gamma : \gamma \in C_1 \setminus \{\hat{\alpha}\}, \hat{\alpha} \notin C_1(\gamma), \text{ and } |\hat{\alpha} \cap \gamma| \leq k\} \cup \{\gamma \setminus \gamma_{\hat{\alpha}} : \hat{\alpha} \in C_1(\gamma) \text{ and } |\hat{\alpha} \cap \gamma| \leq k\}.$$

That is, A is a collection of curves that contains, for each curve in C_1 that intersects $\hat{\alpha}$ in at most k points and has a left endpoint above that of $\hat{\alpha}$, the portion between L_1 and its first intersection with $\hat{\alpha}$. The collection B contains all other curves of C_1 that intersects $\hat{\alpha}$ in at most k points, and it also contains the remaining portions of the curves covered by A (Fig. 2, right). Notice that $|A| \leq |B| < n_1$, and the collection of curves in A is double grounded with the grounds being the line L_1 and the curve $\hat{\alpha}$. Also notice that the number of intersecting pairs $(\gamma_{\hat{\alpha}}, \beta) \in A \times (A \cup B)$ is at least $n_1^2/40$; and so the cardinality of A is at least $n_1/50$.

Let $\mu = n_1^2/(10^6k^2)$. First assume that the number of intersecting pairs in A is at most μ . In this case, the number of intersecting pairs in $A \times B$ must be at least $n_1^2/40 - 2\mu$, which is greater than $20k\sqrt{\mu}n_1$. Applying Lemma 3.2, we find subsets $A' \subset A$ and $B' \subset B$ such that

$$|A'| \geq \frac{\sqrt{\mu}}{7} = \frac{1}{7} \sqrt{\frac{n_1^2}{10^6k^2}} > \frac{\varepsilon^2 n}{10^5 k} \geq 2t, \quad \text{and}$$

$$|B'| \geq \frac{k\mu|B|}{|A|^2} = \left(\frac{n_1^2}{10^6k^2}\right) \frac{k|B|}{|A|^2} \geq \left(\frac{n_1^2}{10^6k^2}\right) \frac{k}{n_1} = \frac{n_1}{10^6k} \geq \frac{\varepsilon^2 n}{5 \cdot 10^6 k} \geq 2t,$$

and every curve in A' intersects every curve in B' . Even though a curve of C_1 may have a portion in A and a disjoint portion in B , nevertheless, the intersection graph of C still contains $K_{t,t}$.

Next assume that the number of intersecting pairs in A is larger than $\mu = n_1^2/(10^6k^2) \geq |A|^2/(10^6k^2)$. Let us estimate the number of pairs $\beta_{\hat{\alpha}}, \gamma_{\hat{\alpha}} \in A$ that intersect in at least k points. Clearly, $|\beta_{\hat{\alpha}} \cap \gamma_{\hat{\alpha}}| \leq |\beta \cap \gamma|$, and there are at most δn^2 pairs with $|\beta \cap \gamma| > k$. If $|\beta_{\hat{\alpha}} \cap \gamma_{\hat{\alpha}}| = k$ and $|\beta \cap \gamma| = k$, then $(\hat{\alpha}, \beta, \gamma)$ is a bad triple of T . Hence, the number of pairs in A that intersect in at least k points cannot exceed

$$\delta n^2 + 60tn_1 \leq \left(\frac{25\delta}{\varepsilon^4} + \frac{300t}{\varepsilon^2 n}\right) n_1^2 \leq \left(\frac{25\delta}{\varepsilon^4} + \frac{300t}{\varepsilon^2 n}\right) (50|A|)^2 \leq 10^6 \left(\frac{\delta}{\varepsilon^4} + \frac{t}{\varepsilon^2 n}\right) |A|^2.$$

In the first case we showed that G contains the bi-clique $K_{t,t}$, and in the second case the intersection graph of A contains a bi-clique with parts of size $f(\varepsilon', \delta', k-1, n')$. We conclude that $f(\varepsilon, \delta, k, n) \geq \min(t, f(\varepsilon', \delta', k-1, n'))$. \square

5. Proof of Theorem 1.1

The aim of this section is to prove Theorem 1.1. In order to use our results on $[0, 1]$ -curves and on double-grounded curves, we choose one or two curves to act as grounds. By Observation 2, the intersection graph of the portions of the curves between the chosen ground curves can be realized by $[0, 1]$ -curves.

OBSERVATION 2. The following two statements are equivalent:

- (i) G is the intersection graph of a k -intersecting collection of $[0, 1]$ -curves.
- (ii) G is the intersection graph of a k -intersecting collection C of grounded curves with the following property: We are given a ground g and a curve α with one endpoint incident to g and the rest of α lying in the exterior of g . Every curve of C has one endpoint incident to g , the other endpoint incident to α , and the rest of the curve is disjoint from α and g . (See Fig. 3.) \square




FIGURE 3. (a) A collection of $[0, 1]$ -curves. (b) We may assume that all curves lie in a rectangle R between vertical lines L_1 and L_2 . (c) The boundary of R can be continuously deformed to a circular arc and two segments along radii. (d) The segments along the radii can be identified by a further deformation, and an inversion transforms the curves into a collection described in Observation 2(ii). The above mapping of a collection of curves in (a) to the collection in (d) is invertible.

THEOREM 5.1. *Let C be a k -intersecting collection of n curves. Suppose that $C_1 \subseteq C$ is a collection of grounded curves with ground g , $C_2 = C \setminus C_1$ is a collection of curves disjoint from g , and there are at least εn^2 intersecting (nonidentical) pairs in $C_1 \times C$.*

- (i) *If $C = C_1$ (hence $C_2 = \emptyset$), then the intersection graph of C contains a bi-clique of size at least $c_k \varepsilon^8 n$, where $c_k > 0$ is a constant depending on k only.*
- (ii) *Otherwise, the intersection graph of C contains a bi-clique of size at least $c'_k \varepsilon^{32} n$, where $c'_k > 0$ is a constant depending on k only.*

Proof. For any curve $\gamma \in C_1$, let $N(\gamma)$ denote the set of curves in C that intersect γ , and let $d(\gamma) = |N(\gamma)|$. Label the curves $\alpha \in N(\gamma)$ with integers from 1 to $d(\gamma)$ according to their closest intersection points to the ground along γ , that is, the label $\ell_\gamma(\alpha)$ of $\alpha \in N(\gamma)$ is the number of curves that intersect the portion of γ between g and its first intersection with α . The sum of the labels is $\binom{d(\gamma)+1}{2}$ for a given $\gamma \in C_1$. Since $\sum_{\gamma \in C_1} d(\gamma) \geq \varepsilon n^2$, the sum of all the labels is

$$\sum_{\gamma \in C_1} \sum_{\alpha \in N(\gamma)} \ell_\gamma(\alpha) = \sum_{\gamma \in C_1} \binom{d(\gamma)+1}{2} > \sum_{\gamma \in C_1} d(\gamma)^2/2 \geq \frac{1}{2n} \left(\sum_{\gamma \in C_1} d(\gamma) \right)^2 \geq \frac{\varepsilon^2}{2} n^3,$$

by Jensen's inequality. Let the *weight* $w(\alpha)$ of a curve $\alpha \in C$ be the sum of its labels, so $w(\alpha) = \sum_{\gamma: \alpha \in N(\gamma)} \ell_\gamma(\alpha)$. Pick a curve $\hat{\alpha}$ of at least average weight, which is at least $\frac{\varepsilon^2}{2} n^2$.

Let A be the set containing the portion of every curve $\gamma \in N(\hat{\alpha})$ between g and its first intersection with $\hat{\alpha}$. Let B be the set containing all curves $\gamma \in C \setminus N(\hat{\alpha})$, $\gamma \neq \hat{\alpha}$; and the portion of every curve $\gamma \in N(\hat{\alpha})$ beyond its first intersection with $\hat{\alpha}$. The number of intersecting pairs in $A \times (A \cup B)$ is at least the weight of $\hat{\alpha}$, which is at least $\frac{\varepsilon^2}{2} n^2$. Since $|B| = |C \setminus \{\hat{\alpha}\}| = n - 1$, we have $|A| \geq \varepsilon^2 n/4$.

Let $m = \varepsilon^4 n^2 / (10^4 k^2)$. Assume first that the number of intersecting pairs in A is at most m . This means that the number of intersecting pairs in $A \times B$ must be at least $\frac{\varepsilon^2}{4} n^2 - 2m$, which is greater than $20k\sqrt{m}|B|$. By Lemma 3.2, there are subsets $A' \subset A$ and $B' \subset B$ of size

$$|A'| \geq \frac{\sqrt{m}}{7} = \frac{1}{7} \sqrt{\frac{\varepsilon^4 n^2}{10^4 k^2}} > \frac{\varepsilon^2 n}{10^3 k}, \quad \text{and}$$

$$|B'| \geq km|B|/|A|^2 = k \cdot \frac{\varepsilon^4 n^2}{10^4 k^2} \cdot \frac{|B|}{|A|^2} \geq \frac{\varepsilon^4 n}{10^4 k}$$

such that every curve in A' intersects every curve in B' . Even though a single curve of C may have a portion in A and a portion in B , the intersection graph of C still contains a bi-clique of size at least $c_k \varepsilon^4 n^2$ for some constant $c_k > 0$ depending only on k .

Assume now that the number of intersecting pairs in A is more than m . To prove (i), we assume that $C = C_1$ and so $\hat{\alpha} \in C_1$. By Observation 2, the intersection graph of A is the intersection graph of $[0, 1]$ -curves. By Theorem 4.1, there is a constant b_k depending on k only such that the intersection graph of A (and hence, the intersection graph of C) contains a bi-clique of size at least $b_k m^2 / |A|^3 \geq b_k m^2 / n^3 \geq c_k \varepsilon^8 n$ for some constant $c_k > 0$ depending only on k . This completes the proof of part (i).

Now we prove (ii). As before, if the number of intersecting pairs in A is at most m , then we are done. So we may assume that the number of intersecting pairs in A is larger than m . By applying (i) to the collection A , it follows that the intersection graph of A (and hence, the intersection graph of C) contains a bi-clique of size at least

$$c_k \left(\frac{m}{|A|^2} \right)^8 |A| \geq c_k m^8 |A|^{-15} \geq c_k \left(\frac{\varepsilon^4 n^2}{10^4 k^2} \right)^8 n^{-15} \geq c'_k \varepsilon^{32} n$$

for some constant $c'_k > 0$ depending on k . \square

We are almost ready to prove Theorem 1.1. In the proof, we choose a curve of a k -intersecting family as a *ground* and apply Theorem 5.1(ii). Since each curve intersects any other curve in at most k points, but a ground may intersect each curve in only one point, we keep only a piece of each curve, using the following proposition.




FIGURE 4. If a curve α has k intersection points with γ , we can split α into $k + 1$ disjoint subcurves such that each subcurve intersect γ in exactly one point (namely, at an endpoint), and any intersection point with other curves lies in one of the subcurves.

PROPOSITION 5.2. *Let C be a k -intersecting collection of n curves, and let M be a set of m intersecting pairs $(\alpha, \beta) \in C \times C$. For every curve $g \in C$, we can choose a subcurve α' of each curve $\alpha \in C \setminus \{g\}$ such that if α intersects g , then $\alpha' \cap g$ is an endpoint of α' , and there are at least $m / (k + 1)^2$ intersecting pairs (α', β') with $(\alpha, \beta) \in M$.*

Proof. Every curve $\alpha \in C$ that intersects g can be split into at most $k + 1$ disjoint subcurves, each intersecting g in exactly one point (namely, at an endpoint), see Fig. 4. Let C' contain one subcurve of each curve of C chosen uniformly at random. Each intersection point of two curves of C is the intersection point of two subcurves in C' with probability at least $1 / (k + 1)^2$. The expected number of intersecting pairs $(\alpha', \beta') \in C' \times C'$ with $(\alpha, \beta) \in M$ is at least $m / (k + 1)^2$, hence there is a choice of subcurves C' that gives at least $m / (k + 1)^2$ such intersecting pairs. \square

We now have all the ingredients to prove Theorem 1.1.

Proof of Theorem 1.1: Let C be a k -intersecting collection of n curves in the plane with at least εn^2 intersecting pairs. Successively delete every curve of C that intersect fewer than $\frac{\varepsilon}{2}n$ other curves. We are left with a collection $C_0 \subseteq C$ of curves, each of which intersects at least $\varepsilon n/2$ other curves of C_0 . Since the deletion of these curves removes fewer than $\varepsilon n^2/2$ intersecting pairs, there are still at least $\varepsilon n^2/2$ intersecting pairs in C_0 .

Choose an arbitrary curve $g \in C_0$ and let $C_1 \subset C_0$ be the set of curves that intersect g . Since each curve in C_0 intersects at least $\varepsilon n/2$ others, there are at least $\varepsilon^2 n^2/4$ intersecting pairs $(\alpha, \beta) \in C_1 \times C_0$.

By Proposition 5.2, there is a collection C'_0 of subcurves of curves in $C_0 \setminus \{g\}$ such that the subcurves $C'_1 \subseteq C'_0$ of the curves of C_1 are grounded curves with ground g and there are at least $\frac{\varepsilon^2}{4}n^2/(k+1)^2$ intersecting pairs in $C'_1 \times C'_0$. Applying Theorem 5.1(ii), we conclude that the intersection graph of C'_0 , hence that of C , contains a bi-clique of size at least $c_k \varepsilon^{64} n^2$, for some constant $c_k > 0$ depending only on k . \square

6. Proof of Theorem 1.3

We need the following separator theorem from [14], mentioned in the introduction. Recall that a crossing between a pair of Jordan regions is defined as an intersection point of their boundaries or a containment between them.

LEMMA 6.1. [14] *The intersection graph of any finite collection of Jordan regions in the plane with a total of m crossings has a separator of size $O(\sqrt{m})$.*

Applying this result successively to a collection C of Jordan regions in the plane, we obtain a number of small clusters of C such that any two regions belonging to distinct clusters are disjoint.

THEOREM 6.2. *Let C be a collection of Jordan regions with a total of m crossings. For any $d \geq 1$, there exists a partition $C = \bigcup_{i=0}^s V_i$, for some $s \leq 2d + 1$ such that $|V_0| = O((dm)^{1/2})$, $|V_i| \leq |C|/d$ for $i = 1, 2, \dots, s$, and no element in V_i intersects any element in V_j for $1 \leq i < j \leq s$.*

Proof. The proof is by a simple iteration of Lemma 6.1. At step i , we have a collection S_i of disjoint subsets of C . For a subcollection $W \subseteq C$, let $m(W)$ denote the total number of crossings between the Jordan regions in W . Let $q = \lceil \log_{3/2} d \rceil$.

- (i) Let $i = 0$ and $S_0 = \{C\}$.
- (ii) Repeat until $i = q$.
 - Set $i := i + 1$ and $S_i := \emptyset$. For every $W \in S_{i-1}$, do
 - * If $|W| \geq (\frac{2}{3})^i n$, then apply Theorem 6.1 to partition $W = W_0 \cup W_1 \cup W_2$ such that $|W_0| = O(\sqrt{m(W)})$, $|W_1|, |W_2| \leq \frac{2}{3}|W|$, and no set in W_1 intersects a set in W_2 . In this case, let $S_i := S_i \cup \{W_1, W_2\}$.
 - * Otherwise, let $S_i := S_i \cup \{W\}$.
- (iii) Return S_q .

Every S_i is a collection of disjoint subsets of C . Therefore, we have $\sum_{W \in S_i} m(W) \leq m$. Since each set partitioned at step i has at least $(\frac{2}{3})^i n$ elements, the number of subsets partitioned at step i is at most $(\frac{3}{2})^i$. Thus, for $i \in [q]$, the number of sets in C that belong to one of the

subsets in S_{i-1} , but not to any subset in S_i , is

$$\sum_{W \in S_i, |W| \geq \left(\frac{2}{3}\right)^i n} O(\sqrt{m(W)}) = O\left(\left(\frac{3}{2}\right)^i \sqrt{\left(\frac{2}{3}\right)^i m}\right) = O\left(\left(\frac{3}{2}\right)^{i/2} \sqrt{m}\right),$$

by Jensen's inequality, using the concavity of the function $g(x) = \sqrt{x}$. Let V_0 be the set of elements of C that do not belong to any subsets in S_q . Summing the number of deleted sets over all steps, we obtain that $|V_0| \leq O(\sum_{i=1}^q \left(\frac{3}{2}\right)^{i/2} m^{1/2}) = O((dm)^{1/2})$. Denote the elements of S_q be V_1, \dots, V_s , where $s = |S_q|$. In the partition $C = \bigcup_{i=0}^s V_i$, each set $V_i \in S_q$ has at most $|C|/d$ elements, and no Jordan region in V_i intersects any Jordan region in V_j , for $1 \leq i < j \leq s$. By grouping together some small V_i 's, we may assume that $s \leq 2d + 1$ holds. This completes the proof. \square

Proof of Theorem 1.3: Let G be the intersection graph of a k -intersecting collection R of n r -regions. Let C denote the family of all components of elements R , so C is a k -intersecting family of Jordan regions with $n \leq |C| \leq rn$.

Distinguish two cases.

Case 1: *The intersection graph of C has at most $\frac{cn^2}{kr^2}$ edges*, where c is a sufficiently small absolute positive constant determined by the constant implied in Theorem 6.2. In this case, applying Theorem 6.2 with $d = 3r^2$, there is a partition $C = \bigcup_{i=1}^s V_i$ with $s \leq 2d + 1$, $|V_0| \leq n/3$, and $|V_i| \leq \frac{|C|}{d} \leq \frac{n}{3r}$ for $i \in [s]$, such that every set in V_i is disjoint from every set in V_j ($1 \leq i < j \leq s$). Let R_1 be the collection of r -regions in R none of whose components lie in V_0 , so that $|R_1| \geq 2n/3$. Each $\gamma \in R_1$ has at most r components, each of which lies in one of the s subsets V_1, \dots, V_s . Therefore, there are indices $i_1, \dots, i_r \in [s]$ and a subcollection $A \subseteq R_1$ such that $|A| \geq |R_1|/s^r = \Omega(n/(7r)^{2r})$ and, for each $\gamma \in A$, all of the components of γ belong to $V_{i_1} \cup \dots \cup V_{i_r}$. Notice that the number of sets in R_1 that have at least one component in $V_{i_1} \cup \dots \cup V_{i_r}$ is at most $|V_{i_1} \cup \dots \cup V_{i_r}| \leq r \frac{n}{3r} = n/3$. Therefore, letting $B \subset R_1$ be the set of all γ whose components all lie in $\bigcup_{i \in [s] \setminus \{i_1, \dots, i_r\}} V_i$, we have $|B| \geq n/3$ and every set in B is disjoint from every set in A .

Case 2: *The intersection graph of C has at least $\frac{cn^2}{kr^2}$ edges*. In this case, either half of the edges come from containments of components or at least half of them correspond to boundary intersections. If at least half the intersecting pairs correspond to pairs, for which one of the components of one set contains a component of the other, then one of the components is contained in at least $\frac{cn^2}{kr^2|C|} \geq \frac{cn}{kr^3}$ other components. In this case, the intersection graph of R contains a clique of size at least $\frac{cn}{2kr^4}$. If at least half of the intersecting pairs have components with intersecting boundaries, then we can apply Theorem 1.1 to the boundary curves of the components: The intersection graph of C contains a bi-clique of size $c_{k,r}n$, where $c_{k,r} > 0$ depends on k and r only. Since each set of R consists of at most r components, the intersection graph of R contains a bi-clique of size at least $\frac{c_{k,r}}{r}n$. \square

7. Applications and Open Problems

First, we mention some applications of Theorem 1.1 and Corollary 1.2. We conclude by listing a few open problems.

In [15], we use Theorem 1.1 to prove the following result. The chromatic number of every K_t -free intersection graph of $n \geq t$ curves in the plane, with no pair of curves intersecting in more than k points, is at most $(c_k \frac{\log n}{\log t})^{c \log t}$, where c is an absolute constant and $c_k > 0$ depends only on k . In particular, this implies that, for any $\varepsilon > 0$ and for any positive integer k , there is a $\delta = \delta(\varepsilon, k) > 0$ such that the intersection graph of every k -intersecting collection

of n curves in the plane contains a clique of size at least n^δ or an independent set of size at least $n^{1-\varepsilon}$. This result is a strengthening of the Erdős-Hajnal property for intersection graphs of k -intersecting families of curves. This stronger version was not known even for intersection graphs of straight line segments.

The well-known *crossing lemma*, proved independently by Ajtai *et al.* [1] and Leighton [17], states that every graph of n vertices and $m \geq 4n$ edges, drawn in the plane so that the vertices are represented by points and the edges by continuous arcs connecting the corresponding points, determines $\Omega(m^3/n^2)$ crossings. This is easily seen to be equivalent to saying that there is always an edge that participates in $\Omega(m^2/n^2)$ crossings. In [13], we use Theorem 1.1 to prove a strengthening of the crossing lemma under the assumption that the edges form a k -intersecting collection of curves, for some constant k . We prove that, for any drawing of a graph $G = (V, E)$ with n vertices and $m \geq 4n$ edges, with no pair of edges intersecting in more than k points, there are disjoint subsets $E_1, E_2 \subset E$, each of size $\Omega_k(m^2/n^2)$, such that every edge in E_1 crosses every edge in E_2 .

Our arguments made essential use of the separator theorem for intersection graphs of families of curves (or Jordan regions) such that the total number of intersections (crossings) between them is relatively small. In particular, we used the fact (Lemma 6.1) that, for any fixed k , the intersection graph G of any k -intersecting collection of n curves in the plane has a separator of size $O_k(\sqrt{m})$, where $m = |E(G)|$. We do not know whether this statement remains true if we drop the assumption that any pair of curves can intersect in at most a constant number of points. In the literature, intersection graphs of continuous curves in the plane are usually called *string graphs*.

QUESTION 1. Does every string graph with m edges have a separator of size $O(\sqrt{m})$?

It is shown in [15] that the answer is yes for “outerstring graphs,” that is, for intersection graphs of grounded curves.

Corollary 1.2 states that the family of string graphs representable as intersection graphs of curves in the plane, any two of which meet in at most a fixed number of points, has the strong Erdős-Hajnal property, and hence the Erdős-Hajnal property. As we have pointed out in the introduction, the family of string graphs does not have the strong Erdős-Hajnal property. We have been unable to answer the following

QUESTION 2. Does the family of string graphs have the Erdős-Hajnal property?

Acknowledgments

The authors would like to thank Peter Keevash for helpful discussion that led to the separator theorem for curves. We are also grateful to the anonymous referee for many helpful comments.

References

1. M. Ajtai, V. Chvátal, M. Newborn, and E. Szemerédi, Crossing-free subgraphs, in *Theory and Practice of Combinatorics*, vol. 60, Mathematical Studies, North-Holland, Amsterdam, 1982, pp. 9–12.
2. N. Alon, J. Pach, R. Pinchasi, R. Radoičić, and M. Sharir, Crossing patterns of semi-algebraic sets, *J. Combin. Theory Ser. A* **111** (2) (2005), 310–326.
3. N. Alon, J. Pach, and J. Solymosi, Ramsey-type theorems with forbidden subgraphs, *Combinatorica* **21** (2001), 155–170.
4. S. Basu, Combinatorial complexity in o-minimal geometry, in *Proc. 39th Sympos. Theory of Computing (San Diego, CA)*, ACM Press, 2007, pp. 47–56.
5. M. Chudnovsky and S. Safra, The Erdős-Hajnal conjecture for bull-free graphs, *J. Combin. Theory, Ser. B* (2007), to appear.
6. G. Ehrlich, S. Even, and R. E. Tarjan, Intersection graphs of curves in the plane, *J. Combin. Theory Ser. B* **21** (1) (1976), 8–20.

7. P. Erdős, Some remarks on the theory of graphs, *Bulletin of the Amer. Math. Soc.* **53** (1947), 292–294.
8. P. Erdős and A. Hajnal, Ramsey-type theorems, *Discrete Appl. Math.* **25** (1989), 37–52.
9. P. Erdős, A. Hajnal, and J. Pach, Ramsey-type theorem for bipartite graphs, *Geombinatorics* **10** (2000), 64–68.
10. P. Erdős and G. Szekeres, A combinatorial problem in geometry, *Compositio Mathematica* **2** (1935), 463–470.
11. J. Fox, A bipartite analogue of Dilworth’s theorem, *Order* **23** (2-3) (2006), 197–209.
12. J. Fox, J. Pach, and Cs. D. Tóth, Turán-type results for partial orders and intersection graphs of convex sets, *Israel J. of Math.* (2007), to appear.
13. J. Fox, J. Pach, and Cs. D. Tóth, A bipartite strengthening of the Crossing Lemma, in *Proc. 15th Sympos. on Graph Drawing (Sydney, 2007)*, vol. 4875 of LNCS, Springer, 2008, pp. 13–24.
14. J. Fox and J. Pach, Separator theorems and Turán-type results for planar intersection graphs, submitted, 2007.
15. J. Fox and J. Pach, Coloring planar intersection graphs and the number of edges of k -quasi-planar topological graphs, manuscript, 2007.
16. M. C. Golumbic, D. Rotem, and J. Urrutia, Comparability graphs and intersection graphs. *Discrete Math.* **43** (1983), 37–46.
17. T. Leighton, New lower bound techniques for VLSI, *Math. Systems Theory* **17** (1984), 47–70.
18. R. J. Lipton and R. E. Tarjan, A separator theorem for planar graphs, *SIAM J. Appl. Math.* **36** (2) (1979), 177–189.
19. B. Mohar and C. Thomassen, *Graphs on Surfaces*, The John Hopkins University Press, Baltimore and London, 2001.
20. J. Pach and G. Tóth, Comment of Fox News, *Geombinatorics* **15** (2006), 150–154.
21. J. B. Sidney, S. J. Sidney, and J. Urrutia, Circle orders, n -gon orders and the crossing number, *Order* **5** (1) (1988), 1–10.

Jacob Fox
Department of Mathematics
Princeton University
Fine Hall, Washington Road
Princeton, NJ 08544-1000
USA

jacobfox@math.princeton.edu

János Pach
City College, CUNY and
Courant Institute, NYU
251 Mercer Street
New York, NY 10012
USA

pach@cims.nyu.edu

Csaba D. Tóth
Department of Mathematics
University of Calgary
2500 University Drive NW
Calgary, AB, T2P 1N4
Canada

cdtoth@ucalgary.ca