18.440 Final Exam: 100 points Carefully and clearly show your work on each problem (without writing anything that is technically not true) and put a box around each of your final computations. 1. (10 points) Let X be the number on a standard die roll (i.e., each of  $\{1, 2, 3, 4, 5, 6\}$  is equally likely) and Y the number on an independent standard die roll. Write Z = X + Y.

1. Compute the condition probability P[X = 4|Z = 6].

2. Compute the conditional expectation E[Z|Y] as a function of Y.

2. (10 points) Janet is standing outside at time zero when it starts to drizzle. The times at which raindrops hit her are a Poisson point process with parameter  $\lambda = 2$ . In expectation, she is hit by 2 raindrops in each given second.

(a) What is the expected amount of time until she is first hit by a raindrop?

(b) What is the probability that she is hit by exactly 4 raindrops during the first 2 seconds of time?

3. (10 points) Let X be a random variable with density function f, cumulative distribution function F, variance V and mean M.

(a) Compute the mean and variance of 3X + 3 in terms of V and M.

(b) If  $X_1, \ldots, X_n$  are independent copies of X. Compute (in terms of F) the cumulative distribution function for the largest of the  $X_i$ .

4. (10 points) Suppose that  $X_i$  are i.i.d. random variables, each uniform on [0, 1]. Compute the moment generating function for the sum  $\sum_{i=1}^{n} X_i$ .

5. (10 points) Suppose that X and Y are outcomes of independent standard die rolls (each equal to  $\{1, 2, 3, 4, 5, 6\}$  with equal probability). Write Z = X + Y.

(a) Compute the entropies H(X) and H(Y).

(b) Compute H(X, Z).

(c) Compute H(10X + Y).

(d) Compute  $H(Z) + H_Z(Y)$ . (Hint: you shouldn't need to do any more calculations.)

6. (10 points) Elaine's not-so-trusty old car has three states: broken (in Elaine's possession), working (in Elaine's possession), and in the shop. Denote these states B, W, and S.

- (i) Each morning the car starts out B, it has a .5 chance of staying B and a .5 chance of switching to S by the next morning.
- (ii) Each morning the car starts out W, it has .5 chance of staying W, and a .5 chance of switching to B by the next morning.
- (iii) Each morning the car starts out S, it has a .5 chance of staying S and a .5 chance of switching to W by the next morning.

Answer the following

(a) Write the three-by-three Markov transition matrix for this problem.

(b) If the car starts out B on one morning, what is the probability that it will start out B two days later?

(c) Over the long term, what fraction of mornings does the car start out in each of the three states, B, S, and W?

7. Suppose that  $X_1, X_2, X_3, \ldots$  is an infinite sequence of independent random variables which are each equal to 2 with probability 1/3 and .5 with probability 2/3. Let  $Y_0 = 1$  and  $Y_n = \prod_{i=1}^n X_i$  for  $n \ge 1$ .

(a) What is the probability that  $Y_n$  reaches 8 before the first time that it reaches  $\frac{1}{8}$ ?

(b) Find the mean and variance of  $\log Y_{10000}$ .

(c) Use the central limit theorem to approximate the probability that  $\log Y_{10000}$  (and hence  $Y_{10000}$ ) is greater than its median value.

8. (10 points) Eight people toss their hats into a bin and the hats are redistributed, with all of the 8! hat permutations being equally likely. Let N be the number of people who get their own hat. Compute the following:

(a)  $\mathbb{E}[N]$ 

(b)  $\operatorname{Var}[N]$ 

9. (10 points) Let X be a normal random variable with mean  $\mu$  and variance  $\sigma^2.$ 

(a)  $\mathbb{E}e^X$ .

(b) Find  $\mu$ , assuming that  $\sigma^2 = 3$  and  $E[e^X] = 1$ .

## 10. (10 points)

- 1. Let  $X_1, X_2, \ldots$  be independent random variables, each equal to 1 with probability 1/2 and -1 with probability 1/2. In which of the cases below is the sequence  $Y_n$  a martingale? (Just circle the corresponding letters.)
  - (a)  $Y_n = X_n$
  - (b)  $Y_n = 1 + X_n$
  - (c)  $Y_n = 7$
  - (d)  $Y_n = \sum_{i=1}^n iX_i$
  - (e)  $Y_n = \prod_{i=1}^n (1 + X_i)$
- 2. Let  $Y_n = \sum_{i=1}^n X_i$ . Which of the following is necessarily a stopping time for  $Y_n$ ?
  - (a) The smallest *n* for which  $|Y_n| = 5$ .
  - (b) The largest n for which  $Y_n = 12$  and n < 100.
  - (c) The smallest value n for which n > 100 and  $Y_n = 12$ .