

LECTURE 8: PUPPE SEQUENCES

Let $f : X \rightarrow Y$ be a map of pointed spaces. Taking iterated cofibers, we obtain an infinite sequence

$$X \xrightarrow{f} Y \xrightarrow{j} C(f) \xrightarrow{\pi} \Sigma X \xrightarrow{-\Sigma f} \Sigma Y \xrightarrow{-\Sigma j} \Sigma C(f) \xrightarrow{-\Sigma \pi} \Sigma^2 X \xrightarrow{\Sigma^2 f} \dots$$

The minus signs means take the degree -1 map on the suspension coordinate. This sequence $\{Z_i\}$ has the property that at each stage there is a homotopy commutative diagram

$$\begin{array}{ccccc} Z_i & \xrightarrow{g} & Z_{i+1} & \longrightarrow & C(g) \\ \parallel & & \parallel & & \simeq \downarrow \\ Z_i & \xrightarrow{g} & Z_{i+1} & \longrightarrow & Z_{i+2} \end{array}$$

whose vertical maps are homotopy equivalences.

We deduce that for a pointed space Z , there is a LES

$$[X, Z]_* \xleftarrow{f^*} [Y, Z]_* \leftarrow [C(f), Z]_* \leftarrow [\Sigma X, Z]_* \leftarrow \dots$$

Remark 0.1. The set $[\Sigma^k X, Z]_*$ is a group for $k \geq 1$, and is an abelian group for $k \geq 2$.

Remark 0.2. Taking $Z = K(\pi, n)$ for n large, we recover the LES for a pair

$$\tilde{H}^n(X, \pi) \xleftarrow{f^*} \tilde{H}^n(Y, \pi) \leftarrow H^n(Y, X, \pi) \leftarrow \tilde{H}^{n-1}(X, \pi) \leftarrow \dots$$

Dually, taking iterated fibers, we obtain an infinite sequence

$$\dots \xrightarrow{\Omega^2 f} \Omega^2 Y \xrightarrow{-\Omega \pi} \Omega F(f) \xrightarrow{-\Omega j} \Omega X \xrightarrow{-\Omega f} \Omega Y \xrightarrow{\pi} F(f) \xrightarrow{j} X \xrightarrow{f} Y$$

For a pointed space Z this induces a long exact sequence

$$\dots \rightarrow [Z, \Omega Y]_* \rightarrow [Z, F(f)]_* \rightarrow [Z, X]_* \xrightarrow{f_*} [Z, Y]_*$$

Specializing to $Z = S^0$, we recover the long exact sequence for the homotopy fiber.