

**FINAL EXAM**  
**MATH 18.022, MIT, AUTUMN 10**

You have three hours. This test is closed book, closed notes, no calculators.

Name: MODEL ANSWERS

Signature: \_\_\_\_\_

Recitation Time: \_\_\_\_\_

There are 10 problems, and the total number of points is 200. Show all your work. *Please make your work as clear and easy to follow as possible.*

Problem	Points	Score
1	20	
2	20	
3	20	
4	20	
5	20	
6	20	
7	20	
8	20	
9	20	
10	20	
Total	200	

1. (20pts) Find the shortest distance between the plane  $\Pi$  given by the equation  $2x - y + 3z = 3$  and the point of intersection of the two lines  $l_1$  and  $l_2$  given parametrically by


$$(x, y, z) = (2t - 3, t, 1 - t) \quad \text{and} \quad (x, y, z) = (1, 1 - t, t).$$

- Point where the two lines intersect:

$$\begin{cases} 2t_1 - 3 = 1 \\ t_1 = 1 - t_2 \\ 1 - t_1 = t_2 \end{cases} \Leftrightarrow \begin{cases} t_1 = 2 \\ t_2 = -1 \end{cases}$$

$$\text{so } (x, y, z) = (1, 2, -1) = P$$

- Arbitrary point on the plane  $\Pi$ ; for example  $Q = (0, 0, 1)$
- Normal vector to the plane  $\Pi$ :  $\vec{n} = (2, -1, 3)$



$$\vec{QP} = (1, 2, -1) - (0, 0, 1) = (1, 2, -2)$$

$$\text{distance} = \left\| \text{proj}_{\vec{n}} \vec{QP} \right\| = \left\| \frac{\vec{QP} \cdot \vec{n}}{\vec{n} \cdot \vec{n}} \vec{n} \right\|$$


$$= \left\| \frac{2-2-6}{4+1+9} (2, -1, 3) \right\| = \frac{6}{\sqrt{14}} \sqrt{4+1+9} =$$

$$\boxed{\frac{6}{\sqrt{14}}}$$

2. (20pts) Let  $W$  be the solid bounded by the paraboloid  $z = 9 - x^2 - y^2$ , the  $xy$ -plane, and the cylinder  $x^2 + y^2 = 4$ .

(a) Set up an integral in cylindrical coordinates for evaluating the volume of  $W$ .

$$\text{vol } W = \int_0^{2\pi} \int_0^2 \int_0^{9-r^2} r \, dz \, dr \, d\theta$$



(b) Evaluate this integral.

$$\text{vol } W = 2\pi \int_0^2 r(9-r^2) \, dr$$

$$= 2\pi \int_0^2 (9r - r^3) \, dr$$

$$= 2\pi \left[ \frac{9r^2}{2} - \frac{r^4}{4} \right] \Big|_{r=0}^2$$

$$= 2\pi \left( 9 \frac{4}{2} - \frac{16}{4} \right)$$


$$= 2\pi (18 - 4)$$

$$= \boxed{28\pi}$$

3. (20pts) (a) Change the order of integration of the integral

$$\int_0^1 \int_{-x}^x y^2 \cos(xy) dy dx.$$

$$= \int_{-1}^0 \int_{-y}^1 y^2 \cos(xy) dx dy + \int_0^1 \int_y^1 y^2 \cos(xy) dx dy$$



(b) Evaluate this integral.

$$= \int_{-1}^0 (y \sin(xy)) \Big|_{x=-y}^1 dy + \int_0^1 (y \sin(xy)) \Big|_{x=y}^1 dy$$

$$= \int_{-1}^0 (y \sin y - y \sin(-y^2)) dy + \int_0^1 (y \sin y - y \sin(y^2)) dy$$

$$= \underbrace{\int_{-1}^0 y \sin y dy}_{\text{Integration by parts}} + \underbrace{\int_{-1}^0 -y \sin(-y^2) dy}_{\text{Integration by parts}} + \underbrace{\int_0^1 -y \sin(y^2) dy}_{\text{Integration by parts}} = \textcircled{*}$$

$$= 2 \int_0^1 y \sin y dy$$

$$= \left[ \frac{(\cos(-y))^2}{-2} \right] \Big|_{y=-1}^0$$

$$= \left( \frac{\cos(-y^2)}{2} \right) \Big|_{y=0}^1$$

$$= 2 \left[ (-y \cos y) \Big|_{y=0}^1 - \int_0^1 -\cos y dy \right]$$

$$= \frac{1 - \cos 1}{-2}$$

$$= \frac{\cos 1 - 1}{2}$$

$$= 2 \left[ -\cos 1 - (-\sin y) \Big|_{y=0}^1 \right]$$

$$= \frac{\cos 1 - 1}{2}$$

$$= 2 \left[ -\cos 1 + \sin 1 \right]$$

$$\textcircled{*} = -2 \cos 1 + 2 \sin 1 + \cos 1 - 1 = \boxed{2(\sin 1) - (\cos 1) - 1}$$

Integration  
by parts

$$u = y$$

$$v' = \sin y$$

$$(uv' - u'v)$$

4. (20pts) Let  $\vec{r}: I \rightarrow \mathbb{R}^3$  be a regular smooth curve parametrised by arclength. Let  $a \in I$  and suppose that

$$\vec{T}(a) = \frac{2}{7}\hat{i} - \frac{6}{7}\hat{j} - \frac{3}{7}\hat{k}, \quad \vec{B}(a) = \frac{3}{7}\hat{i} - \frac{2}{7}\hat{j} + \frac{6}{7}\hat{k}, \quad \frac{d\vec{N}}{ds}(a) = -\frac{13}{7}\hat{i} + \frac{18}{7}\hat{j} - \frac{12}{7}\hat{k}.$$

Find:

(i) the unit normal vector  $\vec{N}(a)$ .

$$B = T \times N, \text{ and } \begin{array}{c} \uparrow B \\ \nearrow T \\ \downarrow N \end{array} \quad N = B \times T$$

$$\vec{N}(a) = B(a) \times T(a) = \begin{vmatrix} i & j & k \\ \frac{3}{7} & -\frac{2}{7} & \frac{6}{7} \\ \frac{2}{7} & -\frac{4}{7} & -\frac{3}{7} \end{vmatrix} = \frac{1}{49} (6+36, 12+9, -18+4) = \boxed{\left(\frac{6}{7}, \frac{3}{7}, -\frac{2}{7}\right)}$$

(ii) the curvature  $\kappa(a)$ .

$$N' = -\kappa T + \tau B \rightsquigarrow (\text{dot with } T \text{ on both sides}) \rightsquigarrow N \cdot T = -\kappa$$

$$\kappa(a) = -N'(a) \cdot T(a) = -\left(-\frac{13}{7}, \frac{18}{7}, -\frac{12}{7}\right) \cdot \left(\frac{2}{7}, -\frac{4}{7}, -\frac{3}{7}\right) = \frac{1}{49} (26+108-36) = \frac{98}{49} = \boxed{2}$$

(iii) the torsion  $\tau(a)$ .

$$N' = -\kappa T + \tau B \rightsquigarrow (\text{dot with } B \text{ on both sides}) \rightsquigarrow N \cdot B = \tau$$

$$\tau(a) = N'(a) \cdot B(a) = \frac{1}{49} (-13, 18, -12) \cdot (3, -2, 6)$$

$$= \frac{1}{49} (-39 - 36 - 72)$$

$$= -\frac{147}{49} = \boxed{-3}$$

5. (20pts) Let

$$C = \{(x, y, z) \in \mathbb{R}^3 \mid x^2z^3 - x^3z^2 = 0, x^2y + xy^3 = 2\}.$$

(a) Show that in a neighbourhood of the point  $P = (1, 1, 1)$ ,  $C$  is a smooth curve with a parametrisation of the form

$$\bar{g}(x) = (x, g_1(x), g_2(x)).$$

$$\text{Let } F(u, y, z) = (u^2z^3 - u^3z^2, u^2y + uy^3 - 2)$$

$$\left| \frac{\partial F}{\partial (y, z)}(1, 1, 1) \right| = \det \begin{bmatrix} 0 & 3u^2z^2 - 2u^3z \\ u^2 + 3uy^2 & 0 \end{bmatrix} \Big|_{(u, y, z)=(1, 1, 1)} = \det \begin{bmatrix} 0 & 1 \\ 4 & 0 \end{bmatrix} = -4 \neq 0$$

Then, by the implicit function theorem, in a neighborhood of  $P=(1, 1, 1)$ ,

(b) Find a parametrisation of the tangent line to  $C$  at  $P$ .  $\underline{[can be parametrized by x.]}$

Take the derivative with respect to  $x$  on both sides of the

equation  $F(u, g_1(u), g_2(u)) = (0, 0)$ , and evaluate at  $u=1$ :

$$\begin{aligned} \frac{\partial F}{\partial u} + \frac{\partial F}{\partial y} g'_1(u) + \frac{\partial F}{\partial z} g'_2(u) &= (0, 0) && \leftarrow \text{actually two equations!} \\ \Leftrightarrow \begin{cases} (2uz^3 - 3u^2z^2) + 0 + (3u^2z^2 - 2u^3z)g'_2(u) = 0 \\ (2uy + y^3) + (u^2 + 3uy^2)g'_1(u) = 0 \end{cases} & \Leftrightarrow \begin{cases} -1 + g'_2(1) = 0 \\ 3 + 4g'_1(1) = 0 \end{cases} & \Rightarrow \begin{cases} g'_2(1) = 1 \\ g'_1(1) = -\frac{3}{4} \end{cases} \end{aligned}$$

Vector tangent to curve  $C$  at  $P=(1, 1, 1)$ :  $(1, g'_1(1), g'_2(1)) = (1, -\frac{3}{4}, 1)$

Point on tangent line:  $P = (1, 1, 1)$

5

Equation of the line:  $(u, y, z) = (1, 1, 1) + t(1, -\frac{3}{4}, 1)$

$$\text{or } \boxed{(u, y, z) = (1+t, 1-\frac{3}{4}t, 1+t)}$$

6. (20pts) Let  $f: \mathbb{R}^2 \rightarrow \mathbb{R}$  be the function  $f(x, y) = xy$ .

(a) Show that  $f$  has a global maximum on the ellipse  $9x^2 + 4y^2 = 36$ .

$f$  is a continuous function  
the ellipse is a compact set  
(bounded + closed)

$\Rightarrow f$  has a global maximum  
on the ellipse

(b) Find this global maximum value of  $f$ .

$$\nabla f = \lambda \nabla g \Leftrightarrow (y, x) = \lambda (18x, 8y) \Leftrightarrow \begin{cases} y = 18\lambda x \\ x = 8\lambda y \end{cases}$$

$$y = 18 \cdot 8 \lambda^2 y \quad \begin{array}{l} y=0 \\ \lambda = \pm \frac{1}{12} \end{array} \quad \rightarrow x=y=0 \quad (\text{not on ellipse})$$

$$\lambda = \frac{1}{12} : \quad y = \frac{18}{12} x = \frac{3}{2} x, \quad \text{so} \quad 9x^2 + 4 \left(\frac{3}{2}x\right)^2 = 36 \Leftrightarrow 18x^2 = 36 \Leftrightarrow x = \pm\sqrt{2} \quad \Rightarrow y = \frac{3}{2}x = \pm\frac{3}{2}\sqrt{2}$$

$$f\left(\pm\sqrt{2}, \pm\frac{3}{2}\sqrt{2}\right) = 3$$

$$\lambda = -\frac{1}{12} : \quad y = -\frac{18}{12} x = -\frac{3}{2} x, \quad \text{so} \quad x = \pm\sqrt{2} \Rightarrow y = \mp\frac{3}{2}\sqrt{2}$$

$$f\left(\pm\sqrt{2}, \mp\frac{3}{2}\sqrt{2}\right) = -3$$

Maximum value of  $f$  is 3 (at points  $(\sqrt{2}, \frac{3}{2}\sqrt{2})$  and  $(-\sqrt{2}, -\frac{3}{2}\sqrt{2})$ )

7. (20pts) Let  $D$  be the region bounded by the four curves  $x^2 - y^2 = 1$ ,  $x^2 - y^2 = 4$ ,  $x^2/4 + y^2 = 1$  and  $x^2/16 + y^2/4 = 1$ .

(a) Compute  $dx dy$  in terms of  $du dv$ , where  $u = x^2 - y^2$  and  $v = x^2/4 + y^2$ .

$$\frac{\partial(u, v)}{\partial(x, y)} = \left| \det \begin{bmatrix} 2x & -2y \\ \frac{2x}{4} & 2y \end{bmatrix} \right| = |4xy + 2y| = 5|2y|$$

$$\text{So } \boxed{dx dy = \frac{1}{5|2y|} du dv}$$

(b) Evaluate the integral

$$\iint_D \frac{xy}{y^2 - x^2} dx dy.$$

The function  $\frac{xy}{y^2 - x^2}$  is odd


with respect to  $x$  (and  $y$  too),


so its integral over the first and fourth

quadrants will cancel out with the integral over the second and

third quadrants:

$$\iint_D \frac{xy}{y^2 - x^2} dx dy = 0$$





8. (20pts) (a) Find the area of the region that lies inside the closed curve defined by the equation  $r = 2a(1 + \sin 2\theta)$  in polar coordinates

$$\begin{aligned}
 \text{area} &= \int_0^{2\pi} \int_0^{2a(1+\sin 2\theta)} r dr d\theta = \int_0^{2\pi} \frac{4a^2(1+\sin 2\theta)^2}{2} d\theta \\
 &= 2a^2 \int_0^{2\pi} (1+\sin 2\theta)^2 d\theta = 2a^2 \int_0^{2\pi} (1 + (\sin 2\theta)^2 + 2\sin 2\theta) d\theta \\
 &= 2a^2 \left( 2\pi + \pi + (-\cos 2\theta) \Big|_{\theta=0}^{2\pi} \right) = \boxed{6\pi a^2}
 \end{aligned}$$

$\int_0^{2\pi} (\sin 2\theta)^2 d\theta = \frac{1}{2} \int_0^{2\pi} (\sin 2\theta)^2 + (\cos 2\theta)^2 d\theta = \frac{1}{2} 2\pi = \pi$

(b) Find the line integral of  $\vec{F} = -y\hat{i} + x\hat{j}$  along the curve, oriented counter-clockwise.

∫ Green's theorem in vector form / Stokes theorem

$$\oint_C \vec{F} \cdot d\vec{s} = \iint_D (\nabla \times \vec{F}) \cdot \hat{k} dA$$

$\int_C 2 dA = 2 \cdot 6\pi a^2 = \boxed{12\pi a^2}$

$$\nabla \times \vec{F} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \partial_x & \partial_y & \partial_z \\ -y & x & 0 \end{vmatrix} = (0, 0, 2)$$

9. (20pts) Let  $S$  be the circle with centre  $(2, 3, -1)$  and radius 3 lying in the plane with normal vector  $\hat{n} = (\frac{1}{3}, \frac{2}{3}, -\frac{2}{3})$ . Find the flux of the vector field  $\vec{F}(x, y, z) = y\hat{j} + z\hat{j} + x\hat{k}$  through  $S$  in the direction of  $\hat{n}$ .

First find two vectors  $\hat{a}, \hat{b}$  orthogonal to  $\hat{n}$ .

$$\hat{a} = \left(-\frac{2}{3}, \frac{2}{3}, \frac{1}{3}\right) \quad \hat{a} \cdot \hat{n} = 0$$

$$\hat{b} = \hat{n} \times \hat{a} = \begin{vmatrix} \hat{i} & \hat{j} & \hat{k} \\ \frac{1}{3} & \frac{2}{3} & -\frac{2}{3} \\ -\frac{2}{3} & \frac{2}{3} & \frac{1}{3} \end{vmatrix} = \frac{2}{3}\hat{i} + \frac{1}{3}\hat{j} + \frac{1}{3}\hat{k}$$

$(\hat{a}, \hat{b}, \hat{n})$  right handed set.

$$U = [0, 2] \times (0, \pi)$$

$$\hat{g}: U \rightarrow \mathbb{R}^3$$

$$(r, \theta) \mapsto \vec{OP} + r \cos \theta \hat{a} + r \sin \theta \hat{b}$$

$$\frac{d\hat{g}}{dr} = \cos \theta \hat{a} + \sin \theta \hat{b}$$

$$\frac{d\hat{g}}{d\theta} = -r \sin \theta \hat{a} + r \cos \theta \hat{b}$$

$$\frac{d\hat{g}}{dr} \times \frac{d\hat{g}}{d\theta} = (\cos \theta \hat{a} + \sin \theta \hat{b}) \times (-r \sin \theta \hat{a} + r \cos \theta \hat{b})$$

$$= r(\cos^2 \theta + \sin^2 \theta) \hat{n} = r \hat{n}$$

$$\iint_S \vec{F} \cdot d\vec{s} = \iint_U \vec{F} \left( \frac{\partial \hat{g}}{\partial r} \times \frac{\partial \hat{g}}{\partial \theta} \right) ds = \iint_U (\vec{F}(r, \theta) \cdot \hat{n}) dr d\theta$$

$$= \int_0^{2\pi} \left[ \int_0^r \left[ -r^2 \right] dr \right] d\theta = \int_0^{2\pi} \int_0^r (r(r + p \cos \theta + q \sin \theta)) dr d\theta$$

$$\alpha = \frac{3}{3} - \frac{2}{3} - \frac{2}{3} = -1$$

$$\int_0^{2\pi} \cos \theta d\theta = \int_0^{2\pi} \sin \theta d\theta = 0$$

ignore  $p, q$ .

10. (20pts) Let  $S_a(P)$  denote the sphere centred at  $P$  of radius  $a$  and oriented outwards. A smooth vector field  $\vec{F}$  is defined on all of  $\mathbb{R}^3$  except the three points  $P_1 = (0, 0, 0)$ ,  $P_2 = (4, 0, 0)$  and  $P_3 = (8, 0, 0)$ . Suppose that the divergence of  $\vec{F}$  is zero and that

$$\iint_{S_1(P_1)} \vec{F} \cdot d\vec{S} = 1, \quad \iint_{S_6(P_1)} \vec{F} \cdot d\vec{S} = 3 \quad \text{and} \quad \iint_{S_6(P_3)} \vec{F} \cdot d\vec{S} = 5.$$

Find the following flux integrals:

(a)

$$\iint_{S_1(P_2)} \vec{F} \cdot d\vec{S} = 2$$

Let  $M$  be the closed ball of radius 6 containing  $P_1$ , minus the interior of the balls of radius 1 about  $P_1$  and  $P_2$ . Then  $\partial M = S_6(P_1) + S'_1(P_1) + S'_1(P_2)$ .  
and  $M$  is a smooth 2-manifold with boundary

(b)

same reasoning:

$$\begin{aligned}
 & \iint_{S_1(P_3)} \vec{F} \cdot d\vec{S} \\
 & 5 - 2 \\
 & \quad || \\
 & \quad 3. \\
 & \left| \begin{array}{l} \text{Gauss divergence theorem:} \\ 0 = \iiint_M \nabla \cdot \vec{F} dV = \iint_{\partial M} \vec{F} \cdot d\vec{S} \\ = \iint_{S_6(P_1)} \vec{F} \cdot d\vec{S} + \iint_{S'_1(P_1)} \vec{F} \cdot d\vec{S} + \iint_{S'_1(P_2)} \vec{F} \cdot d\vec{S} \\ \iint_{S_6(P_1)} \vec{F} \cdot d\vec{S} = \iint_{S_1(P_1)} \vec{F} \cdot d\vec{S} - \iint_{S'_1(P_1)} \vec{F} \cdot d\vec{S} \\ = 3 - 1 \\ = 2 \end{array} \right.
 \end{aligned}$$