

22. DOUBLE INTEGRALS

Definition 22.1. Let $R = [a, b] \times [c, d] \subset \mathbb{R}^2$ be a rectangle in the plane. A **partition** \mathcal{P} of R is a pair of sequences:

$$\begin{aligned} a &= x_0 < x_1 < \cdots < x_n = b \\ c &= y_0 < y_1 < \cdots < y_n = d. \end{aligned}$$

The **mesh** of \mathcal{P} is

$$m(\mathcal{P}) = \max\{x_i - x_{i-1}, y_i - y_{i-1} \mid 1 \leq i \leq n\}.$$

Now suppose we are given a function

$$f: R \longrightarrow \mathbb{R}$$

Pick

$$\vec{c}_{ij} \in R_{ij} = [x_{i-1}, x_i] \times [y_{j-1}, y_j].$$

Definition 22.2. The sum

$$S = \sum_{i=1}^n \sum_{j=1}^n f(\vec{c}_{ij})(x_i - x_{i-1})(y_j - y_{j-1}),$$

is called a **Riemann sum**.

We will use the short hand notation

$$\Delta x_i = x_i - x_{i-1} \quad \text{and} \quad \Delta y_j = y_j - y_{j-1}.$$

Definition 22.3. The function $f: R \longrightarrow \mathbb{R}$ is called **integrable**, with integral I , if for every $\epsilon > 0$, we may find a $\delta > 0$ such that for every mesh \mathcal{P} whose mesh size is less than δ , we have

$$|I - S| < \epsilon,$$

where S is any Riemann sum associated to \mathcal{P} .

We write

$$\iint_R f(x, y) \, dx \, dy = I,$$

to mean that f is integrable with integral I .

We use a sneaky trick to integrate over regions other than rectangles. Suppose that D is a bounded subset of the plane. Then we can find a rectangle R which completely contains D .

Definition 22.4. The **indicator function** of $D \subset R$ is the function

$$i_D: R \longrightarrow \mathbb{R},$$

given by

$$i_D(x) = \begin{cases} 1 & \text{if } x \in D \\ 0 & \text{if } x \notin D. \end{cases}$$

If i_D is integrable, then we say that the **area of D** is the integral

$$\iint_R i_D \, dx \, dy.$$

If i_D is not integrable, then D does not have an area.

Example 22.5. Let

$$D = \{ (x, y) \in [0, 1] \times [0, 1] \mid x, y \in \mathbb{Q} \}.$$

Then D does not have an area.

Definition 22.6. If $f: D \rightarrow \mathbb{R}$ is a function and D is bounded, then pick $D \subset R \subset \mathbb{R}^2$ a rectangle. Define

$$\tilde{f}: R \rightarrow \mathbb{R},$$

by the rule

$$\tilde{f}(x) = \begin{cases} f(x) & \text{if } x \in D \\ 0 & \text{otherwise.} \end{cases}$$

We say that f is **integrable** over D if \tilde{f} is integrable over R . In this case

$$\iint_D f(x, y) \, dx \, dy = \iint_R \tilde{f}(x, y) \, dx \, dy.$$

Proposition 22.7. Let $D \subset \mathbb{R}^2$ be a bounded subset and let $f: D \rightarrow \mathbb{R}$ and $g: D \rightarrow \mathbb{R}$ be two integrable functions. Let λ be a scalar.

Then

(1) $f + g$ is integrable over D and

$$\iint_D f(x, y) + g(x, y) \, dx \, dy = \iint_D f(x, y) \, dx \, dy + \iint_D g(x, y) \, dx \, dy.$$

(2) λf is integrable over D and

$$\iint_D \lambda f(x, y) \, dx \, dy = \lambda \iint_D f(x, y) \, dx \, dy.$$

(3) If $f(x, y) \leq g(x, y)$ for any $(x, y) \in D$, then

$$\iint_D f(x, y) \, dx \, dy \leq \iint_D g(x, y) \, dx \, dy.$$

(4) $|f|$ is integrable over D and

$$\left| \iint_D f(x, y) \, dx \, dy \right| \leq \iint_D |f(x, y)| \, dx \, dy.$$

It is straightforward to integrate continuous functions over regions of three special types:

Definition 22.8. A bounded subset $D \subset \mathbb{R}^2$ is an **elementary region** if it is one of three types:

Type 1:

$$D = \{ (x, y) \in \mathbb{R}^2 \mid a \leq x \leq b, \gamma(x) \leq y \leq \delta(x) \},$$

where $\gamma: [a, b] \rightarrow \mathbb{R}$ and $\delta: [a, b] \rightarrow \mathbb{R}$ are continuous functions.

Type 2:

$$D = \{ (x, y) \in \mathbb{R}^2 \mid c \leq y \leq d, \alpha(y) \leq x \leq \beta(y) \},$$

where $\alpha: [c, d] \rightarrow \mathbb{R}$ and $\beta: [c, d] \rightarrow \mathbb{R}$ are continuous functions.

Type 3: D is both type 1 and 2.

Theorem 22.9. Let $D \subset \mathbb{R}^2$ be an elementary region and let $f: D \rightarrow \mathbb{R}$ be a continuous function.

Then

(1) If D is of type 1, then

$$\iint_D f(x, y) \, dx \, dy = \int_a^b \left(\int_{\gamma(x)}^{\delta(x)} f(x, y) \, dy \right) dx.$$

(2) If D is of type 2, then

$$\iint_D f(x, y) \, dx \, dy = \int_c^d \left(\int_{\alpha(y)}^{\beta(y)} f(x, y) \, dx \right) dy.$$

Example 22.10. Let D be the region bounded by the lines $x = 0$, $y = 4$ and the parabola $y = x^2$. Let $f: D \rightarrow \mathbb{R}$ be the function given by $f(x, y) = x^2 + y^2$.

If we view D as a region of type 1, then we get

$$\begin{aligned}
 \iint_D f(x, y) \, dx \, dy &= \int_0^2 \left(\int_{x^2}^4 x^2 + y^2 \, dy \right) dx \\
 &= \int_0^2 \left[x^2 y + \frac{y^3}{3} \right]_{x^2}^4 dx \\
 &= \int_0^2 4x^2 + \frac{2^6}{3} - x^4 - \frac{x^6}{3} dx \\
 &= \left[\frac{4x^3}{3} + \frac{2^6 x}{3} - \frac{x^5}{5} - \frac{x^7}{3 \cdot 7} \right]_0^2 \\
 &= \frac{2^5}{3} + \frac{2^7}{3} - \frac{2^5}{5} - \frac{2^7}{3 \cdot 7} \\
 &= \frac{2^6}{3 \cdot 5} + \frac{2^8}{7} \\
 &= 2^6 \left(\frac{1}{3 \cdot 5} + \frac{2^2}{7} \right).
 \end{aligned}$$

On the other hand, if we view D as a region of type 2, then we get

$$\begin{aligned}
 \iint_D f(x, y) \, dx \, dy &= \int_0^4 \left(\int_0^{\sqrt{y}} x^2 + y^2 \, dx \right) dy \\
 &= \int_0^4 \left[\frac{x^3}{3} + xy^2 \right]_0^{\sqrt{y}} dy \\
 &= \int_0^4 \frac{y^{3/2}}{3} + y^{5/2} dy \\
 &= \left[\frac{2y^{5/2}}{3 \cdot 5} + \frac{2y^{7/2}}{7} \right]_0^4 \\
 &= \frac{2^6}{3 \cdot 5} + \frac{2^8}{7} \\
 &= 2^6 \left(\frac{1}{3 \cdot 5} + \frac{2^2}{7} \right).
 \end{aligned}$$