

## 26. CHANGE OF COORDINATES: II

**Example 26.1.** Let  $D$  be the region bounded by the cardioid,

$$r = 1 - \cos \theta.$$

If we multiply both sides by  $r$ , take  $r \cos \theta$  over the other side, then we get

$$(x^2 + y^2 + x)^2 = x^2 + y^2.$$

We have

$$\begin{aligned} \text{area}(D) &= \iint_D dx dy \\ &= \iint_{D^*} r dr d\theta \\ &= \int_{-\pi}^{\pi} \left( \int_0^{1-\cos\theta} r dr \right) d\theta \\ &= \int_{-\pi}^{\pi} \left[ \frac{r^2}{2} \right]_0^{1-\cos\theta} d\theta \\ &= \int_{-\pi}^{\pi} \frac{(1 - \cos \theta)^2}{2} d\theta \\ &= \int_{-\pi}^{\pi} \frac{1}{2} - \cos \theta + \frac{\cos^2 \theta}{2} d\theta \\ &= \left[ \frac{\theta}{2} - \sin \theta \right]_{-\pi}^{\pi} \frac{\pi}{2} \\ &= \frac{\pi}{2}. \end{aligned}$$

In  $\mathbb{R}^3$ , we can either use cylindrical or spherical coordinates, instead of Cartesian coordinates.

Let's first do the case of cylindrical coordinates. Recall that

$$\begin{aligned} x &= r \cos \theta \\ y &= r \sin \theta \\ z &= z. \end{aligned}$$

So the Jacobian is given by

$$\frac{\partial(x, y, z)}{\partial(r, \theta, z)}(r, \theta, z) = \begin{vmatrix} \cos \theta & -r \sin \theta & 0 \\ \sin \theta & r \cos \theta & 0 \\ 0 & 0 & 1 \end{vmatrix} = r.$$

So,

$$\iiint_W f(x, y, z) dx dy dz = \iiint_{W^*} f(r, \theta, z) dr d\theta dz.$$

**Example 26.2.** Consider a cone of height  $b$  and base radius  $a$ . Put the vertex of the cone at the point  $(0, 0, b)$ , so that the base of the cone is the circle of radius  $a$ , centred at the origin, in the  $xy$ -plane. Note that at height  $z$ , we have a circle of radius

$$a \left(1 - \frac{z}{b}\right).$$

$$\begin{aligned} \text{vol}(W) &= \iiint_W dx dy dz \\ &= \iiint_{W^*} r dr d\theta dz \\ &= \int_0^b \left( \int_0^{2\pi} \left( \int_0^{a(1-z/b)} r dr \right) d\theta \right) dz \\ &= \frac{1}{2} \int_0^b \left( \int_0^{2\pi} [r^2]_0^{a(1-z/b)} d\theta \right) dz \\ &= \frac{1}{2} \int_0^b \left( \int_0^{2\pi} a^2 \left(1 - \frac{z}{b}\right)^2 d\theta \right) dz \\ &= \pi a^2 \int_{-a}^a \left(1 - \frac{z}{b}\right)^2 dz \\ &= -\pi a^2 b \int_1^0 u^2 du \\ &= \pi a^2 b \int_0^1 u^2 du \\ &= \frac{\pi a^2 b}{3}. \end{aligned}$$

**Example 26.3.** Consider a ball of radius  $a$ . Put the centre of the ball at the point  $(0, 0, 0)$ . Note that

$$x^2 + y^2 + z^2 = a^2,$$

translates to the equation

$$r^2 + z^2 = a^2,$$

so that

$$r = \sqrt{a^2 - z^2}.$$

$$\begin{aligned}
\text{vol}(W) &= \iiint_W dx dy dz \\
&= \iint_{W^*} r dr d\theta dz \\
&= \int_{-a}^a \left( \int_0^{2\pi} \left( \int_0^{\sqrt{a^2-z^2}} r dr \right) d\theta \right) dz \\
&= \frac{1}{2} \int_{-a}^a \left( \int_0^{2\pi} [r^2]_0^{\sqrt{a^2-z^2}} d\theta \right) dz \\
&= \frac{1}{2} \int_{-a}^a \left( \int_0^{2\pi} a^2 - z^2 d\theta \right) dz \\
&= \pi \int_{-a}^a a^2 - z^2 dz \\
&= \pi \left[ a^2 z - \frac{z^3}{3} \right]_{-a}^a \\
&= \frac{4\pi a^3}{3}.
\end{aligned}$$

Now consider using spherical coordinates. Recall that

$$\begin{aligned}
x &= \rho \sin \phi \cos \theta \\
y &= \rho \sin \phi \sin \theta \\
z &= \rho \cos \phi.
\end{aligned}$$

So

$$\begin{aligned}
\frac{\partial(x, y, z)}{\partial(\rho, \phi, \theta)}(\rho, \phi, \theta) &= \begin{vmatrix} \sin \phi \cos \theta & \rho \cos \phi \cos \theta & -\rho \sin \phi \sin \theta \\ \sin \phi \sin \theta & \rho \cos \phi \sin \theta & \rho \sin \phi \cos \theta \\ \cos \phi & -\rho \sin \phi & 0 \end{vmatrix} \\
&= \rho^2 \cos^2 \phi \sin \phi + \rho^2 \sin^3 \phi = \rho^2 \sin \phi.
\end{aligned}$$

Notice that this is greater than zero, if  $0 < \phi < \pi$ . So,

$$\iiint_W f(x, y, z) dx dy dz = \iiint_{W^*} f(\rho, \phi, \theta) \rho^2 \sin \phi dr d\theta dz.$$

**Example 26.4.** Consider a ball of radius  $a$ . Put the centre of the ball at the point  $(0, 0, 0)$ .

$$\begin{aligned}
\text{vol}(W) &= \iiint_W dx dy dz \\
&= \iiint_{W^*} \rho^2 \sin \phi d\rho d\phi d\theta \\
&= \int_0^{2\pi} \left( \int_0^\pi \left( \int_0^a \rho^2 \sin \phi d\rho \right) d\phi \right) d\theta \\
&= \int_0^{2\pi} \left( \int_0^\pi \sin \phi \left[ \frac{\rho^3}{3} \right]_0^a d\phi \right) d\theta \\
&= \int_0^{2\pi} \left( \int_0^\pi \sin \phi \frac{a^3}{3} d\phi \right) d\theta \\
&= \frac{a^3}{3} \int_0^{2\pi} [-\cos \phi]_0^\pi d\theta \\
&= \frac{2a^3}{3} \int_0^{2\pi} d\theta \\
&= \frac{4\pi a^3}{3}.
\end{aligned}$$