

## HWK #6, DUE WEDNESDAY 03/16

1. Let  $\pi: X \rightarrow B$  be a projective and surjective morphism with connected fibres of dimension  $n$ , where  $X$  and  $B$  are quasi-projective and  $X$  is irreducible. Let  $f: X \rightarrow Y$  be a morphism of quasi-projective varieties.

If there is a point  $b_0 \in B$  such that  $f(\pi^{-1}(b_0))$  is a point, then  $f(\pi^{-1}(b))$  is a point for every  $b \in B$ . This result is known as *the rigidity lemma*. (*Hint: consider the morphism  $f \times \pi: X \rightarrow Y \times B$* ).

2. Recall that an abelian variety  $A$  is a connected and projective algebraic group (you may assume that a connected algebraic group is irreducible). Show that every abelian variety is a commutative group. (*Hint: consider the morphism  $A \times A \rightarrow A$  given by conjugation*).

If  $A$  is a commutative algebraic group and  $a \in A$  then the action of  $A$  on itself by left (or right) translation defines a morphism  $\tau_a: A \rightarrow A$ ,  $\tau_a(x) = x + a$ . We will refer to any such morphism as a *translation*.

3. Show that if  $\pi: A \rightarrow B$  is a morphism of abelian varieties then  $\pi$  is the composition of a translation and a group homomorphism.

4. Show that if  $\pi: G \rightarrow H$  is a morphism of algebraic tori then  $\pi$  is the composition of a translation and a group homomorphism. In particular, if  $G = \mathbb{G}_m^n$  and  $H = \mathbb{G}_m^n$  and  $\pi$  sends the identity to the identity then there are integers  $a_1, a_2, \dots, a_n$  such that  $\pi(t) = (t^{a_1}, t^{a_2}, \dots, t^{a_n})$ . (*Hint: consider the map of group algebras*).

5. Let  $A$  be an abelian variety. Show that every rational map  $f: \mathbb{P}^1 \dashrightarrow A$  is constant. You may use the fact that every morphism  $\pi: G \rightarrow A$  is a composition of a translation and a group homomorphism, where  $G$  is a group isomorphic to either  $\mathbb{G}_a$  or  $\mathbb{G}_m$ .

(*Just for fun: For those who know some of the theory of complex manifolds, note that if the underlying field is  $\mathbb{C}$ , then every abelian variety is a complex torus. Give another proof that  $f$  is constant in this case*).

6. Let  $X$  and  $Y$  be two projective varieties in  $\mathbb{P}^n$  of dimensions  $d$  and  $e$ .

(i) Show that if  $X$  and  $Y$  belong to linear spaces which don't intersect then the dimension of the join of  $X$  and  $Y$  is equal to  $d + e + 1$ .

(ii) Show that if  $X$  and  $Y$  don't intersect then the dimension of the join of  $X$  and  $Y$  is equal to  $d + e + 1$  (*Hint: reduce to the case above, by realising  $X$  and  $Y$  as the projection of  $\tilde{X}$  and  $\tilde{Y}$  in  $\mathbb{P}^{2n+1}$* ).

(iii) Show that if  $d + e \geq n$  then  $X$  and  $Y$  must intersect.