

### 13. COHOMOLOGY OF AFFINE SCHEME

**Proposition 13.1.** *Let  $I$  be an injective module over a noetherian ring  $A$ .*

*Then the sheaf  $\tilde{I}$  on  $X = \text{Spec } A$  is flasque.*

**Corollary 13.2.** *Let  $X$  be a noetherian scheme.*

*Then ever quasi-coherent sheaf  $\mathcal{F}$  on  $X$  can be embedded in a flasque, quasi-coherent sheaf  $\mathcal{G}$ .*

*Proof.* Let  $U_i = \text{Spec } A_i$  be a finite open affine cover of  $X$  and let  $\mathcal{F}|_{U_i} = \tilde{M}_i$  for each  $i$ . Pick an embedding of  $M_i$  into an injective  $A_i$ -module  $I_i$ . Let  $f_i: U_i \rightarrow X$  be the inclusion and let

$$\mathcal{G} = \bigoplus_i f_{i*} \tilde{I}_i.$$

Now for each  $i$  there is an injective map  $\mathcal{F}|_{U_i} \rightarrow \tilde{I}_i$ , which induces a map  $\mathcal{F} \rightarrow f_{i*} \tilde{I}_i$ . This induces a map  $\mathcal{F} \rightarrow \mathcal{G}$ , which is clearly injective.

But  $\tilde{I}_i$  is flasque and quasi-coherent on  $U_i$ , so that  $f_{i*} \tilde{I}_i$  is flasque and quasi-coherent on  $X$ . But then  $\mathcal{G}$  is flasque and quasi-coherent.  $\square$

**Theorem 13.3** (Serre). *Let  $X$  be a Noetherian scheme.*

*TFAE*

- (1)  $X$  is affine,
- (2)  $H^i(X, \mathcal{F}) = 0$  for all  $i > 0$  and all quasi-coherent sheaves,
- (3)  $H^1(X, \mathcal{I}) = 0$  for all coherent sheaves of ideals  $\mathcal{I}$ .

*Proof.* Suppose  $X$  is affine. Let  $M = H^0(X, \mathcal{F})$  and take an injective resolution  $I^\bullet$  of  $M$  in the category of  $A$ -modules. Then  $\tilde{I}^\bullet$  is a flasque resolution of  $\mathcal{F}$  on  $X$ . If we take global sections we get back the original injective resolution of  $A$ , so that  $H^i(X, \mathcal{F}) = 0$  for all  $i > 0$ . Thus (i) implies (ii).

(ii) implies (iii) is easy. Suppose that  $H^1(X, \mathcal{I}) = 0$  for all coherent sheaves of ideals  $\mathcal{I}$ .

Fix a closed point  $p$  of  $X$  together with an open neighbourhood  $U$  of  $p$  and let  $Y = X - U$ . Then there is a short exact sequence

$$0 \rightarrow \mathcal{I}_{Y \cup \{P\}} \rightarrow \mathcal{I}_Y \rightarrow k(P) \rightarrow 0.$$

This gives us an exact sequence

$$H^0(X, \mathcal{I}_Y) \rightarrow H^0(X, k(P)) \rightarrow H^1(X, \mathcal{I}_{Y \cup \{P\}}) \rightarrow 0.$$

But then there is regular function  $f \in A = H^0(U, \mathcal{O}_U)$  which is not zero at  $p$ , so that  $p \in X_f \subset U$  is an open neighbourhood of  $p$ . As  $X_f = U_f$  it follows that  $X_f$  is affine.

As  $X$  is noetherian, it is compact, so that we can cover  $X$  by finitely many open affines,  $X_{f_i}$ , where  $f_1, f_2, \dots, f_r \in A$ .

Finally we check that  $f_1, f_2, \dots, f_r$  generate the unit ideal. There is a short exact sequence of sheaves

$$0 \longrightarrow \mathcal{F} \longrightarrow \mathcal{O}_X^r \longrightarrow \mathcal{O}_X \longrightarrow 0.$$

The last map  $\alpha$  sends  $(a_1, a_2, \dots, a_r)$  to  $\sum a_i f_i$ . It is surjective as it is surjective on stalks.  $\mathcal{F}$  is then the kernel of  $\alpha$ .

There is a filtration of  $\mathcal{F}$  as follows:

$$\mathcal{F} \cap \mathcal{O}_X \subset \mathcal{F} \cap \mathcal{O}_X^2 \subset \mathcal{F} \cap \mathcal{O}_X^3 \subset \mathcal{F} \cap \mathcal{O}_X^r = \mathcal{F}.$$

The quotients are naturally  $\mathcal{O}_X$ -submodules of  $\mathcal{O}_X$ , that is, the quotients are coherent sheaves of ideals. Taking the long exact sequence of cohomology ( $r$  times), we get that  $H^1(X, \mathcal{F}) = 0$ . Taking the long exact sequence of cohomology of the sequence above, we get that  $\alpha$  is surjective. But then

$$1 = \alpha(a_1, a_2, \dots, a_r) = \sum a_i f_i,$$

in the ideal generated by  $f_1, f_2, \dots, f_r$ . (II.2.17) shows that  $X$  is affine.  $\square$