

**THE PROBLEMS FOR THE SECOND TEST FOR 18.102  
WILL BE SELECTED FROM THIS LIST**

RICHARD MELROSE

Question 1

Show that if  $A \in \mathcal{B}(H)$ , where  $H$  is a separable Hilbert space, has the property that  $u_n \rightharpoonup u$  (converges weakly) implies that  $Au_n$  is Cauchy. Show that  $A(\overline{\{u \in H; \|u\| \leq 1\}})$  is compact.

Question 2

If  $H$  is an infinite-dimensional separable Hilbert space, define a set of operators  $T(H)$  by the property

$$A \in T(H) \text{ iff } A \in \mathcal{B}(H) \text{ and } |A| \in \mathcal{K}(H), \sum_i \lambda_i(|A|) < \infty$$

where the  $\lambda_i(|A|)$  are the eigenvalues of  $|A|$  repeated with multiplicity. Show that  $T(H)$  is a 2-sided \*-closed ideal.

Question 3

Suppose  $A \in \mathcal{B}(H)$  is a bounded operator on a Hilbert space such that  $A^*A$  is a compact operator. Show that  $A$  is a compact operator.

Question 4

If  $H$  is a separable, infinite dimensional, Hilbert space set

$$(1) \quad l^2(H) = \{u : \mathbb{N} \longrightarrow H; \|u\|_{l^2(H)}^2 = \sum_i \|u_i\|_H^2 < \infty\}.$$

Show that  $l^2(H)$  has a Hilbert space structure and construct an explicit isometric (norm-preserving) isomorphism (bijection) from  $l^2(H)$  to  $H$ .

Question 5

Starting from the definition of weak convergence of a sequence in a separable Hilbert space,  $v_n \rightharpoonup v$ , that  $\langle v_n, w \rangle \rightarrow \langle v, w \rangle$  in  $\mathbb{C}$  for each  $w \in H$ , show that a weakly convergent sequence  $\{v_n\}$ , is (strongly) convergent if and only if

$$(2) \quad \|v\|_H = \lim_{n \rightarrow \infty} \|v_n\|_H.$$

## Question 6

Let  $e_k, k \in \mathbb{N}$ , be an orthonormal basis in a separable Hilbert space,  $H$ . Show that there is a unique bounded linear operator  $T : H \rightarrow H$  satisfying

$$(3) \quad Te_j = e_{j-1} \quad \forall j \geq 2, \quad Te_1 = 0,$$

and that if  $B \in \mathcal{B}(H)$  has  $\|B\| < 1$  then  $T + B$  has one-dimensional null space.

## Question 7

Suppose  $H$  is an infinite dimensional separable Hilbert space with an orthonormal basis  $\{e_k\}_{k=1}^{\infty}$ . Show that a continuous function  $K : [0, 1] \rightarrow H$  has the property that the Fourier-Bessel series of  $K(x) \in H$ , for  $x \in [0, 1]$ , converges uniformly in the sense that if  $K_n(x) = \sum_{k \leq n} \langle K(x), e_k \rangle e_k$  then  $K_n : [0, 1] \rightarrow H$  is also continuous and

$$(4) \quad \sup_{x \in [0, 1]} \|K(x) - K_n(x)\|_H \rightarrow 0.$$

## Question 8

If  $A \in \mathcal{B}(H)$  is a bounded operator on a separable, infinite-dimensional, Hilbert space  $H$ , explain why  $|A| = (A^*A)^{1/2} \in \mathcal{B}(H)$  is well-defined. Suppose that  $|A|$  is compact and for a orthonormal basis of  $H$  consisting of eigenvectors of  $|A|$ ,

$$(5) \quad \sum_i \| |A| e_i \|^2 < \infty.$$

Show that

$$\sum_i \| A f_j \|^2 < \infty$$

for any orthonormal sequence  $f_j$ .

## Question 9

Show that a separable Hilbert space  $H$  with the property that every bounded operator from  $H$  to itself is compact is necessarily finite dimensional.

## Question 10

Show that if  $B$  is a compact operator on a separable Hilbert space  $H$  and  $A$  is an invertible operator then

$$(6) \quad \{u \in H; Bu = Au\}$$

is finite dimensional.

## Question 11

Assuming the Stone-Weierstrass theorem, show that there is a complete orthonormal basis of  $L^2([0, 2\pi])$  consisting of polynomials.

### Question 12

Let  $H$  be a separable Hilbert space and let  $\mathcal{C}_c(\mathbb{R}; H)$  be the linear space of continuous maps from  $\mathbb{R}$  to  $H$  which vanish outside some interval  $[-R, R]$  depending on the function. Show that

$$(7) \quad \|u\|^2 = \int_{\mathbb{R}} \|u(x)\|_H^2$$

defines a norm which comes from a pre-Hilbert structure on  $\mathcal{C}_c(\mathbb{R}; H)$ . Show that if  $u_n$  is a Cauchy sequence in this pre-Hilbert space and  $h \in H$  then  $\langle u_n(x), h \rangle_H$  converges in  $L^2(\mathbb{R})$ .

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY

*Email address:* `rbm@math.mit.edu`