

**PROBLEM SET 10 FOR 18.102, SPRING 2016  
DUE FRIDAY 6 MAY.**

RICHARD MELROSE

This is the last problem set. You may freely use the fact that the Fourier transform extends from an isomorphism on Schwartz space  $\mathcal{S}(\mathbb{R})$ , which is a dense subspace of  $L^2(\mathbb{R})$ , to an isomorphism of  $L^2(\mathbb{R})$ .

Define  $H^2(\mathbb{R}) \subset L^2(\mathbb{R})$  by the condition

$$u \in H^2(\mathbb{R}) \iff u \in L^2(\mathbb{R}) \text{ and } \xi^2 \hat{u}(\xi) \in L^2(\mathbb{R}).$$

P10.1 Show that  $H^2(\mathbb{R})$  is a Hilbert space with the norm  $(\|u\|_{L^2}^2 + \|D^2 u\|_{L^2}^2)^{\frac{1}{2}}$  where  $\widehat{D^2 u}(\xi) = \xi^2 \hat{u}(\xi)$ .

Hint: For a Cauchy sequence in  $H^2(\mathbb{R})$  both  $u_n \rightarrow u$  and  $D^2 u_n \rightarrow v$  converge in  $L^2$  so you only need show that  $\hat{v} = \xi^2 \hat{u}$  and this follows from Monotonicity/LDC.

P10.2 Show that if  $u \in H^2(\mathbb{R})$  then  $u$  ‘is’ continuously differentiable (meaning, since we value precision, has a representative which is a continuously differentiable function on  $\mathbb{R}$ ).

Hint: Since  $\hat{u}$  and  $\xi^2 \hat{u} \in L^2$  it follows that  $\hat{u}$  and  $\xi \hat{u} \in L^1$  by Cauchy-Schwartz, so they have bounded continuous inverse FTs,  $u, v$ . Apply LDC to the integral for the IFT giving  $u$  to see that the difference quotient converges to  $v$ .

P10.3 Show that  $D^2 + 1$  is an isomorphism from  $H^2(\mathbb{R})$  to  $L^2(\mathbb{R})$

Hint: The inverse is  $\times(1 + \xi^2)^{-1}$  on the FT side.

P10.4 Show that  $(D^2 + 1)^{-1}$  is a self-adjoint operator on  $L^2(\mathbb{R})$  and that it has spectrum precisely the interval  $[0, 1]$ .

P10.5 Prove that if  $V \geq 0$  is a bounded continuous function on  $\mathbb{R}$  then

$$(1) \quad (D^2 + 1 + V) : H^2(\mathbb{R}) \longrightarrow L^2(\mathbb{R})$$

is a topological isomorphism, i.e. a bijection with a bounded inverse.

Hint: Recall the discussion of the Dirichlet problem. If  $A^2 = (1 + D^2)^{-1}$  is given by the functional calculus then  $\text{Id} + AVA$  is invertible on  $L^2$  and is of the form  $\text{Id} + AEA$  with  $E$  bounded; the inverse to (1) is  $A(\text{Id} + AVA)^{-1}A$  and you need to check that this maps  $L^2$  to  $H^2$  and is indeed the inverse.

P10.6 – extra Show that if  $f \in \mathcal{C}_c(\mathbb{R})$  is a continuous function of compact support then (under the same hypotheses as above)

$$(2) \quad -\frac{d^2 u}{dx^2} + u + Vu = f$$

has a unique twice continuously differentiable solution which is in  $L^2(\mathbb{R})$ .

Hint: Show by integration (making sure of the behaviour at infinity) that the equation has a unique solution  $u$  which is  $\mathcal{C}^2$  and in  $L^2$  for each  $f \in \mathcal{C}_c(\mathbb{R})$ .

P10.7 – extra Show that this solution to (2) defines a self-adjoint operator on  $L^2(\mathbb{R})$  which has spectrum contained in  $[0, 1]$ .

Hint: Then show that the solution in the previous question is actually  $(D^2 + 1)^{-1}f$  and using the  $L^2$  inverse and a regularity argument.

DEPARTMENT OF MATHEMATICS, MASSACHUSETTS INSTITUTE OF TECHNOLOGY  
*E-mail address:* `rbm@math.mit.edu`