

**PROBLEM SET 7, 18.155**  
**DUE 8:30AM NOVEMBER 11, 2017**

This week I ask you to think about *unbounded self-adjoint operators* which you might well encounter. By definition this means a linear map

$$A : D(A) \longrightarrow H$$

where  $D(A) \subset H$  is a dense linear subspace of a (separable) Hilbert space  $H$  and in addition we assume/require symmetry:

$$(Au, v) = (u, Av) \quad \forall u, v \in D(A)$$

and a maximality condition on  $D(A)$  :

if  $v \in H$  and  $D \ni u \longmapsto (Au, v)$  extends to be continuous on  $H$   
then  $v \in D(A)$ .

Q7.1

Show that if  $P(\xi)$  is a real and elliptic polynomial of degree  $m$  in  $n$  variables and  $V \in \mathcal{C}_c^\infty(\mathbb{R}^n)$  is real-valued then

$$P(D) + V(x) : H^m(\mathbb{R}^n) \longrightarrow L^2(\mathbb{R}^n)$$

is an unbounded self-adjoint operator (with  $H^m(\mathbb{R}^n)$  as domain).

Hint: Use the density of  $\mathcal{S}(\mathbb{R}^n)$  in  $H^m(\mathbb{R}^n)$  to get the symmetry condition. If  $v \in L^2(\mathbb{R}^n)$  satisfies the adjoint condition then  $(P(D)+V)v \in L^2(\mathbb{R}^n)$  and hence  $v \in H^m(\mathbb{R}^n)$  by elliptic regularity.

Q7.2

If  $A$  is unbounded self-adjoint show that

- (1) The graph of  $A$  is closed in  $H \times H$ .
- (2) The spectrum, defined as

$$\text{Spec}(A) = \{z \in \mathbb{C}; A - z \text{Id} : D(A) \longrightarrow H \text{ is not a bijection}\}$$

is a closed subset of  $\mathbb{R}$ .

- (3)  $R(A, z) = (A - z \text{Id})^{-1} \in \mathcal{B}(H)$  for  $z \in \mathbb{C} \setminus \text{Spec}(A)$  is holomorphic in  $z$ .

Hint:

- (1) If  $(v_n, A(v_n))$  in the graph and  $(v_n, Av_n) \rightarrow (v, f)$  in  $H \times H$  and  $u \in D(A)$  then by the symmetry condition

$$(1) \quad (Au, v) = \lim_n (Au, v_n) = \lim_n (u, Av_n) = (u, f)$$

extends by continuity to  $u \in H$  so  $v \in D(A)$  and  $Av = f$  (by symmetry) showing that the graph is closed.

- (2) By symmetry if  $u \in D(A)$  then  $(Au, u)$  is real. Certainly  $A + z : D(A) \rightarrow H$  and  $\operatorname{Im}((A + z)u, u) = \operatorname{Im} z \|u\|^2$  so  $A + z$  is injective. If  $u_n \in D(A)$  and  $f_n = Au_n \rightarrow f$  in  $H$  then  $|\operatorname{Im} z| \|u_n\|^2 = \operatorname{Im}(f_n, u_n) \leq \|f_n\| \|u_n\|$ . Applying the same inequality to  $u_n - u_m$  (and assuming  $z$  is not real) it follows that  $u_n$  is Cauchy and hence converges and so  $(A + z)$  is a bijection from  $D(A)$  to  $H$ . Thus the spectrum is contained in  $\mathbb{R}$ . Whenever  $A + z$  is a bijection it has an ‘algebraic’ inverse,  $R(z)$ , the graph of which is the ‘reverse’ of the graph of  $A + z$  so as a map  $H \rightarrow H$   $R(z)$  is bounded since its graph is closed. If  $z \notin \operatorname{Spec}(A)$  then  $R(z)(A + z + t) = \operatorname{Id} + tR(z)$  and similarly for the opposite composition so for  $|t|$  small  $A + z + t$  is also a bijection. Thus the spectrum is closed and real.
- (3) The Neumann series for  $\operatorname{Id} + tR(z)$  converges for small  $t$  and shows that  $R(z)$  is holomorphic on the open set where it is defined.

### Q7.3

If  $A$  is unbounded self-adjoint let  $R(A, i) = BU$  be the polar decomposition of  $R(A, i)$ .

- (1) Show that  $D(A) = \operatorname{Ran}(B)$ .
- (2) If  $E$  is the spectral subspace for  $B$  corresponding to the interval  $(-\infty, \frac{1}{2}]$  (the  $\frac{1}{2}$  is chosen somewhat randomly in  $(0, 1)$ ) then

$$D(A) = E^\perp + \operatorname{Ran}(B|_E),$$

$$A : E^\perp \rightarrow E^\perp \text{ is bounded and self-adjoint}$$

$$A : \operatorname{Ran}(B|_E) \rightarrow E \text{ is a bijection and}$$

$$A^{-1} : E \rightarrow E \text{ is bounded and self-adjoint.}$$

Hint:

- (1) Since  $R(i)$  has range  $D(A)$  it follows from the proof of the polar decomposition that  $U$  is unitary – it is a bijection on  $H$ . Taking the adjoint  $R(-i) = BU^*$  shows that the range of  $B$  is equal to

the range of  $R(-i)$  is  $D(A)$ . Now  $R(i)$  commutes with  $A$  and hence with  $A - i$  and so with  $R(-i) = R(i)^*$  (so  $R(i)$  is normal) and hence  $R(i)$  commutes with  $B^2 = R(-i)R(i)$  and so with  $B$ . [Maybe there is an easier way to see this!]

- (2) Any bounded operator that commutes with  $B$  maps  $E$  to itself and  $E^\perp$  to itself (from the definition of the  $E$  as a spectral subspace for  $B$ ). It follows that  $R(i)$  does this and is invertible on  $E^\perp$  with inverse  $A + i$ ; it follows that  $E^\perp \subset D(A)$  and that  $A : E^\perp \rightarrow E^\perp$  is bounded and self-adjoint.
- (3) I leave the last part to you!

## Q7.4

Show how to define  $f(A) \in \mathcal{B}(H)$  where  $A$  is unbounded self-adjoint and  $f \in \mathcal{C}_0(\mathbb{R}; \mathbb{R})$  is real valued and vanishes at infinity in such a way as to give a continuous linear map

$$\mathcal{C}_0(\mathbb{R}; \mathbb{R}) \ni f \rightarrow f(A) \in \mathcal{B}(H) \text{ s.t.}$$

$$f(A)g(A) = (fg)(A), \quad B = (A^2 + 1)^{-\frac{1}{2}}$$

where  $B$  is given in (Q3). Hint: Define it separately on  $E$  and  $E^\perp$ .

## Q7.5

For the special case of (Q1),  $\Delta + V$ ,  $P(\xi) = |\xi|^2$ , show that

$$\text{Spec}(\Delta + V) \subset [M, \infty), \quad M = \inf_{x \in \mathbb{R}^n} V(x).$$

Hint: Just use the fact that  $V - M \geq 0$  to show that  $A = \Delta - M + V$  is unbounded self-adjoint with domain  $H^2(\mathbb{R}^n)$  and  $(Au, u) \geq 0$  so  $A$  has no spectrum in  $(-\infty, 0)$ .

Q7.6-Opt

Continue (Q5) to show that the part of the spectrum below 0 (if any) consists of a finite number of eigenvalues of finite multiplicity.

## Q7.7-Opt

Under the same conditions as (Q5) show that  $\text{Spec}(\Delta + V) \supset [0, \infty)$ .