

**PROBLEM SET 7 FOR 18.102, SPRING 2013  
DUE FRIDAY 3 MAY (I.E. 4AM SATURDAY 4 MAY).**

RICHARD MELROSE

Problem 7.1

Let  $A \in \mathcal{B}(H)$ ,  $H$  a separable Hilbert space, be such that for some orthonormal basis  $\{e_i\}$

$$(5.1) \quad \sum_i \|Ae_i\|_H^2 < \infty.$$

Show that the same inequality is true for any other orthonormal basis and that  $A^*$  satisfies the same inequality.

Problem 7.2

The elements of  $A \in \mathcal{B}(H)$  as in Problem 7.1 are called ‘Hilbert-Schmidt operators’. Show that these form a 2-sided  $*$ -closed ideal [more precisely it is a 2-sided ideal in  $\mathcal{B}(H)$  consisting of compact operators and the adjoint of any Hilbert-Schmidt operator is also Hilbert-Schmidt]  $\text{HS}(H)$ , inside the compact operators and that

$$(5.2) \quad \langle A, B \rangle = \sum_i \langle Ae_i, Be_i \rangle,$$

for any choice of orthonormal basis, makes this into a Hilbert space.

Problem 7.3

Suppose  $E$  is a compact, self-adjoint and injective operator on a separable infinite-dimensional Hilbert space  $H$  and that it is positive in the sense that  $(Eu, u) \geq 0$  for all  $u \in H$ . Show that there is a decreasing sequence of positive eigenvalues given by the minimax formula:-

$$s_j(E) = \max_{F \subset H; \dim F=j} \left( \min_{u \in F; \|u\|=1} (Eu, u) \right).$$

Problem 7.4

With  $E$  as above, suppose that  $D \in \mathcal{B}(H)$  is a bounded self-adjoint injective operator. Show that

$$s_j(DED) \leq \|D\|^2 s_j(E) \quad \forall j.$$

Problem 7.5

Recall what we have shown in class:- If  $V \in \mathcal{C}([0, 2\pi])$  is real-valued and non-negative then  $\lambda \in \mathbb{R}$  is an eigenvalue for the Dirichlet problem

$$\left(-\frac{d^2}{dx^2} + V(x)\right)u(x) = \lambda u(x) \text{ on } [0, 2\pi], \quad u(0) = 0 = u(2\pi)$$

with a twice-continuously differentiable eigenfunction if and only if  $s = 1/\lambda$  is an eigenvalue of the operator

$$(\text{Id} + AVA)^{-\frac{1}{2}} A^2 (\text{Id} + AVA)^{-\frac{1}{2}}$$

where  $A$  is positive, self-adjoint and compact with eigenvalues  $\frac{2}{\pi k}$ ,  $k \in \mathbb{N}_0$ .

Show that the eigenvalues of the Dirichlet problem, with  $V$  real-valued and continuous, repeated according to multiplicity (the dimension of the eigenspace) and arranged as a non-decreasing sequence,

$$\lambda_1 \leq \lambda_2 \leq \lambda_N \rightarrow \infty$$

satisfy

$$\lambda_k \geq \pi^2 k^2 / 4 + \min_{[0, 2\pi]} V.$$

Problem 7.6-extra

An operator  $T$  on a separable Hilbert space is said to be ‘of trace class’ (where this is just old-fashioned language) if it can be written as a finite sum

$$T = \sum_{i=1}^N A_i B_i$$

where all the  $A_i$ ,  $B_i$  are Hilbert-Schmidt. Show that these trace class operators form a 2-sided ideal in the bounded operators, closed under passage to adjoints and that

$$\sup_{\{e_i\}, \{f_i\}} \sum_i |\langle T e_i, f_i \rangle| < \infty$$

where the sup is over all pairs of orthonormal bases.

Problem 7.7-extra

Show that the trace functional

$$\text{Tr}(T) = \sum_i \langle T e_i, e_i \rangle$$

is well-defined on trace class operators, independent of the orthonormal basis  $\{e_i\}$  used to compute it and that if  $A$  is self-adjoint, compact and of trace class then

$$\text{Tr}(A) = \sum_i s_i$$

is the (absolutely convergent) sum of the eigenvalues repeated with multiplicity.