

8. CONVOLUTION AND DENSITY

We have defined an inclusion map

$$(8.1) \quad \mathcal{S}(\mathbb{R}^n) \ni \varphi \longmapsto u_\varphi \in \mathcal{S}'(\mathbb{R}^n), \quad u_\varphi(\psi) = \int_{\mathbb{R}^n} \varphi(x)\psi(x) dx \quad \forall \psi \in \mathcal{S}(\mathbb{R}^n).$$

This allows us to ‘think of’ $\mathcal{S}(\mathbb{R}^n)$ as a subspace of $\mathcal{S}'(\mathbb{R}^n)$; that is we habitually identify u_φ with φ . We can do this because we know (8.1) to be injective. We can extend the map (8.1) to include bigger spaces

$$(8.2) \quad \begin{aligned} \mathcal{C}_0^0(\mathbb{R}^n) \ni \varphi &\longmapsto u_\varphi \in \mathcal{S}'(\mathbb{R}^n) \\ L^p(\mathbb{R}^n) \ni \varphi &\longmapsto u_\varphi \in \mathcal{S}'(\mathbb{R}^n) \\ M(\mathbb{R}^n) \ni \mu &\longmapsto u_\mu \in \mathcal{S}'(\mathbb{R}^n) \\ u_\mu(\psi) &= \int_{\mathbb{R}^n} \psi d\mu, \end{aligned}$$

but we need to know that these maps are injective before we can forget about them.

We can see this using *convolution*. This is a sort of ‘product’ of functions. To begin with, suppose $v \in \mathcal{C}_0^0(\mathbb{R}^n)$ and $\psi \in \mathcal{S}(\mathbb{R}^n)$. We define a new function by ‘averaging v with respect to ψ ’

$$(8.3) \quad v * \psi(x) = \int_{\mathbb{R}^n} v(x-y)\psi(y) dy.$$

The integral converges by dominated convergence, namely $\psi(y)$ is integrable and v is bounded,

$$|v(x-y)\psi(y)| \leq \|v\|_{\mathcal{C}_0^0} |\psi(y)|.$$

We can use the same sort of estimates to show that $v * \psi$ is continuous. Fix $x \in \mathbb{R}^n$,

$$(8.4) \quad \begin{aligned} v * \psi(x+x') - v * \psi(x) & \\ &= \int (v(x+x'-y) - v(x-y))\psi(y) dy. \end{aligned}$$

To see that this is small for x' small, we split the integral into two pieces. Since ψ is very small near infinity, given $\epsilon > 0$ we can choose R so large that

$$(8.5) \quad \|v\|_\infty \cdot \int_{|y| \geq R} |\psi(y)| dy \leq \epsilon/4.$$

The set $|y| \leq R$ is compact and if $|x| \leq R'$, $|x'| \leq 1$ then $|x+x'-y| \leq R+R'+1$. A continuous function is *uniformly continuous* on any

compact set, so we can choose $\delta > 0$ such that

$$(8.6) \quad \sup_{\substack{|x'| < \delta \\ |y| \leq R}} |v(x + x' - y) - v(x - y)| \cdot \int_{|y| \leq R} |\psi(y)| dy < \epsilon/2.$$

Combining (8.5) and (8.6) we conclude that $v * \psi$ is continuous. Finally, we conclude that

$$(8.7) \quad v \in \mathcal{C}_0^0(\mathbb{R}^n) \Rightarrow v * \psi \in \mathcal{C}_0^0(\mathbb{R}^n).$$

For this we need to show that $v * \psi$ is small at infinity, which follows from the fact that v is small at infinity. Namely given $\epsilon > 0$ there exists $R > 0$ such that $|v(y)| \leq \epsilon$ if $|y| \geq R$. Divide the integral defining the convolution into two

$$\begin{aligned} |v * \psi(x)| &\leq \int_{|y| > R} u(y)\psi(x - y)dy + \int_{|y| < R} |u(y)\psi(x - y)|dy \\ &\leq \epsilon/2 \|\psi\|_\infty + \|u\|_\infty \sup_{B(x, R)} |\psi|. \end{aligned}$$

Since $\psi \in \mathcal{S}(\mathbb{R}^n)$ the last constant tends to 0 as $|x| \rightarrow \infty$.

We can do much better than this! Assuming $|x'| \leq 1$ we can use Taylor's formula with remainder to write

$$(8.8) \quad \psi(z + x') - \psi(z) = \int_0^1 \frac{d}{dt} \psi(z + tx') dt = \sum_{j=1}^n x_j \cdot \tilde{\psi}_j(z, x').$$

As Problem 23 I ask you to check carefully that

$$(8.9) \quad \psi_j(z; x') \in \mathcal{S}(\mathbb{R}^n) \text{ depends continuously on } x' \text{ in } |x'| \leq 1.$$

Going back to (8.3) we can use the translation and reflection-invariance of Lebesgue measure to rewrite the integral (by changing variable) as

$$(8.10) \quad v * \psi(x) = \int_{\mathbb{R}^n} v(y)\psi(x - y) dy.$$

This reverses the role of v and ψ and shows that if *both* v and ψ are in $\mathcal{S}(\mathbb{R}^n)$ then $v * \psi = \psi * v$.

Using this formula on (8.4) we find

$$(8.11) \quad \begin{aligned} v * \psi(x + x') - v * \psi(x) &= \int v(y)(\psi(x + x' - y) - \psi(x - y)) dy \\ &= \sum_{j=1}^n x_j \int_{\mathbb{R}^n} v(y) \tilde{\psi}_j(x - y, x') dy = \sum_{j=1}^n x_j (v * \psi_j(\cdot; x'))(x). \end{aligned}$$

From (8.9) and what we have already shown, $v * \psi(\cdot; x')$ is continuous in both variables, and is in $\mathcal{C}_0^0(\mathbb{R}^n)$ in the first. Thus

$$(8.12) \quad v \in \mathcal{C}_0^0(\mathbb{R}^n), \psi \in \mathcal{S}(\mathbb{R}^n) \Rightarrow v * \psi \in \mathcal{C}_0^1(\mathbb{R}^n).$$

In fact we also see that

$$(8.13) \quad \frac{\partial}{\partial x_j} v * \psi = v * \frac{\partial \psi}{\partial x_j}.$$

Thus $v * \psi$ inherits its regularity from ψ .

Proposition 8.1. *If $v \in \mathcal{C}_0^0(\mathbb{R}^n)$ and $\psi \in \mathcal{S}(\mathbb{R}^n)$ then*

$$(8.14) \quad v * \psi \in \mathcal{C}_0^\infty(\mathbb{R}^n) = \bigcap_{k \geq 0} \mathcal{C}_0^k(\mathbb{R}^n).$$

Proof. This follows from (8.12), (8.13) and induction. \square

Now, let us make a more special choice of ψ . We have shown the existence of

$$(8.15) \quad \varphi \in \mathcal{C}_c^\infty(\mathbb{R}^n), \varphi \geq 0, \text{supp}(\varphi) \subset \{|x| \leq 1\}.$$

We can also assume $\int_{\mathbb{R}^n} \varphi dx = 1$, by multiplying by a positive constant. Now consider

$$(8.16) \quad \varphi_t(x) = t^{-n} \varphi\left(\frac{x}{t}\right) \quad 1 \geq t > 0.$$

This has all the same properties, except that

$$(8.17) \quad \text{supp } \varphi_t \subset \{|x| \leq t\}, \quad \int \varphi_t dx = 1.$$

Proposition 8.2. *If $v \in \mathcal{C}_0^0(\mathbb{R}^n)$ then as $t \rightarrow 0$, $v_t = v * \varphi_t \rightarrow v$ in $\mathcal{C}_0^0(\mathbb{R}^n)$.*

Proof. using (8.17) we can write the difference as

$$(8.18) \quad |v_t(x) - v(x)| = \left| \int_{\mathbb{R}^n} (v(x-y) - v(x)) \varphi_t(y) dy \right| \\ \leq \sup_{|y| \leq t} |v(x-y) - v(x)| \rightarrow 0.$$

Here we have used the fact that $\varphi_t \geq 0$ has support in $|y| \leq t$ and has integral 1. Thus $v_t \rightarrow v$ uniformly on any set on which v is uniformly continuous, namely \mathbb{R}^n ! \square

Corollary 8.3. $\mathcal{C}_0^k(\mathbb{R}^n)$ is dense in $\mathcal{C}_0^p(\mathbb{R}^n)$ for any $k \geq p$.

Proposition 8.4. $\mathcal{S}(\mathbb{R}^n)$ is dense in $\mathcal{C}_0^k(\mathbb{R}^n)$ for any $k \geq 0$.

Proof. Take $k = 0$ first. The subspace $\mathcal{C}_c^0(\mathbb{R}^n)$ is dense in $\mathcal{C}_0^0(\mathbb{R}^n)$, by cutting off outside a large ball. If $v \in \mathcal{C}_c^0(\mathbb{R}^n)$ has support in $\{|x| \leq R\}$ then

$$v * \varphi_t \in \mathcal{C}_c^\infty(\mathbb{R}^n) \subset \mathcal{S}(\mathbb{R}^n)$$

has support in $\{|x| \leq R + 1\}$. Since $v * \varphi_t \rightarrow v$ the result follows for $k = 0$.

For $k \geq 1$ the same argument works, since $D^\alpha(v * \varphi_t) = (D^\alpha v) * \varphi_t$. \square

Corollary 8.5. *The map from finite Radon measures*

$$(8.19) \quad M_{fn}(\mathbb{R}^n) \ni \mu \longmapsto u_\mu \in \mathcal{S}'(\mathbb{R}^n)$$

is injective.

Now, we want the same result for $L^2(\mathbb{R}^n)$ (and maybe for $L^p(\mathbb{R}^n)$, $1 \leq p < \infty$). I leave the measure-theoretic part of the argument to you.

Proposition 8.6. *Elements of $L^2(\mathbb{R}^n)$ are “continuous in the mean” i.e.,*

$$(8.20) \quad \lim_{|t| \rightarrow 0} \int_{\mathbb{R}^n} |u(x+t) - u(x)|^2 dx = 0.$$

This is Problem 24.

Using this we conclude that

$$(8.21) \quad \mathcal{S}(\mathbb{R}^n) \hookrightarrow L^2(\mathbb{R}^n) \text{ is dense}$$

as before. First observe that the space of L^2 functions of compact support is dense in $L^2(\mathbb{R}^n)$, since

$$\lim_{R \rightarrow \infty} \int_{|x| \geq R} |u(x)|^2 dx = 0 \quad \forall u \in L^2(\mathbb{R}^n).$$

Then look back at the discussion of $v * \varphi$, now v is replaced by $u \in L_c^2(\mathbb{R}^n)$. The compactness of the support means that $u \in L^1(\mathbb{R}^n)$ so in

$$(8.22) \quad u * \varphi(x) = \int_{\mathbb{R}^n} u(x-y)\varphi(y)dy$$

the integral is absolutely convergent. Moreover

$$\begin{aligned} & |u * \varphi(x+x') - u * \varphi(x)| \\ &= \left| \int u(y)(\varphi(x+x'-y) - \varphi(x-y)) dy \right| \\ &\leq C \|u\| \sup_{|y| \leq R} |\varphi(x+x'-y) - \varphi(x-y)| \rightarrow 0 \end{aligned}$$

when $\{|x| \leq R\}$ large enough. Thus $u * \varphi$ is continuous and the same argument as before shows that

$$u * \varphi_t \in \mathcal{S}(\mathbb{R}^n).$$

Now to see that $u * \varphi_t \rightarrow u$, assuming u has compact support (or not) we estimate the integral

$$\begin{aligned} |u * \varphi_t(x) - u(x)| &= \left| \int (u(x-y) - u(x)) \varphi_t(y) dy \right| \\ &\leq \int |u(x-y) - u(x)| \varphi_t(y) dy. \end{aligned}$$

Using the same argument twice

$$\begin{aligned} &\int |u * \varphi_t(x) - u(x)|^2 dx \\ &\leq \iiint |u(x-y) - u(x)| \varphi_t(y) |u(x-y') - u(x)| \varphi_t(y') dx dy dy' \\ &\leq \left(\int |u(x-y) - u(x)|^2 \varphi_t(y) \varphi_t(y') dx dy dy' \right) \\ &\leq \sup_{|y| \leq t} \int |u(x-y) - u(x)|^2 dx. \end{aligned}$$

Note that at the second step here I have used Schwarz's inequality with the integrand written as the product

$$|u(x-y) - u(x)| \varphi_t^{1/2}(y) \varphi_t^{1/2}(y') \cdot |u(x-y') - u(x)| \varphi_t^{1/2}(y) \varphi_t^{1/2}(y').$$

Thus we now know that

$$L^2(\mathbb{R}^n) \hookrightarrow \mathcal{S}'(\mathbb{R}^n) \text{ is injective.}$$

This means that all our usual spaces of functions 'sit inside' $\mathcal{S}'(\mathbb{R}^n)$.

Finally we can use convolution with φ_t to show the existence of *smooth* partitions of unity. If $K \Subset U \subset \mathbb{R}^n$ is a compact set in an open set then we have shown the existence of $\xi \in \mathcal{C}_c^0(\mathbb{R}^n)$, with $\xi = 1$ in some neighborhood of K and $\xi = 0$ in some neighborhood of K^c and $\text{supp}(\xi) \Subset U$.

Then consider $\xi * \varphi_t$ for t small. In fact

$$\text{supp}(\xi * \varphi_t) \subset \{p \in \mathbb{R}^n; \text{dist}(p, \text{supp} \xi) \leq 2t\}$$

and similarly, $0 \leq \xi * \varphi_t \leq 1$ and

$$\xi * \varphi_t = 1 \text{ at } p \text{ if } \xi = 1 \text{ on } B(p, 2t).$$

Using this we get:

Proposition 8.7. *If $U_a \subset \mathbb{R}^n$ are open for $a \in A$ and $K \Subset \bigcup_{a \in A} U_a$ then there exist finitely many $\varphi_i \in \mathcal{C}_c^\infty(\mathbb{R}^n)$, with $0 \leq \varphi_i \leq 1$, $\text{supp}(\varphi_i) \subset U_{a_i}$ such that $\sum_i \varphi_i = 1$ in a neighbourhood of K .*

Proof. By the compactness of K we may choose a finite open subcover. Using Lemma 1.8 we may choose a continuous partition, ϕ'_i , of unity subordinate to this cover. Using the convolution argument above we can replace ϕ'_i by $\phi'_i * \varphi_t$ for $t > 0$. If t is sufficiently small then this is again a partition of unity subordinate to the cover, but now smooth. \square

Next we can make a simple ‘cut off argument’ to show

Lemma 8.8. *The space $\mathcal{C}_c^\infty(\mathbb{R}^n)$ of \mathcal{C}^∞ functions of compact support is dense in $\mathcal{S}(\mathbb{R}^n)$.*

Proof. Choose $\varphi \in \mathcal{C}_c^\infty(\mathbb{R}^n)$ with $\varphi(x) = 1$ in $|x| \leq 1$. Then given $\psi \in \mathcal{S}(\mathbb{R}^n)$ consider the sequence

$$\psi_n(x) = \varphi(x/n)\psi(x).$$

Clearly $\psi_n = \psi$ on $|x| \leq n$, so if it converges in $\mathcal{S}(\mathbb{R}^n)$ it must converge to ψ . Suppose $m \geq n$ then by Leibniz’s formula¹³

$$\begin{aligned} D_x^\alpha(\psi_n(x) - \psi_m(x)) &= \sum_{\beta \leq \alpha} \binom{\alpha}{\beta} D_x^\beta \left(\varphi\left(\frac{x}{n}\right) - \varphi\left(\frac{x}{m}\right) \right) \cdot D_x^{\alpha-\beta} \psi(x). \end{aligned}$$

All derivatives of $\varphi(x/n)$ are bounded, independent of n and $\psi_n = \psi_m$ in $|x| \leq n$ so for any p

$$|D_x^\alpha(\psi_n(x) - \psi_m(x))| \leq \begin{cases} 0 & |x| \leq n \\ C_{\alpha,p} \langle x \rangle^{-2p} & |x| \geq n \end{cases}.$$

Hence ψ_n is Cauchy in $\mathcal{S}(\mathbb{R}^n)$. \square

Thus every element of $\mathcal{S}'(\mathbb{R}^n)$ is determined by its restriction to $\mathcal{C}_c^\infty(\mathbb{R}^n)$. The support of a tempered distribution was defined above to be

$$(8.23) \quad \text{supp}(u) = \{x \in \mathbb{R}^n; \exists \varphi \in \mathcal{S}(\mathbb{R}^n), \varphi(x) \neq 0, \varphi u = 0\}^c.$$

Using the preceding lemma and the construction of smooth partitions of unity we find

Proposition 8.9. *$f u \in \mathcal{S}'(\mathbb{R}^n)$ and $\text{supp}(u) = \emptyset$ then $u = 0$.*

¹³Problem 25.

Proof. From (8.23), if $\psi \in \mathcal{S}(\mathbb{R}^n)$, $\text{supp}(\psi u) \subset \text{supp}(u)$. If $x \in \text{supp}(u)$ then, by definition, $\varphi u = 0$ for some $\varphi \in \mathcal{S}(\mathbb{R}^n)$ with $\varphi(x) \neq 0$. Thus $\varphi \neq 0$ on $B(x, \epsilon)$ for $\epsilon > 0$ sufficiently small. If $\psi \in \mathcal{C}_c^\infty(\mathbb{R}^n)$ has support in $B(x, \epsilon)$ then $\psi u = \tilde{\psi} \varphi u = 0$, where $\tilde{\psi} \in \mathcal{C}_c^\infty(\mathbb{R}^n)$:

$$\tilde{\psi} = \begin{cases} \psi/\varphi & \text{in } B(x, \epsilon) \\ 0 & \text{elsewhere.} \end{cases}$$

Thus, given $K \Subset \mathbb{R}^n$ we can find $\varphi_j \in \mathcal{C}_c^\infty(\mathbb{R}^n)$, supported in such balls, so that $\sum_j \varphi_j \equiv 1$ on K but $\varphi_j u = 0$. For given $\mu \in \mathcal{C}_c^\infty(\mathbb{R}^n)$ apply this to $\text{supp}(\mu)$. Then

$$\mu = \sum_j \varphi_j \mu \Rightarrow u(\mu) = \sum_j (\varphi_j u)(\mu) = 0.$$

Thus $u = 0$ on $\mathcal{C}_c^\infty(\mathbb{R}^n)$, so $u = 0$. \square

The linear space of distributions of compact support will be denoted $\mathcal{C}_c^{-\infty}(\mathbb{R}^n)$; it is often written $\mathcal{E}'(\mathbb{R}^n)$.

Now let us give a characterization of the ‘delta function’

$$\delta(\varphi) = \varphi(0) \quad \forall \varphi \in \mathcal{S}(\mathbb{R}^n),$$

or at least the one-dimensional subspace of $\mathcal{S}'(\mathbb{R}^n)$ it spans. This is based on the simple observation that $(x_j \varphi)(0) = 0$ if $\varphi \in \mathcal{S}(\mathbb{R}^n)$!

Proposition 8.10. *If $u \in \mathcal{S}'(\mathbb{R}^n)$ satisfies $x_j u = 0$, $j = 1, \dots, n$ then $u = c\delta$.*

Proof. The main work is in characterizing the null space of δ as a linear functional, namely in showing that

$$(8.24) \quad \mathcal{H} = \{\varphi \in \mathcal{S}(\mathbb{R}^n); \varphi(0) = 0\}$$

can also be written as

$$(8.25) \quad \mathcal{H} = \left\{ \varphi \in \mathcal{S}(\mathbb{R}^n); \varphi = \sum_{j=1}^n x_j \psi_j, \psi_j \in \mathcal{S}(\mathbb{R}^n) \right\}.$$

Clearly the right side of (8.25) is contained in the left. To see the converse, suppose first that

$$(8.26) \quad \varphi \in \mathcal{S}(\mathbb{R}^n), \quad \varphi = 0 \text{ in } |x| < 1.$$

Then define

$$\psi = \begin{cases} 0 & |x| < 1 \\ \varphi/|x|^2 & |x| \geq 1. \end{cases}$$

All the derivatives of $1/|x|^2$ are bounded in $|x| \geq 1$, so from Leibniz's formula it follows that $\psi \in \mathcal{S}(\mathbb{R}^n)$. Since

$$\varphi = \sum_j x_j (x_j \psi)$$

this shows that φ of the form (8.26) is in the right side of (8.25). In general suppose $\varphi \in \mathcal{S}(\mathbb{R}^n)$. Then

$$(8.27) \quad \begin{aligned} \varphi(x) - \varphi(0) &= \int_0^1 \frac{d}{dt} \varphi(tx) dt \\ &= \sum_{j=1}^n x_j \int_0^1 \frac{\partial \varphi}{\partial x_j}(tx) dt. \end{aligned}$$

Certainly these integrals are \mathcal{C}^∞ , but they may not decay rapidly at infinity. However, choose $\mu \in \mathcal{C}_c^\infty(\mathbb{R}^n)$ with $\mu = 1$ in $|x| \leq 1$. Then (8.27) becomes, if $\varphi(0) = 0$,

$$\begin{aligned} \varphi &= \mu \varphi + (1 - \mu) \varphi \\ &= \sum_{j=1}^n x_j \psi_j + (1 - \mu) \varphi, \quad \psi_j = \mu \int_0^1 \frac{\partial \varphi}{\partial x_j}(tx) dt \in \mathcal{S}(\mathbb{R}^n). \end{aligned}$$

Since $(1 - \mu)\varphi$ is of the form (8.26), this proves (8.25).

Our assumption on u is that $x_j u = 0$, thus

$$u(\varphi) = 0 \quad \forall \varphi \in \mathcal{H}$$

by (8.25). Choosing μ as above, a general $\varphi \in \mathcal{S}(\mathbb{R}^n)$ can be written

$$\varphi = \varphi(0) \cdot \mu + \varphi', \quad \varphi' \in \mathcal{H}.$$

Then

$$u(\varphi) = \varphi(0)u(\mu) \Rightarrow u = c\delta, \quad c = u(\mu).$$

□

This result is quite powerful, as we shall soon see. The Fourier transform of an element $\varphi \in \mathcal{S}(\mathbb{R}^n)$ is¹⁴

$$(8.28) \quad \hat{\varphi}(\xi) = \int e^{-ix \cdot \xi} \varphi(x) dx, \quad \xi \in \mathbb{R}^n.$$

¹⁴Normalizations vary, but it doesn't matter much.

The integral certainly converges, since $|\varphi| \leq C\langle x \rangle^{-n-1}$. In fact it follows easily that $\hat{\varphi}$ is continuous, since

$$\begin{aligned} |\hat{\varphi}(\xi) - \hat{\varphi}(\xi')| &\in \int \left| e^{ix-\xi} - e^{-ix-\xi'} \right| |\varphi| dx \\ &\rightarrow 0 \text{ as } \xi' \rightarrow \xi. \end{aligned}$$

In fact

Proposition 8.11. *Fourier transformation, (8.28), defines a continuous linear map*

$$(8.29) \quad \mathcal{F} : \mathcal{S}(\mathbb{R}^n) \rightarrow \mathcal{S}(\mathbb{R}^n), \mathcal{F}\varphi = \hat{\varphi}.$$

Proof. Differentiating under the integral¹⁵ sign shows that

$$\partial_{\xi_j} \hat{\varphi}(\xi) = -i \int e^{-ix \cdot \xi} x_j \varphi(x) dx.$$

Since the integral on the right is absolutely convergent that shows that (remember the i 's)

$$(8.30) \quad D_{\xi_j} \hat{\varphi} = -\widehat{x_j \varphi}, \quad \forall \varphi \in \mathcal{S}(\mathbb{R}^n).$$

Similarly, if we multiply by ξ_j and observe that $\xi_j e^{-ix \cdot \xi} = i \frac{\partial}{\partial x_j} e^{-ix \cdot \xi}$ then integration by parts shows

$$\begin{aligned} (8.31) \quad \xi_j \hat{\varphi} &= i \int \left(\frac{\partial}{\partial x_j} e^{-ix \cdot \xi} \right) \varphi(x) dx \\ &= -i \int e^{-ix \cdot \xi} \frac{\partial \varphi}{\partial x_j} dx \\ \widehat{D_j \varphi} &= \xi_j \hat{\varphi}, \quad \forall \varphi \in \mathcal{S}(\mathbb{R}^n). \end{aligned}$$

Since $x_j \varphi, D_j \varphi \in \mathcal{S}(\mathbb{R}^n)$ these results can be iterated, showing that

$$(8.32) \quad \xi^\alpha D_\xi^\beta \hat{\varphi} = \mathcal{F} \left((-1)^{|\beta|} D_x^\alpha x^\beta \varphi \right).$$

Thus $\left| \xi^\alpha D_\xi^\beta \hat{\varphi} \right| \leq C_{\alpha\beta} \sup | \langle x \rangle^{+n+1} D_x^\alpha x^\beta \varphi | \leq C \| \langle x \rangle^{n+1+|\beta|} \varphi \|_{C^{|\alpha|}}$, which shows that \mathcal{F} is continuous as a map (8.32). □

Suppose $\varphi \in \mathcal{S}(\mathbb{R}^n)$. Since $\hat{\varphi} \in \mathcal{S}(\mathbb{R}^n)$ we can consider the distribution $u \in \mathcal{S}'(\mathbb{R}^n)$

$$(8.33) \quad u(\varphi) = \int_{\mathbb{R}^n} \hat{\varphi}(\xi) d\xi.$$

¹⁵See [5]

The continuity of u follows from the fact that integration is continuous and (8.29). Now observe that

$$\begin{aligned} u(x_j\varphi) &= \int_{\mathbb{R}^n} \widehat{x_j\varphi}(\xi) d\xi \\ &= - \int_{\mathbb{R}^n} D_{\xi_j} \hat{\varphi} d\xi = 0 \end{aligned}$$

where we use (8.30). Applying Proposition 8.10 we conclude that $u = c\delta$ for some (universal) constant c . By definition this means

$$(8.34) \quad \int_{\mathbb{R}^n} \hat{\varphi}(\xi) d\xi = c\varphi(0).$$

So what is the constant? To find it we need to work out an example. The simplest one is

$$\varphi = \exp(-|x|^2/2).$$

Lemma 8.12. *The Fourier transform of the Gaussian $\exp(-|x|^2/2)$ is the Gaussian $(2\pi)^{n/2} \exp(-|\xi|^2/2)$.*

Proof. There are two obvious methods — one uses complex analysis (Cauchy's theorem) the other, which I shall follow, uses the uniqueness of solutions to ordinary differential equations.

First observe that $\exp(-|x|^2/2) = \prod_j \exp(-x_j^2/2)$. Thus¹⁶

$$\hat{\varphi}(\xi) = \prod_{j=1}^n \hat{\psi}(\xi_j), \quad \psi(x) = e^{-x^2/2},$$

being a function of one variable. Now ψ satisfies the differential equation

$$(\partial_x + x)\psi = 0,$$

and is the *only* solution of this equation up to a constant multiple. By (8.30) and (8.31) its Fourier transform satisfies

$$\widehat{\partial_x \psi} + \widehat{x\psi} = i\xi \hat{\psi} + i \frac{d}{d\xi} \hat{\varphi} = 0.$$

This is the same equation, but in the ξ variable. Thus $\hat{\psi} = ce^{-|\xi|^2/2}$. Again we need to find the constant. However,

$$\hat{\psi}(0) = c = \int e^{-x^2/2} dx = (2\pi)^{1/2}$$

¹⁶Really by Fubini's theorem, but here one can use Riemann integrals.

by the standard use of polar coordinates:

$$c^2 = \int_{\mathbb{R}^n} e^{-(x^2+y^2)/2} dx dy = \int_0^\infty \int_0^{2\pi} e^{-r^2/2} r dr d\theta = 2\pi.$$

This proves the lemma. □

Thus we have shown that for any $\varphi \in \mathcal{S}(\mathbb{R}^n)$

$$(8.35) \quad \int_{\mathbb{R}^n} \hat{\varphi}(\xi) d\xi = (2\pi)^n \varphi(0).$$

Since this is true for $\varphi = \exp(-|x|^2/2)$. The identity allows us to *invert* the Fourier transform.