

## REFERENCES

- [1] M. F. Atiyah, V. K. Patodi, and I. M. Singer, *Spectral asymmetry and Riemannian geometry. III*, Math. Proc. Cambridge Philos. Soc. **79** (1976), no. 1, 71–99. MR 53 #1655c
- [2] J.-L. Brylinski, *Loop spaces, characteristic classes and geometric quantization*, Progress in Mathematics, vol. 107, Birkhäuser Boston, Inc., Boston, MA, 1993.
- [3] V.W. Guillemin, *A new proof of Weyl's formula on the asymptotic distribution of eigenvalues*, Adv. Math. **55** (1985), 131–160.
- [4] L. Hörmander, *The Weyl calculus of pseudo-differential operators*, Comm. Pure Appl. Math. **32** (1979), 359–443.
- [5] V. Mathai, R.B. Melrose, and I.M. Singer, *The index of projective families of elliptic operators: The decomposable case*, Arxiv:0809.0028.
- [6] Eckhard Meinrenken, *The basic gerbe over a compact simple lie group*, L'Enseignement Mathématique **49** (2003), 307–333.
- [7] Richard B. Melrose, *The eta invariant and families of pseudodifferential operators*, Math. Res. Lett. **2** (1995), no. 5, 541–561. MR 96h:58169
- [8] M.K. Murray, *Bundle gerbes*, J. London Math. Soc. **54** (1996), 403–416.
- [9] A. Pressley and G. Segal, *Loop groups*, Oxford Mathematical Monographs, The Clarendon Press, Oxford University Press, New York, 1986.
- [10] D. Quillen, *Paper on determinant*, Funct. Anal. and its Appl. **19** (1985), 37–41.
- [11] R. T. Seeley, *Complex powers of an elliptic operator*, Singular Integrals (Proc. Sympos. Pure Math., Chicago, Ill., 1966), Amer. Math. Soc., Providence, R.I., 1967, pp. 288–307. MR 38 #6220
- [12] M. Wodzicki, *Noncommutative residue. I. Fundamentals*, *K*-theory, arithmetic and geometry (Moscow, 1984–1986), Lecture Notes in Math., Springer, Berlin-New York, 1987, pp. 320–399.

MIT, MSRI AND CHANCELLOR'S PROFESSOR AT U.C. BERKELEY, FALL 2008

*E-mail address:* `rbm@math.berkeley.edu`