

18.100B, FALL 2002
FINAL EXAM: DECEMBER 16

Try each of the questions; the first seven are worth 12 points each, the last one is harder and is worth 16. You may use theorems from class, or the book, provided you can recall them correctly! This includes standard properties of the exponential and trigonometric functions. No books or papers are permitted.

PROBLEM 1

Show that the set $\{z \in \mathbb{C}; z = \exp(it^3) \text{ for some } t \in \mathbb{R}\}$ is connected.

PROBLEM 2

Let $\{x_n\}$ be a sequence in a metric space X and suppose that there is a point $p \in X$ with the property that every subsequence of $\{x_n\}$ has a subsequence which converges to p . Show that $\{x_n\}$ converges to p .

PROBLEM 3

- (1) Why is the function $f(x) = \exp(\frac{x^3-15}{x^2+x+1})$ continuously differentiable on $[0, 1]$?
- (2) Why does f have a minimum value on this interval?

PROBLEM 4

Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be twice differentiable and suppose that 0 is a local maximum of f , i.e. for some $\epsilon > 0$ $f(x) \leq f(0)$ for all $x \in (-\epsilon, \epsilon)$. Show that $f''(0) \leq 0$.

PROBLEM 5

If $g : \mathbb{R} \rightarrow \mathbb{R}$ is a differentiable function and g' is bounded on \mathbb{R} , show that g is uniformly continuous.

PROBLEM 6

Using standard properties of the cosine function explain why the formula

$$f(x) = \sum_{n=0}^{\infty} \frac{1}{n^3} \cos(nx)$$

defines a continuously differentiable function on the real line.

PROBLEM 7

Explain carefully why the Riemann-Stieltjes integral

$$\int_0^1 \exp(3x^2) d\alpha$$

exists for any increasing function $\alpha : [0, 1] \rightarrow \mathbb{R}$.

PROBLEM 8

Let $A : [0, 1]^2 \rightarrow \mathbb{R}$ be a continuous function satisfying

$$\sup_{[0,1]^2} |A(x, y)| \leq \frac{1}{2}.$$

(1) Show that if $f \in \mathcal{C}([0, 1])$ then

$$g(x) = \int_0^1 A(x, y)f(y)dy \in \mathcal{C}([0, 1]).$$

(2) Estimate $\|g\| = \sup_{[0,1]} |g(x)|$ in terms of $\|f\|$.

(3) If $h \in \mathcal{C}([0, 1])$ is a fixed function show that

$$(Gf)(x) = h(x) + \int_0^1 A(x, y)f(y)dy$$

defines a contraction G on $\mathcal{C}([0, 1])$ sending f to Gf .

(4) Show that there exists a unique $f \in \mathcal{C}([0, 1])$ such that

$$f(x) = h(x) + \int_0^1 A(x, y)f(y)dy \quad \forall x \in [0, 1].$$