

The Local Langlands Correspondence for tamely ramified groups

David Roe

Department of Mathematics
Harvard University

Quebec-Maine 2010

Outline

- 1 Local Langlands for GL_n
 - GL_1
 - GL_n
- 2 Beyond GL_n
 - Statements
 - Construction

What is the Langlands Correspondence?

- A generalization of class field theory to non-abelian extensions.
- A tool for studying L-functions.
- A correspondence between representations of Galois groups and representations of algebraic groups.

The Global Correspondence

Irreducible n -dim'l complex representations of $\text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$



Automorphic representations of $GL_n(\mathbb{A})$

The local-global principal suggests that we try to break this down to local pieces.

$$\text{Gal}(\bar{\mathbb{Q}}_p/\mathbb{Q}_p) \subset \text{Gal}(\bar{\mathbb{Q}}/\mathbb{Q})$$

$$GL_n(\mathbb{A}) = \prod'_v GL_n(\mathbb{Q}_v)$$

Local Class Field Theory

The one dimensional case of local Langlands is local class field theory.

Irreducible representations of $\text{Gal}(\bar{\mathbb{Q}}_p/\mathbb{Q}_p)^{\text{ab}}$



Irreducible representations of \mathbb{Q}_p^\times

The Weil Group

Write $G_{\mathbb{Q}_p} = \text{Gal}(\bar{\mathbb{Q}}_p/\mathbb{Q}_p)$ and recall $G_{\mathbb{F}_p} \cong \hat{\mathbb{Z}}$, topologically generated by Frobenius.

$$\begin{array}{ccccccc}
 1 & \longrightarrow & I_p & \longrightarrow & G_{\mathbb{Q}_p} & \longrightarrow & \hat{\mathbb{Z}} \longrightarrow 1 \\
 & & \parallel & & \uparrow & & \uparrow \\
 1 & \longrightarrow & I_p & \longrightarrow & W_{\mathbb{Q}_p} & \longrightarrow & \mathbb{Z} \longrightarrow 1
 \end{array}$$

$W_{\mathbb{Q}_p}$ is dense in $G_{\mathbb{Q}_p}$ so $\{\text{irreps of } G_{\mathbb{Q}_p}\} \hookrightarrow \{\text{irreps of } W_{\mathbb{Q}_p}\}$.

Theorem (Local Class Field Theory)

The artin map $\mathbb{Q}_p^\times \rightarrow W_{\mathbb{Q}_p}^{\text{ab}}$ is an isomorphism and induces isomorphisms $\widehat{\mathbb{Q}_p^\times} \rightarrow G_{\mathbb{Q}_p}^{\text{ab}}$ and $\mathbb{Z}_p^\times \rightarrow I_p^{\text{ab}}$.

Corollary

Irreducible 1-dimensional representations of $\mathcal{W}_{\mathbb{Q}_p}$



Irreducible representations of $GL_1(\mathbb{Q}_p)$

Conjecture

Irreducible n -dimensional representations of $\mathcal{W}_{\mathbb{Q}_p}$



Irreducible representations of $GL_n(\mathbb{Q}_p)$

Smooth Representations

For $n > 1$, the representations of $GL_n(\mathbb{Q}_p)$ that appear are usually infinite dimensional.

Definition

A *smooth \mathbb{C} -representation* of $GL_n(\mathbb{Q}_p)$ is a pair (π, V) , where

- V is a \mathbb{C} -vector space (possibly infinite dimensional),
- $\pi: GL_n(\mathbb{Q}_p) \rightarrow GL(V)$ is a homomorphism,
- The stabilizer of each $v \in V$ is open in $GL_n(\mathbb{Q}_p)$.

The only finite-dimensional irreducible smooth π are

$$g \mapsto \chi(\det(g))$$

for some character $\chi: \mathbb{Q}_p^\times \rightarrow \mathbb{C}^\times$.

Langlands Parameters

We also need to clarify what kinds of representations of $\mathcal{W}_{\mathbb{Q}_p}$ to focus on.

Definition

A *Langlands parameter* is a pair (φ, V) with

$$\varphi: \mathcal{W}_{\mathbb{Q}_p} \rightarrow \mathrm{GL}(V) \quad \dim_{\mathbb{C}} V = n$$

such that φ is continuous and semisimple.

Parabolic Subgroups

Given a number of Langlands parameters $\varphi_i: \mathbf{W}_{\mathbb{Q}_p} \rightarrow GL(V_i)$, one can form their direct sum. There should be a corresponding operation on the $GL_n(\mathbb{Q}_p)$ side.

Definition

A *parabolic subgroup* of GL_n is a subgroup P conjugate to one consisting of block triangular matrices of a given pattern. For example:

$$\begin{pmatrix} * & * & * & * & * \\ 0 & * & * & * & * \\ 0 & * & * & * & * \\ 0 & 0 & 0 & * & * \\ 0 & 0 & 0 & * & * \end{pmatrix}$$

Such a subgroup has a Levi decomposition $P = M \ltimes N$, where M is conjugate to the corresponding subgroup of block diagonal matrices, and N consists of the subgroup of P with identity blocks on the diagonal.

Parabolic Induction

Since each Levi subgroup M is just a direct product of GL_{n_i} , a collection of representations $\pi_i: GL_{n_i}(\mathbb{Q}_p) \rightarrow GL(V_i)$ yields a representation $\boxtimes_i \pi_i$ of M . We can pull this back to P and then induce to obtain

$$\pi = \text{Ind}_P^{GL_n(\mathbb{Q}_p)} \boxtimes_i \pi_i.$$

Definition

We say that π is the *parabolic induction* of the π_i . We say that π is *supercuspidal* if π is not parabolically induced from any proper parabolic subgroup of $GL_n(\mathbb{Q}_p)$.

The Weil-Deligne Group

There is a natural bijection

Supercuspidal
representations of $GL_n(\mathbb{Q}_p)$

\leftrightarrow

n -dimensional irreducible
representations of $\mathcal{W}_{\mathbb{Q}_p}$.

But the parabolic induction of irreducible representations does not always remain irreducible. To extend this bijection from supercuspidal representations of $GL_n(\mathbb{Q}_p)$ to all smooth irreducible representations of $GL_n(\mathbb{Q}_p)$, one enlarges the right hand side using the following group:

$$WD_{\mathbb{Q}_p} := \mathcal{W}_{\mathbb{Q}_p} \times SL_2(\mathbb{C}).$$

Local Langlands for GL_n

Theorem (Harris-Taylor, Henniart)

There is a unique system of bijections

*Irreducible representations
of $GL_n(\mathbb{Q}_p)$*

$\xrightarrow{\text{rec}_n}$

*n -dimensional
representations of $WD_{\mathbb{Q}_p}$*

- rec_1 is induced by the Artin map of local class field theory.
- rec_n is compatible with 1-dimensional characters:
 $\text{rec}_n(\pi \otimes \chi \circ \det) = \text{rec}_n(\pi) \otimes \text{rec}_1(\chi)$.
- The central character ω_π of π corresponds to $\det \circ \text{rec}_n$:
 $\text{rec}_1(\omega_\pi) = \det(\text{rec}_n(\pi))$.
- $\text{rec}_n(\pi^\vee) = \text{rec}_n(\pi)^\vee$
- rec_n respects natural invariants associated to each side, namely L -factors and ϵ -factors of pairs.

A First Guess

Now suppose \mathbf{G} is some other connected reductive group defined over \mathbb{Q}_p , such as SO_n , Sp_n or U_n . We'd like to use a Langlands correspondence to understand representations of $\mathbf{G}(\mathbb{Q}_p)$ in terms of Galois representations. Something like

Homomorphisms
 $\varphi: \mathrm{WD}_{\mathbb{Q}_p} \rightarrow \mathbf{G}(\mathbb{C})$

\leftrightarrow

Irreducible representations
of $\mathbf{G}(\mathbb{Q}_p)$.

We need to modify this guess in three ways:

- change $\mathbf{G}(\mathbb{C})$ to a related group, ${}^L\mathbf{G}(\mathbb{C})$,
- account for the fact that our correspondence is no longer a bijection,
- account for the fact that our representations are no longer all of $\mathbf{G}(\mathbb{Q}_p)$, but instead of pure inner forms of $\mathbf{G}(\mathbb{Q}_p)$.

Root Data

Reductive groups over algebraically closed fields are classified by root data

$$(X^*(\mathbf{S}), \Phi(\mathbf{G}, \mathbf{S}), X_*(\mathbf{S}), \Phi^\vee(\mathbf{G}, \mathbf{S})),$$

where

- $\mathbf{S} \subset \mathbf{G}$ is a maximal torus,
- $X^*(\mathbf{S})$ is the lattice of characters $\chi: \mathbf{S} \rightarrow \mathbb{G}_m$,
- $X_*(\mathbf{S})$ is the lattice of cocharacters $\lambda: \mathbb{G}_m \rightarrow \mathbf{S}$,
- $\Phi(\mathbf{G}, \mathbf{S})$ is the set of roots (eigenvalues of the adjoint action of \mathbf{S} on \mathfrak{g}),
- $\Phi^\vee(\mathbf{G}, \mathbf{S})$ is the set of coroots ($\langle \alpha, \alpha^\vee \rangle = 2$).

Connected Langlands Dual

Given $\mathbf{G} \supset \mathbf{S}$, the connected Langlands dual group $\hat{\mathbf{G}}$ is defined to be the algebraic group over \mathbb{C} with root datum

$$(X_*(\mathbf{S}), \Phi^\vee(\mathbf{G}, \mathbf{S}), X^*(\mathbf{S}), \Phi(\mathbf{G}, \mathbf{S})).$$

For semisimple groups, this has the effect of exchanging the long and short roots (as well as interchanging the simply connected and adjoint forms).

\mathbf{G}	GL_n	SL_n	PGL_n	Sp_{2n}	SO_{2n}	U_n
$\hat{\mathbf{G}}$	GL_n	PGL_n	SL_n	SO_{2n+1}	SO_{2n}	GL_n

Langlands Dual Group

For non-split \mathbf{G} , such as U_n , we need to work a little harder. Suppose that \mathbf{G} is quasi-split with Borel $\mathbf{B} \supset \mathbf{S}$, splitting over a finite extension E/\mathbb{Q}_p . Then $\text{Gal}(E/\mathbb{Q}_p)$ acts on the root datum, and we get an action on $\hat{\mathbf{G}}$ via pinned automorphisms. Define

$${}^L\mathbf{G} := \hat{\mathbf{G}} \rtimes \text{Gal}(E/\mathbb{Q}_p).$$

Unitary Groups

- E/\mathbb{Q}_p a quadratic extension (so for $p \neq 2$ there are three possibilities),
- $\tau \in \text{Gal}(E/\mathbb{Q}_p)$ the nontrivial element,
- V an n -dimensional E -vector space,
- Non-degenerate Hermitian form \langle, \rangle (so $\langle x, y \rangle = \tau \langle y, x \rangle$).

Then $U(V)$ is the group of automorphisms of V preserving \langle, \rangle . Over $\bar{\mathbb{Q}}_p$, U becomes isomorphic to GL_n , so \hat{U}_n is GL_n , but ${}^L\mathbf{G}$ is non-connected: τ acts on $GL_n(\mathbb{C})$ by the outer automorphism

$$g \mapsto (g^{-1})^T.$$

Pure Inner Forms

- Pure inner forms in general are parameterized by $H^1(\mathbb{Q}_p, \mathbf{G})$ (as opposed to forms, which are parameterized by $H^1(\mathbb{Q}_p, \text{Aut}(\mathbf{G}))$ or inner forms, by $H^1(\mathbb{Q}_p, \mathbf{G}_{\text{ad}})$). For unitary groups for the same quadratic field E/\mathbb{Q}_p , there are two: one for each isomorphism class of Hermitian space of dimension n .
- Pure inner forms have the same Langlands dual group ${}^L\mathbf{G}$, thus entangling the representations corresponding to a given Langlands parameter.
- Every \mathbf{G} over \mathbb{Q}_p has a quasi-split pure inner form, so we assume from now on that \mathbf{G} is quasi-split.

Langlands Parameters

A Langlands parameter is now an equivalence class of homomorphisms

$$\varphi: \mathrm{WD}_{\mathbb{Q}_p} \rightarrow {}^L\mathbf{G}.$$

- We require that the composition of φ with the projection ${}^L\mathbf{G} \rightarrow \mathrm{Gal}(E/\mathbb{Q}_p)$ agrees with the standard projection $\mathcal{W}_{\mathbb{Q}_p} \rightarrow \mathrm{Gal}(E/\mathbb{Q}_p)$.
- We consider two parameters to be equivalent if they are conjugate by an element of $\hat{\mathbf{G}}$. This definition of equivalence is chosen to match up with the notion of isomorphic representations on the $\mathbf{G}(\mathbb{Q}_p)$ side.

A Map

There is a natural map

Irreducible representations
of pure inner forms of \mathbf{G}

\rightarrow

Langlands parameters
 $\varphi: \mathrm{WD}_{\mathbb{Q}_p} \rightarrow {}^L\mathbf{G}$

It is surjective and finite-to-one; the fibers are called *L-packets*.

L-packets

Moreover, we can naturally parameterize these fibers. Given a Langlands parameter φ , let $Z_{\hat{\mathbf{G}}}(\varphi)$ be the centralizer in $\hat{\mathbf{G}}$ of φ , and let ${}^L Z$ be the center of ${}^L \mathbf{G}$. Define

$$A_\varphi = \pi_0(Z_{\hat{\mathbf{G}}}(\varphi)).$$

The fibers should be in bijection with

$$A_\varphi^\vee = \{\text{irreducible representations of } A_\varphi\}.$$

So we get a natural bijection

Irreducible representations
of pure inner forms of \mathbf{G}

\leftrightarrow

(φ, ρ) with $\varphi: \text{WD}_{\mathbb{Q}_p} \rightarrow {}^L \mathbf{G}$
and $\rho \in A_\varphi^\vee$

Which Pure Inner Form?

The choice of $\rho \in A_\varphi^\vee$ determines which pure inner form arises on the other side of the correspondence. Kottwitz showed

$$H^1(\mathbb{Q}_p, \mathbf{G}) = \text{Hom}(\pi_0({}^L Z), \mathbb{C}^\times).$$

Using this canonical bijection, together with the fact that ${}^L Z \subset Z_{\hat{\mathbf{G}}}(\varphi)$, the restriction of $\rho \in A_\varphi^\vee$ to $\pi_0({}^L Z)$ determines a pure inner form of \mathbf{G} ; for any φ the representation associated to (φ, ρ) should be a representation of this group.

Restrictions on φ

From now on we fix a totally ramified quadratic extension E/\mathbb{Q}_p and set $\mathbf{G} = \mathbf{U}(V)$ for V a quasi-split Hermitian space over E . We say that a Langlands parameter φ is

- *discrete* if $Z_{\hat{\mathbf{G}}}(\varphi)$ is finite,
- *tame* if φ factors through the maximal tame quotient (and thus $p \neq 2$).
- *regular* if $Z_{\hat{\mathbf{G}}}(\varphi(\tilde{\tau}))$ is connected and minimum dimensional (here $\tilde{\tau}$ is a procyclic generator of tame inertia).

We will construct an L-packet of supercuspidal representations of pure inner forms of $\mathbf{G}(\mathbb{Q}_p)$ given a tame, discrete regular parameter.

Filtrations

$\mathbf{G}(\mathbb{Q}_p)$ acts on the Bruhat-Tits building $\mathcal{B}(\mathbf{G})$, and we can classify the compact subgroups of $\mathbf{G}(\mathbb{Q}_p)$ as stabilizers of convex subsets of $\mathcal{B}(\mathbf{G})$

- Each such compact \mathbf{H} has the structure of a \mathbb{Z}_p -scheme.
- There is a decreasing filtration on each \mathbf{H} .
- \mathbf{H}^0 is just the connected component of the identity (as a \mathbb{Z}_p -scheme) and is of finite index in \mathbf{H} .
- The special fiber $\mathbf{H}(\mathbb{F}_p)$ is given by $\mathbf{H}^0/\mathbf{H}^{0+}$.
- The filtration on \mathbf{T} is the one given by Moy and Prasad, coming from the filtration on \mathbb{Q}_p^\times .

We can thus obtain representations of compact subgroups of \mathbf{G} by pulling back representations of reductive groups over finite fields.

Outline

Our plan for constructing an L-packet from φ is as follows. We construct:

- A maximal unramified anisotropic torus \mathbf{T} , which embeds into \mathbf{G} in various ways,
- A character χ_φ on \mathbf{T}^0 that vanishes on \mathbf{T}^{0+} ,
- For each $\rho \in A_\varphi^\vee$, an embedding of \mathbf{T} into a maximal compact subgroup $\mathbf{H} \subset \mathbf{G}'$ for \mathbf{G}' a pure inner form of \mathbf{G} .
- We get a Deligne-Lusztig representation of $\mathbf{H}^0(\mathbb{F}_\rho) = \mathbf{H}^0/\mathbf{H}^{0+}$ associated to the torus $\mathbf{T}^0(\mathbb{F}_\rho) = \mathbf{T}^0/\mathbf{T}^{0+}$ and the character χ_φ .
- We induce this representation up to a representation of \mathbf{G}' .

An Unramified Anisotropic Torus

We have a specified torus $\hat{\mathbf{S}} \subset \hat{\mathbf{G}}$.

- We can conjugate φ so that $\varphi(\tilde{\tau}) \in \hat{\mathbf{S}} \rtimes \text{Gal}(E/\mathbb{Q}_p)$.
- $\varphi(F)$ then lies in the normalizer of $\hat{\mathbf{S}}$. It's projection onto the Weyl group gives an element of

$$H^1(\langle F \rangle, W^I) \hookrightarrow H^1(\mathbb{Q}_p, W),$$

which is exactly the data we need to define a maximal unramified torus as a twist of \mathbf{S} .

- Discreteness of φ implies that

$$(\hat{\mathfrak{g}}^I)^F = 0 \iff \hat{\mathfrak{g}}^I = \hat{\mathfrak{s}}^I \text{ and } X_*(\mathbf{S})^F = 0$$

and thus that \mathbf{T} is anisotropic.

A Character

- The image of φ lands inside some finite extension of $\hat{\mathbf{S}}$. If it were a semidirect product, we could just use the local Langlands correspondence for tori:

$$H^1(\mathbb{Q}_p, \hat{\mathbf{T}}) \cong \text{Hom}(\mathbf{T}(\mathbb{Q}_p), \mathbb{C}^\times).$$

- In general this extension is not a semidirect product. But we can use a modification of the Langlands correspondence for tori to obtain a character χ_φ of $\mathbf{T}^0(\mathbb{Q}_p)$, where \mathbf{T}^0 is the connected component in the Néron model of \mathbf{T} .
- The depth preservation feature of the Langlands correspondence for tori, together with the fact that φ is tamely ramified, implies that χ_φ vanishes on \mathbf{T}^{0+} .

Construction of χ_φ

Let $D \subset {}^L\mathbf{G}$ be the group generated by $\hat{\mathbf{S}} \rtimes \text{Gal}(E/\mathbb{Q}_p)$ and $\varphi(\mathbf{F})$. Denote by $M = \mathbb{Q}_{p^s} \cdot E$ the splitting field of \mathbf{T} .


$$1 \rightarrow \hat{\mathbf{T}} \rightarrow D \rightarrow \text{Gal}(M/\mathbb{Q}_p) \rightarrow 1.$$

Let $P_K(D, \mathbf{T})$ be the set of homomorphisms from $\text{Gal}(\bar{K}/K)$ to D that project correctly onto $\text{Gal}(M/\mathbb{Q}_p)$, modulo conjugacy by $\hat{\mathbf{T}}$. Set D_s as the preimage in D of $\text{Gal}(M/\mathbb{Q}_{p^s})$ and let $\Gamma = \text{Gal}(\mathbb{Q}_{p^s}/\mathbb{Q}_p)$. Then χ_φ is the image of φ under

$$\begin{aligned} P_{\mathbb{Q}_p}(D, \mathbf{T}) &\xrightarrow{\text{res}} P_{\mathbb{Q}_{p^s}}(D_s, \mathbf{T})^\Gamma \cong H^1(\mathbb{Q}_{p^s}, \hat{\mathbf{T}})^\Gamma \\ &\cong \text{Hom}(\mathbf{T}(\mathbb{Q}_{p^s})_\Gamma, \mathbb{C}^\times) \rightarrow \text{Hom}(\mathbf{T}^0(\mathbb{Q}_{p^s})_\Gamma, \mathbb{C}^\times) \\ &\cong \text{Hom}(\mathbf{T}^0(\mathbb{Q}_p), \mathbb{C}^\times). \end{aligned}$$

Basic Tori

We classify unramified anisotropic twists of the “quasi-split” torus \mathbf{S} . Essentially, they are products of basic tori. For each $s = 2r$, define $\mathbf{T}_s = \{x \in E_s : \text{Nm}_{E_s/L_r} x = 1\}$,



Embeddings of Basic Tori

In order to get Deligne-Lustig representations, we need to embed \mathbf{T} into maximal compacts of \mathbf{G} . We do so by building a Hermitian space around each basic torus in the product decomposition of \mathbf{T} .

For each $\kappa \in L_r^\times$, we define a Hermitian product on E_s

$$\phi_\kappa(x, y) = \mathrm{Tr}_{E_s/E} \left(\frac{\kappa}{\pi_L} x \cdot \eta_s(y) \right)$$

This Hermitian space is quasi-split if and only if $v_L(\kappa)$ is even.

Embeddings of General Tori

In general, we choose a κ_j for each basic torus in the decomposition of \mathbf{T} . This choice corresponds to a choice of $\rho \in A_\varphi^\vee$, and the pure inner form \mathbf{G}' of \mathbf{G} we obtain depends on the sum of the valuations of the κ_j modulo 2.

We prove \mathbf{T} fixes a unique point on the building $\mathcal{B}(\mathbf{G}')$ and thus embeds in a unique maximal compact $\mathbf{H} \subset \mathbf{G}' = \mathbf{U}(V')$.
The reduction of \mathbf{H} is

$$\mathbf{O}(m) \times \mathbf{Sp}(m'),$$

where m is the sum of the dimensions of basic tori whose κ_j has even valuation and m' is the sum of those with $v(\kappa_j)$ odd.

Constructing a representation of $\mathbf{G}(\mathbb{Q}_p)$

Modulo p , we have a maximal torus $\mathbf{T}^0(\mathbb{F}_p)$ sitting in a connected reductive group $\mathbf{H}^0(\mathbb{F}_p)$ and a character χ_φ of $\mathbf{T}^0(\mathbb{F}_p)$. This situation was studied by Deligne and Lusztig, and they produce a representation from étale cohomology. The irreducibility of this representation follows from the regularity condition on φ . We pull back to \mathbf{H}^0 and the only wrinkle in the induction process occurs between \mathbf{H}^0 and \mathbf{H} . Once we have a representation of \mathbf{H} , we define a representation on all of $\mathbf{G}(\mathbb{Q}_p)$ by compact induction.

A Finite Induction

There are three cases for the induction from \mathbf{H}^0 to \mathbf{H} .

- n even, $\mathbf{H}(\mathbb{F}_p) = \mathrm{Sp}(n)$. Here $\mathbf{H} = \mathbf{H}^0$ and there is no induction.
- n even, otherwise. The fact that the normalizer of $\mathbf{T}^0(\mathbb{F}_p)$ in $\mathbf{H}(\mathbb{F}_p)$ contains the normalizer in $\mathbf{H}^0(\mathbb{F}_p)$ with index 2 implies that the induction remains irreducible.
- n odd. Now the induction from \mathbf{H}^0 to \mathbf{H} splits into two irreducible components. We can pick one using a recipe for the central character, together with the fact that in the case that n is odd the center of $O(m)$ is not contained in $SO(m)$.

Summary

$$\begin{array}{ccccc}
 \mathbf{T}(\mathbb{Q}_p) & \longrightarrow & \mathbf{H}(\mathbb{Q}_p) & \longrightarrow & \mathbf{G}'(\mathbb{Q}_p) \\
 \uparrow & & \uparrow & \searrow & \searrow \\
 \mathbf{T}^0(\mathbb{Q}_p) & \longrightarrow & \mathbf{H}^0(\mathbb{Q}_p) & \xrightarrow{\text{Ind}(\text{DL}(\mathbf{T}^0, \chi_\varphi))} & \text{GL}(\text{Ind } V) \\
 \downarrow & & \downarrow & \searrow & \searrow \\
 \mathbf{T}^0(\mathbb{F}_p) & \longrightarrow & \mathbf{H}^0(\mathbb{F}_p) & \xrightarrow{\text{DL}(\mathbf{T}^0, \chi_\varphi)} & \text{GL}(V) \\
 & \searrow & \searrow & \searrow & \\
 & \chi_\varphi & & & \\
 & & \mathbb{C}^\times & &
 \end{array}$$

$\xrightarrow{c\text{Ind}(\text{DL}(\mathbf{T}^0, \chi_\varphi))} \text{GL}(c\text{Ind } V)$

Further Work

- There are only a few instances where I depend on the fact that $\mathbf{G} = \mathbf{U}(V)$. I'd like to come up with general arguments that work for any reductive group splitting over a tame extension.
- There are various properties we expect the correspondence to satisfy. All of the properties that Reeder and DeBacker prove about their L-packets (including stability and an analysis of which representations are generic) we should be able to do for L-packets in the tame case.
- I want to build a computational framework within Sage to experiment with these L-packets.