partition λ of $n \geq 0$:

$$\lambda = (\lambda_1, \lambda_2, \ldots)$$

$$\lambda_1 \geq \lambda_2 \geq \cdots \geq 0$$

$$\sum \lambda_i = n$$

Young diagram of $\lambda = (4, 4, 1)$:



standard Young tableau (**SYT**) of shape $(4, 4, 1)$:

$$<$$

	1	3	4	7
$\wedge$	2	6	8	9
	5			

$\boldsymbol{f^{\lambda}}$ = # of SYT of shape λ

$$\begin{array}{lll} 1\,2\,3\,4 & 1\,2\,3\,5 & 1\,2\,3\,6 \\ 5\,6 & 4\,6 & 4\,5 \\ & & \\ 1\,2\,4\,5 & 1\,2\,4\,6 & 1\,2\,5\,6 \\ 3\,6 & 3\,5 & 3\,4 \\ & & \\ 1\,3\,4\,5 & 1\,3\,4\,6 & 1\,3\,5\,6 \\ 2\,6 & 2\,5 & 2\,4 \end{array}$$

$$f^{4,2} = 9$$



$\boldsymbol{H(u)}$: **hook** at (or of) u

$\boldsymbol{h(u)} = \#H(u)$: **hook length** at u

Frame-Robinson-Thrall hook length formula (1954):

$$f^\lambda = \frac{n!}{\prod_{u\in\lambda} h(u)}$$

$$\text{hook lengths}: \qquad \begin{matrix} 5&4&2&1\\ 2&1 \end{matrix}$$

$$f^{4,2} = \frac{6!}{5\cdot 4\cdot 2\cdot 2\cdot 1\cdot 1} = 9$$

"nice" bijective proof by Novelli-Pak-Stoyanovskii (1997)

Robinson-Schensted-Knuth (RSK) algorithm: $w \stackrel{\text{rsk}}{\rightarrow} (P,Q)$, where $w \in \mathfrak{S}_n$ and P,Q are SYT of same shape $\lambda \vdash n$

Note. Schensted = Ea Ea

(ea.ea.home.mindspring.com)

$$w = 4273615$$

$$\begin{array}{lll} \mathbf{4} & \qquad & 1 \end{array}$$

$$\begin{array}{lll} \mathbf{2} & \qquad & 1 \\ \mathbf{4} & & 2 \end{array}$$

$$\begin{array}{lll} 2\,\mathbf{7} & \qquad & 1\,3 \\ 4 & & 2 \end{array}$$

$$\begin{array}{lll} 2\,\mathbf{3} & \qquad & 1\,3 \\ 4\,\mathbf{7} & & 2\,4 \end{array}$$

$$\begin{array}{lll} 2\,3\,\mathbf{6} & \qquad & 1\,3\,5 \\ 4\,7 & & 2\,4 \end{array}$$

$$\begin{array}{lll} \mathbf{1}\,3\,6 & \qquad & 1\,3\,5 \\ \mathbf{2}\,7 & & 2\,4 \\ \mathbf{4} & & 6 \end{array}$$

$$\begin{array}{lll} 1\,3\,\mathbf{5} & \qquad & 1\,3\,5 \\ 2\,\mathbf{6} & & 2\,4 \\ 4\,\mathbf{7} & & 6\,7 \end{array}$$

An element j bumps the smallest $i > j$.

$\boldsymbol{\chi^\lambda}$: irreducible character of $\mathfrak{S}_n$
indexed by $\lambda \vdash n$

$$f^\lambda = \chi^\lambda(1) = \dim \chi^\lambda$$

$$\sum_{\lambda \vdash n} \left(f^\lambda\right)^2 = n!$$

First symmetry property.

$$w = 4273615 \xrightarrow{\text{rsk}} \begin{array}{lll} 1&3&5 \\ 2&6 \\ 4&7 \end{array} \quad \begin{array}{lll} 1&3&5 \\ 2&4 \\ 6&7 \end{array}$$

$$w^{-1} = 6241753 \xrightarrow{\text{rsk}} \begin{array}{lll} 1&3&5 \\ 2&4 \\ 6&7 \end{array} \quad \begin{array}{lll} 1&3&5 \\ 2&6 \\ 4&7 \end{array}$$

Theorem (Schützenberger) *If* $w \stackrel{\text{rsk}}{\rightarrow} (P, Q)$, *then*

$$w^{-1} \xrightarrow{\text{rsk}} (Q, P).$$

Corollary. *Let* $t(n)$ *denote the number of SYT with* n *squares. Then*

$$t(n) = \#\{w \in \mathfrak{S}_n \,:\, w^2 = 1\}$$

$$\sum_{n \geq 0} t(n) \frac{x^n}{n!} = \exp\left(x + \frac{x^2}{2}\right).$$

$$t(n) = \sum_{\lambda \vdash n} f^\lambda = \sum_{\lambda \vdash n} \chi^\lambda(1)$$

Theorem (Frobenius). *Let G be a finite group and $\hat{G}$ its set of (complex) irreducible characters. Then*

$$\sum_{\chi \in \hat{G}} \chi(1) = \#\{w \in G \,:\, w^2 = 1\}$$

if and only if every representation of G is equivalent to a real representation (true for $\mathfrak{S}_n$).

$$w = 3\mathbf{1}8\mathbf{4}9\mathbf{67}25$$

$$\boldsymbol{is}(w) := \text{length of longest increasing subsequence of } w \in \mathfrak{S}_n$$

$$\mathrm{is}(318496725) = 4$$

$$318496725 \xrightarrow{\text{rsk}} \begin{array}{llll} 1 & 2 & 5 & 7 \\ 3 & 4 & 6 & \\ 8 & 9 & & \end{array} \qquad \begin{array}{llll} 1 & 3 & 5 & 7 \\ 2 & 4 & 6 & \\ 8 & 9 & & \end{array}$$

Theorem. *Let* $w \xrightarrow{\text{rsk}} (P, Q)$,

$$\text{shape}(P) = (\lambda_1, \lambda_2, \ldots).$$

Then

$$\text{is}(w) = \lambda_1.$$

Proof (sketch). Let the first row be

$$b_1, b_2, \ldots, b_k.$$

Straightforward to show by induction on n that b_i is the rightmost element j of w for which the longest increasing subsequence of w ending at j has length i. $\square$

Second symmetry property.

$$w = a_1 a_2 \cdots a_n, \quad \overline{w} := a_n \cdots a_2 a_1$$

Theorem (Schensted). *If* $w \stackrel{\text{rsk}}{\to} (P, Q)$, *then*

$$\overline{w} \stackrel{\text{rsk}}{\to} (P^t, \operatorname{evac}(Q)^t).$$

Corollary. *Let* $w \xrightarrow{\text{rsk}} (P, Q)$, $\operatorname{shape}(P) = (\lambda_1, \lambda_2, \ldots)$. *Then*

$$\operatorname{ds}(w) = \lambda_1' = \ell(\lambda).$$

Corollary (Erdős-Szekeres). *Let* $w \in \mathfrak{S}_{pq+1}$. *Then either*

$$\operatorname{is}(w) > p \ \text{ or } \ \operatorname{ds}(w) > q.$$

Corollary. *Let $p \le q$ (say). Then*

$$\#\{w \in \mathfrak{S}_{pq} \,:\, \mathrm{is}(w) = p,\ \mathrm{ds}(w) = q\}$$

$$= f^{q\times p}$$

$$= \frac{(pq)!}{1^1 2^2 \cdots p^p (p+1)^p \cdots q^p (q+1)^{p-1} \cdots (p+q-1)^1}.$$

7	6	5	4	3
6	5	4	3	2
5	4	3	2	1

$$\begin{tikzcd}
\begin{array}{c}\begin{bmatrix} 0&0&0&1&0\\ 1&0&0&0&0\\ 0&1&0&0&0\\ 0&0&0&0&1\\ 0&0&1&0&0 \end{bmatrix}\\ w = 41253\end{array} \arrow[r] \arrow[d] & \begin{array}{c}\begin{bmatrix} 0&1&0&0&0\\ 0&0&1&0&0\\ 0&0&0&0&1\\ 1&0&0&0&0\\ 0&0&0&1&0 \end{bmatrix}\\ w^{-1} = 23514\end{array} \\
\begin{array}{c}\begin{bmatrix} 0&0&1&0&0\\ 0&0&0&0&1\\ 0&1&0&0&0\\ 1&0&0&0&0\\ 0&0&0&1&0 \end{bmatrix}\\ \overline{w} = 35214\end{array} &
\end{tikzcd}$$

These two reflections generate the dihedral group D_4 (symmetries of a square).

A **reverse semistandard tableau** T of shape $(5, 4, 3)$:

$$\begin{array}{lllll} 6 & 6 & 4 & 2 & 2 \\ 4 & 4 & 3 & 1 & \\ 2 & 1 & 1 & & \end{array}$$

$$x^T = x_1^3 x_2^3 x_3 x_4^3 x_6^2$$

(weakly decreasing in rows, strictly decreasing in columns)

$$A = \begin{bmatrix} 0 & 0 & 0 & 2 \\ 0 & 2 & 1 & 0 \\ 1 & 0 & 1 & 0 \end{bmatrix}$$

$$w_A = \begin{pmatrix} 3 & 3 & 2 & 2 & 2 & 1 & 1 \\ 3 & 1 & 3 & 2 & 2 & 4 & 4 \end{pmatrix}$$

$$\begin{array}{l} \mathbf{3} \end{array} \qquad \begin{array}{l} 3 \end{array}$$

$$\begin{array}{ll} 3 & \mathbf{1} \end{array} \qquad \begin{array}{ll} 3 & 3 \end{array}$$

$$\begin{array}{ll} 3 & \mathbf{3} \\ \mathbf{1} & \end{array} \qquad \begin{array}{ll} 3 & 3 \\ 2 & \end{array}$$

$$\begin{array}{lll} 3 & 3 & \mathbf{2} \\ 1 & & \end{array} \qquad \begin{array}{lll} 3 & 3 & 2 \\ 2 & & \end{array}$$

$$\begin{array}{llll} 3 & 3 & 2 & \mathbf{2} \\ 1 & & & \end{array} \qquad \begin{array}{llll} 3 & 3 & 2 & 2 \\ 2 & & & \end{array}$$

$$\begin{array}{llll} \mathbf{4} & 3 & 2 & 2 \\ \mathbf{3} & & & \\ \mathbf{1} & & & \end{array} \qquad \begin{array}{llll} 3 & 3 & 2 & 2 \\ 2 & & & \\ 1 & & & \end{array}$$

$$\begin{array}{llll} 4 & \mathbf{4} & 2 & 2 \\ 3 & \mathbf{3} & & \\ 1 & & & \end{array} \qquad \begin{array}{llll} 3 & 3 & 2 & 2 \\ 2 & 1 & & \\ 1 & & & \end{array}$$

An element j bumps the largest element $i < j$.

Lemma (simple). Let $A \xrightarrow{\text{rsk}} (P, Q)$. Equal elements of Q are inserted left-to-right (allowing the construction of the inverse map $(P, Q) \to A$).

Schur function:

$$\boldsymbol{s_\lambda} = \sum_{\substack{\text{RSSYT } T \\ \text{shape}(T)=\lambda}} x^T$$

3 3	3 3	3 2	3 1
2	1	2	1

2 2	2 1	3 2	3 1
1	1	1	2

$$s_{2,1}(x_1, x_2, x_3) = x_2x_3^2 + x_1x_3^2 + x_2^2x_3 + x_1^2x_3 \\ + x_1x_2^2 + x_1^2x_2 + 2x_1x_2x_3$$

Cauchy identity:

$$\prod_{i,j}(1 - x_iy_j)^{-1} = \sum_\lambda s_\lambda(x)s_\lambda(y)$$

(analytic formulation of RSK for (reverse) SSYT)

Plane partition of $n \geq 0$:

$$\pi = (\pi_{ij})_{i,j\geq 1}$$

$$\pi_{ij} \geq \pi_{i+1,j}, \quad \pi_{ij} \geq \pi_{i,j+1}, \quad \sum \pi_{ij} = n$$

$$\boldsymbol{a(n)} = \#\ \text{plane partitions of}\ n$$

$$\begin{array}{llllll} 5 & 5 & 3 & 2 & 2 & 1 \\ 5 & 3 & 3 & 1 & 1 & \\ 5 & 2 & 2 & 1 & 1 & \\ 2 & 2 & 1 & & & \end{array}$$

$$3 \qquad 21 \qquad 111 \qquad \begin{array}{l} 2 \\ 1 \end{array} \qquad \begin{array}{l} 1 \\ 1 \\ 1 \end{array} \qquad \begin{array}{ll} 1 & 1 \\ 1 & \end{array}$$

$$a(3) = 6$$

MacMahon:

$$\sum_{n\geq 0} a(n)x^n = \prod_{i\geq 1}(1-x^i)^{-i}.$$

$$A = \begin{bmatrix} 0 & 0 & 1 \\ 0 & 1 & 2 \\ 1 & 0 & 1 \\ 0 & 2 & 0 \end{bmatrix}$$

$$\xrightarrow{\text{rsk}} \begin{array}{l} 3\,3\,3\,3\,2 \\ 2\,2\,2 \\ 1 \end{array} \quad \begin{array}{l} 4\,4\,3\,2\,1 \\ 3\,2\,2 \\ 1 \end{array}$$

$$\xrightarrow{\text{merge}} \begin{array}{l} 3\,3\,3\,3\,2 \\ 3\,3\,3\,1 \\ 3\,3\,2 \\ 2\,1 \end{array}$$

$$\xrightarrow{\text{rowconj}} \begin{array}{l} 5\,5\,4 \\ 4\,3\,3 \\ 3\,3\,2 \\ 2\,1 \end{array} = \pi_A$$

Merge column-by-column.

merge of $(5, 4, 2)$ and $(6, 3, 1)$ is $(5, 5, 4, 2, 1, 1)$:



$$|\pi_A| = \sum_{i,j}(i+j-1)a_{ij}$$

$$\#(\text{rows of } \pi_A) = \#(\text{rows of } A)$$

$$\#(\text{columns of } \pi_A) = \#(\text{columns of } A)$$

$$\begin{aligned}
\Rightarrow \sum_{\pi} x^{|\pi|} &= \sum_{A} x^{(i+j-1)a_{ij}} \\
&= \prod_{i,j\geq 1}\left(\sum_{a_{ij}\geq 0} x^{(i+j-1)a_{ij}}\right) \\
&= \prod_{i,j\geq 1}\left(1-x^{i+j-1}\right)^{-1} \\
&= \prod_{i\geq 1}(1-x^{i})^{-i}
\end{aligned}$$

RSK $\Rightarrow$

$$\sum_{\substack{\pi \\ \leq r \text{ rows} \\ \leq s \text{ cols}}} x^{|\pi|} = \prod_{i=1}^{r} \prod_{j=1}^{s} (1 - x^{i+j-1})^{-1}$$

More difficult (MacMahon):

$$\sum_{\substack{\pi \\ \leq r \text{ rows} \\ \leq s \text{ cols} \\ \max \leq t}} x^{|\pi|} = \prod_{i=1}^{r} \prod_{j=1}^{s} \prod_{k=1}^{t} \frac{1 - x^{i+j+k-1}}{1 - x^{i+j+k-2}}$$

Oscillating tableaux: start with $\emptyset$, add or remove a square at each step.



shape $(3,1)$, length 8

$$\tilde{f}_n^{\lambda} = \#\{\text{osc. tab. of shape } \lambda, \text{ length } n\}$$





Hence

$$\begin{aligned}\tilde{f}^{\emptyset}_{2n} &= \sum_{\lambda} \left(\tilde{f}^{\lambda}_{n}\right)^2 \\ &= (2n-1)!! \\ &:= (2n-1)(2n-3)\cdots 1.\end{aligned}$$

$\tilde{f}^{\lambda}_{n}$ is the dimension of an irreducible representation of the **Brauer algebra** $\mathfrak{B}_n$, a semisimple algebra of dimension $(2n-1)!!$.

$$s_i = (i, i+1) \in \mathfrak{S}_n$$

reduced decomposition $(a_1, \ldots, a_p)$ of $w \in \mathfrak{S}_n$:

$$w = s_{a_1} \cdots s_{a_p},$$

where p is minimal, i.e.,

$$p = \ell(w) = \#\{(i,j)\ :\ i < j,\ w(i) > w(j)\}.$$

$\boldsymbol{R(w)}$: set of reduced decomps. of w

$$\boldsymbol{r(w)} := \#R(w)$$

E.g., $w = 321$, $R(w) = \{(1,2,1), (2,1,2)\}$, $r(w) = 2$.

$$w_0 = n, n-1, \ldots, 1 \in \mathfrak{S}_n$$

Theorem.

$$r(w_0) = f^{(n-1,n-2,\ldots,1)} = \frac{\binom{n}{2}!}{1^{n-1}\,3^{n-2}\,5^{n-3}\cdots(2n-3)}$$

Proof (Edelman-Greene).



$$(2,3,1,2,1,3) \in R(4321)$$

Inverse:



Example. $312132 \in R(4321)$

$$\begin{array}{l} \mathbf{3} \end{array} \qquad \begin{array}{l} 1 \end{array}$$

$$\begin{array}{l} \mathbf{1} \\ \mathbf{3} \end{array} \qquad \begin{array}{l} 1 \\ 2 \end{array}$$

$$\begin{array}{ll} 1 & \mathbf{2} \\ 3 & \end{array} \qquad \begin{array}{ll} 1 & 3 \\ 2 & \end{array}$$

$$\begin{array}{ll} 1 & \mathbf{2} \\ \mathbf{2} & \\ \mathbf{3} & \end{array} \qquad \begin{array}{ll} 1 & 3 \\ 2 & \\ 4 & \end{array}$$

$$\begin{array}{lll} 1 & 2 & \mathbf{3} \\ 2 & & \\ 3 & & \end{array} \qquad \begin{array}{lll} 1 & 3 & 5 \\ 2 & & \\ 4 & & \end{array}$$

$$\begin{array}{lll} 1 & 2 & \mathbf{3} \\ 2 & \mathbf{3} & \\ 3 & & \end{array} \qquad \begin{array}{lll} 1 & 3 & 5 \\ 2 & 6 & \\ 4 & & \end{array}$$

P is always the same, and Q runs through all SYT of shape $(n-1, n-2, \ldots, 1)$.

Variant:

$$f(w) = \sum_{(a_1,\ldots,a_p)\in R(w)} a_1 a_2 \cdots a_p$$

E.g., $w = 321$, $R(321) = \{121, 212\}$,

$$f(321) = 1\cdot 2\cdot 1 + 2\cdot 1\cdot 2 = 6 = 3!$$

Theorem (Macdonald, Fomin-S).

$$f(w_0) = \binom{n}{2}!$$

More generally, $f(w) = \mathfrak{S}_w(1,1,\ldots,1)\ell(w)!$, where $\mathfrak{S}_w$ is a **Schubert polynomial**.

Corollary. $f(w) = \ell(w)!$ if and only if there never holds

$$i < j < k \Rightarrow w(i) < w(k) < w(j).$$

Number of such $w \in \mathfrak{S}_n$ is the Catalan number $C_n = \frac{1}{n+1}\binom{2n}{n}$.