

# 18.600: Lecture 25

## Lectures 15-24 Review

Scott Sheffield

MIT

# Outline

Continuous random variables

Problems motivated by coin tossing

Random variable properties

Continuous random variables

Problems motivated by coin tossing

Random variable properties

## Continuous random variables

- ▶ Say  $X$  is a **continuous random variable** if there exists a **probability density function**  $f = f_X$  on  $\mathbb{R}$  such that 
$$P\{X \in B\} = \int_B f(x)dx := \int 1_B(x)f(x)dx.$$

# Continuous random variables

- ▶ Say  $X$  is a **continuous random variable** if there exists a **probability density function**  $f = f_X$  on  $\mathbb{R}$  such that 
$$P\{X \in B\} = \int_B f(x)dx := \int 1_B(x)f(x)dx.$$
- ▶ We may assume  $\int_{\mathbb{R}} f(x)dx = \int_{-\infty}^{\infty} f(x)dx = 1$  and  $f$  is non-negative.

# Continuous random variables

- ▶ Say  $X$  is a **continuous random variable** if there exists a **probability density function**  $f = f_X$  on  $\mathbb{R}$  such that  $P\{X \in B\} = \int_B f(x)dx := \int 1_B(x)f(x)dx$ .
- ▶ We may assume  $\int_{\mathbb{R}} f(x)dx = \int_{-\infty}^{\infty} f(x)dx = 1$  and  $f$  is non-negative.
- ▶ Probability of interval  $[a, b]$  is given by  $\int_a^b f(x)dx$ , the area under  $f$  between  $a$  and  $b$ .

# Continuous random variables

- ▶ Say  $X$  is a **continuous random variable** if there exists a **probability density function**  $f = f_X$  on  $\mathbb{R}$  such that  $P\{X \in B\} = \int_B f(x)dx := \int 1_B(x)f(x)dx$ .
- ▶ We may assume  $\int_{\mathbb{R}} f(x)dx = \int_{-\infty}^{\infty} f(x)dx = 1$  and  $f$  is non-negative.
- ▶ Probability of interval  $[a, b]$  is given by  $\int_a^b f(x)dx$ , the area under  $f$  between  $a$  and  $b$ .
- ▶ Probability of any single point is zero.

# Continuous random variables

- ▶ Say  $X$  is a **continuous random variable** if there exists a **probability density function**  $f = f_X$  on  $\mathbb{R}$  such that  $P\{X \in B\} = \int_B f(x)dx := \int 1_B(x)f(x)dx$ .
- ▶ We may assume  $\int_{\mathbb{R}} f(x)dx = \int_{-\infty}^{\infty} f(x)dx = 1$  and  $f$  is non-negative.
- ▶ Probability of interval  $[a, b]$  is given by  $\int_a^b f(x)dx$ , the area under  $f$  between  $a$  and  $b$ .
- ▶ Probability of any single point is zero.
- ▶ Define **cumulative distribution function**  
 $F(a) = F_X(a) := P\{X < a\} = P\{X \leq a\} = \int_{-\infty}^a f(x)dx$ .



## Expectations of continuous random variables

- ▶ Recall that when  $X$  was a discrete random variable, with  $p(x) = P\{X = x\}$ , we wrote

$$E[X] = \sum_{x:p(x)>0} p(x)x.$$

# Expectations of continuous random variables

- ▶ Recall that when  $X$  was a discrete random variable, with  $p(x) = P\{X = x\}$ , we wrote

$$E[X] = \sum_{x:p(x)>0} p(x)x.$$

- ▶ How should we define  $E[X]$  when  $X$  is a continuous random variable?

# Expectations of continuous random variables

- ▶ Recall that when  $X$  was a discrete random variable, with  $p(x) = P\{X = x\}$ , we wrote

$$E[X] = \sum_{x:p(x)>0} p(x)x.$$

- ▶ How should we define  $E[X]$  when  $X$  is a continuous random variable?
- ▶ Answer:  $E[X] = \int_{-\infty}^{\infty} f(x)x dx.$

# Expectations of continuous random variables

- ▶ Recall that when  $X$  was a discrete random variable, with  $p(x) = P\{X = x\}$ , we wrote

$$E[X] = \sum_{x:p(x)>0} p(x)x.$$

- ▶ How should we define  $E[X]$  when  $X$  is a continuous random variable?
- ▶ Answer:  $E[X] = \int_{-\infty}^{\infty} f(x)x dx$ .
- ▶ Recall that when  $X$  was a discrete random variable, with  $p(x) = P\{X = x\}$ , we wrote

$$E[g(X)] = \sum_{x:p(x)>0} p(x)g(x).$$

# Expectations of continuous random variables

- ▶ Recall that when  $X$  was a discrete random variable, with  $p(x) = P\{X = x\}$ , we wrote

$$E[X] = \sum_{x:p(x)>0} p(x)x.$$

- ▶ How should we define  $E[X]$  when  $X$  is a continuous random variable?
- ▶ Answer:  $E[X] = \int_{-\infty}^{\infty} f(x)x dx$ .
- ▶ Recall that when  $X$  was a discrete random variable, with  $p(x) = P\{X = x\}$ , we wrote

$$E[g(X)] = \sum_{x:p(x)>0} p(x)g(x).$$

- ▶ What is the analog when  $X$  is a continuous random variable?

# Expectations of continuous random variables

- ▶ Recall that when  $X$  was a discrete random variable, with  $p(x) = P\{X = x\}$ , we wrote

$$E[X] = \sum_{x:p(x)>0} p(x)x.$$

- ▶ How should we define  $E[X]$  when  $X$  is a continuous random variable?

- ▶ Answer:  $E[X] = \int_{-\infty}^{\infty} f(x)x dx$ .

- ▶ Recall that when  $X$  was a discrete random variable, with  $p(x) = P\{X = x\}$ , we wrote

$$E[g(X)] = \sum_{x:p(x)>0} p(x)g(x).$$

- ▶ What is the analog when  $X$  is a continuous random variable?
- ▶ Answer: we will write  $E[g(X)] = \int_{-\infty}^{\infty} f(x)g(x) dx$ .

## Variance of continuous random variables

- ▶ Suppose  $X$  is a continuous random variable with mean  $\mu$ .

## Variance of continuous random variables

- ▶ Suppose  $X$  is a continuous random variable with mean  $\mu$ .
- ▶ We can write  $\text{Var}[X] = E[(X - \mu)^2]$ , same as in the discrete case.



## Variance of continuous random variables

- ▶ Suppose  $X$  is a continuous random variable with mean  $\mu$ .
- ▶ We can write  $\text{Var}[X] = E[(X - \mu)^2]$ , same as in the discrete case.
- ▶ Next, if  $g = g_1 + g_2$  then

$$E[g(X)] = \int g_1(x)f(x)dx + \int g_2(x)f(x)dx = \int (g_1(x) + g_2(x))f(x)dx = E[g_1(X)] + E[g_2(X)].$$

## Variance of continuous random variables

- ▶ Suppose  $X$  is a continuous random variable with mean  $\mu$ .
- ▶ We can write  $\text{Var}[X] = E[(X - \mu)^2]$ , same as in the discrete case.
- ▶ Next, if  $g = g_1 + g_2$  then
$$E[g(X)] = \int g_1(x)f(x)dx + \int g_2(x)f(x)dx = \int (g_1(x) + g_2(x))f(x)dx = E[g_1(X)] + E[g_2(X)].$$
- ▶ Furthermore,  $E[ag(X)] = aE[g(X)]$  when  $a$  is a constant.

## Variance of continuous random variables

- ▶ Suppose  $X$  is a continuous random variable with mean  $\mu$ .
- ▶ We can write  $\text{Var}[X] = E[(X - \mu)^2]$ , same as in the discrete case.
- ▶ Next, if  $g = g_1 + g_2$  then
$$E[g(X)] = \int g_1(x)f(x)dx + \int g_2(x)f(x)dx = \int (g_1(x) + g_2(x))f(x)dx = E[g_1(X)] + E[g_2(X)].$$
- ▶ Furthermore,  $E[ag(X)] = aE[g(X)]$  when  $a$  is a constant.
- ▶ Just as in the discrete case, we can expand the variance expression as  $\text{Var}[X] = E[X^2 - 2\mu X + \mu^2]$  and use additivity of expectation to say that
$$\text{Var}[X] = E[X^2] - 2\mu E[X] + E[\mu^2] = E[X^2] - 2\mu^2 + \mu^2 = E[X^2] - E[X]^2.$$

## Variance of continuous random variables

- ▶ Suppose  $X$  is a continuous random variable with mean  $\mu$ .
- ▶ We can write  $\text{Var}[X] = E[(X - \mu)^2]$ , same as in the discrete case.
- ▶ Next, if  $g = g_1 + g_2$  then
$$E[g(X)] = \int g_1(x)f(x)dx + \int g_2(x)f(x)dx = \int (g_1(x) + g_2(x))f(x)dx = E[g_1(X)] + E[g_2(X)].$$
- ▶ Furthermore,  $E[ag(X)] = aE[g(X)]$  when  $a$  is a constant.
- ▶ Just as in the discrete case, we can expand the variance expression as  $\text{Var}[X] = E[X^2 - 2\mu X + \mu^2]$  and use additivity of expectation to say that
$$\text{Var}[X] = E[X^2] - 2\mu E[X] + E[\mu^2] = E[X^2] - 2\mu^2 + \mu^2 = E[X^2] - E[X]^2.$$
- ▶ Expectation of square minus square of expectation.

## Variance of continuous random variables

- ▶ Suppose  $X$  is a continuous random variable with mean  $\mu$ .
- ▶ We can write  $\text{Var}[X] = E[(X - \mu)^2]$ , same as in the discrete case.
- ▶ Next, if  $g = g_1 + g_2$  then
$$E[g(X)] = \int g_1(x)f(x)dx + \int g_2(x)f(x)dx = \int (g_1(x) + g_2(x))f(x)dx = E[g_1(X)] + E[g_2(X)].$$
- ▶ Furthermore,  $E[ag(X)] = aE[g(X)]$  when  $a$  is a constant.
- ▶ Just as in the discrete case, we can expand the variance expression as  $\text{Var}[X] = E[X^2 - 2\mu X + \mu^2]$  and use additivity of expectation to say that
$$\text{Var}[X] = E[X^2] - 2\mu E[X] + E[\mu^2] = E[X^2] - 2\mu^2 + \mu^2 = E[X^2] - E[X]^2.$$
- ▶ Expectation of square minus square of expectation.
- ▶ This formula is often useful for calculations.

# Outline

Continuous random variables

Problems motivated by coin tossing

Random variable properties

# Outline

Continuous random variables

Problems motivated by coin tossing

Random variable properties

## It's the coins, stupid

- ▶ Much of what we have done in this course can be motivated by the i.i.d. sequence  $X_i$  where each  $X_i$  is 1 with probability  $p$  and 0 otherwise. Write  $S_n = \sum_{i=1}^n X_n$ .



## It's the coins, stupid

- ▶ Much of what we have done in this course can be motivated by the i.i.d. sequence  $X_i$  where each  $X_i$  is 1 with probability  $p$  and 0 otherwise. Write  $S_n = \sum_{i=1}^n X_i$ .
- ▶ **Binomial** ( $S_n$  — number of heads in  $n$  tosses), **geometric** (steps required to obtain one heads), **negative binomial** (steps required to obtain  $n$  heads).

## It's the coins, stupid

- ▶ Much of what we have done in this course can be motivated by the i.i.d. sequence  $X_i$  where each  $X_i$  is 1 with probability  $p$  and 0 otherwise. Write  $S_n = \sum_{i=1}^n X_i$ .
- ▶ **Binomial** ( $S_n$  — number of heads in  $n$  tosses), **geometric** (steps required to obtain one heads), **negative binomial** (steps required to obtain  $n$  heads).
- ▶ **Standard normal** approximates law of  $\frac{S_n - E[S_n]}{SD(S_n)}$ . Here  $E[S_n] = np$  and  $SD(S_n) = \sqrt{\text{Var}(S_n)} = \sqrt{npq}$  where  $q = 1 - p$ .

## It's the coins, stupid

- ▶ Much of what we have done in this course can be motivated by the i.i.d. sequence  $X_i$  where each  $X_i$  is 1 with probability  $p$  and 0 otherwise. Write  $S_n = \sum_{i=1}^n X_i$ .
- ▶ **Binomial** ( $S_n$  — number of heads in  $n$  tosses), **geometric** (steps required to obtain one heads), **negative binomial** (steps required to obtain  $n$  heads).
- ▶ **Standard normal** approximates law of  $\frac{S_n - E[S_n]}{SD(S_n)}$ . Here  $E[S_n] = np$  and  $SD(S_n) = \sqrt{\text{Var}(S_n)} = \sqrt{npq}$  where  $q = 1 - p$ .
- ▶ **Poisson** is limit of binomial as  $n \rightarrow \infty$  when  $p = \lambda/n$ .

## It's the coins, stupid

- ▶ Much of what we have done in this course can be motivated by the i.i.d. sequence  $X_i$  where each  $X_i$  is 1 with probability  $p$  and 0 otherwise. Write  $S_n = \sum_{i=1}^n X_i$ .
- ▶ **Binomial** ( $S_n$  — number of heads in  $n$  tosses), **geometric** (steps required to obtain one heads), **negative binomial** (steps required to obtain  $n$  heads).
- ▶ **Standard normal** approximates law of  $\frac{S_n - E[S_n]}{SD(S_n)}$ . Here  $E[S_n] = np$  and  $SD(S_n) = \sqrt{\text{Var}(S_n)} = \sqrt{npq}$  where  $q = 1 - p$ .
- ▶ **Poisson** is limit of binomial as  $n \rightarrow \infty$  when  $p = \lambda/n$ .
- ▶ **Poisson point process**: toss one  $\lambda/n$  coin during each length  $1/n$  time increment, take  $n \rightarrow \infty$  limit.

## It's the coins, stupid

- ▶ Much of what we have done in this course can be motivated by the i.i.d. sequence  $X_i$  where each  $X_i$  is 1 with probability  $p$  and 0 otherwise. Write  $S_n = \sum_{i=1}^n X_i$ .
- ▶ **Binomial** ( $S_n$  — number of heads in  $n$  tosses), **geometric** (steps required to obtain one heads), **negative binomial** (steps required to obtain  $n$  heads).
- ▶ **Standard normal** approximates law of  $\frac{S_n - E[S_n]}{SD(S_n)}$ . Here  $E[S_n] = np$  and  $SD(S_n) = \sqrt{\text{Var}(S_n)} = \sqrt{npq}$  where  $q = 1 - p$ .
- ▶ **Poisson** is limit of binomial as  $n \rightarrow \infty$  when  $p = \lambda/n$ .
- ▶ **Poisson point process**: toss one  $\lambda/n$  coin during each length  $1/n$  time increment, take  $n \rightarrow \infty$  limit.
- ▶ **Exponential**: time till first event in  $\lambda$  Poisson point process.

## It's the coins, stupid

- ▶ Much of what we have done in this course can be motivated by the i.i.d. sequence  $X_i$  where each  $X_i$  is 1 with probability  $p$  and 0 otherwise. Write  $S_n = \sum_{i=1}^n X_i$ .
- ▶ **Binomial** ( $S_n$  — number of heads in  $n$  tosses), **geometric** (steps required to obtain one heads), **negative binomial** (steps required to obtain  $n$  heads).
- ▶ **Standard normal** approximates law of  $\frac{S_n - E[S_n]}{\text{SD}(S_n)}$ . Here  $E[S_n] = np$  and  $\text{SD}(S_n) = \sqrt{\text{Var}(S_n)} = \sqrt{npq}$  where  $q = 1 - p$ .
- ▶ **Poisson** is limit of binomial as  $n \rightarrow \infty$  when  $p = \lambda/n$ .
- ▶ **Poisson point process**: toss one  $\lambda/n$  coin during each length  $1/n$  time increment, take  $n \rightarrow \infty$  limit.
- ▶ **Exponential**: time till first event in  $\lambda$  Poisson point process.
- ▶ **Gamma distribution**: time till  $n$ th event in  $\lambda$  Poisson point process.

# Discrete random variable properties derivable from coin toss intuition

- ▶ **Sum of two independent binomial random variables** with parameters  $(n_1, p)$  and  $(n_2, p)$  is itself binomial  $(n_1 + n_2, p)$ .

## Discrete random variable properties derivable from coin toss intuition

- ▶ **Sum of two independent binomial random variables** with parameters  $(n_1, p)$  and  $(n_2, p)$  is itself binomial  $(n_1 + n_2, p)$ .
- ▶ **Sum of  $n$  independent geometric random variables** with parameter  $p$  is negative binomial with parameter  $(n, p)$ .



# Discrete random variable properties derivable from coin toss intuition

- ▶ **Sum of two independent binomial random variables** with parameters  $(n_1, p)$  and  $(n_2, p)$  is itself binomial  $(n_1 + n_2, p)$ .
- ▶ **Sum of  $n$  independent geometric random variables** with parameter  $p$  is negative binomial with parameter  $(n, p)$ .
- ▶ **Expectation of geometric random variable** with parameter  $p$  is  $1/p$ .

# Discrete random variable properties derivable from coin toss intuition

- ▶ **Sum of two independent binomial random variables** with parameters  $(n_1, p)$  and  $(n_2, p)$  is itself binomial  $(n_1 + n_2, p)$ .
- ▶ **Sum of  $n$  independent geometric random variables** with parameter  $p$  is negative binomial with parameter  $(n, p)$ .
- ▶ **Expectation of geometric random variable** with parameter  $p$  is  $1/p$ .
- ▶ **Expectation of binomial random variable** with parameters  $(n, p)$  is  $np$ .

# Discrete random variable properties derivable from coin toss intuition

- ▶ **Sum of two independent binomial random variables** with parameters  $(n_1, p)$  and  $(n_2, p)$  is itself binomial  $(n_1 + n_2, p)$ .
- ▶ **Sum of  $n$  independent geometric random variables** with parameter  $p$  is negative binomial with parameter  $(n, p)$ .
- ▶ **Expectation of geometric random variable** with parameter  $p$  is  $1/p$ .
- ▶ **Expectation of binomial random variable** with parameters  $(n, p)$  is  $np$ .
- ▶ **Variance of binomial random variable** with parameters  $(n, p)$  is  $np(1 - p) = npq$ .

## Continuous random variable properties derivable from coin toss intuition

- ▶ **Sum of  $n$  independent exponential random variables** each with parameter  $\lambda$  is gamma with parameters  $(n, \lambda)$ .

## Continuous random variable properties derivable from coin toss intuition

- ▶ **Sum of  $n$  independent exponential random variables** each with parameter  $\lambda$  is gamma with parameters  $(n, \lambda)$ .
- ▶ **Memoryless properties:** given that exponential random variable  $X$  is greater than  $T > 0$ , the conditional law of  $X - T$  is the same as the original law of  $X$ .

# Continuous random variable properties derivable from coin toss intuition

- ▶ **Sum of  $n$  independent exponential random variables** each with parameter  $\lambda$  is gamma with parameters  $(n, \lambda)$ .
- ▶ **Memoryless properties:** given that exponential random variable  $X$  is greater than  $T > 0$ , the conditional law of  $X - T$  is the same as the original law of  $X$ .
- ▶ Write  $p = \lambda/n$ . **Poisson random variable expectation** is  $\lim_{n \rightarrow \infty} np = \lim_{n \rightarrow \infty} n \frac{\lambda}{n} = \lambda$ . **Variance** is  $\lim_{n \rightarrow \infty} np(1 - p) = \lim_{n \rightarrow \infty} n(1 - \lambda/n)\lambda/n = \lambda$ .

## Continuous random variable properties derivable from coin toss intuition

- ▶ **Sum of  $n$  independent exponential random variables** each with parameter  $\lambda$  is gamma with parameters  $(n, \lambda)$ .
- ▶ **Memoryless properties:** given that exponential random variable  $X$  is greater than  $T > 0$ , the conditional law of  $X - T$  is the same as the original law of  $X$ .
- ▶ Write  $p = \lambda/n$ . **Poisson random variable expectation** is  $\lim_{n \rightarrow \infty} np = \lim_{n \rightarrow \infty} n \frac{\lambda}{n} = \lambda$ . **Variance** is  $\lim_{n \rightarrow \infty} np(1 - p) = \lim_{n \rightarrow \infty} n(1 - \lambda/n)\lambda/n = \lambda$ .
- ▶ **Sum of  $\lambda_1$  Poisson and independent  $\lambda_2$  Poisson** is a  $\lambda_1 + \lambda_2$  Poisson.

## Continuous random variable properties derivable from coin toss intuition

- ▶ **Sum of  $n$  independent exponential random variables** each with parameter  $\lambda$  is gamma with parameters  $(n, \lambda)$ .
- ▶ **Memoryless properties:** given that exponential random variable  $X$  is greater than  $T > 0$ , the conditional law of  $X - T$  is the same as the original law of  $X$ .
- ▶ Write  $p = \lambda/n$ . **Poisson random variable expectation** is  $\lim_{n \rightarrow \infty} np = \lim_{n \rightarrow \infty} n \frac{\lambda}{n} = \lambda$ . **Variance** is  $\lim_{n \rightarrow \infty} np(1 - p) = \lim_{n \rightarrow \infty} n(1 - \lambda/n)\lambda/n = \lambda$ .
- ▶ **Sum of  $\lambda_1$  Poisson and independent  $\lambda_2$  Poisson** is a  $\lambda_1 + \lambda_2$  Poisson.
- ▶ **Times between successive events** in  $\lambda$  Poisson process are independent exponentials with parameter  $\lambda$ .



## Continuous random variable properties derivable from coin toss intuition

- ▶ **Sum of  $n$  independent exponential random variables** each with parameter  $\lambda$  is gamma with parameters  $(n, \lambda)$ .
- ▶ **Memoryless properties:** given that exponential random variable  $X$  is greater than  $T > 0$ , the conditional law of  $X - T$  is the same as the original law of  $X$ .
- ▶ Write  $p = \lambda/n$ . **Poisson random variable expectation** is  $\lim_{n \rightarrow \infty} np = \lim_{n \rightarrow \infty} n \frac{\lambda}{n} = \lambda$ . **Variance** is  $\lim_{n \rightarrow \infty} np(1 - p) = \lim_{n \rightarrow \infty} n(1 - \lambda/n)\lambda/n = \lambda$ .
- ▶ **Sum of  $\lambda_1$  Poisson and independent  $\lambda_2$  Poisson** is a  $\lambda_1 + \lambda_2$  Poisson.
- ▶ **Times between successive events** in  $\lambda$  Poisson process are independent exponentials with parameter  $\lambda$ .
- ▶ **Minimum of independent exponentials** with parameters  $\lambda_1$  and  $\lambda_2$  is itself exponential with parameter  $\lambda_1 + \lambda_2$ .

- ▶ **DeMoivre-Laplace limit theorem (special case of central limit theorem):**

$$\lim_{n \rightarrow \infty} P\left\{a \leq \frac{S_n - np}{\sqrt{npq}} \leq b\right\} \rightarrow \Phi(b) - \Phi(a).$$

# DeMoivre-Laplace Limit Theorem

- ▶ **DeMoivre-Laplace limit theorem (special case of central limit theorem):**

$$\lim_{n \rightarrow \infty} P\left\{a \leq \frac{S_n - np}{\sqrt{npq}} \leq b\right\} \rightarrow \Phi(b) - \Phi(a).$$

- ▶ This is  $\Phi(b) - \Phi(a) = P\{a \leq X \leq b\}$  when  $X$  is a standard normal random variable.

# Problems

- ▶ Toss a million fair coins. Approximate the probability that I get more than 501,000 heads.

# Problems

- ▶ Toss a million fair coins. Approximate the probability that I get more than 501,000 heads.
- ▶ Answer: well,  $\sqrt{npq} = \sqrt{10^6 \times .5 \times .5} = 500$ . So we're asking for probability to be over two SDs above mean. This is approximately  $1 - \Phi(2) = \Phi(-2)$ .

# Problems

- ▶ Toss a million fair coins. Approximate the probability that I get more than 501,000 heads.
- ▶ Answer: well,  $\sqrt{npq} = \sqrt{10^6 \times .5 \times .5} = 500$ . So we're asking for probability to be over two SDs above mean. This is approximately  $1 - \Phi(2) = \Phi(-2)$ .
- ▶ Roll 60000 dice. Expect to see 10000 sixes. What's the probability to see more than 9800?

# Problems

- ▶ Toss a million fair coins. Approximate the probability that I get more than 501,000 heads.
- ▶ Answer: well,  $\sqrt{npq} = \sqrt{10^6 \times .5 \times .5} = 500$ . So we're asking for probability to be over two SDs above mean. This is approximately  $1 - \Phi(2) = \Phi(-2)$ .
- ▶ Roll 60000 dice. Expect to see 10000 sixes. What's the probability to see more than 9800?
- ▶ Here  $\sqrt{npq} = \sqrt{60000 \times \frac{1}{6} \times \frac{5}{6}} \approx 91.28$ .

# Problems

- ▶ Toss a million fair coins. Approximate the probability that I get more than 501,000 heads.
- ▶ Answer: well,  $\sqrt{npq} = \sqrt{10^6 \times .5 \times .5} = 500$ . So we're asking for probability to be over two SDs above mean. This is approximately  $1 - \Phi(2) = \Phi(-2)$ .
- ▶ Roll 60000 dice. Expect to see 10000 sixes. What's the probability to see more than 9800?
- ▶ Here  $\sqrt{npq} = \sqrt{60000 \times \frac{1}{6} \times \frac{5}{6}} \approx 91.28$ .
- ▶ And  $200/91.28 \approx 2.19$ . Answer is about  $1 - \Phi(-2.19)$ .



# Properties of normal random variables

- ▶ Say  $X$  is a (standard) **normal random variable** if
$$f(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}.$$

# Properties of normal random variables

- ▶ Say  $X$  is a (standard) **normal random variable** if
$$f(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}.$$
- ▶ Mean zero and variance one.

# Properties of normal random variables

- ▶ Say  $X$  is a (standard) **normal random variable** if
$$f(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}.$$
- ▶ Mean zero and variance one.
- ▶ The random variable  $Y = \sigma X + \mu$  has variance  $\sigma^2$  and expectation  $\mu$ .

# Properties of normal random variables

- ▶ Say  $X$  is a (standard) **normal random variable** if  $f(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}$ .
- ▶ Mean zero and variance one.
- ▶ The random variable  $Y = \sigma X + \mu$  has variance  $\sigma^2$  and expectation  $\mu$ .
- ▶  $Y$  is said to be normal with parameters  $\mu$  and  $\sigma^2$ . Its density function is  $f_Y(x) = \frac{1}{\sqrt{2\pi\sigma}} e^{-(x-\mu)^2/2\sigma^2}$ .

# Properties of normal random variables

- ▶ Say  $X$  is a (standard) **normal random variable** if  $f(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}$ .
- ▶ Mean zero and variance one.
- ▶ The random variable  $Y = \sigma X + \mu$  has variance  $\sigma^2$  and expectation  $\mu$ .
- ▶  $Y$  is said to be normal with parameters  $\mu$  and  $\sigma^2$ . Its density function is  $f_Y(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-(x-\mu)^2/2\sigma^2}$ .
- ▶ Function  $\Phi(a) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^a e^{-x^2/2} dx$  can't be computed explicitly.

# Properties of normal random variables

- ▶ Say  $X$  is a (standard) **normal random variable** if  $f(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}$ .
- ▶ Mean zero and variance one.
- ▶ The random variable  $Y = \sigma X + \mu$  has variance  $\sigma^2$  and expectation  $\mu$ .
- ▶  $Y$  is said to be normal with parameters  $\mu$  and  $\sigma^2$ . Its density function is  $f_Y(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-(x-\mu)^2/2\sigma^2}$ .
- ▶ Function  $\Phi(a) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^a e^{-x^2/2} dx$  can't be computed explicitly.
- ▶ Values:  $\Phi(-3) \approx .0013$ ,  $\Phi(-2) \approx .023$  and  $\Phi(-1) \approx .159$ .

# Properties of normal random variables

- ▶ Say  $X$  is a (standard) **normal random variable** if  $f(x) = \frac{1}{\sqrt{2\pi}} e^{-x^2/2}$ .
- ▶ Mean zero and variance one.
- ▶ The random variable  $Y = \sigma X + \mu$  has variance  $\sigma^2$  and expectation  $\mu$ .
- ▶  $Y$  is said to be normal with parameters  $\mu$  and  $\sigma^2$ . Its density function is  $f_Y(x) = \frac{1}{\sqrt{2\pi}\sigma} e^{-(x-\mu)^2/2\sigma^2}$ .
- ▶ Function  $\Phi(a) = \frac{1}{\sqrt{2\pi}} \int_{-\infty}^a e^{-x^2/2} dx$  can't be computed explicitly.
- ▶ Values:  $\Phi(-3) \approx .0013$ ,  $\Phi(-2) \approx .023$  and  $\Phi(-1) \approx .159$ .
- ▶ Rule of thumb: "two thirds of time within one SD of mean, 95 percent of time within 2 SDs of mean."

## Properties of exponential random variables

- ▶ Say  $X$  is an **exponential random variable of parameter  $\lambda$**  when its probability distribution function is  $f(x) = \lambda e^{-\lambda x}$  for  $x \geq 0$  (and  $f(x) = 0$  if  $x < 0$ ).



## Properties of exponential random variables

- ▶ Say  $X$  is an **exponential random variable of parameter  $\lambda$**  when its probability distribution function is  $f(x) = \lambda e^{-\lambda x}$  for  $x \geq 0$  (and  $f(x) = 0$  if  $x < 0$ ).
- ▶ For  $a > 0$  have

$$F_X(a) = \int_0^a f(x) dx = \int_0^a \lambda e^{-\lambda x} dx = -e^{-\lambda x} \Big|_0^a = 1 - e^{-\lambda a}.$$

## Properties of exponential random variables

- ▶ Say  $X$  is an **exponential random variable of parameter  $\lambda$**  when its probability distribution function is  $f(x) = \lambda e^{-\lambda x}$  for  $x \geq 0$  (and  $f(x) = 0$  if  $x < 0$ ).
- ▶ For  $a > 0$  have

$$F_X(a) = \int_0^a f(x) dx = \int_0^a \lambda e^{-\lambda x} dx = -e^{-\lambda x} \Big|_0^a = 1 - e^{-\lambda a}.$$

- ▶ Thus  $P\{X < a\} = 1 - e^{-\lambda a}$  and  $P\{X > a\} = e^{-\lambda a}$ .

# Properties of exponential random variables

- ▶ Say  $X$  is an **exponential random variable of parameter  $\lambda$**  when its probability distribution function is  $f(x) = \lambda e^{-\lambda x}$  for  $x \geq 0$  (and  $f(x) = 0$  if  $x < 0$ ).
- ▶ For  $a > 0$  have

$$F_X(a) = \int_0^a f(x) dx = \int_0^a \lambda e^{-\lambda x} dx = -e^{-\lambda x} \Big|_0^a = 1 - e^{-\lambda a}.$$

- ▶ Thus  $P\{X < a\} = 1 - e^{-\lambda a}$  and  $P\{X > a\} = e^{-\lambda a}$ .
- ▶ Formula  $P\{X > a\} = e^{-\lambda a}$  is very important in practice.

# Properties of exponential random variables

- ▶ Say  $X$  is an **exponential random variable of parameter  $\lambda$**  when its probability distribution function is  $f(x) = \lambda e^{-\lambda x}$  for  $x \geq 0$  (and  $f(x) = 0$  if  $x < 0$ ).
- ▶ For  $a > 0$  have

$$F_X(a) = \int_0^a f(x) dx = \int_0^a \lambda e^{-\lambda x} dx = -e^{-\lambda x} \Big|_0^a = 1 - e^{-\lambda a}.$$

- ▶ Thus  $P\{X < a\} = 1 - e^{-\lambda a}$  and  $P\{X > a\} = e^{-\lambda a}$ .
- ▶ Formula  $P\{X > a\} = e^{-\lambda a}$  is very important in practice.
- ▶ Repeated integration by parts gives  $E[X^n] = n!/\lambda^n$ .

# Properties of exponential random variables

- ▶ Say  $X$  is an **exponential random variable of parameter  $\lambda$**  when its probability distribution function is  $f(x) = \lambda e^{-\lambda x}$  for  $x \geq 0$  (and  $f(x) = 0$  if  $x < 0$ ).

- ▶ For  $a > 0$  have

$$F_X(a) = \int_0^a f(x) dx = \int_0^a \lambda e^{-\lambda x} dx = -e^{-\lambda x} \Big|_0^a = 1 - e^{-\lambda a}.$$

- ▶ Thus  $P\{X < a\} = 1 - e^{-\lambda a}$  and  $P\{X > a\} = e^{-\lambda a}$ .
- ▶ Formula  $P\{X > a\} = e^{-\lambda a}$  is very important in practice.
- ▶ Repeated integration by parts gives  $E[X^n] = n!/\lambda^n$ .
- ▶ If  $\lambda = 1$ , then  $E[X^n] = n!$ . Value  $\Gamma(n) := E[X^{n-1}]$  defined for real  $n > 0$  and  $\Gamma(n) = (n-1)!$ .

## Defining $\Gamma$ distribution

- ▶ Say that random variable  $X$  has gamma distribution with parameters  $(\alpha, \lambda)$  if  $f_X(x) = \begin{cases} \frac{(\lambda x)^{\alpha-1} e^{-\lambda x} \lambda}{\Gamma(\alpha)} & x \geq 0 \\ 0 & x < 0 \end{cases}$ .

## Defining $\Gamma$ distribution

- ▶ Say that random variable  $X$  has gamma distribution with parameters  $(\alpha, \lambda)$  if  $f_X(x) = \begin{cases} \frac{(\lambda x)^{\alpha-1} e^{-\lambda x} \lambda}{\Gamma(\alpha)} & x \geq 0 \\ 0 & x < 0 \end{cases}$ .
- ▶ Same as exponential distribution when  $\alpha = 1$ . Otherwise, multiply by  $x^{\alpha-1}$  and divide by  $\Gamma(\alpha)$ . The fact that  $\Gamma(\alpha)$  is what you need to divide by to make the total integral one just follows from the definition of  $\Gamma$ .

## Defining $\Gamma$ distribution

- ▶ Say that random variable  $X$  has gamma distribution with parameters  $(\alpha, \lambda)$  if  $f_X(x) = \begin{cases} \frac{(\lambda x)^{\alpha-1} e^{-\lambda x} \lambda}{\Gamma(\alpha)} & x \geq 0 \\ 0 & x < 0 \end{cases}$ .
- ▶ Same as exponential distribution when  $\alpha = 1$ . Otherwise, multiply by  $x^{\alpha-1}$  and divide by  $\Gamma(\alpha)$ . The fact that  $\Gamma(\alpha)$  is what you need to divide by to make the total integral one just follows from the definition of  $\Gamma$ .
- ▶ Waiting time interpretation makes sense only for integer  $\alpha$ , but distribution is defined for general positive  $\alpha$ .



# Outline

Continuous random variables

Problems motivated by coin tossing

Random variable properties

# Outline

Continuous random variables

Problems motivated by coin tossing

Random variable properties

# Properties of uniform random variables

- ▶ Suppose  $X$  is a random variable with probability density

$$\text{function } f(x) = \begin{cases} \frac{1}{\beta - \alpha} & x \in [\alpha, \beta] \\ 0 & x \notin [\alpha, \beta]. \end{cases}$$

# Properties of uniform random variables

- ▶ Suppose  $X$  is a random variable with probability density function  $f(x) = \begin{cases} \frac{1}{\beta-\alpha} & x \in [\alpha, \beta] \\ 0 & x \notin [\alpha, \beta]. \end{cases}$
- ▶ Then  $E[X] = \frac{\alpha+\beta}{2}$ .

# Properties of uniform random variables

- ▶ Suppose  $X$  is a random variable with probability density function  $f(x) = \begin{cases} \frac{1}{\beta - \alpha} & x \in [\alpha, \beta] \\ 0 & x \notin [\alpha, \beta]. \end{cases}$
- ▶ Then  $E[X] = \frac{\alpha + \beta}{2}$ .
- ▶ And  $\text{Var}[X] = \text{Var}[(\beta - \alpha)Y + \alpha] = \text{Var}[(\beta - \alpha)Y] = (\beta - \alpha)^2 \text{Var}[Y] = (\beta - \alpha)^2 / 12$ .

## Distribution of function of random variable

- ▶ Suppose  $P\{X \leq a\} = F_X(a)$  is known for all  $a$ . Write  $Y = X^3$ . What is  $P\{Y \leq 27\}$ ?

## Distribution of function of random variable

- ▶ Suppose  $P\{X \leq a\} = F_X(a)$  is known for all  $a$ . Write  $Y = X^3$ . What is  $P\{Y \leq 27\}$ ?
- ▶ Answer: note that  $Y \leq 27$  if and only if  $X \leq 3$ . Hence  $P\{Y \leq 27\} = P\{X \leq 3\} = F_X(3)$ .

## Distribution of function of random variable

- ▶ Suppose  $P\{X \leq a\} = F_X(a)$  is known for all  $a$ . Write  $Y = X^3$ . What is  $P\{Y \leq 27\}$ ?
- ▶ Answer: note that  $Y \leq 27$  if and only if  $X \leq 3$ . Hence  $P\{Y \leq 27\} = P\{X \leq 3\} = F_X(3)$ .
- ▶ Generally  $F_Y(a) = P\{Y \leq a\} = P\{X \leq a^{1/3}\} = F_X(a^{1/3})$



## Distribution of function of random variable

- ▶ Suppose  $P\{X \leq a\} = F_X(a)$  is known for all  $a$ . Write  $Y = X^3$ . What is  $P\{Y \leq 27\}$ ?
- ▶ Answer: note that  $Y \leq 27$  if and only if  $X \leq 3$ . Hence  $P\{Y \leq 27\} = P\{X \leq 3\} = F_X(3)$ .
- ▶ Generally  $F_Y(a) = P\{Y \leq a\} = P\{X \leq a^{1/3}\} = F_X(a^{1/3})$
- ▶ This is a general principle. If  $X$  is a continuous random variable and  $g$  is a strictly increasing function of  $x$  and  $Y = g(X)$ , then  $F_Y(a) = F_X(g^{-1}(a))$ .

## Joint probability mass functions: discrete random variables

- ▶ If  $X$  and  $Y$  assume values in  $\{1, 2, \dots, n\}$  then we can view  $A_{i,j} = P\{X = i, Y = j\}$  as the entries of an  $n \times n$  matrix.

## Joint probability mass functions: discrete random variables

- ▶ If  $X$  and  $Y$  assume values in  $\{1, 2, \dots, n\}$  then we can view  $A_{i,j} = P\{X = i, Y = j\}$  as the entries of an  $n \times n$  matrix.
- ▶ Let's say I don't care about  $Y$ . I just want to know  $P\{X = i\}$ . How do I figure that out from the matrix?

## Joint probability mass functions: discrete random variables

- ▶ If  $X$  and  $Y$  assume values in  $\{1, 2, \dots, n\}$  then we can view  $A_{i,j} = P\{X = i, Y = j\}$  as the entries of an  $n \times n$  matrix.
- ▶ Let's say I don't care about  $Y$ . I just want to know  $P\{X = i\}$ . How do I figure that out from the matrix?
- ▶ Answer:  $P\{X = i\} = \sum_{j=1}^n A_{i,j}$ .

## Joint probability mass functions: discrete random variables

- ▶ If  $X$  and  $Y$  assume values in  $\{1, 2, \dots, n\}$  then we can view  $A_{i,j} = P\{X = i, Y = j\}$  as the entries of an  $n \times n$  matrix.
- ▶ Let's say I don't care about  $Y$ . I just want to know  $P\{X = i\}$ . How do I figure that out from the matrix?
- ▶ Answer:  $P\{X = i\} = \sum_{j=1}^n A_{i,j}$ .
- ▶ Similarly,  $P\{Y = j\} = \sum_{i=1}^n A_{i,j}$ .

## Joint probability mass functions: discrete random variables

- ▶ If  $X$  and  $Y$  assume values in  $\{1, 2, \dots, n\}$  then we can view  $A_{i,j} = P\{X = i, Y = j\}$  as the entries of an  $n \times n$  matrix.
- ▶ Let's say I don't care about  $Y$ . I just want to know  $P\{X = i\}$ . How do I figure that out from the matrix?
- ▶ Answer:  $P\{X = i\} = \sum_{j=1}^n A_{i,j}$ .
- ▶ Similarly,  $P\{Y = j\} = \sum_{i=1}^n A_{i,j}$ .
- ▶ In other words, the probability mass functions for  $X$  and  $Y$  are the row and column sums of  $A_{i,j}$ .

## Joint probability mass functions: discrete random variables

- ▶ If  $X$  and  $Y$  assume values in  $\{1, 2, \dots, n\}$  then we can view  $A_{i,j} = P\{X = i, Y = j\}$  as the entries of an  $n \times n$  matrix.
- ▶ Let's say I don't care about  $Y$ . I just want to know  $P\{X = i\}$ . How do I figure that out from the matrix?
- ▶ Answer:  $P\{X = i\} = \sum_{j=1}^n A_{i,j}$ .
- ▶ Similarly,  $P\{Y = j\} = \sum_{i=1}^n A_{i,j}$ .
- ▶ In other words, the probability mass functions for  $X$  and  $Y$  are the row and column sums of  $A_{i,j}$ .
- ▶ Given the joint distribution of  $X$  and  $Y$ , we sometimes call distribution of  $X$  (ignoring  $Y$ ) and distribution of  $Y$  (ignoring  $X$ ) the **marginal** distributions.

## Joint probability mass functions: discrete random variables

- ▶ If  $X$  and  $Y$  assume values in  $\{1, 2, \dots, n\}$  then we can view  $A_{i,j} = P\{X = i, Y = j\}$  as the entries of an  $n \times n$  matrix.
- ▶ Let's say I don't care about  $Y$ . I just want to know  $P\{X = i\}$ . How do I figure that out from the matrix?
- ▶ Answer:  $P\{X = i\} = \sum_{j=1}^n A_{i,j}$ .
- ▶ Similarly,  $P\{Y = j\} = \sum_{i=1}^n A_{i,j}$ .
- ▶ In other words, the probability mass functions for  $X$  and  $Y$  are the row and column sums of  $A_{i,j}$ .
- ▶ Given the joint distribution of  $X$  and  $Y$ , we sometimes call distribution of  $X$  (ignoring  $Y$ ) and distribution of  $Y$  (ignoring  $X$ ) the **marginal** distributions.
- ▶ In general, when  $X$  and  $Y$  are jointly defined discrete random variables, we write  $p(x, y) = p_{X,Y}(x, y) = P\{X = x, Y = y\}$ .



## Joint distribution functions: continuous random variables

- ▶ Given random variables  $X$  and  $Y$ , define  $F(a, b) = P\{X \leq a, Y \leq b\}$ .

## Joint distribution functions: continuous random variables

- ▶ Given random variables  $X$  and  $Y$ , define  $F(a, b) = P\{X \leq a, Y \leq b\}$ .
- ▶ The region  $\{(x, y) : x \leq a, y \leq b\}$  is the lower left “quadrant” centered at  $(a, b)$ .

## Joint distribution functions: continuous random variables

- ▶ Given random variables  $X$  and  $Y$ , define  $F(a, b) = P\{X \leq a, Y \leq b\}$ .
- ▶ The region  $\{(x, y) : x \leq a, y \leq b\}$  is the lower left “quadrant” centered at  $(a, b)$ .
- ▶ Refer to  $F_X(a) = P\{X \leq a\}$  and  $F_Y(b) = P\{Y \leq b\}$  as **marginal** cumulative distribution functions.

## Joint distribution functions: continuous random variables

- ▶ Given random variables  $X$  and  $Y$ , define  $F(a, b) = P\{X \leq a, Y \leq b\}$ .
- ▶ The region  $\{(x, y) : x \leq a, y \leq b\}$  is the lower left “quadrant” centered at  $(a, b)$ .
- ▶ Refer to  $F_X(a) = P\{X \leq a\}$  and  $F_Y(b) = P\{Y \leq b\}$  as **marginal** cumulative distribution functions.
- ▶ Question: if I tell you the two parameter function  $F$ , can you use it to determine the marginals  $F_X$  and  $F_Y$ ?

## Joint distribution functions: continuous random variables

- ▶ Given random variables  $X$  and  $Y$ , define  $F(a, b) = P\{X \leq a, Y \leq b\}$ .
- ▶ The region  $\{(x, y) : x \leq a, y \leq b\}$  is the lower left “quadrant” centered at  $(a, b)$ .
- ▶ Refer to  $F_X(a) = P\{X \leq a\}$  and  $F_Y(b) = P\{Y \leq b\}$  as **marginal** cumulative distribution functions.
- ▶ Question: if I tell you the two parameter function  $F$ , can you use it to determine the marginals  $F_X$  and  $F_Y$ ?
- ▶ Answer: Yes.  $F_X(a) = \lim_{b \rightarrow \infty} F(a, b)$  and  $F_Y(b) = \lim_{a \rightarrow \infty} F(a, b)$ .

## Joint distribution functions: continuous random variables

- ▶ Given random variables  $X$  and  $Y$ , define  $F(a, b) = P\{X \leq a, Y \leq b\}$ .
- ▶ The region  $\{(x, y) : x \leq a, y \leq b\}$  is the lower left “quadrant” centered at  $(a, b)$ .
- ▶ Refer to  $F_X(a) = P\{X \leq a\}$  and  $F_Y(b) = P\{Y \leq b\}$  as **marginal** cumulative distribution functions.
- ▶ Question: if I tell you the two parameter function  $F$ , can you use it to determine the marginals  $F_X$  and  $F_Y$ ?
- ▶ Answer: Yes.  $F_X(a) = \lim_{b \rightarrow \infty} F(a, b)$  and  $F_Y(b) = \lim_{a \rightarrow \infty} F(a, b)$ .
- ▶ Density:  $f(x, y) = \frac{\partial}{\partial x} \frac{\partial}{\partial y} F(x, y)$ .

# Independent random variables

- ▶ We say  $X$  and  $Y$  are independent if for any two (measurable) sets  $A$  and  $B$  of real numbers we have

$$P\{X \in A, Y \in B\} = P\{X \in A\}P\{Y \in B\}.$$

# Independent random variables

- ▶ We say  $X$  and  $Y$  are independent if for any two (measurable) sets  $A$  and  $B$  of real numbers we have

$$P\{X \in A, Y \in B\} = P\{X \in A\}P\{Y \in B\}.$$

- ▶ When  $X$  and  $Y$  are discrete random variables, they are independent if  $P\{X = x, Y = y\} = P\{X = x\}P\{Y = y\}$  for all  $x$  and  $y$  for which  $P\{X = x\}$  and  $P\{Y = y\}$  are non-zero.



# Independent random variables

- ▶ We say  $X$  and  $Y$  are independent if for any two (measurable) sets  $A$  and  $B$  of real numbers we have

$$P\{X \in A, Y \in B\} = P\{X \in A\}P\{Y \in B\}.$$

- ▶ When  $X$  and  $Y$  are discrete random variables, they are independent if  $P\{X = x, Y = y\} = P\{X = x\}P\{Y = y\}$  for all  $x$  and  $y$  for which  $P\{X = x\}$  and  $P\{Y = y\}$  are non-zero.
- ▶ When  $X$  and  $Y$  are continuous, they are independent if  $f(x, y) = f_X(x)f_Y(y)$ .

## Summing two random variables

- ▶ Say we have independent random variables  $X$  and  $Y$  and we know their density functions  $f_X$  and  $f_Y$ .

## Summing two random variables

- ▶ Say we have independent random variables  $X$  and  $Y$  and we know their density functions  $f_X$  and  $f_Y$ .
- ▶ Now let's try to find  $F_{X+Y}(a) = P\{X + Y \leq a\}$ .

## Summing two random variables

- ▶ Say we have independent random variables  $X$  and  $Y$  and we know their density functions  $f_X$  and  $f_Y$ .
- ▶ Now let's try to find  $F_{X+Y}(a) = P\{X + Y \leq a\}$ .
- ▶ This is the integral over  $\{(x, y) : x + y \leq a\}$  of  $f(x, y) = f_X(x)f_Y(y)$ . Thus,

## Summing two random variables

- ▶ Say we have independent random variables  $X$  and  $Y$  and we know their density functions  $f_X$  and  $f_Y$ .
- ▶ Now let's try to find  $F_{X+Y}(a) = P\{X + Y \leq a\}$ .
- ▶ This is the integral over  $\{(x, y) : x + y \leq a\}$  of  $f(x, y) = f_X(x)f_Y(y)$ . Thus,



$$\begin{aligned} P\{X + Y \leq a\} &= \int_{-\infty}^{\infty} \int_{-\infty}^{a-y} f_X(x)f_Y(y)dx dy \\ &= \int_{-\infty}^{\infty} F_X(a - y)f_Y(y)dy. \end{aligned}$$

## Summing two random variables

- ▶ Say we have independent random variables  $X$  and  $Y$  and we know their density functions  $f_X$  and  $f_Y$ .
- ▶ Now let's try to find  $F_{X+Y}(a) = P\{X + Y \leq a\}$ .
- ▶ This is the integral over  $\{(x, y) : x + y \leq a\}$  of  $f(x, y) = f_X(x)f_Y(y)$ . Thus,



$$\begin{aligned}P\{X + Y \leq a\} &= \int_{-\infty}^{\infty} \int_{-\infty}^{a-y} f_X(x)f_Y(y)dx dy \\ &= \int_{-\infty}^{\infty} F_X(a-y)f_Y(y)dy.\end{aligned}$$

- ▶ Differentiating both sides gives  $f_{X+Y}(a) = \frac{d}{da} \int_{-\infty}^{\infty} F_X(a-y)f_Y(y)dy = \int_{-\infty}^{\infty} f_X(a-y)f_Y(y)dy$ .

## Summing two random variables

- ▶ Say we have independent random variables  $X$  and  $Y$  and we know their density functions  $f_X$  and  $f_Y$ .
- ▶ Now let's try to find  $F_{X+Y}(a) = P\{X + Y \leq a\}$ .
- ▶ This is the integral over  $\{(x, y) : x + y \leq a\}$  of  $f(x, y) = f_X(x)f_Y(y)$ . Thus,



$$\begin{aligned}P\{X + Y \leq a\} &= \int_{-\infty}^{\infty} \int_{-\infty}^{a-y} f_X(x)f_Y(y)dx dy \\ &= \int_{-\infty}^{\infty} F_X(a-y)f_Y(y)dy.\end{aligned}$$

- ▶ Differentiating both sides gives  $f_{X+Y}(a) = \frac{d}{da} \int_{-\infty}^{\infty} F_X(a-y)f_Y(y)dy = \int_{-\infty}^{\infty} f_X(a-y)f_Y(y)dy$ .
- ▶ Latter formula makes some intuitive sense. We're integrating over the set of  $x, y$  pairs that add up to  $a$ .

# Conditional distributions

- ▶ Let's say  $X$  and  $Y$  have joint probability density function  $f(x, y)$ .



# Conditional distributions

- ▶ Let's say  $X$  and  $Y$  have joint probability density function  $f(x, y)$ .
- ▶ We can *define* the conditional probability density of  $X$  given that  $Y = y$  by  $f_{X|Y=y}(x) = \frac{f(x,y)}{f_Y(y)}$ .

# Conditional distributions

- ▶ Let's say  $X$  and  $Y$  have joint probability density function  $f(x, y)$ .
- ▶ We can *define* the conditional probability density of  $X$  given that  $Y = y$  by  $f_{X|Y=y}(x) = \frac{f(x,y)}{f_Y(y)}$ .
- ▶ This amounts to restricting  $f(x, y)$  to the line corresponding to the given  $y$  value (and dividing by the constant that makes the integral along that line equal to 1).

## Maxima: pick five job candidates at random, choose best

- ▶ Suppose I choose  $n$  random variables  $X_1, X_2, \dots, X_n$  uniformly at random on  $[0, 1]$ , independently of each other.

## Maxima: pick five job candidates at random, choose best

- ▶ Suppose I choose  $n$  random variables  $X_1, X_2, \dots, X_n$  uniformly at random on  $[0, 1]$ , independently of each other.
- ▶ The  $n$ -tuple  $(X_1, X_2, \dots, X_n)$  has a constant density function on the  $n$ -dimensional cube  $[0, 1]^n$ .

## Maxima: pick five job candidates at random, choose best

- ▶ Suppose I choose  $n$  random variables  $X_1, X_2, \dots, X_n$  uniformly at random on  $[0, 1]$ , independently of each other.
- ▶ The  $n$ -tuple  $(X_1, X_2, \dots, X_n)$  has a constant density function on the  $n$ -dimensional cube  $[0, 1]^n$ .
- ▶ What is the probability that the *largest* of the  $X_i$  is less than  $a$ ?

## Maxima: pick five job candidates at random, choose best

- ▶ Suppose I choose  $n$  random variables  $X_1, X_2, \dots, X_n$  uniformly at random on  $[0, 1]$ , independently of each other.
- ▶ The  $n$ -tuple  $(X_1, X_2, \dots, X_n)$  has a constant density function on the  $n$ -dimensional cube  $[0, 1]^n$ .
- ▶ What is the probability that the *largest* of the  $X_i$  is less than  $a$ ?
- ▶ ANSWER:  $a^n$ .

## Maxima: pick five job candidates at random, choose best

- ▶ Suppose I choose  $n$  random variables  $X_1, X_2, \dots, X_n$  uniformly at random on  $[0, 1]$ , independently of each other.
- ▶ The  $n$ -tuple  $(X_1, X_2, \dots, X_n)$  has a constant density function on the  $n$ -dimensional cube  $[0, 1]^n$ .
- ▶ What is the probability that the *largest* of the  $X_i$  is less than  $a$ ?
- ▶ ANSWER:  $a^n$ .
- ▶ So if  $X = \max\{X_1, \dots, X_n\}$ , then what is the probability density function of  $X$ ?

## Maxima: pick five job candidates at random, choose best

- ▶ Suppose I choose  $n$  random variables  $X_1, X_2, \dots, X_n$  uniformly at random on  $[0, 1]$ , independently of each other.
- ▶ The  $n$ -tuple  $(X_1, X_2, \dots, X_n)$  has a constant density function on the  $n$ -dimensional cube  $[0, 1]^n$ .
- ▶ What is the probability that the *largest* of the  $X_i$  is less than  $a$ ?
- ▶ ANSWER:  $a^n$ .
- ▶ So if  $X = \max\{X_1, \dots, X_n\}$ , then what is the probability density function of  $X$ ?

▶ Answer:  $F_X(a) = \begin{cases} 0 & a < 0 \\ a^n & a \in [0, 1] \\ 1 & a > 1 \end{cases}$ . And

$$f_X(a) = F'_X(a) = na^{n-1}.$$



## General order statistics

- ▶ Consider i.i.d random variables  $X_1, X_2, \dots, X_n$  with continuous probability density  $f$ .

## General order statistics

- ▶ Consider i.i.d random variables  $X_1, X_2, \dots, X_n$  with continuous probability density  $f$ .
- ▶ Let  $Y_1 < Y_2 < Y_3 \dots < Y_n$  be list obtained by *sorting* the  $X_j$ .

## General order statistics

- ▶ Consider i.i.d random variables  $X_1, X_2, \dots, X_n$  with continuous probability density  $f$ .
- ▶ Let  $Y_1 < Y_2 < Y_3 \dots < Y_n$  be list obtained by *sorting* the  $X_j$ .
- ▶ In particular,  $Y_1 = \min\{X_1, \dots, X_n\}$  and  $Y_n = \max\{X_1, \dots, X_n\}$  is the maximum.

## General order statistics

- ▶ Consider i.i.d random variables  $X_1, X_2, \dots, X_n$  with continuous probability density  $f$ .
- ▶ Let  $Y_1 < Y_2 < Y_3 \dots < Y_n$  be list obtained by *sorting* the  $X_j$ .
- ▶ In particular,  $Y_1 = \min\{X_1, \dots, X_n\}$  and  $Y_n = \max\{X_1, \dots, X_n\}$  is the maximum.
- ▶ What is the joint probability density of the  $Y_i$ ?

## General order statistics

- ▶ Consider i.i.d random variables  $X_1, X_2, \dots, X_n$  with continuous probability density  $f$ .
- ▶ Let  $Y_1 < Y_2 < Y_3 \dots < Y_n$  be list obtained by *sorting* the  $X_j$ .
- ▶ In particular,  $Y_1 = \min\{X_1, \dots, X_n\}$  and  $Y_n = \max\{X_1, \dots, X_n\}$  is the maximum.
- ▶ What is the joint probability density of the  $Y_i$ ?
- ▶ Answer:  $f(x_1, x_2, \dots, x_n) = n! \prod_{i=1}^n f(x_i)$  if  $x_1 < x_2 \dots < x_n$ , zero otherwise.

## General order statistics

- ▶ Consider i.i.d random variables  $X_1, X_2, \dots, X_n$  with continuous probability density  $f$ .
- ▶ Let  $Y_1 < Y_2 < Y_3 \dots < Y_n$  be list obtained by *sorting* the  $X_j$ .
- ▶ In particular,  $Y_1 = \min\{X_1, \dots, X_n\}$  and  $Y_n = \max\{X_1, \dots, X_n\}$  is the maximum.
- ▶ What is the joint probability density of the  $Y_i$ ?
- ▶ Answer:  $f(x_1, x_2, \dots, x_n) = n! \prod_{i=1}^n f(x_i)$  if  $x_1 < x_2 \dots < x_n$ , zero otherwise.
- ▶ Let  $\sigma : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  be the permutation such that  $X_j = Y_{\sigma(j)}$

## General order statistics

- ▶ Consider i.i.d random variables  $X_1, X_2, \dots, X_n$  with continuous probability density  $f$ .
- ▶ Let  $Y_1 < Y_2 < Y_3 \dots < Y_n$  be list obtained by *sorting* the  $X_j$ .
- ▶ In particular,  $Y_1 = \min\{X_1, \dots, X_n\}$  and  $Y_n = \max\{X_1, \dots, X_n\}$  is the maximum.
- ▶ What is the joint probability density of the  $Y_i$ ?
- ▶ Answer:  $f(x_1, x_2, \dots, x_n) = n! \prod_{i=1}^n f(x_i)$  if  $x_1 < x_2 \dots < x_n$ , zero otherwise.
- ▶ Let  $\sigma : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  be the permutation such that  $X_j = Y_{\sigma(j)}$
- ▶ Are  $\sigma$  and the vector  $(Y_1, \dots, Y_n)$  independent of each other?

## General order statistics

- ▶ Consider i.i.d random variables  $X_1, X_2, \dots, X_n$  with continuous probability density  $f$ .
- ▶ Let  $Y_1 < Y_2 < Y_3 \dots < Y_n$  be list obtained by *sorting* the  $X_j$ .
- ▶ In particular,  $Y_1 = \min\{X_1, \dots, X_n\}$  and  $Y_n = \max\{X_1, \dots, X_n\}$  is the maximum.
- ▶ What is the joint probability density of the  $Y_i$ ?
- ▶ Answer:  $f(x_1, x_2, \dots, x_n) = n! \prod_{i=1}^n f(x_i)$  if  $x_1 < x_2 \dots < x_n$ , zero otherwise.
- ▶ Let  $\sigma : \{1, 2, \dots, n\} \rightarrow \{1, 2, \dots, n\}$  be the permutation such that  $X_j = Y_{\sigma(j)}$
- ▶ Are  $\sigma$  and the vector  $(Y_1, \dots, Y_n)$  independent of each other?
- ▶ Yes.



# Properties of expectation

- ▶ Several properties we derived for discrete expectations continue to hold in the continuum.

# Properties of expectation

- ▶ Several properties we derived for discrete expectations continue to hold in the continuum.
- ▶ If  $X$  is discrete with mass function  $p(x)$  then  $E[X] = \sum_x p(x)x$ .

## Properties of expectation

- ▶ Several properties we derived for discrete expectations continue to hold in the continuum.
- ▶ If  $X$  is discrete with mass function  $p(x)$  then 
$$E[X] = \sum_x p(x)x.$$
- ▶ Similarly, if  $X$  is continuous with density function  $f(x)$  then 
$$E[X] = \int f(x)x dx.$$

# Properties of expectation

- ▶ Several properties we derived for discrete expectations continue to hold in the continuum.
- ▶ If  $X$  is discrete with mass function  $p(x)$  then 
$$E[X] = \sum_x p(x)x.$$
- ▶ Similarly, if  $X$  is continuous with density function  $f(x)$  then 
$$E[X] = \int f(x)x dx.$$
- ▶ If  $X$  is discrete with mass function  $p(x)$  then 
$$E[g(x)] = \sum_x p(x)g(x).$$

## Properties of expectation

- ▶ Several properties we derived for discrete expectations continue to hold in the continuum.
- ▶ If  $X$  is discrete with mass function  $p(x)$  then
$$E[X] = \sum_x p(x)x.$$
- ▶ Similarly, if  $X$  is continuous with density function  $f(x)$  then
$$E[X] = \int f(x)x dx.$$
- ▶ If  $X$  is discrete with mass function  $p(x)$  then
$$E[g(x)] = \sum_x p(x)g(x).$$
- ▶ Similarly,  $X$  if is continuous with density function  $f(x)$  then
$$E[g(X)] = \int f(x)g(x) dx.$$

# Properties of expectation

- ▶ Several properties we derived for discrete expectations continue to hold in the continuum.
- ▶ If  $X$  is discrete with mass function  $p(x)$  then
$$E[X] = \sum_x p(x)x.$$
- ▶ Similarly, if  $X$  is continuous with density function  $f(x)$  then
$$E[X] = \int f(x)x dx.$$
- ▶ If  $X$  is discrete with mass function  $p(x)$  then
$$E[g(x)] = \sum_x p(x)g(x).$$
- ▶ Similarly,  $X$  if is continuous with density function  $f(x)$  then
$$E[g(X)] = \int f(x)g(x) dx.$$
- ▶ If  $X$  and  $Y$  have joint mass function  $p(x, y)$  then
$$E[g(X, Y)] = \sum_y \sum_x g(x, y)p(x, y).$$

## Properties of expectation

- ▶ Several properties we derived for discrete expectations continue to hold in the continuum.
- ▶ If  $X$  is discrete with mass function  $p(x)$  then
$$E[X] = \sum_x p(x)x.$$
- ▶ Similarly, if  $X$  is continuous with density function  $f(x)$  then
$$E[X] = \int f(x)x dx.$$
- ▶ If  $X$  is discrete with mass function  $p(x)$  then
$$E[g(x)] = \sum_x p(x)g(x).$$
- ▶ Similarly,  $X$  if is continuous with density function  $f(x)$  then
$$E[g(X)] = \int f(x)g(x) dx.$$
- ▶ If  $X$  and  $Y$  have joint mass function  $p(x, y)$  then
$$E[g(X, Y)] = \sum_y \sum_x g(x, y)p(x, y).$$
- ▶ If  $X$  and  $Y$  have joint probability density function  $f(x, y)$  then
$$E[g(X, Y)] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x, y)f(x, y) dx dy.$$

# Properties of expectation

- ▶ For both discrete and continuous random variables  $X$  and  $Y$  we have  $E[X + Y] = E[X] + E[Y]$ .



# Properties of expectation

- ▶ For both discrete and continuous random variables  $X$  and  $Y$  we have  $E[X + Y] = E[X] + E[Y]$ .
- ▶ In both discrete and continuous settings,  $E[aX] = aE[X]$  when  $a$  is a constant. And  $E[\sum a_i X_i] = \sum a_i E[X_i]$ .

# Properties of expectation

- ▶ For both discrete and continuous random variables  $X$  and  $Y$  we have  $E[X + Y] = E[X] + E[Y]$ .
- ▶ In both discrete and continuous settings,  $E[aX] = aE[X]$  when  $a$  is a constant. And  $E[\sum a_i X_i] = \sum a_i E[X_i]$ .
- ▶ But what about that delightful “area under  $1 - F_X$ ” formula for the expectation?

# Properties of expectation

- ▶ For both discrete and continuous random variables  $X$  and  $Y$  we have  $E[X + Y] = E[X] + E[Y]$ .
- ▶ In both discrete and continuous settings,  $E[aX] = aE[X]$  when  $a$  is a constant. And  $E[\sum a_i X_i] = \sum a_i E[X_i]$ .
- ▶ But what about that delightful “area under  $1 - F_X$ ” formula for the expectation?
- ▶ When  $X$  is non-negative with probability one, do we always have  $E[X] = \int_0^\infty P\{X > x\}$ , in both discrete and continuous settings?

# Properties of expectation

- ▶ For both discrete and continuous random variables  $X$  and  $Y$  we have  $E[X + Y] = E[X] + E[Y]$ .
- ▶ In both discrete and continuous settings,  $E[aX] = aE[X]$  when  $a$  is a constant. And  $E[\sum a_i X_i] = \sum a_i E[X_i]$ .
- ▶ But what about that delightful “area under  $1 - F_X$ ” formula for the expectation?
- ▶ When  $X$  is non-negative with probability one, do we always have  $E[X] = \int_0^\infty P\{X > x\}$ , in both discrete and continuous settings?
- ▶ Define  $g(y)$  so that  $1 - F_X(g(y)) = y$ . (Draw horizontal line at height  $y$  and look where it hits graph of  $1 - F_X$ .)

## Properties of expectation

- ▶ For both discrete and continuous random variables  $X$  and  $Y$  we have  $E[X + Y] = E[X] + E[Y]$ .
- ▶ In both discrete and continuous settings,  $E[aX] = aE[X]$  when  $a$  is a constant. And  $E[\sum a_i X_i] = \sum a_i E[X_i]$ .
- ▶ But what about that delightful “area under  $1 - F_X$ ” formula for the expectation?
- ▶ When  $X$  is non-negative with probability one, do we always have  $E[X] = \int_0^\infty P\{X > x\}$ , in both discrete and continuous settings?
- ▶ Define  $g(y)$  so that  $1 - F_X(g(y)) = y$ . (Draw horizontal line at height  $y$  and look where it hits graph of  $1 - F_X$ .)
- ▶ Choose  $Y$  uniformly on  $[0, 1]$  and note that  $g(Y)$  has the same probability distribution as  $X$ .

# Properties of expectation

- ▶ For both discrete and continuous random variables  $X$  and  $Y$  we have  $E[X + Y] = E[X] + E[Y]$ .
- ▶ In both discrete and continuous settings,  $E[aX] = aE[X]$  when  $a$  is a constant. And  $E[\sum a_i X_i] = \sum a_i E[X_i]$ .
- ▶ But what about that delightful “area under  $1 - F_X$ ” formula for the expectation?
- ▶ When  $X$  is non-negative with probability one, do we always have  $E[X] = \int_0^\infty P\{X > x\}$ , in both discrete and continuous settings?
- ▶ Define  $g(y)$  so that  $1 - F_X(g(y)) = y$ . (Draw horizontal line at height  $y$  and look where it hits graph of  $1 - F_X$ .)
- ▶ Choose  $Y$  uniformly on  $[0, 1]$  and note that  $g(Y)$  has the same probability distribution as  $X$ .
- ▶ So  $E[X] = E[g(Y)] = \int_0^1 g(y) dy$ , which is indeed the area under the graph of  $1 - F_X$ .

## A property of independence

- ▶ If  $X$  and  $Y$  are independent then  
 $E[g(X)h(Y)] = E[g(X)]E[h(Y)]$ .

## A property of independence

- ▶ If  $X$  and  $Y$  are independent then  $E[g(X)h(Y)] = E[g(X)]E[h(Y)]$ .
- ▶ Just write  $E[g(X)h(Y)] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x)h(y)f(x, y)dx dy$ .



## A property of independence

- ▶ If  $X$  and  $Y$  are independent then  $E[g(X)h(Y)] = E[g(X)]E[h(Y)]$ .
- ▶ Just write  $E[g(X)h(Y)] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x)h(y)f(x, y)dx dy$ .
- ▶ Since  $f(x, y) = f_X(x)f_Y(y)$  this factors as  $\int_{-\infty}^{\infty} h(y)f_Y(y)dy \int_{-\infty}^{\infty} g(x)f_X(x)dx = E[h(Y)]E[g(X)]$ .

## Defining covariance and correlation

- ▶ Now define covariance of  $X$  and  $Y$  by  
$$\text{Cov}(X, Y) = E[(X - E[X])(Y - E[Y])].$$

## Defining covariance and correlation

- ▶ Now define covariance of  $X$  and  $Y$  by
$$\text{Cov}(X, Y) = E[(X - E[X])(Y - E[Y])].$$
- ▶ Note: by definition  $\text{Var}(X) = \text{Cov}(X, X)$ .

## Defining covariance and correlation

- ▶ Now define covariance of  $X$  and  $Y$  by 
$$\text{Cov}(X, Y) = E[(X - E[X])(Y - E[Y])].$$
- ▶ Note: by definition  $\text{Var}(X) = \text{Cov}(X, X)$ .
- ▶ Covariance formula  $E[XY] - E[X]E[Y]$ , or “expectation of product minus product of expectations” is frequently useful.

## Defining covariance and correlation

- ▶ Now define covariance of  $X$  and  $Y$  by 
$$\text{Cov}(X, Y) = E[(X - E[X])(Y - E[Y])].$$
- ▶ Note: by definition  $\text{Var}(X) = \text{Cov}(X, X)$ .
- ▶ Covariance formula  $E[XY] - E[X]E[Y]$ , or “expectation of product minus product of expectations” is frequently useful.
- ▶ If  $X$  and  $Y$  are independent then  $\text{Cov}(X, Y) = 0$ .

## Defining covariance and correlation

- ▶ Now define covariance of  $X$  and  $Y$  by 
$$\text{Cov}(X, Y) = E[(X - E[X])(Y - E[Y])].$$
- ▶ Note: by definition  $\text{Var}(X) = \text{Cov}(X, X)$ .
- ▶ Covariance formula  $E[XY] - E[X]E[Y]$ , or “expectation of product minus product of expectations” is frequently useful.
- ▶ If  $X$  and  $Y$  are independent then  $\text{Cov}(X, Y) = 0$ .
- ▶ Converse is not true.

## Basic covariance facts

- ▶  $\text{Cov}(X, Y) = \text{Cov}(Y, X)$

## Basic covariance facts

- ▶  $\text{Cov}(X, Y) = \text{Cov}(Y, X)$
- ▶  $\text{Cov}(X, X) = \text{Var}(X)$



## Basic covariance facts

- ▶  $\text{Cov}(X, Y) = \text{Cov}(Y, X)$
- ▶  $\text{Cov}(X, X) = \text{Var}(X)$
- ▶  $\text{Cov}(aX, Y) = a\text{Cov}(X, Y)$ .

## Basic covariance facts

- ▶  $\text{Cov}(X, Y) = \text{Cov}(Y, X)$
- ▶  $\text{Cov}(X, X) = \text{Var}(X)$
- ▶  $\text{Cov}(aX, Y) = a\text{Cov}(X, Y)$ .
- ▶  $\text{Cov}(X_1 + X_2, Y) = \text{Cov}(X_1, Y) + \text{Cov}(X_2, Y)$ .

## Basic covariance facts

- ▶  $\text{Cov}(X, Y) = \text{Cov}(Y, X)$
- ▶  $\text{Cov}(X, X) = \text{Var}(X)$
- ▶  $\text{Cov}(aX, Y) = a\text{Cov}(X, Y)$ .
- ▶  $\text{Cov}(X_1 + X_2, Y) = \text{Cov}(X_1, Y) + \text{Cov}(X_2, Y)$ .
- ▶ **General statement of bilinearity of covariance:**

$$\text{Cov}\left(\sum_{i=1}^m a_i X_i, \sum_{j=1}^n b_j Y_j\right) = \sum_{i=1}^m \sum_{j=1}^n a_i b_j \text{Cov}(X_i, Y_j).$$

## Basic covariance facts

- ▶  $\text{Cov}(X, Y) = \text{Cov}(Y, X)$
- ▶  $\text{Cov}(X, X) = \text{Var}(X)$
- ▶  $\text{Cov}(aX, Y) = a\text{Cov}(X, Y)$ .
- ▶  $\text{Cov}(X_1 + X_2, Y) = \text{Cov}(X_1, Y) + \text{Cov}(X_2, Y)$ .
- ▶ **General statement of bilinearity of covariance:**

$$\text{Cov}\left(\sum_{i=1}^m a_i X_i, \sum_{j=1}^n b_j Y_j\right) = \sum_{i=1}^m \sum_{j=1}^n a_i b_j \text{Cov}(X_i, Y_j).$$

- ▶ Special case:

$$\text{Var}\left(\sum_{i=1}^n X_i\right) = \sum_{i=1}^n \text{Var}(X_i) + 2 \sum_{(i,j): i < j} \text{Cov}(X_i, X_j).$$

## Defining correlation

- ▶ Again, by definition  $\text{Cov}(X, Y) = E[XY] - E[X]E[Y]$ .

## Defining correlation

- ▶ Again, by definition  $\text{Cov}(X, Y) = E[XY] - E[X]E[Y]$ .
- ▶ **Correlation** of  $X$  and  $Y$  defined by

$$\rho(X, Y) := \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}}.$$

## Defining correlation

- ▶ Again, by definition  $\text{Cov}(X, Y) = E[XY] - E[X]E[Y]$ .
- ▶ **Correlation** of  $X$  and  $Y$  defined by

$$\rho(X, Y) := \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}}.$$

- ▶ Correlation doesn't care what units you use for  $X$  and  $Y$ . If  $a > 0$  and  $c > 0$  then  $\rho(aX + b, cY + d) = \rho(X, Y)$ .

## Defining correlation

- ▶ Again, by definition  $\text{Cov}(X, Y) = E[XY] - E[X]E[Y]$ .
- ▶ **Correlation** of  $X$  and  $Y$  defined by

$$\rho(X, Y) := \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}}.$$

- ▶ Correlation doesn't care what units you use for  $X$  and  $Y$ . If  $a > 0$  and  $c > 0$  then  $\rho(aX + b, cY + d) = \rho(X, Y)$ .
- ▶ Satisfies  $-1 \leq \rho(X, Y) \leq 1$ .



## Defining correlation

- ▶ Again, by definition  $\text{Cov}(X, Y) = E[XY] - E[X]E[Y]$ .
- ▶ **Correlation** of  $X$  and  $Y$  defined by

$$\rho(X, Y) := \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}}.$$

- ▶ Correlation doesn't care what units you use for  $X$  and  $Y$ . If  $a > 0$  and  $c > 0$  then  $\rho(aX + b, cY + d) = \rho(X, Y)$ .
- ▶ Satisfies  $-1 \leq \rho(X, Y) \leq 1$ .
- ▶ If  $a$  and  $b$  are positive constants and  $a > 0$  then  $\rho(aX + b, X) = 1$ .

## Defining correlation

- ▶ Again, by definition  $\text{Cov}(X, Y) = E[XY] - E[X]E[Y]$ .
- ▶ **Correlation** of  $X$  and  $Y$  defined by

$$\rho(X, Y) := \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}}.$$

- ▶ Correlation doesn't care what units you use for  $X$  and  $Y$ . If  $a > 0$  and  $c > 0$  then  $\rho(aX + b, cY + d) = \rho(X, Y)$ .
- ▶ Satisfies  $-1 \leq \rho(X, Y) \leq 1$ .
- ▶ If  $a$  and  $b$  are positive constants and  $a > 0$  then  $\rho(aX + b, X) = 1$ .
- ▶ If  $a$  and  $b$  are positive constants and  $a < 0$  then  $\rho(aX + b, X) = -1$ .

## Conditional probability distributions

- ▶ It all starts with the definition of conditional probability:  
 $P(A|B) = P(AB)/P(B)$ .

## Conditional probability distributions

- ▶ It all starts with the definition of conditional probability:  
 $P(A|B) = P(AB)/P(B)$ .
- ▶ If  $X$  and  $Y$  are jointly discrete random variables, we can use this to define a probability mass function for  $X$  *given*  $Y = y$ .

## Conditional probability distributions

- ▶ It all starts with the definition of conditional probability:  
 $P(A|B) = P(AB)/P(B)$ .
- ▶ If  $X$  and  $Y$  are jointly discrete random variables, we can use this to define a probability mass function for  $X$  *given*  $Y = y$ .
- ▶ That is, we write  $p_{X|Y}(x|y) = P\{X = x|Y = y\} = \frac{p(x,y)}{p_Y(y)}$ .

## Conditional probability distributions

- ▶ It all starts with the definition of conditional probability:  
 $P(A|B) = P(AB)/P(B)$ .
- ▶ If  $X$  and  $Y$  are jointly discrete random variables, we can use this to define a probability mass function for  $X$  given  $Y = y$ .
- ▶ That is, we write  $p_{X|Y}(x|y) = P\{X = x|Y = y\} = \frac{p(x,y)}{p_Y(y)}$ .
- ▶ In words: first restrict sample space to pairs  $(x, y)$  with given  $y$  value. Then divide the original mass function by  $p_Y(y)$  to obtain a probability mass function on the restricted space.

## Conditional probability distributions

- ▶ It all starts with the definition of conditional probability:  
 $P(A|B) = P(AB)/P(B)$ .
- ▶ If  $X$  and  $Y$  are jointly discrete random variables, we can use this to define a probability mass function for  $X$  given  $Y = y$ .
- ▶ That is, we write  $p_{X|Y}(x|y) = P\{X = x|Y = y\} = \frac{p(x,y)}{p_Y(y)}$ .
- ▶ In words: first restrict sample space to pairs  $(x, y)$  with given  $y$  value. Then divide the original mass function by  $p_Y(y)$  to obtain a probability mass function on the restricted space.
- ▶ We do something similar when  $X$  and  $Y$  are continuous random variables. In that case we write  $f_{X|Y}(x|y) = \frac{f(x,y)}{f_Y(y)}$ .

## Conditional probability distributions

- ▶ It all starts with the definition of conditional probability:  
 $P(A|B) = P(AB)/P(B)$ .
- ▶ If  $X$  and  $Y$  are jointly discrete random variables, we can use this to define a probability mass function for  $X$  given  $Y = y$ .
- ▶ That is, we write  $p_{X|Y}(x|y) = P\{X = x|Y = y\} = \frac{p(x,y)}{p_Y(y)}$ .
- ▶ In words: first restrict sample space to pairs  $(x, y)$  with given  $y$  value. Then divide the original mass function by  $p_Y(y)$  to obtain a probability mass function on the restricted space.
- ▶ We do something similar when  $X$  and  $Y$  are continuous random variables. In that case we write  $f_{X|Y}(x|y) = \frac{f(x,y)}{f_Y(y)}$ .
- ▶ Often useful to think of sampling  $(X, Y)$  as a two-stage process. First sample  $Y$  from its marginal distribution, obtain  $Y = y$  for some particular  $y$ . Then sample  $X$  from its probability distribution given  $Y = y$ .



## Example

- ▶ Let  $X$  be a random variable of variance  $\sigma_X^2$  and  $Y$  an independent random variable of variance  $\sigma_Y^2$  and write  $Z = X + Y$ . Assume  $E[X] = E[Y] = 0$ .

## Example

- ▶ Let  $X$  be a random variable of variance  $\sigma_X^2$  and  $Y$  an independent random variable of variance  $\sigma_Y^2$  and write  $Z = X + Y$ . Assume  $E[X] = E[Y] = 0$ .
- ▶ What are the covariances  $\text{Cov}(X, Y)$  and  $\text{Cov}(X, Z)$ ?

## Example

- ▶ Let  $X$  be a random variable of variance  $\sigma_X^2$  and  $Y$  an independent random variable of variance  $\sigma_Y^2$  and write  $Z = X + Y$ . Assume  $E[X] = E[Y] = 0$ .
- ▶ What are the covariances  $\text{Cov}(X, Y)$  and  $\text{Cov}(X, Z)$ ?
- ▶ How about the correlation coefficients  $\rho(X, Y)$  and  $\rho(X, Z)$ ?

# Examples

- ▶ If  $X$  is binomial with parameters  $(p, n)$  then  
 $M_X(t) = (pe^t + 1 - p)^n$ .

# Examples

- ▶ If  $X$  is binomial with parameters  $(p, n)$  then  $M_X(t) = (pe^t + 1 - p)^n$ .
- ▶ If  $X$  is Poisson with parameter  $\lambda > 0$  then  $M_X(t) = \exp[\lambda(e^t - 1)]$ .

# Examples

- ▶ If  $X$  is binomial with parameters  $(p, n)$  then  $M_X(t) = (pe^t + 1 - p)^n$ .
- ▶ If  $X$  is Poisson with parameter  $\lambda > 0$  then  $M_X(t) = \exp[\lambda(e^t - 1)]$ .
- ▶ If  $X$  is normal with mean 0, variance 1, then  $M_X(t) = e^{t^2/2}$ .

# Examples

- ▶ If  $X$  is binomial with parameters  $(p, n)$  then  $M_X(t) = (pe^t + 1 - p)^n$ .
- ▶ If  $X$  is Poisson with parameter  $\lambda > 0$  then  $M_X(t) = \exp[\lambda(e^t - 1)]$ .
- ▶ If  $X$  is normal with mean 0, variance 1, then  $M_X(t) = e^{t^2/2}$ .
- ▶ If  $X$  is normal with mean  $\mu$ , variance  $\sigma^2$ , then  $M_X(t) = e^{\sigma^2 t^2/2 + \mu t}$ .

# Examples

- ▶ If  $X$  is binomial with parameters  $(p, n)$  then  $M_X(t) = (pe^t + 1 - p)^n$ .
- ▶ If  $X$  is Poisson with parameter  $\lambda > 0$  then  $M_X(t) = \exp[\lambda(e^t - 1)]$ .
- ▶ If  $X$  is normal with mean 0, variance 1, then  $M_X(t) = e^{t^2/2}$ .
- ▶ If  $X$  is normal with mean  $\mu$ , variance  $\sigma^2$ , then  $M_X(t) = e^{\sigma^2 t^2/2 + \mu t}$ .
- ▶ If  $X$  is exponential with parameter  $\lambda > 0$  then  $M_X(t) = \frac{\lambda}{\lambda - t}$ .



# Cauchy distribution

- ▶ A standard **Cauchy random variable** is a random real number with probability density  $f(x) = \frac{1}{\pi} \frac{1}{1+x^2}$ .

# Cauchy distribution

- ▶ A standard **Cauchy random variable** is a random real number with probability density  $f(x) = \frac{1}{\pi} \frac{1}{1+x^2}$ .
- ▶ There is a “spinning flashlight” interpretation. Put a flashlight at  $(0, 1)$ , spin it to a uniformly random angle in  $[-\pi/2, \pi/2]$ , and consider point  $X$  where light beam hits the  $x$ -axis.

# Cauchy distribution

- ▶ A standard **Cauchy random variable** is a random real number with probability density  $f(x) = \frac{1}{\pi} \frac{1}{1+x^2}$ .
- ▶ There is a “spinning flashlight” interpretation. Put a flashlight at  $(0, 1)$ , spin it to a uniformly random angle in  $[-\pi/2, \pi/2]$ , and consider point  $X$  where light beam hits the  $x$ -axis.
- ▶  $F_X(x) = P\{X \leq x\} = P\{\tan \theta \leq x\} = P\{\theta \leq \tan^{-1} x\} = \frac{1}{2} + \frac{1}{\pi} \tan^{-1} x$ .

# Cauchy distribution

- ▶ A standard **Cauchy random variable** is a random real number with probability density  $f(x) = \frac{1}{\pi} \frac{1}{1+x^2}$ .
- ▶ There is a “spinning flashlight” interpretation. Put a flashlight at  $(0, 1)$ , spin it to a uniformly random angle in  $[-\pi/2, \pi/2]$ , and consider point  $X$  where light beam hits the  $x$ -axis.
- ▶  $F_X(x) = P\{X \leq x\} = P\{\tan \theta \leq x\} = P\{\theta \leq \tan^{-1} x\} = \frac{1}{2} + \frac{1}{\pi} \tan^{-1} x$ .
- ▶ Find  $f_X(x) = \frac{d}{dx} F(x) = \frac{1}{\pi} \frac{1}{1+x^2}$ .

# Cauchy distribution

- ▶ A standard **Cauchy random variable** is a random real number with probability density  $f(x) = \frac{1}{\pi} \frac{1}{1+x^2}$ .
- ▶ There is a “spinning flashlight” interpretation. Put a flashlight at  $(0, 1)$ , spin it to a uniformly random angle in  $[-\pi/2, \pi/2]$ , and consider point  $X$  where light beam hits the  $x$ -axis.
- ▶  $F_X(x) = P\{X \leq x\} = P\{\tan \theta \leq x\} = P\{\theta \leq \tan^{-1} x\} = \frac{1}{2} + \frac{1}{\pi} \tan^{-1} x$ .
- ▶ Find  $f_X(x) = \frac{d}{dx} F(x) = \frac{1}{\pi} \frac{1}{1+x^2}$ .
- ▶ Cool fact: if  $X_1, X_2, \dots, X_n$  are i.i.d. Cauchy then their average  $A = \frac{X_1 + X_2 + \dots + X_n}{n}$  is also Cauchy.

# Beta distribution

- ▶ Two part experiment: first let  $p$  be uniform random variable  $[0, 1]$ , then let  $X$  be binomial  $(n, p)$  (number of heads when we toss  $n$   $p$ -coins).

# Beta distribution

- ▶ Two part experiment: first let  $p$  be uniform random variable  $[0, 1]$ , then let  $X$  be binomial  $(n, p)$  (number of heads when we toss  $n$   $p$ -coins).
- ▶ **Given** that  $X = a - 1$  and  $n - X = b - 1$  the conditional law of  $p$  is called the  $\beta$  distribution.

# Beta distribution

- ▶ Two part experiment: first let  $p$  be uniform random variable  $[0, 1]$ , then let  $X$  be binomial  $(n, p)$  (number of heads when we toss  $n$   $p$ -coins).
- ▶ **Given** that  $X = a - 1$  and  $n - X = b - 1$  the conditional law of  $p$  is called the  $\beta$  distribution.
- ▶ The density function is a constant (that doesn't depend on  $x$ ) times  $x^{a-1}(1-x)^{b-1}$ .



# Beta distribution

- ▶ Two part experiment: first let  $p$  be uniform random variable  $[0, 1]$ , then let  $X$  be binomial  $(n, p)$  (number of heads when we toss  $n$   $p$ -coins).
- ▶ **Given** that  $X = a - 1$  and  $n - X = b - 1$  the conditional law of  $p$  is called the  $\beta$  distribution.
- ▶ The density function is a constant (that doesn't depend on  $x$ ) times  $x^{a-1}(1-x)^{b-1}$ .
- ▶ That is  $f(x) = \frac{1}{B(a,b)}x^{a-1}(1-x)^{b-1}$  on  $[0, 1]$ , where  $B(a, b)$  is constant chosen to make integral one. Can show 
$$B(a, b) = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}.$$

# Beta distribution

- ▶ Two part experiment: first let  $p$  be uniform random variable  $[0, 1]$ , then let  $X$  be binomial  $(n, p)$  (number of heads when we toss  $n$   $p$ -coins).
- ▶ **Given** that  $X = a - 1$  and  $n - X = b - 1$  the conditional law of  $p$  is called the  $\beta$  distribution.
- ▶ The density function is a constant (that doesn't depend on  $x$ ) times  $x^{a-1}(1-x)^{b-1}$ .
- ▶ That is  $f(x) = \frac{1}{B(a,b)}x^{a-1}(1-x)^{b-1}$  on  $[0, 1]$ , where  $B(a, b)$  is constant chosen to make integral one. Can show 
$$B(a, b) = \frac{\Gamma(a)\Gamma(b)}{\Gamma(a+b)}.$$
- ▶ Turns out that  $E[X] = \frac{a}{a+b}$  and the mode of  $X$  is  $\frac{(a-1)}{(a-1)+(b-1)}$ .