

# **18.600: Lecture 25**

## **Covariance and some conditional expectation exercises**

Scott Sheffield

MIT

Covariance and correlation

Paradoxes: getting ready to think about conditional expectation

Covariance and correlation

Paradoxes: getting ready to think about conditional expectation

## A property of independence

- ▶ If  $X$  and  $Y$  are independent then
$$E[g(X)h(Y)] = E[g(X)]E[h(Y)].$$

## A property of independence

- ▶ If  $X$  and  $Y$  are independent then
$$E[g(X)h(Y)] = E[g(X)]E[h(Y)].$$
- ▶ Just write  $E[g(X)h(Y)] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x)h(y)f(x, y)dx dy$ .

## A property of independence

- ▶ If  $X$  and  $Y$  are independent then
$$E[g(X)h(Y)] = E[g(X)]E[h(Y)].$$
- ▶ Just write  $E[g(X)h(Y)] = \int_{-\infty}^{\infty} \int_{-\infty}^{\infty} g(x)h(y)f(x, y)dxdy$ .
- ▶ Since  $f(x, y) = f_X(x)f_Y(y)$  this factors as
$$\int_{-\infty}^{\infty} h(y)f_Y(y)dy \int_{-\infty}^{\infty} g(x)f_X(x)dx = E[h(Y)]E[g(X)].$$

## Defining covariance and correlation

- ▶ Now define covariance of  $X$  and  $Y$  by  
$$\text{Cov}(X, Y) = E[(X - E[X])(Y - E[Y])].$$

## Defining covariance and correlation

- ▶ Now define covariance of  $X$  and  $Y$  by
$$\text{Cov}(X, Y) = E[(X - E[X])(Y - E[Y])].$$
- ▶ Note: by definition  $\text{Var}(X) = \text{Cov}(X, X)$ .



# Defining covariance and correlation

- ▶ Now define covariance of  $X$  and  $Y$  by
$$\text{Cov}(X, Y) = E[(X - E[X])(Y - E[Y])].$$
- ▶ Note: by definition  $\text{Var}(X) = \text{Cov}(X, X)$ .
- ▶ Covariance (like variance) can also be written a different way. Write  $\mu_X = E[X]$  and  $\mu_Y = E[Y]$ . If laws of  $X$  and  $Y$  are known, then  $\mu_X$  and  $\mu_Y$  are just constants.

# Defining covariance and correlation

- ▶ Now define covariance of  $X$  and  $Y$  by  
$$\text{Cov}(X, Y) = E[(X - E[X])(Y - E[Y])].$$
- ▶ Note: by definition  $\text{Var}(X) = \text{Cov}(X, X)$ .
- ▶ Covariance (like variance) can also be written a different way. Write  $\mu_X = E[X]$  and  $\mu_Y = E[Y]$ . If laws of  $X$  and  $Y$  are known, then  $\mu_X$  and  $\mu_Y$  are just constants.
- ▶ Then

$$\begin{aligned}\text{Cov}(X, Y) &= E[(X - \mu_X)(Y - \mu_Y)] = E[XY - \mu_X Y - \mu_Y X + \mu_X \mu_Y] = \\ &E[XY] - \mu_X E[Y] - \mu_Y E[X] + \mu_X \mu_Y = E[XY] - E[X]E[Y].\end{aligned}$$

# Defining covariance and correlation

- ▶ Now define covariance of  $X$  and  $Y$  by  
$$\text{Cov}(X, Y) = E[(X - E[X])(Y - E[Y])].$$
- ▶ Note: by definition  $\text{Var}(X) = \text{Cov}(X, X)$ .
- ▶ Covariance (like variance) can also be written a different way. Write  $\mu_X = E[X]$  and  $\mu_Y = E[Y]$ . If laws of  $X$  and  $Y$  are known, then  $\mu_X$  and  $\mu_Y$  are just constants.
- ▶ Then

$$\begin{aligned}\text{Cov}(X, Y) &= E[(X - \mu_X)(Y - \mu_Y)] = E[XY - \mu_X Y - \mu_Y X + \mu_X \mu_Y] = \\ &E[XY] - \mu_X E[Y] - \mu_Y E[X] + \mu_X \mu_Y = E[XY] - E[X]E[Y].\end{aligned}$$

- ▶ Covariance formula  $E[XY] - E[X]E[Y]$ , or “expectation of product minus product of expectations” is frequently useful.

# Defining covariance and correlation

- ▶ Now define covariance of  $X$  and  $Y$  by
$$\text{Cov}(X, Y) = E[(X - E[X])(Y - E[Y])].$$
- ▶ Note: by definition  $\text{Var}(X) = \text{Cov}(X, X)$ .
- ▶ Covariance (like variance) can also be written a different way. Write  $\mu_X = E[X]$  and  $\mu_Y = E[Y]$ . If laws of  $X$  and  $Y$  are known, then  $\mu_X$  and  $\mu_Y$  are just constants.
- ▶ Then

$$\begin{aligned}\text{Cov}(X, Y) &= E[(X - \mu_X)(Y - \mu_Y)] = E[XY - \mu_X Y - \mu_Y X + \mu_X \mu_Y] = \\ &E[XY] - \mu_X E[Y] - \mu_Y E[X] + \mu_X \mu_Y = E[XY] - E[X]E[Y].\end{aligned}$$

- ▶ Covariance formula  $E[XY] - E[X]E[Y]$ , or “expectation of product minus product of expectations” is frequently useful.
- ▶ Note: if  $X$  and  $Y$  are independent then  $\text{Cov}(X, Y) = 0$ .

## Basic covariance facts

- ▶ Using  $\text{Cov}(X, Y) = E[XY] - E[X]E[Y]$  as a definition, certain facts are immediate.

# Basic covariance facts

- ▶ Using  $\text{Cov}(X, Y) = E[XY] - E[X]E[Y]$  as a definition, certain facts are immediate.
- ▶  $\text{Cov}(X, Y) = \text{Cov}(Y, X)$

# Basic covariance facts

- ▶ Using  $\text{Cov}(X, Y) = E[XY] - E[X]E[Y]$  as a definition, certain facts are immediate.
- ▶  $\text{Cov}(X, Y) = \text{Cov}(Y, X)$
- ▶  $\text{Cov}(X, X) = \text{Var}(X)$

# Basic covariance facts

- ▶ Using  $\text{Cov}(X, Y) = E[XY] - E[X]E[Y]$  as a definition, certain facts are immediate.
- ▶  $\text{Cov}(X, Y) = \text{Cov}(Y, X)$
- ▶  $\text{Cov}(X, X) = \text{Var}(X)$
- ▶  $\text{Cov}(aX, Y) = a\text{Cov}(X, Y)$ .



# Basic covariance facts

- ▶ Using  $\text{Cov}(X, Y) = E[XY] - E[X]E[Y]$  as a definition, certain facts are immediate.
- ▶  $\text{Cov}(X, Y) = \text{Cov}(Y, X)$
- ▶  $\text{Cov}(X, X) = \text{Var}(X)$
- ▶  $\text{Cov}(aX, Y) = a\text{Cov}(X, Y)$ .
- ▶  $\text{Cov}(X_1 + X_2, Y) = \text{Cov}(X_1, Y) + \text{Cov}(X_2, Y)$ .

# Basic covariance facts

- ▶ Using  $\text{Cov}(X, Y) = E[XY] - E[X]E[Y]$  as a definition, certain facts are immediate.
- ▶  $\text{Cov}(X, Y) = \text{Cov}(Y, X)$
- ▶  $\text{Cov}(X, X) = \text{Var}(X)$
- ▶  $\text{Cov}(aX, Y) = a\text{Cov}(X, Y)$ .
- ▶  $\text{Cov}(X_1 + X_2, Y) = \text{Cov}(X_1, Y) + \text{Cov}(X_2, Y)$ .
- ▶ **General statement of bilinearity of covariance:**

$$\text{Cov}\left(\sum_{i=1}^m a_i X_i, \sum_{j=1}^n b_j Y_j\right) = \sum_{i=1}^m \sum_{j=1}^n a_i b_j \text{Cov}(X_i, Y_j).$$

# Basic covariance facts

- ▶ Using  $\text{Cov}(X, Y) = E[XY] - E[X]E[Y]$  as a definition, certain facts are immediate.
- ▶  $\text{Cov}(X, Y) = \text{Cov}(Y, X)$
- ▶  $\text{Cov}(X, X) = \text{Var}(X)$
- ▶  $\text{Cov}(aX, Y) = a\text{Cov}(X, Y)$ .
- ▶  $\text{Cov}(X_1 + X_2, Y) = \text{Cov}(X_1, Y) + \text{Cov}(X_2, Y)$ .
- ▶ **General statement of bilinearity of covariance:**

$$\text{Cov}\left(\sum_{i=1}^m a_i X_i, \sum_{j=1}^n b_j Y_j\right) = \sum_{i=1}^m \sum_{j=1}^n a_i b_j \text{Cov}(X_i, Y_j).$$

- ▶ Special case:

$$\text{Var}\left(\sum_{i=1}^n X_i\right) = \sum_{i=1}^n \text{Var}(X_i) + 2 \sum_{(i,j): i < j} \text{Cov}(X_i, X_j).$$

# Defining correlation

- ▶ Again, by definition  $\text{Cov}(X, Y) = E[XY] - E[X]E[Y]$ .

# Defining correlation

- ▶ Again, by definition  $\text{Cov}(X, Y) = E[XY] - E[X]E[Y]$ .
- ▶ **Correlation** of  $X$  and  $Y$  defined by

$$\rho(X, Y) := \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}}.$$

# Defining correlation

- ▶ Again, by definition  $\text{Cov}(X, Y) = E[XY] - E[X]E[Y]$ .
- ▶ **Correlation** of  $X$  and  $Y$  defined by

$$\rho(X, Y) := \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}}.$$

- ▶ Correlation doesn't care what units you use for  $X$  and  $Y$ . If  $a > 0$  and  $c > 0$  then  $\rho(aX + b, cY + d) = \rho(X, Y)$ .

# Defining correlation

- ▶ Again, by definition  $\text{Cov}(X, Y) = E[XY] - E[X]E[Y]$ .
- ▶ **Correlation** of  $X$  and  $Y$  defined by

$$\rho(X, Y) := \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}}.$$

- ▶ Correlation doesn't care what units you use for  $X$  and  $Y$ . If  $a > 0$  and  $c > 0$  then  $\rho(aX + b, cY + d) = \rho(X, Y)$ .
- ▶ Satisfies  $-1 \leq \rho(X, Y) \leq 1$ .

# Defining correlation

- ▶ Again, by definition  $\text{Cov}(X, Y) = E[XY] - E[X]E[Y]$ .
- ▶ **Correlation** of  $X$  and  $Y$  defined by

$$\rho(X, Y) := \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}}.$$

- ▶ Correlation doesn't care what units you use for  $X$  and  $Y$ . If  $a > 0$  and  $c > 0$  then  $\rho(aX + b, cY + d) = \rho(X, Y)$ .
- ▶ Satisfies  $-1 \leq \rho(X, Y) \leq 1$ .
- ▶ Why is that? Something to do with  $E[(X + Y)^2] \geq 0$  and  $E[(X - Y)^2] \geq 0$ ?



# Defining correlation

- ▶ Again, by definition  $\text{Cov}(X, Y) = E[XY] - E[X]E[Y]$ .
- ▶ **Correlation** of  $X$  and  $Y$  defined by

$$\rho(X, Y) := \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}}.$$

- ▶ Correlation doesn't care what units you use for  $X$  and  $Y$ . If  $a > 0$  and  $c > 0$  then  $\rho(aX + b, cY + d) = \rho(X, Y)$ .
- ▶ Satisfies  $-1 \leq \rho(X, Y) \leq 1$ .
- ▶ Why is that? Something to do with  $E[(X + Y)^2] \geq 0$  and  $E[(X - Y)^2] \geq 0$ ?
- ▶ If  $a$  and  $b$  are constants and  $a > 0$  then  $\rho(aX + b, X) = 1$ .

# Defining correlation

- ▶ Again, by definition  $\text{Cov}(X, Y) = E[XY] - E[X]E[Y]$ .
- ▶ **Correlation** of  $X$  and  $Y$  defined by

$$\rho(X, Y) := \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var}(X)\text{Var}(Y)}}.$$

- ▶ Correlation doesn't care what units you use for  $X$  and  $Y$ . If  $a > 0$  and  $c > 0$  then  $\rho(aX + b, cY + d) = \rho(X, Y)$ .
- ▶ Satisfies  $-1 \leq \rho(X, Y) \leq 1$ .
- ▶ Why is that? Something to do with  $E[(X + Y)^2] \geq 0$  and  $E[(X - Y)^2] \geq 0$ ?
- ▶ If  $a$  and  $b$  are constants and  $a > 0$  then  $\rho(aX + b, X) = 1$ .
- ▶ If  $a$  and  $b$  are constants and  $a < 0$  then  $\rho(aX + b, X) = -1$ .

## Important point

- ▶ Say  $X$  and  $Y$  are uncorrelated when  $\rho(X, Y) = 0$ .

## Important point

- ▶ Say  $X$  and  $Y$  are uncorrelated when  $\rho(X, Y) = 0$ .
- ▶ Are independent random variables  $X$  and  $Y$  always uncorrelated?

## Important point

- ▶ Say  $X$  and  $Y$  are uncorrelated when  $\rho(X, Y) = 0$ .
- ▶ Are independent random variables  $X$  and  $Y$  always uncorrelated?
- ▶ Yes, assuming variances are finite (so that correlation is defined).

# Important point

- ▶ Say  $X$  and  $Y$  are uncorrelated when  $\rho(X, Y) = 0$ .
- ▶ Are independent random variables  $X$  and  $Y$  always uncorrelated?
- ▶ Yes, assuming variances are finite (so that correlation is defined).
- ▶ Are uncorrelated random variables always independent?

# Important point

- ▶ Say  $X$  and  $Y$  are uncorrelated when  $\rho(X, Y) = 0$ .
- ▶ Are independent random variables  $X$  and  $Y$  always uncorrelated?
- ▶ Yes, assuming variances are finite (so that correlation is defined).
- ▶ Are uncorrelated random variables always independent?
- ▶ No. Uncorrelated just means  $E[(X - E[X])(Y - E[Y])] = 0$ , i.e., the outcomes where  $(X - E[X])(Y - E[Y])$  is positive (the upper right and lower left quadrants, if axes are drawn centered at  $(E[X], E[Y])$ ) balance out the outcomes where this quantity is negative (upper left and lower right quadrants). This is a much weaker statement than independence.

# Examples

- ▶ Suppose that  $X_1, \dots, X_n$  are i.i.d. random variables with variance 1. For example, maybe each  $X_j$  takes values  $\pm 1$  according to a fair coin toss.



# Examples

- ▶ Suppose that  $X_1, \dots, X_n$  are i.i.d. random variables with variance 1. For example, maybe each  $X_j$  takes values  $\pm 1$  according to a fair coin toss.
- ▶ Compute  $\text{Cov}(X_1 + X_2 + X_3, X_2 + X_3 + X_4)$ .

# Examples

- ▶ Suppose that  $X_1, \dots, X_n$  are i.i.d. random variables with variance 1. For example, maybe each  $X_j$  takes values  $\pm 1$  according to a fair coin toss.
- ▶ Compute  $\text{Cov}(X_1 + X_2 + X_3, X_2 + X_3 + X_4)$ .
- ▶ Compute the correlation coefficient  $\rho(X_1 + X_2 + X_3, X_2 + X_3 + X_4)$ .

# Examples

- ▶ Suppose that  $X_1, \dots, X_n$  are i.i.d. random variables with variance 1. For example, maybe each  $X_j$  takes values  $\pm 1$  according to a fair coin toss.
- ▶ Compute  $\text{Cov}(X_1 + X_2 + X_3, X_2 + X_3 + X_4)$ .
- ▶ Compute the correlation coefficient  $\rho(X_1 + X_2 + X_3, X_2 + X_3 + X_4)$ .
- ▶ Can we generalize this example?

# Examples

- ▶ Suppose that  $X_1, \dots, X_n$  are i.i.d. random variables with variance 1. For example, maybe each  $X_j$  takes values  $\pm 1$  according to a fair coin toss.
- ▶ Compute  $\text{Cov}(X_1 + X_2 + X_3, X_2 + X_3 + X_4)$ .
- ▶ Compute the correlation coefficient  $\rho(X_1 + X_2 + X_3, X_2 + X_3 + X_4)$ .
- ▶ Can we generalize this example?
- ▶ What is variance of number of people who get their own hat in the hat problem?

# Examples

- ▶ Suppose that  $X_1, \dots, X_n$  are i.i.d. random variables with variance 1. For example, maybe each  $X_j$  takes values  $\pm 1$  according to a fair coin toss.
- ▶ Compute  $\text{Cov}(X_1 + X_2 + X_3, X_2 + X_3 + X_4)$ .
- ▶ Compute the correlation coefficient  $\rho(X_1 + X_2 + X_3, X_2 + X_3 + X_4)$ .
- ▶ Can we generalize this example?
- ▶ What is variance of number of people who get their own hat in the hat problem?
- ▶ Define  $X_i$  to be 1 if  $i$ th person gets own hat, zero otherwise.

# Examples

- ▶ Suppose that  $X_1, \dots, X_n$  are i.i.d. random variables with variance 1. For example, maybe each  $X_j$  takes values  $\pm 1$  according to a fair coin toss.
- ▶ Compute  $\text{Cov}(X_1 + X_2 + X_3, X_2 + X_3 + X_4)$ .
- ▶ Compute the correlation coefficient  $\rho(X_1 + X_2 + X_3, X_2 + X_3 + X_4)$ .
- ▶ Can we generalize this example?
- ▶ What is variance of number of people who get their own hat in the hat problem?
- ▶ Define  $X_i$  to be 1 if  $i$ th person gets own hat, zero otherwise.
- ▶ Recall formula
$$\text{Var}(\sum_{i=1}^n X_i) = \sum_{i=1}^n \text{Var}(X_i) + 2 \sum_{(i,j): i < j} \text{Cov}(X_i, X_j).$$

# Examples

- ▶ Suppose that  $X_1, \dots, X_n$  are i.i.d. random variables with variance 1. For example, maybe each  $X_j$  takes values  $\pm 1$  according to a fair coin toss.
- ▶ Compute  $\text{Cov}(X_1 + X_2 + X_3, X_2 + X_3 + X_4)$ .
- ▶ Compute the correlation coefficient  $\rho(X_1 + X_2 + X_3, X_2 + X_3 + X_4)$ .
- ▶ Can we generalize this example?
- ▶ What is variance of number of people who get their own hat in the hat problem?
- ▶ Define  $X_i$  to be 1 if  $i$ th person gets own hat, zero otherwise.
- ▶ Recall formula
$$\text{Var}(\sum_{i=1}^n X_i) = \sum_{i=1}^n \text{Var}(X_i) + 2 \sum_{(i,j): i < j} \text{Cov}(X_i, X_j).$$
- ▶ Reduces problem to computing  $\text{Cov}(X_i, X_j)$  (for  $i \neq j$ ) and  $\text{Var}(X_i)$ .

Covariance and correlation

Paradoxes: getting ready to think about conditional expectation



Covariance and correlation

Paradoxes: getting ready to think about conditional expectation

# Famous paradox

- ▶ Certain corrupt and amoral banker dies, instructed to spend some number  $n$  (of banker's choosing) days in hell.

# Famous paradox

- ▶ Certain corrupt and amoral banker dies, instructed to spend some number  $n$  (of banker's choosing) days in hell.
- ▶ At the end of this period, a (biased) coin will be tossed. Banker will be assigned to hell forever with probability  $1/n$  and heaven forever with probability  $1 - 1/n$ .

# Famous paradox

- ▶ Certain corrupt and amoral banker dies, instructed to spend some number  $n$  (of banker's choosing) days in hell.
- ▶ At the end of this period, a (biased) coin will be tossed. Banker will be assigned to hell forever with probability  $1/n$  and heaven forever with probability  $1 - 1/n$ .
- ▶ After 10 days, banker reasons, "If I wait another day I reduce my odds of being here forever from  $1/10$  to  $1/11$ . That's a reduction of  $1/110$ . A  $1/110$  chance at infinity has infinite value. Worth waiting one more day."

# Famous paradox

- ▶ Certain corrupt and amoral banker dies, instructed to spend some number  $n$  (of banker's choosing) days in hell.
- ▶ At the end of this period, a (biased) coin will be tossed. Banker will be assigned to hell forever with probability  $1/n$  and heaven forever with probability  $1 - 1/n$ .
- ▶ After 10 days, banker reasons, "If I wait another day I reduce my odds of being here forever from  $1/10$  to  $1/11$ . That's a reduction of  $1/110$ . A  $1/110$  chance at infinity has infinite value. Worth waiting one more day."
- ▶ Repeats this reasoning every day, stays in hell forever.

# Famous paradox

- ▶ Certain corrupt and amoral banker dies, instructed to spend some number  $n$  (of banker's choosing) days in hell.
- ▶ At the end of this period, a (biased) coin will be tossed. Banker will be assigned to hell forever with probability  $1/n$  and heaven forever with probability  $1 - 1/n$ .
- ▶ After 10 days, banker reasons, "If I wait another day I reduce my odds of being here forever from  $1/10$  to  $1/11$ . That's a reduction of  $1/110$ . A  $1/110$  chance at infinity has infinite value. Worth waiting one more day."
- ▶ Repeats this reasoning every day, stays in hell forever.
- ▶ Standard punch line: this is actually what banker deserved.

# Famous paradox

- ▶ Certain corrupt and amoral banker dies, instructed to spend some number  $n$  (of banker's choosing) days in hell.
- ▶ At the end of this period, a (biased) coin will be tossed. Banker will be assigned to hell forever with probability  $1/n$  and heaven forever with probability  $1 - 1/n$ .
- ▶ After 10 days, banker reasons, "If I wait another day I reduce my odds of being here forever from  $1/10$  to  $1/11$ . That's a reduction of  $1/110$ . A  $1/110$  chance at infinity has infinite value. Worth waiting one more day."
- ▶ Repeats this reasoning every day, stays in hell forever.
- ▶ Standard punch line: this is actually what banker deserved.
- ▶ Fairly dark as math humor goes (and no offense intended to anyone...) but dilemma is interesting.

- ▶ **Paradox:** decisions seem sound individually but together yield worst possible outcome. Why? Can we demystify this?



- ▶ **Paradox:** decisions seem sound individually but together yield worst possible outcome. Why? Can we demystify this?
- ▶ **Variant without probability:** Instead of tossing  $(1/n)$ -coin, person deterministically spends  $1/n$  fraction of future days (every  $n$ th day, say) in hell.

- ▶ **Paradox:** decisions seem sound individually but together yield worst possible outcome. Why? Can we demystify this?
- ▶ **Variant without probability:** Instead of tossing  $(1/n)$ -coin, person deterministically spends  $1/n$  fraction of future days (every  $n$ th day, say) in hell.

- ▶ **Paradox:** decisions seem sound individually but together yield worst possible outcome. Why? Can we demystify this?
- ▶ **Variant without probability:** Instead of tossing  $(1/n)$ -coin, person deterministically spends  $1/n$  fraction of future days (every  $n$ th day, say) in hell.

- ▶ **Paradox:** decisions seem sound individually but together yield worst possible outcome. Why? Can we demystify this?
- ▶ **Variant without probability:** Instead of tossing  $(1/n)$ -coin, person deterministically spends  $1/n$  fraction of future days (every  $n$ th day, say) in hell.
- ▶ **Even simpler variant:** infinitely many identical money sacks have labels  $1, 2, 3, \dots$ . I have sack 1. You have all others.

- ▶ **Paradox:** decisions seem sound individually but together yield worst possible outcome. Why? Can we demystify this?
- ▶ **Variant without probability:** Instead of tossing  $(1/n)$ -coin, person deterministically spends  $1/n$  fraction of future days (every  $n$ th day, say) in hell.
- ▶ **Even simpler variant:** infinitely many identical money sacks have labels  $1, 2, 3, \dots$ . I have sack 1. You have all others.
- ▶ You offer me a deal. I give you sack 1, you give me sacks 2 and 3. I give you sack 2 and you give me sacks 4 and 5. On the  $n$ th stage, I give you sack  $n$  and you give me sacks  $2n$  and  $2n + 1$ . Continue until I say stop.

- ▶ **Paradox:** decisions seem sound individually but together yield worst possible outcome. Why? Can we demystify this?
- ▶ **Variant without probability:** Instead of tossing  $(1/n)$ -coin, person deterministically spends  $1/n$  fraction of future days (every  $n$ th day, say) in hell.
- ▶ **Even simpler variant:** infinitely many identical money sacks have labels  $1, 2, 3, \dots$ . I have sack 1. You have all others.
- ▶ You offer me a deal. I give you sack 1, you give me sacks 2 and 3. I give you sack 2 and you give me sacks 4 and 5. On the  $n$ th stage, I give you sack  $n$  and you give me sacks  $2n$  and  $2n + 1$ . Continue until I say stop.
- ▶ Lets me get arbitrarily rich. But if I go on forever, I return every sack given to me. If  $n$ th sack confers right to spend  $n$ th day in heaven, leads to hell-forever paradox.

- ▶ **Paradox:** decisions seem sound individually but together yield worst possible outcome. Why? Can we demystify this?
- ▶ **Variant without probability:** Instead of tossing  $(1/n)$ -coin, person deterministically spends  $1/n$  fraction of future days (every  $n$ th day, say) in hell.
- ▶ **Even simpler variant:** infinitely many identical money sacks have labels  $1, 2, 3, \dots$ . I have sack 1. You have all others.
- ▶ You offer me a deal. I give you sack 1, you give me sacks 2 and 3. I give you sack 2 and you give me sacks 4 and 5. On the  $n$ th stage, I give you sack  $n$  and you give me sacks  $2n$  and  $2n + 1$ . Continue until I say stop.
- ▶ Lets me get arbitrarily rich. But if I go on forever, I return every sack given to me. If  $n$ th sack confers right to spend  $n$ th day in heaven, leads to hell-forever paradox.
- ▶ I make infinitely many good trades and end up with less than I started with. “Paradox” is really just existence of 2-to-1 map from (smaller set)  $\{2, 3, \dots\}$  to (bigger set)  $\{1, 2, \dots\}$ .

# Money pile paradox

- ▶ You have an infinite collection of money piles with labeled  $0, 1, 2, \dots$  from left to right.



# Money pile paradox

- ▶ You have an infinite collection of money piles with labeled  $0, 1, 2, \dots$  from left to right.
- ▶ Precise details not important, but let's say you have  $1/4$  in the 0th pile and  $\frac{3}{8}5^j$  in the  $j$ th pile for each  $j > 0$ . Important thing is that pile size is increasing exponentially in  $j$ .

# Money pile paradox

- ▶ You have an infinite collection of money piles with labeled  $0, 1, 2, \dots$  from left to right.
- ▶ Precise details not important, but let's say you have  $1/4$  in the 0th pile and  $\frac{3}{8}5^j$  in the  $j$ th pile for each  $j > 0$ . Important thing is that pile size is increasing exponentially in  $j$ .
- ▶ Banker proposes to transfer a fraction (say  $2/3$ ) of each pile to the pile on its left and remainder to the pile on its right. Do this simultaneously for all piles.

# Money pile paradox

- ▶ You have an infinite collection of money piles with labeled  $0, 1, 2, \dots$  from left to right.
- ▶ Precise details not important, but let's say you have  $1/4$  in the 0th pile and  $\frac{3}{8}5^j$  in the  $j$ th pile for each  $j > 0$ . Important thing is that pile size is increasing exponentially in  $j$ .
- ▶ Banker proposes to transfer a fraction (say  $2/3$ ) of each pile to the pile on its left and remainder to the pile on its right. Do this simultaneously for all piles.
- ▶ Every pile is bigger after transfer (and this can be true even if banker takes a portion of each pile as a fee).

# Money pile paradox

- ▶ You have an infinite collection of money piles with labeled  $0, 1, 2, \dots$  from left to right.
- ▶ Precise details not important, but let's say you have  $1/4$  in the 0th pile and  $\frac{3}{8}5^j$  in the  $j$ th pile for each  $j > 0$ . Important thing is that pile size is increasing exponentially in  $j$ .
- ▶ Banker proposes to transfer a fraction (say  $2/3$ ) of each pile to the pile on its left and remainder to the pile on its right. Do this simultaneously for all piles.
- ▶ Every pile is bigger after transfer (and this can be true even if banker takes a portion of each pile as a fee).
- ▶ Banker seemed to make you richer (every pile got bigger) but really just reshuffled your infinite wealth.

## Two envelope paradox

- ▶  $X$  is geometric with parameter  $1/2$ . One envelope has  $10^X$  dollars, one has  $10^{X-1}$  dollars. Envelopes shuffled.

## Two envelope paradox

- ▶  $X$  is geometric with parameter  $1/2$ . One envelope has  $10^X$  dollars, one has  $10^{X-1}$  dollars. Envelopes shuffled.
- ▶ You choose an envelope and, after seeing contents, are allowed to choose whether to keep it or switch. (Maybe you have to pay a dollar to switch.)

## Two envelope paradox

- ▶  $X$  is geometric with parameter  $1/2$ . One envelope has  $10^X$  dollars, one has  $10^{X-1}$  dollars. Envelopes shuffled.
- ▶ You choose an envelope and, after seeing contents, are allowed to choose whether to keep it or switch. (Maybe you have to pay a dollar to switch.)
- ▶ Maximizing conditional expectation, it seems it's always better to switch. But if you always switch, why not just choose second-choice envelope first and avoid switching fee?

# Two envelope paradox

- ▶  $X$  is geometric with parameter  $1/2$ . One envelope has  $10^X$  dollars, one has  $10^{X-1}$  dollars. Envelopes shuffled.
- ▶ You choose an envelope and, after seeing contents, are allowed to choose whether to keep it or switch. (Maybe you have to pay a dollar to switch.)
- ▶ Maximizing conditional expectation, it seems it's always better to switch. But if you always switch, why not just choose second-choice envelope first and avoid switching fee?
- ▶ Kind of a disguised version of money pile paradox. But more subtle. One has to replace “ $j$ th pile of money” with “restriction of expectation sum to scenario that first chosen envelop has  $10^j$ ”. Switching indeed makes each pile bigger.



# Two envelope paradox

- ▶  $X$  is geometric with parameter  $1/2$ . One envelope has  $10^X$  dollars, one has  $10^{X-1}$  dollars. Envelopes shuffled.
- ▶ You choose an envelope and, after seeing contents, are allowed to choose whether to keep it or switch. (Maybe you have to pay a dollar to switch.)
- ▶ Maximizing conditional expectation, it seems it's always better to switch. But if you always switch, why not just choose second-choice envelope first and avoid switching fee?
- ▶ Kind of a disguised version of money pile paradox. But more subtle. One has to replace “ $j$ th pile of money” with “restriction of expectation sum to scenario that first chosen envelop has  $10^j$ ”. Switching indeed makes each pile bigger.
- ▶ However, “Higher expectation given amount in first envelope” may not be right notion of “better.” If  $S$  is payout with switching,  $T$  is payout without switching, then  $S$  has same law as  $T - 1$ . In that sense  $S$  is worse.

- ▶ Beware infinite expectations.

- ▶ Beware infinite expectations.
- ▶ Beware unbounded utility functions.

- ▶ Beware infinite expectations.
- ▶ Beware unbounded utility functions.
- ▶ They can lead to strange conclusions, sometimes related to “reshuffling infinite (actual or expected) wealth to create more” paradoxes.

- ▶ Beware infinite expectations.
- ▶ Beware unbounded utility functions.
- ▶ They can lead to strange conclusions, sometimes related to “reshuffling infinite (actual or expected) wealth to create more” paradoxes.
- ▶ Paradoxes can arise even when total transaction is finite with probability one (as in envelope problem).