

## IAP - Math Lecture Series: January 2019

### I. Shape of a 2D meniscus

Consider the meniscus resulting from the interaction between a solid and a free surface (Figure 1). Assume that the contact angle,  $\theta$ , is prescribed. The normal force balance at the interface is expressed by the Young-Laplace equation:

$$\rho g z = \sigma \nabla \cdot \mathbf{n}, \quad (1)$$

which expresses the balance between hydrostatic and curvature pressures. Here  $\rho$  is the fluid density,  $\vec{g} = -g\hat{z}$  is gravitational acceleration,  $z = h(x)$  is the height of the free surface,  $\sigma$  is the surface tension.  $\mathbf{n}$  is the unit normal to the surface pointing towards air, and  $\nabla \cdot \mathbf{n}$ .

a) By defining a functional  $f(x, y, z) = z - h(x, y)$  that vanishes on the free surface, show that the unit normal  $\mathbf{n} = \frac{\nabla f}{|\nabla f|}$  may be expressed

$$\mathbf{n} = \frac{\hat{z} - h_x \hat{x}}{(1 + h_x^2)^{1/2}} \quad , \quad (2)$$

and that the curvature of the surface may be expressed as

$$\nabla \cdot \mathbf{n} = \frac{-h_{xx}}{(1 + h_x^2)^{3/2}} \quad . \quad (3)$$

b) Now by assuming that the interfacial slope is relatively small  $h_x^2 \ll 1$ , one has that  $\nabla \cdot \mathbf{n} = -h_{xx}$ . By substituting this into the Young-Laplace equation (1), solve the resulting differential equation for the interface shape  $h(x)$ . By applying appropriate boundary conditions, i)  $h \rightarrow 0$  as  $x \rightarrow \infty$ , and ii)  $h_x = -\cot\theta$  at  $x = 0$ , show that the meniscus shape is given by:

$$h(x) = L_c \cot\theta \exp(-x/L_c) \quad (4)$$

Note that the capillary length,  $L_c = \sqrt{\frac{\rho g}{\sigma}}$ , prescribes the characteristic length over which the meniscus relaxes to horizontal.

### II. Forces on Floating Bodies

Floating bodies must be supported by some combination of buoyancy and curvature forces. Specifically, since the fluid pressure beneath the interface is given by

$$p = P_0 + \rho g z + \sigma \nabla \cdot \mathbf{n} = P_0 + p_g + p_c \quad ,$$

one may calculate the total hydrostatic force on the body by integrating the hydrostatic pressure  $p_g$  and curvature pressure  $p_c$  over the surface of the body.

The vertical force balance on the body may thus be expressed:

$$Mg = \mathbf{z} \cdot \int_{C_1} -p \mathbf{n} ds = F_b + F_c \quad , \quad (5)$$

where  $F_b$  and  $F_c$  are respectively the buoyancy and curvature forces, defined below.

a) By using the Divergence Theorem, show that the buoyancy force

$$F_b = \mathbf{z} \cdot \int_{C_1} \rho g z \mathbf{n} ds \quad (6)$$

is simply expressible in terms of the volume (per unit length into the page)  $V_c$  of fluid displaced above the object and inside the line of tangency (Figure 2). Thus prove Archimedes Principle holds for a floating body: the buoyancy force is equal to the weight of the fluid displaced by the body:  $F_b = \rho V_c g$ .

b) Proceed by using the Frenet-Serret equation  $(\nabla \cdot \mathbf{n}) \mathbf{n} = \frac{d\mathbf{t}}{ds}$ , that relates the curvature to the derivative of the unit tangent  $\mathbf{t}$  with respect to the arclength  $s$ . By integrating the curvature pressure along the line of contact  $C_1$ , show that the curvature force may be expressed

$$F_c = \mathbf{z} \cdot \int_{C_1} \sigma (\nabla \cdot \mathbf{n}) \mathbf{n} ds = 2\sigma \sin\theta \quad (7)$$

c) Show that the relative magnitude of the buoyancy and curvature forces supporting a floating, non-wetting (2D) body is thus proportional to the relative magnitudes of the characteristic body size  $L$  and the capillary length  $L_c$ :

$$\frac{F_b}{F_c} \sim \frac{L}{L_c} \quad . \quad (8)$$

For an air-water interface,  $g = 980 \text{ cm/s}^2$ ,  $\rho = 1 \text{ g/cc}$ ,  $\sigma = 70 \text{ g/s}^2$  and the capillary length  $L_c \sim 3 \text{ mm}$ . Very small objects ( $L \ll L_c$ ), such as paper clips, pins or insects, are thus supported principally by curvature rather than buoyancy forces. They may reside at rest on a free surface provided the curvature force induced by their deflection of the free surface is sufficient to bear their weight.

d) For a body of contact length  $X$  and total mass  $M$  supported by surface tension, show that static equilibrium on the free surface requires that:

$$\frac{Mg}{\sigma X \sin\theta} < 1 \quad (9)$$

where  $\theta$  is the angle of tangency of the floating body. How big must your feet be for you to walk on water?




Figure 1: A definitional sketch of a planar meniscus at an air-water interface. The free surface is defined by  $z = \eta(x)$ , varying from its maximum elevation at its point of contact with the wall ( $x = 0$ ) to zero at large  $x$ . The shape is prescribed by the Young-Laplace equation.




Figure 2: A non-wetting two-dimensional body of radius  $r$  and mass  $M$  floats on a free surface with surface tension  $\sigma$ . In general, its weight  $Mg$  must be supported by some combination of curvature and buoyancy forces.  $V_c$  and  $V_m$  denote the fluid volumes displaced, respectively, inside and outside the line of tangency.