

**PRACTICE NUMBER 1 FOR TEST 2**  
**18.100B SPRING 2007**

This test is closed book, no books, papers or notes are permitted. You may use theorems, lemmas and propositions from the class and book. Note that where  $\mathbb{R}^k$  is mentioned below the standard metric is assumed.

Problem 1 Put  $B = \{(x, y) \in \mathbb{R}^2; x^2 + y^2 \leq 1\}$  and suppose  $g : B \rightarrow \mathbb{R}$  is a continuous function. Show that the function

$$f(x, y) = x^2 + y^2 + (x^2 + y^2)(1 - (x^2 + y^2))g(x, y)$$

is continuous on  $B$  and prove that there is some point  $(\bar{x}, \bar{y}) \in B$  such that  $f(\bar{x}, \bar{y}) = \frac{1}{2}$ .

Problem 2 If  $f : [0, 1] \rightarrow [0, \infty)$  is increasing and  $f(\frac{1}{2}) > 1$ , show that  $\int_0^1 f(x)dx > \frac{1}{2}$ .

Problem 3 Show that if  $f$  is a continuous function on  $[a, b]$  then there exists a function  $g : [a, b] \rightarrow \mathbb{R}$  such that  $g' = f$ .

Problem 4 If  $f : \mathbb{R} \rightarrow \mathbb{R}$  and  $g : \mathbb{R} \rightarrow \mathbb{R}$  are two functions which are continuous at 0, show that the function

$$h(x) = \max\{f(x), g(x)\}, \quad x \in \mathbb{R}$$

is also continuous at 0.

Problem 5 Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be a differentiable functions which satisfies

$$f(x) = 0, \text{ for } |x| > 10, \quad f(-1) = 1, \quad f(1) = -1.$$

Show that there are at least two values of  $x \in \mathbb{R}$  such that  $f'(x) = 0$ .