

**Massachusetts Institute of Technology**

**Notes for 18.721**

# INTRODUCTION TO ALGEBRAIC GEOMETRY

*(This is a preliminary draft. Please don't reproduce.)*



# TABLE OF CONTENTS

## Chapter 1: PLANE CURVES

- 1.1 The Affine Plane
- 1.2 The Projective Plane
- 1.3 Plane Projective Curves
- 1.4 Tangent Lines
- 1.5 Transcendence Degree
- 1.6 The Dual Curve
- 1.7 Resultants
- 1.8 Nodes and Cusps
- 1.9 Hensel's Lemma
- 1.10 Bézout's Theorem
- 1.11 The Plücker Formulas

## Chapter 2: AFFINE ALGEBRAIC GEOMETRY

- 2.1 Rings and Modules
- 2.2 The Zariski Topology
- 2.3 Some Affine Varieties
- 2.4 The Nullstellensatz
- 2.5 The Spectrum
- 2.6 Localization
- 2.7 Morphisms of Affine Varieties
- 2.8 Finite Group Actions

## Chapter 3: PROJECTIVE ALGEBRAIC GEOMETRY

- 3.1 Projective Varieties
- 3.2 Homogeneous Ideals
- 3.3 Product Varieties
- 3.4 Morphisms and Isomorphisms
- 3.5 Affine Varieties
- 3.6 Lines in Projective Three-Space

## Chapter 4: INTEGRAL MORPHISMS OF AFFINE VARIETIES

- 4.1 The Nakayama Lemma
- 4.2 Integral Extensions
- 4.3 Normalization
- 4.4 Geometry of Integral Morphisms
- 4.5 Finite Group Actions II
- 4.6 Chevalley's Finiteness Theorem
- 4.7 Dimension
- 4.8 Krull's Theorem
- 4.9 Double Planes

## **Chapter 5: STRUCTURE OF VARIETIES IN THE ZARISKI TOPOLOGY**

- 5.1 Modules, review
- 5.2 Valuations
- 5.3 Smooth Curves
- 5.4 Constructible sets
- 5.5 Closed Sets
- 5.6 Fibred Products
- 5.7 Projective Varieties are Proper
- 5.8 Fibre Dimension

## **Chapter 6: MODULES**

- 6.1 The Structure Sheaf
- 6.2  $\mathcal{O}$ -Modules
- 6.3 The Sheaf Property
- 6.4 Some  $\mathcal{O}$ -Modules
- 6.5 Direct Image
- 6.6 Twisting
- 6.7 Proof of Theorem 6.3.2

## **Chapter 7: COHOMOLOGY**

- 7.1 Cohomology of  $\mathcal{O}$ -Modules
- 7.2 Complexes
- 7.3 Characteristic Properties
- 7.4 Construction of Cohomology
- 7.5 Cohomology of the Twisting Modules
- 7.6 Cohomology of Hypersurfaces
- 7.7 Annihilators and Support
- 7.8 Three Theorems about Cohomology
- 7.9 Bézout's Theorem

## **Chapter 8: THE RIEMANN-ROCH THEOREM FOR CURVES**

- 8.1 Branched Coverings
- 8.2 Modules
- 8.3 Divisors
- 8.4 The Riemann-Roch Theorem I
- 8.5 The Birkhoff-Grothendieck Theorem
- 8.6 Differentials
- 8.7 Trace, and proof of Riemann-Roch
- 8.8 The Riemann-Roch Theorem II
- 8.9 Using Riemann-Roch

## Chapter 1 PLANE CURVES

- 1.1 The Affine Plane
- 1.2 The Projective Plane
- 1.3 Plane Projective Curves
- 1.4 Tangent Lines
- 1.5 Transcendence Degree
- 1.6 The Dual Curve
- 1.7 Resultants and Discriminants
- 1.8 Nodes and Cusps
- 1.9 Hensel's Lemma
- 1.10 Bézout's Theorem
- 1.11 The Plücker Formulas

Plane curves were the first algebraic varieties to be studied, so we begin with them. They provide helpful examples, and we will see in Chapter ?? how they control higher dimensional varieties. We will come back to curves in Chapter 8. Chapters 2 - 7 are about varieties of arbitrary dimension.

### 1.1 The Affine Plane


The  $n$ -dimensional *affine space*  $\mathbb{A}^n$  is the space of  $n$ -tuples of complex numbers. The *affine plane*  $\mathbb{A}^2$  is the two-dimensional affine space.

Let  $f(x_1, x_2)$  be an irreducible polynomial in two variables with complex coefficients. The set of points of the affine plane at which  $f$  vanishes, the *locus of zeros* of  $f$ , is called a *plane affine curve*. Let's denote this locus by  $X$ . Using vector notation  $x = (x_1, x_2)$ ,

$$(1.1.1) \quad X = \{x \mid f(x) = 0\}$$

The *degree* of the curve  $X$  is the degree of its irreducible defining polynomial  $f$ .

#### 1.1.2.



**The Cubic Curve  $y^2 = x^3 + x$  (real locus)**

**1.1.3. Note.** In contrast with complex polynomials in one variable, most polynomials in two or more variables are *irreducible* – they cannot be factored. This can be shown by a method called “counting constants”. For instance, quadratic polynomials in  $x_1, x_2$  depend on the six coefficients of the monomials of degree at most two. Linear polynomials  $ax_1 + bx_2 + c$  depend on three coefficients, but the product of two linear polynomials depends on only five parameters, because a scalar factor can be moved from one of the linear polynomials to the other. So the quadratic polynomials cannot all be written as products of linear polynomials. This reasoning is fairly convincing. It can be justified formally in terms of *dimension*, which will be discussed in Chapter ??.

□

We will get an understanding of the geometry of a plane curve as we go along, and we mention just one point here. A plane curve is called a *curve* because it is defined by one equation in two variables. Its *algebraic* dimension is one. But because our scalars are complex numbers, it will be a surface, geometrically. This is analogous to the fact that the *affine line*  $\mathbb{A}^1$  is the plane of complex numbers.

One can see that a plane curve  $X$  is a surface by inspecting its projection to the affine  $x_1$ -line. One writes the defining polynomial as a polynomial in  $x_2$ , whose coefficients  $c_i = c_i(x_1)$  are polynomials in  $x_1$ :

$$f(x_1, x_2) = c_0 x_2^d + c_1 x_2^{d-1} + \cdots + c_d$$

Let's suppose that  $d$  is positive, i.e., that  $f$  isn't a polynomial in  $x_1$  alone (in which case, since it is irreducible, it would be linear).

The *fibre* of a map  $V \rightarrow U$  over a point  $p$  of  $U$  is the inverse image of  $p$ , the set of points of  $V$  that map to  $p$ . The fibre of the projection  $X \rightarrow \mathbb{A}^1$  over the point  $x_1 = a$  is the set of points  $(a, b)$  for which  $b$  is a root of the one-variable polynomial

$$f(a, x_2) = \bar{c}_0 x_2^d + \bar{c}_1 x_2^{d-1} + \cdots + \bar{c}_d$$

with  $\bar{c}_i = c_i(a)$ . There will be finitely many points in this fibre, and the fibre won't be empty unless  $f(a, x_2)$  is a constant. So the curve  $X$  covers most of the  $x_1$ -line, a complex plane, finitely often.

#### (1.1.4) changing coordinates

We allow linear changes of variable and translations in the affine plane  $\mathbb{A}^2$ . When a point  $x$  is written as the column vector  $x = (x_1, x_2)^t$ , the coordinates  $x' = (x'_1, x'_2)^t$  after such a change of variable will be related to  $x$  by the formula

$$(1.1.5) \quad x = Qx' + a$$

where  $Q$  is an invertible  $2 \times 2$  matrix with complex coefficients and  $a = (a_1, a_2)^t$  is a complex translation vector. This changes a polynomial equation  $f(x) = 0$ , to  $f(Qx' + a) = 0$ . One may also multiply a polynomial  $f$  by a nonzero complex scalar without changing the locus  $\{f = 0\}$ . Using these operations, all *lines*, plane curves of degree 1, become equivalent.

An *affine conic* is a plane affine curve of degree two. Every affine conic is equivalent to one of the loci

$$(1.1.6) \quad x_1^2 - x_2^2 = 1 \quad \text{or} \quad x_2 = x_1^2$$

The proof of this is similar to the one used to classify real conics. The two loci might be called a complex 'hyperbola' and 'parabola', respectively. The complex 'ellipse'  $x_1^2 + x_2^2 = 1$  becomes the 'hyperbola' when one multiplies  $x_2$  by  $i$ .

On the other hand, there are infinitely many inequivalent cubic curves. Cubic polynomials in two variables depend on the coefficients of the ten monomials  $1, x_1, x_2, x_1^2, x_1x_2, x_2^2, x_1^3, x_1^2x_2, x_1x_2^2, x_2^3$  of degree at most 3 in  $x$ . Linear changes of variable, translations, and scalar multiplication give us only seven scalars to work with, leaving three essential parameters.

## 1.2 The Projective Plane

The  $n$ -dimensional *projective space*  $\mathbb{P}^n$  is the set of equivalence classes of *nonzero* vectors  $x = (x_0, x_1, \dots, x_n)$ , the equivalence relation being

$$(1.2.1) \quad (x'_0, \dots, x'_n) \sim (x_0, \dots, x_n) \quad \text{if} \quad (x'_0, \dots, x'_n) = (\lambda x_0, \dots, \lambda x_n)$$

for some nonzero complex number  $\lambda$ . The equivalence classes are the *points* of  $\mathbb{P}^n$ , and one often refers to a point by a particular vector in its class.

Points of  $\mathbb{P}^n$  correspond bijectively to one-dimensional subspaces of  $\mathbb{C}^{n+1}$ . When  $x$  is a nonzero vector, the vectors  $\lambda x$ , together with the zero vector, form the one-dimensional subspace of the complex vector space  $\mathbb{C}^{n+1}$  spanned by  $x$ .

The *projective plane*  $\mathbb{P}^2$  is the two-dimensional projective space. Its points are equivalence classes of nonzero vectors  $(x_0, x_1, x_2)$ .

### (1.2.2) the projective line

Points of the *projective line*  $\mathbb{P}^1$  are equivalence classes of nonzero vectors  $(x_0, x_1)$ . If  $x_0$  isn't zero, we may multiply by  $\lambda = x_0^{-1}$  to normalize the first entry of  $(x_0, x_1)$  to 1, and write the point it represents in a unique way as  $(1, u)$ , with  $u = x_1/x_0$ . There is one remaining point, the point represented by the vector  $(0, 1)$ . The projective line  $\mathbb{P}^1$  can be obtained by adding this point, called the *point at infinity*, to the affine  $u$ -line, which is a complex plane. Topologically,  $\mathbb{P}^1$  is a two-dimensional sphere.

### (1.2.3) lines in projective space

A *line* in projective space  $\mathbb{P}^n$  is determined by a pair of distinct points  $p$  and  $q$ . When  $p$  and  $q$  are represented by specific vectors, the set of points  $\{rp + sq\}$ , with  $r, s$  in  $\mathbb{C}$  not both zero is a line. Points of this line correspond bijectively to points of the projective line  $\mathbb{P}^1$ , by

$$(1.2.4) \quad rp + sq \longleftrightarrow (r, s)$$

A line in the projective plane  $\mathbb{P}^2$  can also be described as the locus of solutions of a homogeneous linear equation

$$(1.2.5) \quad s_0x_0 + s_1x_1 + s_2x_2 = 0$$

**1.2.6. Lemma.** *In the projective plane, two distinct lines have exactly one point in common and, in a projective space of any dimension, a pair of distinct points is contained in exactly one line.*  $\square$

### (1.2.7) the standard covering of $\mathbb{P}^2$

If the first entry  $x_0$  of a point  $p = (x_0, x_1, x_2)$  of the projective plane  $\mathbb{P}^2$  isn't zero, we may normalize it to 1 without changing the point:  $(x_0, x_1, x_2) \sim (1, u_1, u_2)$ , where  $u_i = x_i/x_0$ . We did the analogous thing for  $\mathbb{P}^1$  above. The representative vector  $(1, u_1, u_2)$  is uniquely determined by  $p$ , so points with  $x_0 \neq 0$  correspond bijectively to points of an affine plane  $\mathbb{A}^2$  with coordinates  $(u_1, u_2)$ :

$$(x_0, x_1, x_2) \sim (1, u_1, u_2) \longleftrightarrow (u_1, u_2)$$

We regard the affine plane as a subset of  $\mathbb{P}^2$  by this correspondence, and we denote that subset by  $\mathbb{U}^0$ . The points of  $\mathbb{U}^0$ , those with  $x_0 \neq 0$ , are the *points at finite distance*. The *points at infinity* of  $\mathbb{P}^2$ , those of the form  $(0, x_1, x_2)$ , are on the *line at infinity*  $L^0$ , the locus  $\{x_0 = 0\}$ . The projective plane is the union of the two sets  $\mathbb{U}^0$  and  $L^0$ . When a point is given by a coordinate vector, we can assume that the first coordinate is either 1 or 0.

There is an analogous correspondence between points  $(x_0, 1, x_2)$  and points of an affine plane  $\mathbb{A}^2$ , and between points  $(x_0, x_1, 1)$  and points of  $\mathbb{A}^2$ . We denote the subsets  $\{x_1 \neq 0\}$  and  $\{x_2 \neq 0\}$  by  $\mathbb{U}^1$  and  $\mathbb{U}^2$ , respectively. The three sets  $\mathbb{U}^0, \mathbb{U}^1, \mathbb{U}^2$  form the *standard covering* of  $\mathbb{P}^2$  by three *standard affine open sets*. Since the vector  $(0, 0, 0)$  has been ruled out, every point of  $\mathbb{P}^2$  lies in at least one of the standard affine open sets. Points whose three coordinates are nonzero lie in all of them.


**1.2.8. Note.** Which points of  $\mathbb{P}^2$  are at infinity depends on which of the standard affine open sets is taken to be the one at finite distance. When the coordinates are  $(x_0, x_1, x_2)$ , I like to normalize  $x_0$  to 1, as above. Then the points at infinity are those of the form  $(0, x_1, x_2)$ . But when coordinates are  $(x, y, z)$ , I may normalize  $z$  to 1. Then the points at infinity are the points  $(x, y, 0)$ . I hope this won't cause too much confusion.  $\square$

### (1.2.9) digression: the real projective plane

The points of the *real projective plane*  $\mathbb{RP}^2$  are equivalence classes of nonzero real vectors  $x = (x_0, x_1, x_2)$ , the equivalence relation being  $x' \sim x$  if  $x' = \lambda x$  for some nonzero real number  $\lambda$ . The real projective plane can also be thought of as the set of one-dimensional subspaces of the real vector space  $V = \mathbb{R}^3$ .

The plane  $U : \{x_0 = 1\}$  in  $V = \mathbb{R}^3$  is analogous to the standard affine open subset  $\mathbb{U}^0$  in the complex projective plane  $\mathbb{P}^2$ . We can project  $V$  from the origin  $p_0 = (0, 0, 0)$  to  $U$ , sending a point  $x = (x_0, x_1, x_2)$  of  $V$  distinct from  $p_0$  to the point  $(1, u_1, u_2)$ , with  $u_i = x_i/x_0$ . The fibres of this projection are the lines through  $p_0$  and  $x$ , with  $p_0$  omitted. Looking from the origin,  $U$  becomes a “picture plane”.

### 1.2.10.




This illustration is from Dürer's book on perspective

The projection to  $U$  is undefined at the points  $(0, x_1, x_2)$ , which are orthogonal to the  $x_0$ -axis. The line connecting such a point to  $p_0$  doesn't meet  $U$ . Those points correspond to the points at infinity of  $\mathbb{RP}^2$ .

The projection from 3-space to a picture plane goes back to the 16th century, the time of Desargues and Dürer. Projective coordinates were introduced by Möbius, but not until 200 years later.



### 1.2.11.



**A Schematic Representation of the real Projective Plane, with a Conic**

This figure shows the plane  $W: x+y+z=1$  in the real vector space  $\mathbb{R}^3$ . If  $p = (x, y, z)$  is a nonzero vector, the one-dimensional subspace spanned by  $p$  will meet  $W$  in a single point, unless  $p$  is on the line  $x+y+z=0$ . So the plane is a faithful representation of most of  $\mathbb{RP}^2$ .

### (1.2.12) changing coordinates in the projective plane

An invertible  $3 \times 3$  matrix  $P$  determines a linear change of coordinates in  $\mathbb{P}^2$ . With  $x = (x_0, x_1, x_2)^t$  and  $x' = (x'_0, x'_1, x'_2)^t$  represented as column vectors, the coordinate change is given by

$$(1.2.13) \quad Px' = x$$

As the next proposition shows, four special points, the three points  $e_0 = (1, 0, 0)^t, e_1 = (0, 1, 0)^t, e_2 = (0, 0, 1)^t$ , together with the point  $\epsilon = (1, 1, 1)^t$ , determine the coordinates.

**1.2.14. Proposition.** *Let  $p_0, p_1, p_2, q$  be four points of  $\mathbb{P}^2$ , no three of which lie on a line. There is, up to scalar factor, a unique linear coordinate change  $Px' = x$  such that  $Pp_i = e_i$  and  $Pq = \epsilon$ .*

*proof.* The hypothesis that the points  $p_0, p_1, p_2$  don't lie on a line means that the vectors that represent those points are independent. They span  $\mathbb{C}^3$ . So  $q$  will be a combination  $q = c_0p_0 + c_1p_1 + c_2p_2$ , and because no three of the points lie on a line, the coefficients  $c_i$  will be nonzero. We can *scale* the vectors  $p_i$  (multiply them by nonzero scalars) to make  $q = p_0 + p_1 + p_2$  without changing the points. Next, the columns of  $P$  can be an arbitrary set of independent vectors. We let them be  $p_0, p_1, p_2$ . Then  $Pe_i = p_i$ , and  $P\epsilon = q$ . The matrix  $P$  is unique up to scalar factor, as can be verified by looking the reasoning over.  $\square$

### (1.2.15) conics

A polynomial  $f(x_0, x_1, x_2)$  is *homogeneous*, and of *degree*  $d$ , if all monomials that appear with nonzero coefficient have (total) degree  $d$ . For example,  $x_0^3 + x_1^3 - x_0x_1x_2$  is a homogeneous cubic polynomial.

A homogeneous quadratic polynomial is a combination of the six monomials

$$x_0^2, x_1^2, x_2^2, x_0x_1, x_1x_2, x_0x_2$$

A *conic* is the locus of zeros of an irreducible homogeneous quadratic polynomial.

**1.2.16. Proposition.** *For any conic  $C$ , there is a choice of coordinates so that  $C$  becomes the locus*

$$x_0x_1 + x_0x_2 + x_1x_2 = 0$$

*proof.* Since the conic  $C$  isn't a line, it will contain three points that aren't colinear. Let's leave the verification of this fact as an exercise. We choose three non-colinear points on  $C$ , and adjust coordinates so that they become the points  $e_0, e_1, e_2$ . Let  $f$  be the quadratic polynomial in those coordinates whose zero locus is  $C$ . Because  $e_0$  is a point of  $C$ ,  $f(1, 0, 0) = 0$ , and therefore the coefficient of  $x_0^2$  in  $f$  is zero. Similarly, the coefficients of  $x_1^2$  and  $x_2^2$  are zero. So  $f$  has the form

$$f = ax_0x_1 + bx_0x_2 + cx_1x_2$$

Since  $f$  is irreducible,  $a, b, c$  aren't zero. By scaling appropriately, we can make  $a = b = c = 1$ . We will be left with the polynomial  $x_0x_1 + x_0x_2 + x_1x_2$ .  $\square$

### 1.3 Plane Projective Curves

The loci in projective space that are studied in algebraic geometry are the ones that can be defined by systems of *homogeneous* polynomial equations. The reason for homogeneity is that the vectors  $(a_0, \dots, a_n)$  and  $(\lambda a_0, \dots, \lambda a_n)$  represent the same point of  $\mathbb{P}^n$ .

To explain this, we write a polynomial  $f(x_0, \dots, x_n)$  as a sum of its *homogeneous parts*:

$$(1.3.1) \quad f = f_0 + f_1 + \dots + f_d$$

where  $f_0$  is the constant term,  $f_1$  is the linear part, etc., and  $d$  is the degree of  $f$ .

**1.3.2. Lemma.** *Let  $f$  be a polynomial of degree  $d$ , and let  $a = (a_0, \dots, a_n)$  be a nonzero vector. Then  $f(\lambda a) = 0$  for every nonzero complex number  $\lambda$  if and only if  $f_i(a)$  is zero for every  $i = 0, \dots, d$ .*

Thus we may as well work with homogeneous equations.

*proof of Lemma 1.3.2.*  $f(\lambda x_0, \dots, \lambda x_n) = f_0 + \lambda f_1(x) + \lambda^2 f_2(x) + \dots + \lambda^d f_d(x)$ . When we evaluate at some given  $x$ , the right side of this equation becomes a polynomial of degree at most  $d$  in  $\lambda$ . Since a nonzero polynomial of degree at most  $d$  has at most  $d$  roots,  $f(\lambda x)$  won't be zero for *every*  $\lambda$  unless that polynomial is zero.  $\square$

**1.3.3. Lemma.** *If a homogeneous polynomial  $f$  is a product  $gh$  of polynomials, then  $g$  and  $h$  are homogeneous, and the zero locus  $\{f = 0\}$  in projective space is the union of the two loci  $\{g = 0\}$  and  $\{h = 0\}$ .*  $\square$

It is also true that relatively prime homogeneous polynomials  $f$  and  $g$  have only finitely many common zeros. This isn't obvious. It will be proved below, in Proposition 1.3.11.

#### (1.3.4) loci in the projective line

Before going to plane curves, we describe the zero locus in the projective line  $\mathbb{P}^1$  of a homogeneous polynomial in two variables.

**1.3.5. Lemma.** *Every nonzero homogeneous polynomial  $f(x, y) = a_0x^d + a_1x^{d-1}y + \dots + a_dy^d$  with complex coefficients is a product of homogeneous linear polynomials that are unique up to scalar factor.*

To prove this, one uses the fact that the field of complex numbers is algebraically closed. A one-variable complex polynomial factors into linear factors in the polynomial ring  $\mathbb{C}[y]$ . To factor  $f(x, y)$ , one may factor the one-variable polynomial  $f(1, y)$  into linear factors, substitute  $y/x$  for  $y$ , and multiply the result by  $x^d$ . When one adjusts scalar factors, one will obtain the expected factorization of  $f(x, y)$ . For instance, to factor  $f(x, y) = x^2 - 3xy + 2y^2$ , substitute  $x = 1$ :  $2y^2 - 3y + 1 = 2(y - 1)(y - \frac{1}{2})$ . Substituting  $y = y/x$  and multiplying by  $x^2$ ,  $f(x, y) = 2(y - x)(y - \frac{1}{2}x)$ . The scalar 2 can be distributed arbitrarily among the linear factors.  $\square$

Adjusting scalar factors, we may write a homogeneous polynomial as a product of the form

$$(1.3.6) \quad f(x, y) = (v_1x - u_1y)^{r_1} \dots (v_kx - u_ky)^{r_k}$$

where no factor  $v_i x - u_i y$  is a constant multiple of another, and where  $r_1 + \cdots + r_k$  is the degree of  $f$ . The exponent  $r_i$  is the *multiplicity* of the linear factor  $v_i x - u_i y$ .

A linear polynomial  $vx - uy$  determines a point  $(u, v)$  in the projective line  $\mathbb{P}^1$ , the unique *zero* of that polynomial, and changing the polynomial by a scalar factor doesn't change its zero. Thus the linear factors of the homogeneous polynomial (1.3.6) determine points of  $\mathbb{P}^1$ , the *zeros* of  $f$ . The points  $(u_i, v_i)$  are zeros of multiplicity  $r_i$ . The total number of those points, counted with multiplicity, will be the degree of  $f$ .

The zero  $(u_i, v_i)$  of  $f$  corresponds to a root  $x = u_i/v_i$  of multiplicity  $r_i$  of the one-variable polynomial  $f(x, 1)$ , except when the zero is the point  $(1, 0)$ . This happens when the coefficient  $a_0$  of  $f$  is zero, and  $y$  is a factor of  $f$ . One could say that  $f(x, y)$  has a zero at infinity in that case.

This sums up the information contained in an algebraic locus in the projective line. It will be a finite set of points with multiplicities.

### (1.3.7) intersections with a line

Let  $Z$  be the zero locus of a homogeneous polynomial  $f(x_0, \dots, x_n)$  of degree  $d$  in projective space  $\mathbb{P}^n$ , and let  $L$  be a line in  $\mathbb{P}^n$  (1.2.4). Say that  $L$  is the set of points  $rp + sq$ , where  $p = (a_0, \dots, a_n)$  and  $q = (b_0, \dots, b_n)$  are represented by specific vectors, so that  $L$  corresponds to the projective line  $\mathbb{P}^1$  by  $rp + sq \leftrightarrow (r, s)$ . Let's also assume that  $L$  isn't entirely contained in the zero locus  $Z$ . The intersection  $Z \cap L$  corresponds to the zero locus in  $\mathbb{P}^1$  of the polynomial in  $r, s$  that is obtained by substituting  $rp + sq$  into  $f$ . This substitution yields a homogeneous polynomial  $\bar{f}(r, s)$  in  $r, s$ , of degree  $d$ . For example, if  $f = x_0x_1 + x_0x_2 + x_1x_2$ , then with  $p = (a_0, a_1, a_2)$  and  $q = (b_0, b_1, b_2)$ ,  $\bar{f}$  is the following quadratic polynomial in  $r, s$ :

$$\begin{aligned}\bar{f}(r, s) &= f(rp + sq) = (ra_0 + sb_0)(ra_1 + sb_1) + (ra_0 + sb_0)(ra_2 + sb_2) + (ra_1 + sb_1)(ra_2 + sb_2) \\ &= (a_0a_1 + a_0a_2 + a_1a_2)r^2 + \left(\sum_{i \neq j} a_i b_j\right)rs + (b_0b_1 + b_0b_2 + b_1b_2)s^2\end{aligned}$$

The zeros of  $\bar{f}$  in  $\mathbb{P}^1$  correspond to the points of  $Z \cap L$ . There will be  $d$  zeros, when counted with multiplicity.

**1.3.8. Definition.** With notation as above, the *intersection multiplicity* of  $Z$  and  $L$  at a point  $p$  is the multiplicity of zero of the polynomial  $\bar{f}$ .  $\square$

**1.3.9. Corollary.** Let  $Z$  be the zero locus in  $\mathbb{P}^n$  of a homogeneous polynomial  $f$ , and let  $L$  be a line in  $\mathbb{P}^n$  that isn't contained in  $Z$ . The number of intersections of  $Z$  and  $L$ , counted with multiplicity, is equal to the degree of  $f$ .  $\square$

### (1.3.10) loci in the projective plane

**1.3.11. Proposition.** Homogeneous polynomials  $f_1, \dots, f_r$  in  $x, y, z$  with no common factor have finitely many common zeros.

The proof of this proposition is below. It shows that the most interesting type of locus in the projective plane is the zero set of a single equation.

The locus of zeros of an irreducible homogeneous polynomial  $f$  is called a *plane projective curve*. The *degree* of a plane projective curve is the degree of its irreducible defining polynomial.

**1.3.12. Note.** Suppose that a homogeneous polynomial is reducible, say  $f = g_1 \cdots g_k$ , where  $g_i$  are irreducible, and such that  $g_i$  and  $g_j$  don't differ by a scalar factor when  $i \neq j$ . Then the zero locus  $C$  of  $f$  is the union of the zero loci  $V_i$  of the factors  $g_i$ . In this case,  $C$  may be called a *reducible curve*.

When there are multiple factors, say  $f = g_1^{e_1} \cdots g_k^{e_k}$  and some  $e_i$  are greater than 1, it is still true that the locus  $C : \{f = 0\}$  will be the union of the loci  $V_i : \{g_i = 0\}$ , but the connection between the geometry of  $C$  and the algebra is weakened. In this situation, the structure of a *scheme* becomes useful. We won't discuss schemes. The only situation in which we will need to keep track of multiple factors is when counting intersections with another curve  $D$ . For this purpose, one can define the *divisor* of  $f$  to be the integer combination  $e_1V_1 + \cdots + e_kV_k$ .  $\square$

We need a lemma for the proof of Proposition 1.3.11. The ring  $\mathbb{C}[x, y]$  embeds into its field of fractions  $F$ , which is the field of rational functions  $\mathbb{C}(x, y)$  in  $x, y$ . The polynomial ring  $\mathbb{C}[x, y, z]$  is a subring of the one-variable polynomial ring  $F[z]$ . It can be useful to study a problem in the principal ideal domain  $F[z]$  first because its algebra is simpler.

Recall that the *unit ideal* of a ring  $R$  is the ring  $R$  itself.

**1.3.13. Lemma.** *Let  $F = \mathbb{C}(x, y)$  be the field of rational functions in  $x, y$ .*

(i) *Let  $f_1, \dots, f_k$  be homogeneous polynomials in  $x, y, z$  with no common factor. Their greatest common divisor in  $F[z]$  is 1, and therefore  $f_1, \dots, f_k$  generate the unit ideal of  $F[z]$ . There is an equation of the form  $\sum g'_i f_i = 1$  with  $g'_i$  in  $F[z]$ .*

(ii) *Let  $f$  be an irreducible polynomial in  $\mathbb{C}[x, y, z]$  of positive degree in  $z$ , but not divisible by  $z$ . Then  $f$  is also an irreducible element of  $F[z]$ .*

*proof.* (i) Let  $h'$  be an element of  $F[z]$  that isn't a unit of  $F[z]$ , i.e., that isn't an element of  $F$ . Suppose that, for every  $i$ ,  $h'$  divides  $f_i$  in  $F[z]$ , say  $f_i = u'_i h'$ . The coefficients of  $h'$  and  $u'_i$  have denominators that are polynomials in  $x, y$ . We clear denominators from the coefficients, to obtain elements of  $\mathbb{C}[x, y, z]$ . This will give us equations of the form  $d_i f_i = u_i h$ , where  $d_i$  are polynomials in  $x, y$  and  $u_i, h$  are polynomials in  $x, y, z$ .

Since  $h$  isn't in  $F$ , it will have positive degree in  $z$ . Let  $g$  be an irreducible factor of  $h$  of positive degree in  $z$ . Then  $g$  divides  $d_i f_i$  but doesn't divide  $d_i$  which has degree zero in  $z$ . So  $g$  divides  $f_i$ , and this is true for every  $i$ . This contradicts the hypothesis that  $f_1, \dots, f_k$  have no common factor.

(ii) Say that  $f(x, y, z)$  factors in  $F[z]$ ,  $f = g' h'$ , where  $g'$  and  $h'$  are polynomials of positive degree in  $z$  with coefficients in  $F$ . When we clear denominators from  $g'$  and  $h'$ , we obtain an equation of the form  $df = gh$ , where  $g$  and  $h$  are polynomials in  $x, y, z$  of positive degree in  $z$  and  $d$  is a polynomial in  $x, y$ . Then neither  $f$  nor  $g$  divides  $d$ , so  $f$  must be reducible.  $\square$

*proof of Proposition 1.3.11.* We are to show that homogeneous polynomials  $f_1, \dots, f_r$  in  $x, y, z$  with no common factor have finitely many common zeros. Lemma 1.3.13 tells us that we may write  $\sum g'_i f_i = 1$ , with  $g'_i$  in  $F[z]$ . Clearing denominators from  $g'_i$  gives us an equation of the form

$$\sum g_i f_i = d$$

where  $d$  is a polynomial in  $x, y$  and  $g_i$  are polynomials in  $x, y, z$ . Taking suitable homogeneous parts of  $d$  and  $g_i$  produces an equation  $\sum g_i f_i = d$  in which all terms are homogeneous.

Lemma 1.3.5 asserts that  $d$  is a product of linear polynomials, say  $d = \ell_1 \cdots \ell_r$ . A common zero of  $f_1, \dots, f_k$  is also a zero of  $d$ , and therefore it is a zero of  $\ell_j$  for some  $j$ . It suffices to show that, for every  $j$ ,  $f_1, \dots, f_r$  and  $\ell_j$  have finitely many common zeros.

Since  $f_1, \dots, f_k$  have no common factor, there is at least one  $f_i$  that isn't divisible by  $\ell_j$ . Corollary 1.3.9 shows that  $f_i$  and  $\ell_j$  have finitely many common zeros. Therefore  $f_1, \dots, f_k$  and  $\ell_j$  have finitely many common zeros for every  $j$ .  $\square$

**1.3.14. Corollary.** *Every locus in the projective plane  $\mathbb{P}^2$  that can be defined by a system of homogeneous polynomial equations is a finite union of points and curves.*  $\square$

The next corollary is a special case of the Strong Nullstellensatz, which will be proved in the next chapter.

**1.3.15. Corollary.** *Let  $f$  be an irreducible homogeneous polynomial in three variables, that vanishes on an infinite set  $S$  of points of  $\mathbb{P}^2$ . If another homogeneous polynomial  $g$  vanishes on  $S$ , then  $f$  divides  $g$ . Therefore, if an irreducible polynomial vanishes on an infinite set  $S$ , that polynomial is unique up to scalar factor.*

*proof.* If the irreducible polynomial  $f$  doesn't divide  $g$ , then  $f$  and  $g$  have no common factor, and therefore they have finitely many common zeros.  $\square$

### (1.3.16) the classical topology

The usual topology on the affine space  $\mathbb{A}^n$  will be called the *classical topology*. A subset  $U$  of  $\mathbb{A}^n$  is open in the classical topology if, whenever  $U$  contains a point  $p$ , it contains all points sufficiently near to  $p$ . We call this

the classical topology to distinguish it from another topology, the *Zariski topology*, which will be discussed in the next chapter.

The projective space  $\mathbb{P}^n$  also has a classical topology. A subset  $U$  of  $\mathbb{P}^n$  is open if, whenever a point  $p$  of  $U$  is represented by a vector  $(x_0, \dots, x_n)$ , all vectors  $x' = (x'_0, \dots, x'_n)$  sufficiently near to  $x$  represent points of  $U$ .

### (1.3.17) isolated points

A point  $p$  of a topological space  $X$  is *isolated* if both  $\{p\}$  and its complement  $X - \{p\}$  are closed sets, or if  $\{p\}$  is both open and closed. If  $X$  is a subset of  $\mathbb{A}^n$  or  $\mathbb{P}^n$ , a point  $p$  of  $X$  is isolated in the classical topology if  $X$  doesn't contain points  $p'$  distinct from  $p$ , but arbitrarily close to  $p$ .

**1.3.18. Proposition** *Let  $n$  be an integer greater than one. The zero locus of a polynomial in  $\mathbb{A}^n$  or in  $\mathbb{P}^n$  contains no points that are isolated in the classical topology.*

**1.3.19. Lemma.** *Let  $f$  be a polynomial of degree  $d$  in  $n$  variables. When coordinates  $x_1, \dots, x_n$  are chosen suitably,  $f(x)$  will be a monic polynomial of degree  $d$  in the variable  $x_n$ .*

*proof.* We write  $f = f_0 + f_1 + \dots + f_d$ , where  $f_i$  is the homogeneous part of  $f$  of degree  $i$ , and we choose a point  $p$  of  $\mathbb{A}^n$  at which  $f_d$  isn't zero. We change variables so that  $p$  becomes the point  $(0, \dots, 0, 1)$ . We call the new variables  $x_1, \dots, x_n$  and the new polynomial  $f$ . Then  $f_d(0, \dots, 0, x_n)$  will be equal to  $cx_n^d$  for some nonzero constant  $c$ . When we adjust  $x_n$  by a scalar factor to make  $c = 1$ ,  $f$  will be monic.  $\square$

*proof of Proposition 1.3.18.* The proposition is true for loci in affine space and also for loci in projective space. We look at the affine case. Let  $f(x_1, \dots, x_n)$  be a polynomial with zero locus  $Z$ , and let  $p$  be a point of  $Z$ . We adjust coordinates so that  $p$  is the origin  $(0, \dots, 0)$  and  $f$  is monic in  $x_n$ . We relabel  $x_n$  as  $y$ , and write  $f$  as a polynomial in  $y$ . Let's write  $f(x, y) = \tilde{f}(y)$ :

$$\tilde{f}(y) = f(x, y) = y^d + c_{d-1}(x)y^{d-1} + \dots + c_0(x)$$

where  $c_i$  is a polynomial in  $x_1, \dots, x_{n-1}$ . For fixed  $x$ ,  $c_0(x)$  is the product of the roots of  $\tilde{f}(y)$ . Since  $p$  is the origin and  $f(p) = 0$ ,  $c_0(0) = 0$ . So  $c_0(x)$  will tend to zero with  $x$ . Then at least one root  $y$  of  $\tilde{f}(y)$  will tend to zero. This gives us points  $(x, y)$  of  $Z$  that are arbitrarily close to  $p$ .  $\square$

**1.3.20. Corollary.** *Let  $C'$  be the complement of a finite set of points in a plane curve  $C$ . In the classical topology, a continuous function  $g$  on  $C$  that is zero at every point of  $C'$  is identically zero.*  $\square$

## 1.4 Tangent Lines

### (1.4.1) notation for working locally

We will often want to inspect a plane curve  $C : \{f(x_0, x_1, x_2) = 0\}$  in a neighborhood of a particular point  $p$ . To do this we may adjust coordinates so that  $p$  becomes the point  $(1, 0, 0)$ , and look in the standard affine open set  $\mathbb{U}^0 : \{x_0 \neq 0\}$ . There,  $p$  becomes the origin in the affine  $x_1, x_2$ -plane, and  $C$  becomes the zero locus of the non-homogeneous polynomial  $f(1, x_1, x_2)$ .

Of course, it doesn't matter which variable we set to 1. If the variables are  $x, y, z$ , we may prefer to take for  $p$  the point  $(0, 0, 1)$  and work with the polynomial  $f(x, y, 1)$ .

This will be a standard notation for working locally.

**1.4.2. Lemma.** *A homogeneous polynomial  $f(x_0, x_1, x_2)$  not divisible by  $x_0$  is irreducible if and only if its dehomogenization  $f(1, x_1, x_2)$  is irreducible.*  $\square$

### (1.4.3) homogenizing and dehomogenizing

If  $f(x_0, x_1, x_2)$  is a polynomial,  $f(1, x_1, x_2)$  is called the *dehomogenization* of  $f$  with respect to the variable  $x_0$ .

A simple procedure, *homogenization*, inverts dehomogenization. Suppose given a non-homogeneous polynomial  $F(x_1, x_2)$  of degree  $d$ . To homogenize  $F$ , we replace the variables  $x_i$ ,  $i = 1, 2$ , by  $u_i = x_i/x_0$ . Then since  $u_i$  have degree zero in  $x$ , so does  $F(u_1, u_2)$ . When we multiply by  $x_0^d$ , the result will be a homogeneous polynomial of degree  $d$  in  $x_0, x_1, x_2$  that isn't divisible by  $x_0$ ,

We will come back to homogenization in Chapter 2.

#### (1.4.4) smooth points and singular points

Let  $C$  be the plane curve defined by an irreducible homogeneous polynomial  $f(x_0, x_1, x_2)$ , and let  $f_i$  denote the partial derivative  $\frac{\partial f}{\partial x_i}$ , which can be computed by the usual calculus formula. A point of  $C$  at which the partial derivatives  $f_i$  aren't all zero is called a *smooth point* of  $C$ . A point at which all partial derivatives are zero is a *singular point*. A curve is *smooth*, or *nonsingular*, if it contains no singular point; otherwise it is a *singular curve*.

The *Fermat curve*

$$(1.4.5) \quad x_0^d + x_1^d + x_2^d = 0$$

is smooth because the only common zero of the partial derivatives  $dx_0^{d-1}, dx_1^{d-1}, dx_2^{d-1}$ , which is  $(0, 0, 0)$ , doesn't represent a point of  $\mathbb{P}^2$ . The cubic curve  $x_0^3 + x_1^3 - x_0x_1x_2 = 0$  is singular at the point  $(0, 0, 1)$ .

The Implicit Function Theorem explains the meaning of smoothness. Suppose that  $p = (1, 0, 0)$  is a point of  $C$ . We set  $x_0 = 1$  and inspect the locus  $f(1, x_1, x_2) = 0$  in the standard affine open set  $\mathbb{U}^0$ . If  $f_2(p)$  isn't zero, the Implicit Function Theorem tells us that we can solve the equation  $f(1, x_1, x_2) = 0$  for  $x_2$  locally (for small  $x_1$ ) as an analytic function  $\varphi$  of  $x_1$ , with  $\varphi(0) = 0$ . Sending  $x_1$  to  $(1, x_1, \varphi(x_1))$  inverts the projection from  $C$  to the affine  $x_1$ -line, locally. So at a smooth point,  $C$  is locally homeomorphic to the affine line.

**1.4.6. Euler's Formula.** Let  $f$  be a homogeneous polynomial of degree  $d$  in the variables  $x_0, \dots, x_n$ . Then

$$\sum_i x_i \frac{\partial f}{\partial x_i} = d f.$$

*proof.* It is enough to check this formula when  $f$  is a monomial. As an example, let  $f$  be the monomial  $x^2y^3z$ , then

$$xf_x + yf_y + zf_z = x(2xy^3z) + y(3x^2y^2z) + z(x^2y^3) = 6x^2y^3z = 6f \quad \square$$

**1.4.7. Corollary.** (i) If all partial derivatives of an irreducible homogeneous polynomial  $f$  are zero at a point  $p$  of  $\mathbb{P}^2$ , then  $f$  is zero at  $p$ , and therefore  $p$  is a singular point of the curve  $\{f = 0\}$ .

(ii) At a smooth point of a plane curve, at least two partial derivatives will be nonzero.

(iii) The partial derivatives of an irreducible polynomial have no common (nonconstant) factor.

(iv) A plane curve has finitely many singular points.  $\square$

#### (1.4.8) tangent lines and flex points

Let  $C$  be the plane projective curve defined by an irreducible homogeneous polynomial  $f$ . A line  $L$  is *tangent* to  $C$  at a smooth point  $p$  if the intersection multiplicity of  $C$  and  $L$  at  $p$  is at least 2. (See (1.3.8).) There is a unique tangent line at a smooth point.

A smooth point  $p$  of  $C$  is a *flex point* if the intersection multiplicity of  $C$  and its tangent line at  $p$  is at least 3, and  $p$  is an *ordinary flex point* if the intersection multiplicity is equal to 3.

Let  $L$  be a line through a point  $p$  and let  $q$  be a point of  $L$  distinct from  $p$ . We represent  $p$  and  $q$  by specific vectors  $(p_0, p_1, p_2)$  and  $(q_0, q_1, q_2)$ , to write a variable point of  $L$  as  $p + tq$ , and we expand the restriction of  $f$

to  $L$  in a Taylor's Series. The Taylor expansion carries over to complex polynomials because it is an identity. Let  $f_i = \frac{\partial f}{\partial x_i}$  and  $f_{ij} = \frac{\partial^2 f}{\partial x_i \partial x_j}$ . Then

$$(1.4.9) \quad f(p + tq) = f(p) + \left( \sum_i f_i(p) q_i \right) t + \frac{1}{2} \left( \sum_{i,j} q_i f_{ij}(p) q_j \right) t^2 + O(3)$$

where the symbol  $O(3)$  stands for a polynomial in which all terms have degree at least 3 in  $t$ . The point  $q$  is missing from this parametrization, but this won't be important.

We will rewrite this equation twice. Let  $\nabla$  be the *gradient vector*  $(f_0, f_1, f_2)$ , let  $H$  be the *Hessian matrix* of  $f$ , the matrix of second partial derivatives:

$$(1.4.10) \quad H = \begin{pmatrix} f_{00} & f_{01} & f_{02} \\ f_{10} & f_{11} & f_{12} \\ f_{20} & f_{21} & f_{22} \end{pmatrix}$$

and let  $\nabla_p$  and  $H_p$  be the evaluations of  $\nabla$  and  $H$ , respectively, at  $p$ . So  $p$  is a smooth point of  $C$  if  $f(p) = 0$  and  $\nabla_p \neq 0$ . Regarding  $p$  and  $q$  as column vectors, Equation 1.4.9 can be written as

$$(1.4.11) \quad f(p + tq) = f(p) + (\nabla_p q) t + \frac{1}{2} (q^t H_p q) t^2 + O(3)$$

in which  $\nabla_p q$  and  $q^t H_p q$  are to be computed as matrix products.

The intersection multiplicity of  $C$  and  $L$  at  $p$  (1.3.8) is the lowest power of  $t$  that has nonzero coefficient in  $f(p + tq)$ . The intersection multiplicity is at least 1 if  $p$  lies on  $C$ , i.e., if  $f(p) = 0$ .

Suppose that  $p$  is a smooth point of  $C$ . Then  $L$  is tangent to  $C$  at  $p$  if the coefficient  $(\nabla_p q)$  of  $t$  is zero, and  $p$  is a flex point if  $(\nabla_p q)$  and  $(q^t H_p q)$  are both zero. The equation of the tangent line  $L$  at a smooth point  $p$  is  $\nabla_p x = 0$ , or

$$(1.4.12) \quad f_0(p)x_0 + f_1(p)x_1 + f_2(p)x_2 = 0$$

which tells us that a point  $q$  lies on  $L$  if the linear term in  $t$  of (1.4.11) is zero.

By the way, Taylor's formula shows that the restriction of  $f$  to every line through a singular point has a multiple zero. However, we will speak of tangent lines only at smooth points of the curve.

The next lemma is obtained by applying Euler's Formula to the entries of  $H_p$  and  $\nabla_p$ .

**1.4.13. Lemma.**

$$p^t H_p = (d-1) \nabla_p \quad \text{and} \quad \nabla_p p = df(p)$$

□

We rewrite Equation 1.4.9 one more time, using the notation  $\langle u, v \rangle$  to represent the symmetric bilinear form  $u^t H_p v$  on  $V = \mathbb{C}^3$ . It makes sense to say that this form vanishes on a pair of points of  $\mathbb{P}^2$ , because the condition  $\langle u, v \rangle = 0$  doesn't depend on the vectors that represent those points.

**1.4.14. Proposition.** *With notation as above,*

(i) *Equation (1.4.9) can be written as*

$$f(p + tq) = \frac{1}{d(d-1)} \langle p, p \rangle + \frac{1}{d-1} \langle p, q \rangle t + \frac{1}{2} \langle q, q \rangle t^2 + O(3)$$

(ii) *A point  $p$  is a smooth point of  $C$  if and only if  $\langle p, p \rangle = 0$  but  $\langle p, v \rangle$  is not identically zero.*

*proof.* (i) This follows from the formulas

$$(1.4.15) \quad p^t H_p = (d-1) \nabla_p \quad \text{and} \quad \nabla_p p = df(p)$$

which can be obtained by applying Euler's formula to the entries of  $H_p$  and  $\nabla_p$ .

(ii)  $\langle p, v \rangle = (d-1) \nabla_p v$  isn't identically zero at a smooth point  $p$  because  $\nabla_p$  won't be zero. □

**1.4.16. Corollary.** Let  $p$  be a smooth point of  $C$ , let  $q$  be a point of  $\mathbb{P}^2$  distinct from  $p$ , and let  $L$  be the line through  $p$  and  $q$ . Then

- (i)  $L$  is tangent to  $C$  at  $p$  if and only if  $\langle p, p \rangle = \langle p, q \rangle = 0$ , and
- (ii)  $p$  is a flex point of  $C$  with tangent line  $L$  if and only if  $\langle p, p \rangle = \langle p, q \rangle = \langle q, q \rangle = 0$ . □

**1.4.17. Theorem.** A smooth point  $p$  of the curve  $C$  is a flex point if and only if the determinant  $\det H_p$  of the Hessian matrix at  $p$  is zero.

*proof.* Let  $p$  be a smooth point of  $C$ , so that  $\langle p, p \rangle = 0$ . If  $\det H_p = 0$ , the form  $\langle u, v \rangle$  is degenerate. There is a nonzero null vector  $q$ , so that  $\langle p, q \rangle = \langle q, q \rangle = 0$ . But because  $\langle p, v \rangle$  isn't identically zero,  $q$  is distinct from  $p$ . So  $p$  is a flex point.

Conversely, suppose that  $p$  is a flex point and let  $q$  be a point on the tangent line at  $p$  and distinct from  $p$ , so that  $\langle p, p \rangle = \langle p, q \rangle = \langle q, q \rangle = 0$ . The restriction of the form to the two-dimensional space  $W$  spanned by  $p$  and  $q$  is zero, and this implies that the form is degenerate. If  $(p, q, v)$  is a basis of  $V$  with  $p, q$  in  $W$ , the matrix of the form will look like this:

$$\begin{pmatrix} 0 & 0 & * \\ 0 & 0 & * \\ * & * & * \end{pmatrix} \quad \square$$

**1.4.18. Proposition.**

- (i) Let  $f(x, y, z)$  be an irreducible homogeneous polynomial of degree at least two. The **Hessian determinant**  $\det H$  isn't divisible by  $f$ . In particular, it isn't identically zero.
- (ii) A curve that isn't a line has finitely many flex points.

*proof.* (i) Let  $C$  be the curve defined by  $f$ . If  $f$  divides the Hessian determinant, every smooth point of  $C$  will be a flex point. We set  $z = 1$  and look on the standard affine  $\mathbb{U}^2$ , choosing coordinates so that the origin  $p$  is a smooth point of  $C$ , and  $\frac{\partial f}{\partial y} \neq 0$  at  $p$ . The Implicit Function Theorem tells us that we can solve the equation  $f(x, y, 1) = 0$  for  $y$  locally, say  $y = \varphi(x)$ . The graph  $\Gamma : \{y = \varphi(x)\}$  will be equal to  $C$  in a neighborhood of  $p$  (see below). A point of  $\Gamma$  is a flex point if and only if  $\frac{d^2 \varphi}{dx^2}$  is zero there. If this is true for all points near to  $p$ , then  $\frac{d^2 \varphi}{dx^2}$  will be identically zero, and this implies that  $\varphi$  is linear:  $y = ax$ . Then  $y = ax$  solves  $f = 0$ , and therefore  $y - ax$  divides  $f(x, y, 1)$ . But  $f(x, y, z)$  is irreducible, and so is  $f(x, y, 1)$ . Therefore  $f(x, y, 1)$  is linear, contrary to hypothesis.

(ii) This follows from (i) and (1.3.11). The irreducible polynomial  $f$  and the Hessian determinant have finitely many common zeros. □

**1.4.19. Review.** (about the Implicit Function Theorem)

Let  $f(x, y)$  be a polynomial such that  $f(0, 0) = 0$  and  $\frac{df}{dy}(0, 0) \neq 0$ . The Implicit Function Theorem asserts that there is a unique analytic function  $\varphi(x)$ , defined for small  $x$ , such that  $\varphi(0) = 0$  and  $f(x, \varphi(x))$  is identically zero.

We make some further remarks. Let  $\mathcal{R}$  be the ring of functions that are defined and analytic for small  $x$ . In the ring  $\mathcal{R}[y]$  of polynomials in  $y$  with coefficients in  $\mathcal{R}$ , the polynomial  $y - \varphi(x)$ , which is monic in  $y$ , divides  $f(x, y)$ . To see this, we do division with remainder of  $f$  by  $y - \varphi(x)$ :

$$(1.4.20) \quad f(x, y) = (y - \varphi(x))q(x, y) + r(x)$$

The quotient  $q(x, y)$  is in  $\mathcal{R}[y]$ , and the remainder  $r(x)$  has degree zero in  $y$ , so it is in  $\mathcal{R}$ . Setting  $y = \varphi(x)$  in the equation, one sees that  $r(x) = 0$ .

Let  $\Gamma$  be the graph of  $\varphi$  in a suitable neighborhood  $U$  of the origin in  $x, y$ -space. Since  $f(x, y) = (y - \varphi(x))q(x, y)$ , the locus  $f(x, y) = 0$  in  $U$  has the form  $\Gamma \cup \Delta$ , where  $\Gamma$  is the graph of  $\varphi$  and  $\Delta$  is the zero locus of  $q(x, y)$ . Differentiating, we find that  $\frac{\partial f}{\partial y}(0, 0) = q(0, 0)$ . So  $q(0, 0) \neq 0$ . Then  $\Delta$  doesn't contain the origin, while  $\Gamma$  does. This implies that  $\Delta$  is disjoint from  $\Gamma$ , locally. A sufficiently small neighborhood  $U$  of the origin won't contain any of points  $\Delta$ . In such a neighborhood, the locus of zeros of  $f$  will be  $\Gamma$ . □



## 1.5 Transcendence degree

Let  $F \subset K$  be a field extension. A set  $\alpha = \{\alpha_1, \dots, \alpha_n\}$  of elements of  $K$  is *algebraically dependent* over  $F$  if there is a nonzero polynomial  $f(x_1, \dots, x_n)$  with coefficients in  $F$ , such that  $f(\alpha) = 0$ . If there is no such polynomial, the set  $\alpha$  is *algebraically independent* over  $F$ .

An infinite set is called algebraically independent if every finite subset is algebraically independent – if there is no polynomial relation among any finite set of its elements.

The set  $\{\alpha_1\}$  consisting of a single element of  $K$  will be algebraically dependent if  $\alpha_1$  is algebraic over  $F$ . Otherwise, it will be algebraically independent. In this case  $\alpha_1$  is said to be *transcendental* over  $F$ .

An algebraically independent set  $\alpha = \{\alpha_1, \dots, \alpha_n\}$  that isn't contained in a larger algebraically independent set is called a *transcendence basis* for  $K$  over  $F$ . If there is a finite transcendence basis, its order is the *transcendence degree* of the field extension  $K$  of  $F$ . Lemma 1.5.2 below shows that all transcendence basis for  $K$  over  $F$  have the same order, so the transcendence degree is well-defined. If there is no finite transcendence basis, the transcendence degree of  $K$  over  $F$  is infinite.

For example, let  $K = F(x_1, \dots, x_n)$  be the field of rational functions in  $n$  variables. The variables form a transcendence basis of  $K$  over  $F$ , and the transcendence degree of  $K$  over  $F$  is  $n$ .

A *domain* is a nonzero ring with no zero divisors, and a domain that contains a field  $F$  as a subring is called an  $F$ -algebra. We use the customary notation  $F[\alpha_1, \dots, \alpha_n]$  or  $F[\alpha]$  for the  $F$ -algebra generated by a set  $\alpha = \{\alpha_1, \dots, \alpha_n\}$ , and we may denote its field of fractions by  $F(\alpha_1, \dots, \alpha_n)$  or by  $F(\alpha)$ . The set  $\{\alpha_1, \dots, \alpha_n\}$  is algebraically independent over  $F$  if and only if the surjective map from the polynomial algebra  $F[x_1, \dots, x_n]$  to  $F[\alpha_1, \dots, \alpha_n]$  that sends  $x_i$  to  $\alpha_i$  is bijective.

**1.5.1. Lemma.** *Let  $K/F$  be a field extension, let  $\alpha = \{\alpha_1, \dots, \alpha_n\}$  be a set of elements of  $K$  that is algebraically independent over  $F$ , and let  $F(\alpha)$  be the field of fractions of  $F[\alpha]$ .*

- (i) *Every element of the field  $F(\alpha)$  that isn't in  $F$  is transcendental over  $F$ .*
- (ii) *If  $\beta$  is another element of  $K$ , the set  $\{\alpha_1, \dots, \alpha_n, \beta\}$  is algebraically dependent if and only if  $\beta$  is algebraic over  $F(\alpha)$ .*
- (iii) *The algebraically independent set  $\alpha$  is a transcendence basis if and only if every element of  $K$  is algebraic over  $F(\alpha)$ .*

*proof.* (i) We write an element  $z$  of  $F(\alpha)$  as a fraction  $p/q = p(\alpha)/q(\alpha)$ , where  $p(x)$  and  $q(x)$  are relatively prime polynomials. Suppose that  $z$  satisfies a nontrivial polynomial relation  $c_0 z^n + c_1 z^{n-1} + \dots + c_n = 0$  with  $c_i$  in  $F$ . We may assume that  $c_0 = 1$ . Substituting  $z = p/q$  and multiplying by  $q^n$  gives us the equation

$$p^n = -q(c_1 p^{n-1} + \dots + c_n q^{n-1})$$

Because  $\alpha$  is an algebraically independent set, this equation is equivalent with a polynomial equation in  $F[x]$ . It shows that  $q$  divides  $p^n$ , which contradicts the hypothesis that  $p$  and  $q$  are relatively prime.  $\square$

**1.5.2. Lemma.**

- (i) *Let  $K/F$  be a field extension. If  $K$  has a finite transcendence basis, then all algebraically independent subsets of  $K$  are finite, and all transcendence bases have the same order.*
- (ii) *If  $L \supset K \supset F$  are fields and if the degree  $[L : K]$  of  $L$  over  $K$  is finite, then  $K$  and  $L$  have the same transcendence degree over  $F$ .*

*proof.* (i) Let  $\alpha = \{\alpha_1, \dots, \alpha_r\}$  and  $\beta = \{\beta_1, \dots, \beta_s\}$ . Assume that  $K$  is algebraic over  $F(\alpha)$  and that the set  $\beta$  is algebraically independent. We show that  $s \leq r$ . The fact that all transcendence bases have the same order will follow: If both  $\alpha$  and  $\beta$  are transcendence bases, then  $s \leq r$ , and since we can interchange  $\alpha$  and  $\beta$ ,  $r \leq s$ .

The proof that  $s \leq r$  proceeds by reducing to the trivial case that  $\beta$  is a subset of  $\alpha$ . Suppose that some element of  $\beta$ , say  $\beta_s$ , isn't in the set  $\alpha$ . The set  $\beta' = \{\beta_1, \dots, \beta_{s-1}\}$  is algebraically independent, but it isn't a transcendence basis. So  $K$  isn't algebraic over  $F(\beta')$ . Since  $K$  is algebraic over  $F(\alpha)$ , there is at least one element of  $\alpha$ , say  $\alpha_r$ , that isn't algebraic over  $F(\beta')$ . Then  $\gamma = \beta' \cup \{\alpha_s\}$  will be an algebraically independent set of order  $s$ , and it will contain more elements of the set  $\alpha$  than  $\beta$  does. Induction shows that  $s \leq r$ .  $\square$

## 1.6 The Dual Curve

### (1.6.1) the dual plane

Let  $\mathbb{P}$  denote the projective plane with coordinates  $x_0, x_1, x_2$ , and let  $L$  be the line in  $\mathbb{P}$  with the equation

$$(1.6.2) \quad s_0x_0 + s_1x_1 + s_2x_2 = 0$$

The solutions of this equation determine the coefficients  $s_i$  only up to a common nonzero scalar factor, so the line  $L$  determines a point  $(s_0, s_1, s_2)$  in another projective plane  $\mathbb{P}^*$  called the *dual plane*. We denote that point by  $L^*$ . Moreover, a point  $p = (x_0, x_1, x_2)$  in  $\mathbb{P}$  determines a line in the dual plane, the line with the equation (1.6.2), when  $s_i$  are regarded as the variables and  $x_i$  as the scalar coefficients. We denote that line by  $p^*$ . The equation exhibits a duality between  $\mathbb{P}$  and  $\mathbb{P}^*$ . A point  $p$  of  $\mathbb{P}$  lies on the line  $L$  if and only if the equation is satisfied, and this means that, in  $\mathbb{P}^*$ , the point  $L^*$  lies on the line  $p^*$ .

### (1.6.3) the dual curve

Let  $C$  be a plane projective curve of degree at least two, and let  $U$  be the set of its smooth points. This is the complement of a finite subset of  $C$ . We define a map

$$U \xrightarrow{t} \mathbb{P}^*$$

as follows: Let  $p$  be a point of  $U$  and let  $L$  be the tangent line to  $C$  at  $p$ . Then  $t(p) = L^*$ , where  $L^*$  is the point of  $\mathbb{P}^*$  that corresponds to the line  $L$ .

Denoting the partial derivative  $\frac{\partial f}{\partial x_i}$  by  $f_i$  as before, the tangent line  $L$  at a smooth point  $p = (x_0, x_1, x_2)$  of  $C$  has the equation  $f_0x_0 + f_1x_1 + f_2x_2 = 0$  (1.4.12). Therefore  $L^*$  is the point

$$(1.6.4) \quad (s_0, s_1, s_2) = (f_0(x), f_1(x), f_2(x))$$

We'll drop some parentheses, denoting the image  $t(U)$  of  $U$  in  $\mathbb{P}^*$  by  $tU$ . Points of  $tU$  correspond to tangent lines at smooth points of  $C$ . We assume that  $C$  has degree at least two because, if  $C$  were a line,  $tU$  would be a point. Since the partial derivatives have no common factor, the tangent lines aren't constant when the degree is two or more.

??figure??

**1.6.5. Lemma.** *Let  $\varphi(s_0, s_1, s_2)$  be a homogeneous polynomial, and let  $g(x_0, x_1, x_2) = \varphi(f_0(x), f_1(x), f_2(x))$ . Then  $\varphi(s)$  is identically zero on  $tU$  if and only if  $g(x)$  is identically zero on  $U$ . This is true if and only if  $f$  divides  $g$ .*

*proof.* The first assertion follows from the fact that  $(s_0, s_1, s_2)$  and  $(f_0(x), f_1(x), f_2(x))$  represent the same point of  $\mathbb{P}^*$ , and the last one follows from Corollary 1.3.15.  $\square$

**1.6.6. Theorem.** *Let  $C$  be the plane curve defined by an irreducible homogeneous polynomial  $f$  of degree  $d$  at least two. With notation as above, the image  $tU$  is contained in a curve  $C^*$  in the dual space  $\mathbb{P}^*$ .*

The curve  $C^*$  referred to in the theorem is the *dual curve*.

*proof.* If an irreducible homogeneous polynomial  $\varphi(s)$  vanishes on  $tU$ , it will be unique up to scalar factor (Corollary 1.3.15).

Let's use vector notation:  $x = (x_0, x_1, x_2)$ ,  $s = (s_0, s_1, s_2)$ , and  $\nabla f = (f_0, f_1, f_2)$ .

. We show first that there is a nonzero polynomial  $\varphi(s)$ , not necessarily irreducible or homogeneous, that vanishes on  $tU$ . The field  $\mathbb{C}(x_0, x_1, x_2)$  has transcendence degree three over  $\mathbb{C}$ . Therefore the four polynomials  $f_0, f_1, f_2$ , and  $f$  are algebraically dependent. There is a nonzero polynomial  $\psi(s_0, s_1, s_2, t)$  such that  $\psi(f_0(x), f_1(x), f_2(x), f(x)) = \psi(\nabla f(x), f(x))$  is the zero polynomial. We can cancel factors of  $t$ , so we may assume that  $\psi$  isn't divisible by  $t$ . Let  $\varphi(s) = \psi(s_0, s_1, s_2, 0)$ . This isn't the zero polynomial when  $t$  doesn't divide  $\psi$ .

Let  $\bar{x} = (\bar{x}_1, \bar{x}_2, \bar{x}_3)$  be a vector that represents a point of  $U$ . Then  $f(\bar{x}) = 0$ , and therefore

$$\psi(\nabla f(\bar{x}), f(\bar{x})) = \psi(\nabla f(\bar{x}), 0) = \varphi(\nabla f(\bar{x}))$$

Since  $\psi(\nabla f(x), f(x))$  is identically zero,  $\varphi(\nabla f(\bar{x})) = 0$  for all  $\bar{x}$  in  $U$ .

Next, because the vectors  $\bar{x}$  and  $\lambda\bar{x}$  represent the same point of  $U$ ,  $\varphi(\nabla f(\lambda\bar{x})) = 0$ . Since  $f$  has degree  $d$ , the derivatives  $f_i$  are homogeneous of degree  $d - 1$ . Therefore  $\varphi(\nabla f(\lambda\bar{x})) = \varphi(\lambda^{d-1}\nabla f(\bar{x})) = 0$  for all  $\lambda$ . Because the scalar  $\lambda^{d-1}$  can be any complex number, Lemma 1.3.2 tells us that the homogeneous parts of  $\varphi(\nabla f(x))$  vanish for all  $x \in U$ . The homogeneous parts of degree  $r$  of  $\varphi(s)$  correspond to the homogeneous parts of degree  $r(d - 1)$  of  $\varphi(\nabla f(x))$ . So the homogeneous parts of  $\varphi(s)$  vanish on  $tU$ . This shows that there is a homogeneous polynomial  $\varphi(s)$  that vanishes on  $tU$ . We choose such a polynomial  $\varphi(s)$ . Let its degree be  $r$ .

If  $f$  has degree  $d$ , the polynomial  $g(x) = \varphi(\nabla f(x))$  will be homogeneous, of degree  $r(d - 1)$ . It will vanish on  $U$ , and therefore on  $C$  (1.3.20). So  $f$  will divide  $g$ . Finally, if  $\varphi(s)$  factors, then  $g(x)$  factors accordingly, and because  $f$  is irreducible, it will divide one of the factors of  $g$ . The corresponding factor of  $\varphi$  will vanish on  $tU$  (1.6.5). So we may replace the homogeneous polynomial  $\varphi$  by one of its irreducible factors.  $\square$

In principle, the proof of Theorem 1.6.6 gives a method for finding a polynomial that vanishes on the dual curve. Namely, looks for a polynomial relation among  $f_x, f_y, f_z, f$ , and then sets  $f = 0$ . However, it is usually painful to determine the defining polynomial of  $C^*$  explicitly. Most often, the degrees of  $C$  and  $C^*$  will be different, and several points of the dual curve  $C^*$  may correspond to a singular point of  $C$ , and vice versa.

The computation is easy for a conic.

### 1.6.7. Examples.

(i) *(the dual of a conic)* Let  $f = x_0x_1 + x_0x_2 + x_1x_2$  and let  $C$  be the conic  $f = 0$ . Let  $(s_0, s_1, s_2) = (f_0, f_1, f_2) = (x_1 + x_2, x_0 + x_2, x_0 + x_1)$ . Then

$$(1.6.8) \quad s_0^2 + s_1^2 + s_2^2 - 2(x_0^2 + x_1^2 + x_2^2) = 2f \quad \text{and} \quad s_0s_1 + s_1s_2 + s_0s_2 - (x_0^2 + x_1^2 + x_2^2) = 3f$$

We eliminate  $(x_0^2 + x_1^2 + x_2^2)$  from the two equations.

$$(1.6.9) \quad (s_0^2 + s_1^2 + s_2^2) - 2(s_0s_1 + s_1s_2 + s_0s_2) = -4f$$

Setting  $f = 0$  gives us the equation of the dual curve. It is another conic.

(ii) *(the dual of a cuspidal cubic)* The dual of a smooth cubic has degree 6. We compute the dual of a cubic with a cusp instead. The curve  $C$  defined by the irreducible polynomial  $f = y^2z + x^3$  has a cusp at  $(0, 0, 1)$ . The Hessian matrix of  $f$  is

$$H = \begin{pmatrix} 6x & 0 & 0 \\ 0 & 2z & 2y \\ 0 & 2y & 0 \end{pmatrix}$$

and the Hessian determinant  $h = \det H$  is  $-24xy^2$ . The common zeros of  $f$  and  $h$  are the cusp point  $(0, 0, 1)$  and a single flex point  $(0, 1, 0)$ .

We scale the partial derivatives of  $f$  to simplify notation. Let  $u = f_x/3 = x^2$ ,  $v = f_y/2 = yz$ , and  $w = f_z = y^2$ . Then

$$v^2w - u^3 = y^4z^2 - x^6 = (y^2z + x^3)(y^2z - x^3) = f(y^2z - x^3)$$

The zero locus of the irreducible polynomial  $v^2w - u^3$  is the dual curve. It is another cuspidal cubic.  $\square$

### (1.6.10) a local equation for the dual curve

We label the coordinates in  $\mathbb{P}$  and  $\mathbb{P}^*$  as  $x, y, z$  and  $u, v, w$ , respectively, and we work in a neighborhood of a smooth point  $p_0$  of the curve  $C$  defined by a homogeneous polynomial  $f(x, y, z)$ , choosing coordinates so that  $p_0 = (0, 0, 1)$ , and that the tangent line at  $p_0$  is the line  $L_0 : \{y = 0\}$ . The image of  $p_0$  in the dual curve  $C^*$  is  $L_0^* : (u, v, w) = (0, 1, 0)$ .

Let  $\tilde{f}(x, y) = f(x, y, 1)$ . In the affine  $x, y$ -plane, the point  $p_0$  becomes the origin  $p_0 = (0, 0)$ . So  $\tilde{f}(p_0) = 0$ , and since the tangent line is  $L_0$ ,  $\frac{\partial \tilde{f}}{\partial x}(p_0) = 0$ , while  $\frac{\partial \tilde{f}}{\partial y}(p_0) \neq 0$ . We solve the equation  $\tilde{f} = 0$  for  $y$  as an analytic function  $y(x)$  for small  $x$ , with  $y(0) = 0$ . Let  $y'(x)$  denote the derivative  $\frac{dy}{dx}$ . Differentiating the equation  $\tilde{f}(x, y(x)) = 0$  shows that  $y'(0) = 0$ .

Let  $\tilde{p}_1 = (x_1, y_1)$  be a point of  $C_0$  near to  $\tilde{p}_0$ , so that  $y_1 = y(x_1)$ , and let  $y'_1 = y'(x_1)$ . The tangent line  $L_1$  at  $\tilde{p}_1$  has the equation

$$(1.6.11) \quad y - y_1 = y'_1(x - x_1)$$

Putting  $z$  back, the homogeneous equation of the tangent line  $L_1$  at the point  $p_1 = (x_1, y_1, 1)$  is

$$(-y'_1)x + y + (y'_1x_1 - y_1)z = 0$$

The point  $L_1^*$  of the dual plane that corresponds to  $L_1$  is  $(-y'_1, 1, y'_1x_1 - y_1)$ . Let's drop the subscript 1. As  $x$  varies, and writing  $y = y(x)$  and  $y' = y'(x)$ ,

$$(1.6.12) \quad (u, v, w) = (-y', 1, y'x - y),$$

There may be accidents:  $L_0$  might be tangent to  $C$  at distinct smooth points  $q_0$  and  $p_0$ , or it might pass through a singular point of  $C$ . If either of these accidents occurs, we can't analyze the neighborhood of  $L_0^*$  in  $C^*$  by this method. But, provided that there are no accidents, the path (1.6.12) will trace out the dual curve  $C^*$  near to  $L_0^* = (0, 1, 0)$ . (See (1.4.19).)

### (1.6.13) the bidual

The *bidual*  $C^{**}$  of a curve  $C$  is the dual of the curve  $C^*$ . It is a curve in the space  $\mathbb{P}^{**}$ , which is  $\mathbb{P}$ .

**1.6.14. Theorem.** *A plane curve of degree greater than one is equal to its bidual:  $C^{**} = C$ .*

As before,  $U$  will denote the set of smooth points of a curve  $C$ . Let  $V$  be the set of smooth points  $p_0$  of  $C$  such that  $t(p_0)$  is a smooth point of  $C^*$ . Thus  $V \subset U \subset C$ .

### 1.6.15. Lemma.

(i) *The subset  $V$  is the complement of a finite set of points of  $C$ .*

(ii) *Let  $p_1$  be a point near to a smooth point  $p_0$  of a curve  $C$ , let  $L_1$  and  $L_0$  be the tangent line to  $C$  at  $p_1$  and  $p_0$ , respectively, and let  $q$  be intersection point  $L_1 \cap L_0$ . Then  $\lim_{p_1 \rightarrow p_0} q = p_0$ .*

(iii) *If  $p_0$  is a point of  $V$  with tangent line  $L_0$ , the tangent line to  $C^*$  at  $L_0^*$  is  $p_0^*$ .*

*proof.* (i) Let  $S$  and  $S^*$  denote the finite sets of singular points of  $C$ , and  $C^*$ , respectively, the set  $U$  of smooth points of  $C$  is the complement of  $S$  in  $C$ , and  $V$  is obtained from  $U$  by deleting points whose images are in  $S^*$ . The fibre of  $t$  over a point  $L^*$  of  $C^*$  is the set of smooth points  $p$  of  $C$  such that the tangent line at  $p$  is  $L$ . Since  $L$  meets  $C$  in finitely many points, the fibre is finite. So the inverse image of  $S^* \cap U$  will be a finite subset of  $U$ .

(ii) We work analytically in a neighborhood of  $p_0$ , choosing coordinates so that  $p_0 = (0, 0, 1)$  and that  $L_0$  is the line  $\{y = 0\}$ . Let  $(x_q, y_q, 1)$  be the coordinates of  $q = L_0 \cap L_1$ . Since  $q$  is a point of  $L_0$ ,  $y_q = 0$ . The coordinate  $x_q$  can be obtained by substituting  $x = x_q$  and  $y = 0$  into the equation (1.6.11) of  $L_1$ :

$$x_q = x_1 - y_1/y'_1.$$

Now, when a function has an  $n$ th order zero at the point  $x = 0$ , i.e, when it has the form  $y = x^n h(x)$ , where  $n > 0$  and  $h(0) \neq 0$ , the order of zero of its derivative at that point is  $n - 1$ . This is verified by differentiating  $x^n h(x)$ . Since the function  $y(x)$  has a zero of positive order at  $p_0$ ,  $\lim_{p_1 \rightarrow p_0} y_1/y'_1 = 0$ . We also

have  $\lim_{p_1 \rightarrow p_0} x_1 = 0$ . So  $\lim_{p_1 \rightarrow p_0} x_q = 0$  and  $\lim_{p_1 \rightarrow p_0} q = \lim_{p_1 \rightarrow p_0} (x_q, y_q, 1) = (0, 0, 1) = p_0$ .

figure

(iii) Let  $p_1$  be a point of  $C$  near to  $p_0$ , and let  $L_1$  be the tangent line to  $C$  at  $p_1$ . The image of  $p_1$  is  $L_1^* = (f_0(p_1), f_1(p_1), f_2(p_1))$ . Because the partial derivatives  $f_i$  are continuous,

$$\lim_{p_1 \rightarrow p_0} L_1^* = (f_0(p_0), f_1(p_0), f_2(p_0)) = L_0^*$$

With  $q = L_0 \cap L_1$ ,  $q^*$  is the line through the points  $L_0^*$  and  $L_1^*$ . As  $p_1$  approaches  $p_0$ ,  $L_1^*$  approaches  $L_0^*$ , and therefore  $q^*$  approaches the tangent line to  $C^*$  at  $L_0^*$ . On the other hand, (ii) tells us that  $q^*$  approaches  $p_0^*$ . Therefore the tangent line at  $L_0^*$  is  $p_0^*$ .  $\square$

*proof of theorem 1.6.14.* Let  $V$  be the set of smooth points of  $C$  whose images in  $C^*$  are smooth, as in Lemma 1.6.15. Let  $U^*$  denote the set of smooth points of  $C^*$ , and let  $U^* \xrightarrow{t^*} \mathbb{P}^{**} = \mathbb{P}$  be the map analogous to the map  $U \xrightarrow{t} \mathbb{P}^*$ . Recall that  $t$  is defined by  $t(p) = L^*$ . Since the tangent line to  $C^*$  at  $L^*$  is  $p^*$ , the map  $t^*$  is  $t^*(L^*) = (p^*)^* = p$ . So for all points  $p$  of  $V$ ,  $t^*t(p) = t^*(L^*) = p$ . It follows that the restriction of  $t$  to  $V$  is injective, and that it defines a bijective map from  $V$  to its image  $tV$ , whose inverse function is  $t^*$ . So  $V$  is contained in the bidual  $C^{**}$ . Since  $V$  is dense in  $C$  and  $C^{**}$  is a closed set,  $C \subset C^{**}$ . Since  $C$  and  $C^{**}$  are curves,  $C = C^{**}$ .  $\square$

**1.6.16. Corollary.** (i) Let  $U$  be the set of smooth points of a plane curve  $C$ , and let denote the map from  $U$  to the dual curve  $C^*$ . The image of  $U$  is the complement of a finite subset of  $C^*$ .

(ii) If  $C$  is smooth, the map  $C \xrightarrow{t} C^*$ , which is defined at all points of  $C$ , is surjective.

*proof.* (i) Let  $U$  and  $V$  be as above, and let  $U^*$  be the set of smooth points of  $C^*$ . The image  $tV$  of  $V$  is contained in  $U^*$ . Then  $V = t^*tV \subset t^*U^* \subset C^{**} = C$ . Since  $V$  is the complement of a finite subset of  $C$ , so is  $t^*U^*$ . The assertion to be proved follows when we switch  $C$  and  $C^*$ .

(ii) Let  $W$  denote the image of  $C$  in  $C^*$ . The map  $C^* \xrightarrow{t^*} C^{**} = C$  is defined at the smooth points of  $C^*$ , and it inverts  $t$  at those points. Therefore  $W$  contains the smooth points of  $C^*$ . The complement  $S$  of  $W$  in  $C^*$  is a finite set. Since  $C$  is compact, its image  $W$  is compact, and therefore closed in  $C^*$ . Then its complement  $S$  is open, and since it is a finite set,  $S$  is also closed. So  $S$  consists of isolated points of  $C^*$ . Since a plane curve has no isolated point (1.3.18),  $S$  is empty.  $\square$

## 1.7 Resultants and Discriminants

Let  $F$  and  $G$  be monic polynomials in  $x$  with variable coefficients:

$$(1.7.1) \quad F(x) = x^m + a_1x^{m-1} + \cdots + a_m \quad \text{and} \quad G(x) = x^n + b_1x^{n-1} + \cdots + b_n$$

The resultant  $\text{Res}(F, G)$  of  $F$  and  $G$  is a certain polynomial in the coefficients. Its important property is that, when the coefficients of are in a field, the resultant is zero if and only if  $F$  and  $G$  have a common factor.

As an example, suppose that the coefficients  $a_i$  and  $b_i$  in (1.7.1) are polynomials in  $t$ , so that  $F$  and  $G$  become polynomials in two variables. Let  $C$  and  $D$  be (possibly reducible) curves  $F = 0$  and  $G = 0$  in the affine plane  $\mathbb{A}_{t,x}^2$ , and let  $S$  be the set of intersections:  $S = C \cap D$ . The resultant  $\text{Res}(F, G)$ , computed regarding  $x$  as the variable, will be a polynomial in  $t$  whose roots are the  $t$ -coordinates of the set  $S$ .

*figure*

The analogous statement is true when there are more variables. For example, if  $F$  and  $G$  are polynomials in  $x, y, z$ , the loci  $C : \{F = 0\}$  and  $D : \{G = 0\}$  in  $\mathbb{A}^3$  will be surfaces, and  $S = C \cap D$  will be a curve. The resultant  $\text{Res}_z(F, G)$ , computed regarding  $z$  as the variable, is a polynomial in  $x, y$  whose zero locus in the plane  $\mathbb{A}_{xy}^2$  is the projection of  $S$  to the plane.

The formula for the resultant is nicest when one allows leading coefficients different from 1. We work with homogeneous polynomials in two variables to prevent the degrees from dropping when a leading coefficient happens to be zero.

Let  $f$  and  $g$  be homogeneous polynomials in  $x, y$  with complex coefficients:

$$(1.7.2) \quad f(x, y) = a_0x^m + a_1x^{m-1}y + \cdots + a_my^m, \quad g(x, y) = b_0x^n + b_1x^{n-1}y + \cdots + b_ny^n$$

Suppose that they have a common zero  $(x, y) = (u, v)$  in  $\mathbb{P}_{xy}^1$ . Then  $vx - uy$  divides both  $g$  and  $f$ . The polynomial  $h = fg/(vx - uy)$  of degree  $m + n - 1$  will be divisible by  $f$  and by  $g$ , say  $h = pf = qg$ , where  $p$  and  $q$  are homogeneous polynomials of degrees  $n - 1$  and  $m - 1$ , respectively. Then  $h$  will be a linear combination  $pf$  of the polynomials  $x^i y^j f$ , with  $i + j = n - 1$ , and it will also be a linear combination  $qg$  of the polynomials  $x^k y^\ell g$ , with  $k + \ell = m - 1$ . The equation  $pf = qg$  tells us that the  $m + n$  polynomials of degree  $m + n - 1$ ,

$$(1.7.3) \quad x^{n-1}f, x^{n-2}yf, \dots, y^{n-1}f; x^{m-1}g, x^{m-2}yg, \dots, y^{m-1}g$$

will be dependent. For example, suppose that  $f$  has degree 3 and  $g$  has degree 2. If  $f$  and  $g$  have a common zero, the polynomials

$$\begin{aligned} xf &= a_0x^4 + a_1x^3y + a_2x^2y^2 + a_3xy^3 \\ yf &= a_0x^3y + a_1x^2y^2 + a_2xy^3 + a_3y^4 \\ x^2g &= b_0x^4 + b_1x^3y + b_2x^2y^2 \\ xyg &= b_0x^3y + b_1x^2y^2 + b_2xy^3 \\ y^2g &= b_0x^2y^2 + b_1xy^3 + b_2y^4 \end{aligned}$$

will be dependent. Conversely, if the polynomials (1.7.3) are dependent, there will be an equation of the form  $pf = qg$ , with  $p$  of degree  $n - 1$  and  $q$  of degree  $m - 1$ . Then at least one zero of  $g$  must also be a zero of  $f$ .

Let  $r = m + n - 1$ . The polynomials (1.7.3) have degree  $r$ . We form a square  $(r + 1) \times (r + 1)$  matrix  $\mathcal{R}$ , the *resultant matrix*, whose columns are indexed by the monomials  $x^r, x^{r-1}y, \dots, y^r$  of degree  $r$ , and whose rows list the coefficients of the polynomials (1.7.3). The matrix is illustrated below for the cases  $m, n = 3, 2$  and  $m, n = 1, 2$ , with dots representing entries that are zero:

$$(1.7.4) \quad \mathcal{R} = \begin{pmatrix} a_0 & a_1 & a_2 & a_3 & \cdot \\ \cdot & a_0 & a_1 & a_2 & a_3 \\ b_0 & b_1 & b_2 & \cdot & \cdot \\ \cdot & b_0 & b_1 & b_2 & \cdot \\ \cdot & \cdot & b_0 & b_1 & b_2 \end{pmatrix} \quad \text{or} \quad \mathcal{R} = \begin{pmatrix} a_0 & a_1 & \cdot \\ \cdot & a_0 & a_1 \\ b_0 & b_1 & b_2 \end{pmatrix}$$

The *resultant* of  $f$  and  $g$  is defined to be the determinant of  $\mathcal{R}$ .

$$(1.7.5) \quad \text{Res}(f, g) = \det \mathcal{R}$$

The coefficients of  $f$  and  $g$  can be in any ring.

The resultant  $\text{Res}(F, G)$  of the monic, one-variable polynomials  $F(x) = x^m + a_1x^{m-1} + \cdots + a_m$  and  $G(x) = x^n + b_1x^{n-1} + \cdots + b_n$  is the determinant of the matrix  $\mathcal{R}$ , with  $a_0 = b_0 = 1$ .

**1.7.6. Corollary.** *Let  $f$  and  $g$  be homogeneous polynomials in two variables, or monic polynomials in one variable, of degrees  $m$  and  $n$ , respectively, and with coefficients in a field. The resultant  $\text{Res}(f, g)$  is zero if and only if  $f$  and  $g$  have a common factor. If so, there will be polynomials  $p$  and  $q$  of degrees  $n - 1$  and  $m - 1$  respectively, such that  $pf = qg$ .*

*If the coefficients are in  $\mathbb{C}$ , the resultant is zero if and only if  $f$  and  $g$  have a common root.*  $\square$

When the leading coefficients  $a_0$  and  $b_0$  of  $f$  and  $g$  are both zero, the point  $(1, 0)$  of  $\mathbb{P}_{xy}^1$  will be a zero of  $f$  and of  $g$ . In this case, one could say that  $f$  and  $g$  have a common zero at infinity.

#### (1.7.7) weighted degree

When defining the degree of a polynomial, one may assign an integer called a *weight* to each variable. If one assigns weight  $w_i$  to the variable  $x_i$ , the monomial  $x_1^{e_1} \cdots x_n^{e_n}$  gets the *weighted degree*

$$e_1 w_1 + \cdots + e_n w_n$$

For instance, it is natural to assign weight  $k$  to the coefficient  $a_k$  of the polynomial  $f(x) = x^n - a_1 x^{n-1} + a_2 x^{n-2} - \cdots \pm a_n$  because, if  $f$  factors into linear factors,  $f(x) = (x - \alpha_1) \cdots (x - \alpha_n)$ , then  $a_k$  will be the  $k$ th elementary symmetric function in  $\alpha_1, \dots, \alpha_n$ . When written as a polynomial in  $\alpha$ , the degree of  $a_k$  will be  $k$ .

We leave the proof of the next lemma as an exercise.

**1.7.8. Lemma.** *Let  $f(x, y)$  and  $g(x, y)$  be homogeneous polynomials of degrees  $m$  and  $n$  respectively, with variable coefficients  $a_i$  and  $b_i$ , as in (1.7.2). When one assigns weight  $i$  to  $a_i$  and to  $b_i$ , the resultant  $\text{Res}(f, g)$  becomes a weighted homogeneous polynomial of degree  $mn$  in the variables  $\{a_i, b_j\}$ .  $\square$*

**1.7.9. Proposition.** *Let  $F$  and  $G$  be products of monic linear polynomials, say  $F = \prod_i (x - \alpha_i)$  and  $G = \prod_j (x - \beta_j)$ . Then*

$$\text{Res}(F, G) = \prod_{i,j} (\alpha_i - \beta_j) = \prod_i G(\alpha_i)$$

**Note.** Since the resultant vanishes when  $\alpha_i = \beta_j$ , it must be divisible by  $\alpha_i - \beta_j$ . So its weighted degree, though large, is as small as it could be.

*proof.* The equality of the second and third terms is obtained by substituting  $\alpha_i$  for  $x$  into the formula  $G = \prod_j (x - \beta_j)$ . We prove that the first and second terms are equal.

Let the elements  $\alpha_i$  and  $\beta_j$  be variables, let  $R$  denote the resultant  $\text{Res}(F, G)$  and let  $\Pi$  denote the product  $\prod_{i,j} (\alpha_i - \beta_j)$ . When we write the coefficients of  $F$  and  $G$  as symmetric functions in the roots  $\alpha_i$  and  $\beta_j$ ,  $R$  will be homogeneous. Its (unweighted) degree in  $\alpha_i, \beta_j$  will be  $mn$ , the same as the degree of  $\Pi$  (Lemma 1.7.8). To show that  $R = \Pi$ , we choose  $i, j$  and divide  $R$  by the polynomial  $\alpha_i - \beta_j$ , considered as a monic polynomial in  $\alpha_i$ :

$$R = (\alpha_i - \beta_j)q + r,$$

where  $r$  has degree zero in  $\alpha_i$ . The resultant  $R$  vanishes when we substitute  $\alpha_i = \beta_j$ . Looking at this equation, we see that the remainder  $r$  also vanishes when  $\alpha_i = \beta_j$ . On the other hand, the remainder is independent of  $\alpha_i$ . It doesn't change when we set  $\alpha_i = \beta_j$ . Therefore the remainder is zero, and  $\alpha_i - \beta_j$  divides  $R$ . This is true for all  $i$  and all  $j$ , so  $\Pi$  divides  $R$ , and since these two polynomials have the same degree,  $R = c\Pi$  for some scalar  $c$ . To show that  $c = 1$ , one computes  $R$  and  $\Pi$  for some particular polynomials. We suggest using  $F = x^m$  and  $G = x^n - 1$ .  $\square$

**1.7.10. Corollary.** *Let  $F, G, H$  be monic polynomials and let  $c$  be a scalar. Then*

(i)  $\text{Res}(F, GH) = \text{Res}(F, G) \text{Res}(F, H)$ , and

(ii)  $\text{Res}(F(x-c), G(x-c)) = \text{Res}(F(x), G(x))$ .  $\square$

### (1.7.11) the discriminant

The *discriminant*  $\text{Discr}(F)$  of a polynomial  $F = a_0 x^m + a_1 x^{m-1} + \cdots + a_m$  is the resultant of  $F$  and its derivative  $F'$ :

$$(1.7.12) \quad \text{Discr}(F) = \text{Res}(F, F')$$

The computation of the discriminant is made using the formula for the resultant of a polynomial of degree  $m$ . It will be a weighted polynomial of degree  $m(m-1)$ . The definition makes sense when the leading coefficient  $a_0$  is zero, but the discriminant will be zero in that case.

When the coefficients of  $F$  are complex numbers, the discriminant is zero if and only if either  $F$  has a multiple root, which happens when  $F$  and  $F'$  have a common factor, or else  $F$  has degree less than  $m$ .

**Note.** The formula for the discriminant is often normalized by a factor  $\pm a_0^k$ . We won't make this normalization, so our formula of the discriminant is slightly different from the usual one.

Suppose that the coefficients  $a_i$  of  $F$  are polynomials in  $t$ , so that  $F$  becomes a polynomial in two variables. Let  $C$  be the locus  $F = 0$  in the affine plane  $\mathbb{A}_{t,x}^2$ . The discriminant  $\text{Discr}_x(F)$ , computed regarding  $x$  as the variable, will be a polynomial in  $t$ . At a root  $t_0$  of the discriminant, the line  $L_0 : \{t = t_0\}$  is tangent to, or passes through a singular point, of  $C$ .

The discriminant of the quadratic polynomial  $F(x) = ax^2 + bx + c$  is

$$(1.7.13) \quad \det \begin{pmatrix} a & b & c \\ 2a & b & \cdot \\ \cdot & 2a & b \end{pmatrix} = -a(b^2 - 4ac).$$

The discriminant of the monic cubic  $x^3 + px + q$  whose quadratic coefficient is zero is

$$(1.7.14) \quad \det \begin{pmatrix} 1 & \cdot & p & q & \cdot \\ \cdot & 1 & \cdot & p & q \\ 3 & \cdot & p & \cdot & \cdot \\ \cdot & 3 & \cdot & p & \cdot \\ \cdot & \cdot & 3 & \cdot & p \end{pmatrix} = 4p^3 + 27q^2$$

These are the negatives of the usual formulas. The signs are artifacts of our definition. Though it conflicts with our definition, we'll follow tradition and continue writing the discriminant of the polynomial  $ax^2 + bx + c$  as  $b^2 - 4ac$ .

**1.7.15. Proposition.** *Let  $K$  be a field of characteristic zero. The discriminant of an irreducible polynomial  $F$  with coefficients in  $K$  isn't zero. Therefore an irreducible polynomial  $F$  with coefficients in  $K$  has no multiple root.*

*proof.* When  $F$  is irreducible, it cannot have a factor in common with the derivative  $F'$ , which has lower degree.  $\square$

This proposition is false when the characteristic of  $K$  isn't zero. In characteristic  $p$ , the derivative  $F'$  might be the zero polynomial.

**1.7.16. Proposition.** *Let  $F = \prod (x - \alpha_i)$  be a polynomial that is a product of monic linear factors. Then*

$$\text{Discr}(F) = \prod_i F'(\alpha_i) = \prod_{i \neq j} (\alpha_i - \alpha_j) = \pm \prod_{i < j} (\alpha_i - \alpha_j)^2$$

*proof.* The fact that  $\text{Discr}(F) = \prod F'(\alpha_i)$  follows from Proposition 1.7.9. We show that

$$F'(\alpha_i) = \prod_{j, j \neq i} (\alpha_i - \alpha_j) = \prod_i (\alpha_i - \alpha_1) \cdots (\widehat{\alpha_i - \alpha_i}) \cdots (\alpha_i - \alpha_n)$$

where the hat  $\widehat{\phantom{x}}$  indicates that that term is deleted. By the product rule for differentiation,

$$F'(x) = \sum_k (x - \alpha_1) \cdots (\widehat{x - \alpha_k}) \cdots (x - \alpha_n)$$

Substituting  $x = \alpha_i$ , all terms in the sum, except the one with  $i = k$ , become zero.  $\square$

**1.7.17. Corollary.**  $\text{Discr}(F(x)) = \text{Discr}(F(x - c)).$   $\square$

**1.7.18. Proposition.** *Let  $F(x)$  and  $G(x)$  be monic polynomials. Then*

$$\text{Discr}(FG) = \pm \text{Discr}(F) \text{Discr}(G) \text{Res}(F, G)^2$$



*proof.* This proposition follows from Propositions 1.7.9 and 1.7.16 for polynomials with complex coefficients. It is true for polynomials with coefficients in any ring because it is an identity.  $\square$

When  $f$  and  $g$  are polynomials in several variables including a variable  $z$ ,  $\text{Res}_z(f, g)$  and  $\text{Discr}_z(f)$  denote the resultant and the discriminant, computed regarding  $f, g$  as polynomials in  $z$ . They will be polynomials in the other variables.

**1.7.19. Lemma.** *Let  $f$  be an irreducible polynomial in  $\mathbb{C}[x, y, z]$  of positive degree in  $z$ , but not divisible by  $z$ . The discriminant  $\text{Discr}_z(f)$  of  $f$  with respect to the variable  $z$  is a nonzero polynomial in  $x, y$ .*

*proof.* This follows from Lemma 1.3.13 (ii) and Proposition 1.7.15.  $\square$

### (1.7.20) projection to a line

We denote by  $\pi$  the *projection*  $\mathbb{P}^2 \longrightarrow \mathbb{P}^1$  that drops the last coordinate, sending a point  $(x, y, z)$  to  $(x, y)$ . This projection is defined at all points of  $\mathbb{P}^2$  except at the *center of projection*, the point  $q = (0, 0, 1)$ .

The fibre of  $\pi$  over a point  $\tilde{p} = (x_0, y_0)$  of  $\mathbb{P}^1$  is the line  $L_{pq}$  through  $p = (x_0, y_0, 0)$  and  $q = (0, 0, 1)$ , with the point  $q$  omitted – the set of points  $(x_0, y_0, z_0)$ .

*figure*

When a curve  $C$  in the plane doesn't contain the center of projection  $q$ , the projection  $\mathbb{P}^2 \xrightarrow{\pi} \mathbb{P}^1$  will be defined at all points of  $C$ . Say that such a curve  $C$  is defined by an irreducible homogeneous polynomial  $f(x, y, z)$  of degree  $d$ . We write  $f$  as a polynomial in  $z$ ,

$$(1.7.21) \quad f = c_0 z^d + c_1 z^{d-1} + \cdots + c_d$$

with  $c_i$  homogeneous, of degree  $i$  in  $x, y$ . Then  $c_0 = f(0, 0, 1)$  will be a nonzero constant that we normalize to 1, so that  $f$  becomes a monic polynomial of degree  $d$  in  $z$ .

The fibre of  $C$  over a point  $\tilde{p} = (x_0, y_0)$  of  $\mathbb{P}^1$  is the intersection of  $C$  with the line  $L_{pq}$  described above. It consists of the points  $(x_0, y_0, \alpha)$  such that  $\alpha$  is a root of the one-variable polynomial

$$(1.7.22) \quad \tilde{f}(z) = f(x_0, y_0, z)$$

We call  $C$  a *branched covering* of  $\mathbb{P}^1$  of degree  $d$ . All but finitely many fibres of  $C$  over  $\mathbb{P}^1$  consist of  $d$  points (Lemma 1.7.19). The fibres with fewer than  $d$  points are those above the zeros of the discriminant. They are the *branch points* of the covering.

### (1.7.23) the genus of a plane curve

We use the discriminant to describe the topological structure of smooth plane curves in the classical topology.

**1.7.24. Theorem.** *A smooth projective plane curve of degree  $d$  is a compact, orientable and connected manifold of dimension two.*

The fact that a smooth curve is a two-dimensional manifold follows from the Implicit Function Theorem. (See the discussion at (1.4.4)).

*orientability:* A two-dimensional manifold is orientable if one can choose one of its two sides in a continuous, consistent way. A smooth curve  $C$  is orientable because its tangent space at a point is a one-dimensional complex vector space – the affine line with the equation (1.4.11). Multiplication by  $i$  orients the tangent space by defining the counterclockwise rotation. Then the right-hand rule tells us which side of  $C$  is “up”.

*compactness:* A plane projective curve is compact because it is a closed subset of the compact space  $\mathbb{P}^2$ .

The connectedness of a plane curve is a subtle fact whose proof mixes topology and algebra. Unfortunately, I don't know a proof that fits into our discussion here. It will be proved later (see Theorem 8.4.9).

The topological *Euler characteristic*  $e$  of a compact, orientable two-dimensional manifold  $M$  is the alternating sum  $b^0 - b^1 + b^2$  of its Betti numbers. It can be computed using a *topological triangulation*, a subdivision of  $M$  into topological triangles, called *faces*, by the formula

$$(1.7.25) \quad e = |\text{vertices}| - |\text{edges}| + |\text{faces}|$$

For example, a sphere is homeomorphic to a tetrahedron, which has four vertices, six edges, and four faces. Its Euler characteristic is  $4 - 6 + 4 = 2$ . Any other topological triangulation of a sphere, such as the one given by the icosahedron, yields the same Euler characteristic.

Every compact, connected, orientable two-dimensional manifold is homeomorphic to a sphere with a finite number of “handles”. Its *genus* is the number of handles. A torus has one handle. Its genus is one. The projective line  $\mathbb{P}^1$ , which is a two-dimensional sphere, has genus zero.

*Figure*

The Euler characteristic and the genus are related by the formula

$$(1.7.26) \quad e = 2 - 2g$$

The Euler characteristic of a torus is zero, and the Euler characteristic of  $\mathbb{P}^1$  is two.

To compute the Euler characteristic of a smooth curve  $C$  of degree  $d$ , we analyze a generic projection to represent  $C$  as a branched covering of the projective line:  $C \xrightarrow{\pi} \mathbb{P}^1$ .

*figure*

We choose generic coordinates  $x, y, z$  in  $\mathbb{P}^2$  and project from the point  $q = (0, 0, 1)$ . When the defining equation of  $C$  is written as a monic polynomial in  $z$ :  $f = z^d + c_1 z^{d-1} + \cdots + c_d$  where  $c_i$  is a homogeneous polynomial of degree  $i$  in the variables  $x, y$ , the discriminant  $\text{Discr}_z(f)$  with respect to  $z$  will be a homogeneous polynomial of degree  $d(d-1) = d^2 - d$  in  $x, y$ .

Let  $\tilde{p}$  be the image in  $\mathbb{P}^1$  of a point  $p$  of  $C$ . The covering  $C \xrightarrow{\pi} \mathbb{P}^1$  will be branched at  $\tilde{p}$  when the tangent line at  $p$  is the line  $L_{pq}$  through  $p$  and the center of projection  $q$ . When  $q$  is generic, such a point  $p$  will not be a flex point, and then  $C$  and  $L_{pq}$  will have one intersection  $p$  of multiplicity two, and  $d-2$  intersections of multiplicity one (1.9). It is intuitively plausible that the discriminant  $\text{Discr}_z(f)$  will have a simple zero at the image  $\tilde{p}$  of  $p$ . This will be proved below, in Proposition 1.9.13. Assuming this is known, then since the discriminant has degree  $d^2 - d$ , there will be  $d^2 - d$  points  $\tilde{p}$  in  $\mathbb{P}^1$  at which the discriminant vanishes and the fibre contains  $d-1$  points. They are the *branch points* of the covering. All other fibres consist of  $d$  points.

We triangulate the sphere  $\mathbb{P}^1$  in such a way that the branch points are among the vertices, and we use the inverse images of the vertices, edges, and faces to triangulate  $C$ . Then  $C$  will have  $d$  faces and  $d$  edges lying over each face and each edge of  $\mathbb{P}^1$ , respectively. There will also be  $d$  vertices of  $C$  lying over a vertex of  $\mathbb{P}^1$ , except when it is one of the  $d^2 - d$  branch points. In that case the fibre will contain only  $d-1$  vertices. The Euler characteristic of  $C$  is obtained by multiplying the Euler characteristic of  $\mathbb{P}^1$  by  $d$  and subtracting the number of branch points.

$$(1.7.27) \quad e(C) = d e(\mathbb{P}^1) - (d^2 - d) = 2d - (d^2 - d) = 3d - d^2$$

This is the Euler characteristic of any smooth curve of degree  $d$ , so we denote it by  $e_d$ :

$$(1.7.28) \quad e_d = 3d - d^2$$

Formula (1.7.26) shows that the genus  $g_d$  of a smooth curve of degree  $d$  is

$$(1.7.29) \quad g_d = \frac{1}{2}(d^2 - 3d + 2) = \binom{d-1}{2}$$

Thus smooth curves of degrees 1, 2, 3, 4, 5, 6, ... have genus 0, 0, 1, 3, 6, 10, ..., respectively. A smooth plane curve cannot have genus two.

The generic projection to  $\mathbb{P}^1$  also computes the degree of the dual curve  $C^*$  of a smooth curve  $C$  of degree  $d$ . The degree of  $C^*$  is the number of intersections of  $C^*$  with the generic line  $q^*$  in  $\mathbb{P}^*$ . The intersection points have the form  $L^*$ , where  $q$  is a point of  $L$ , and  $L$  is tangent to  $C$  at some point  $p$ . As we have seen, there are  $d(d-1)$  such points.

**1.7.30. Corollary.** *Let  $C$  be a plane curve of degree  $d$ .*

(i) *The degree  $d^*$  of the dual curve  $C^*$  is equal to the number of tangent lines at smooth points of  $C$  that pass through a generic point  $q$  of the plane.*

(ii) *If  $C$  is a smooth curve, the degree  $d^*$  of the dual curve  $C^*$  is  $d(d-1)$ .* □

The formula  $d^* = d(d-1)$  is incorrect when  $C$  is singular. If  $C$  is a smooth curve of degree 3,  $C^*$  will have degree 6, and if  $C^*$  were smooth its dual curve  $C^{**}$ , would have degree 30. But  $C^{**} = C$ .

## 1.8 Nodes and Cusps

Let  $C$  be the projective curve defined by an irreducible homogeneous polynomial  $f(x, y, z)$  of degree  $d$ , and let  $p$  be a point of  $C$ . We choose coordinates so that  $p = (0, 0, 1)$ , and we set  $z = 1$ . This gives us an affine curve  $C_0$  in  $\mathbb{A}_{x,y}^2$ , the zero set of the polynomial  $\tilde{f}(x, y) = f(x, y, 1)$ , and  $p$  becomes the origin  $(0, 0)$ . We write

$$(1.8.1) \quad \tilde{f}(x, y) = f_0 + f_1 + f_2 + \cdots + f_d,$$

where  $f_i$  is the homogeneous part of  $\tilde{f}$  of degree  $i$ , which is also the coefficient of  $z^{d-i}$  in  $f(x, y, z)$ .

If the origin  $p$  is a point of  $C_0$ , the constant term  $f_0$  will be zero. Then the linear term  $f_1$  will define the tangent direction to  $C_0$  at  $p$ . If  $f_0$  and  $f_1$  are both zero,  $p$  will be a singular point of  $C$ .

It seems permissible to drop the tilde and the subscript 0 in what follows, denoting  $f(x, y, 1)$  by  $f(x, y)$ , and  $C_0$  by  $C$ .


### (1.8.2) the multiplicity of a singular point

Let  $f(x, y)$  be an analytic function, defined for small  $x, y$ , and let  $C$  denote the locus of zeros of  $f$  in a neighborhood of  $p = (0, 0)$ . To describe the singularity of  $C$  at  $p$ , we expand  $f$  as a series in  $x, y$  and look at the part of  $f$  of lowest degree. The smallest integer  $r$  such that  $f_r(x, y)$  isn't zero is the *multiplicity* of  $p$ . When the multiplicity of  $p$  is  $r$ ,  $f$  will have the form

$$(1.8.3) \quad f(x, y) = f_r + f_{r+1} + \cdots$$

Let  $L$  be a line  $\{vx = uy\}$  through  $p$ . The intersection multiplicity of  $C$  and  $L$  at  $p$  will be  $r$  unless  $f_r(u, v)$  is zero. It will be greater than  $r$  if  $f_r(u, v) = 0$ . Such a line  $L$  is *special*. The special lines correspond to the zeros of  $f_r$  in  $\mathbb{P}^1$ . Because  $f_r$  has degree  $r$ , there will be at most  $r$  special lines.

### 1.8.4.



a Singular Point, with its Special Lines

### (1.8.5) double points

Suppose that the origin  $p$  is a *double point*, a point of multiplicity 2. Let the quadratic part of  $f$  be

$$(1.8.6) \quad f_2 = ax^2 + bxy + cy^2$$

We may adjust coordinates so that  $c$  isn't zero, and we normalize  $c$  to 1.

To analyze the singularity at the origin  $p$  we blow up the plane. The *blowup* is the map  $W \xrightarrow{\pi} X$  from the  $(x, w)$ -plane  $W$  to the  $(x, y)$ -plane  $X$  defined by  $\pi(x, w) = (x, xw)$ . It is called a “blowup” of  $X$  because the fibre over the origin in  $X$  is the  $w$ -axis  $\{x = 0\}$  in  $W$ . (It might seem more appropriate to call the inverse of  $\pi$  the blowup, but the inverse isn't a map.) The map  $\pi$  is bijective at points at which  $x \neq 0$ , and points  $(x, 0)$  of  $X$  with  $x \neq 0$  aren't in its image.

Writing  $f(x, y) = ax^2 + bxy + y^2 + dx^3 + \dots$ , we make the substitution  $y = xw$  and cancel  $x^2$ . This gives us a polynomial

$$g(x, w) = f(x, xw)/x^2 = a + bw + w^2 + dx + \dots$$

in which the terms represented by  $\dots$  are divisible by  $x$ . Let  $D$  be the locus  $\{g = 0\}$  in  $W$ . The map  $\pi$  restricts to a map  $D \xrightarrow{\pi} C$ . Since  $\pi$  is bijective at points at which  $x \neq 0$ , so is  $\bar{\pi}$ .

Suppose first that the quadratic polynomial  $y^2 + by + a$  has distinct roots  $\alpha, \beta$ , so that  $ax^2 + bxy + y^2 = (y - \alpha x)(y - \beta x)$  and  $g(x, w) = (w - \alpha)(w - \beta) + dx + \dots$ . In this case, the fibre of  $D$  over the origin  $p$  in  $X$  consists of the two points  $p_1 = (0, \alpha)$  and  $p_2 = (0, \beta)$ . The partial derivative  $\frac{\partial g}{\partial t}$  is nonzero at  $p_1$  and  $p_2$ , so those are smooth points of  $D$ . We can solve  $g(x, w) = 0$  for  $w$  as analytic functions of  $x$  near zero, say  $w = u(x)$  and  $w = v(x)$  with  $u(0) = \alpha$  and  $v(0) = \beta$ . The image of  $\pi(D)$  is  $C$ , so  $C$  has two analytic branches  $y = xu(x)$  and  $y = xv(x)$  through the origin with distinct tangent directions  $\alpha$  and  $\beta$ . This singularity is called a *node*. A node is the simplest singularity that a curve can have.

When the discriminant  $b^2 - 4ac$  is zero,  $f_2$  will be a square, and we will have  $f(x, y) = (y - \alpha x)^2 + dx^3 + \dots$ . The blowup substitution  $y = xw$  gives  $g(x, w) = f(x, xw)/x^2 = (w - \alpha)^2 + dx + \dots$ . Here the fibre over  $(x, y) = (0, 0)$  is the point  $(x, w) = (0, \alpha)$ , and  $g_w$  is zero there. However, if  $d \neq 0$ , then  $g_x(0, \alpha) \neq 0$ . In this case,  $D$  is smooth at  $(0, 0)$ , and the equation of  $C$  has the form  $(y - \alpha x)^2 = dx^3 + \dots$ . The singularity of  $C$  at the origin is called a *cusp*. The *standard cusp* is the locus  $y^2 = x^3$ . All cusps are analytically equivalent with the standard cusp.


Cusps have an interesting geometry. The intersection of the standard cusp  $X : \{y^2 = x^3\}$  with a small 3-sphere  $S : \{\bar{x}x + \bar{y}y = \epsilon\}$  in  $\mathbb{C}^2$  is a *trefoil knot*.

To explain this, we parametrize  $X$  by  $(x(t), y(t)) = (t^3, t^2)$ , and we restrict to the unit circle  $t = e^{i\theta}$ . The locus of points of  $X$  of absolute value  $\sqrt{2}$  is  $(x(t), y(t)) = (e^{3i\theta}, e^{2i\theta})$ . To visualize this locus, we embed it in the product of the unit  $x$ -circle and the unit  $y$ -circle, a torus, and we distort that torus, representing it as the usual torus  $T$  in  $\mathbb{R}^3$ . Let the circumference of  $T$  represent the  $x$ -coordinate, and let the loop through the hole represent  $y$ . Then, as  $\theta$  runs from 0 to  $2\pi$ ,  $(x(t), y(t))$  goes around the circumference twice, and it loops through the hole three times, as is illustrated below.

figure

**1.8.7. Corollary.** *A double point  $p$  of a curve  $C$  is a node or a cusp if and only if the blowup of  $C$  is smooth at the points that lie over  $p$ .*  $\square$

The simplest example of a double point that isn't a node or cusp is a *tacnode*, a point at which two smooth branches of a curve intersect with the same tangent direction.




1.8.8.

a Node, a Cusp, and a Tacnode (real locus)

**A note about figures.** In algebraic geometry, the dimensions are too big to allow realistic figures. Even with a plane curve, one is dealing with a locus in the space  $\mathbb{A}^2$ , whose dimension as a real vector space is four. In some cases, such as in the figures above, depicting the real locus can be helpful, but in most cases, even the real locus is too big, and one must make do with a schematic diagram. The figure below is an example. My students have told me that all of my figures look more or less like this:

**1.8.9.**



**A Typical Schematic Figure**

## 1.9 Hensel's Lemma

The resultant matrix (1.7.4) arises in a second context that we explain here.

Suppose given a product  $P = FG$  of two polynomials, say

$$(1.9.1) \quad (c_0x^{m+n} + c_1x^{m+n-1} + \cdots + c_{m+n}) = (a_0x^m + a_1x^{m-1} + \cdots + a_m)(b_0x^n + b_1x^{n-1} + \cdots + b_n)$$

We call the relations among the coefficients implied by this polynomial equation the *product equations*. They are

$$c_i = a_ib_0 + a_{i-1}b_1 + \cdots + a_0b_i$$

for  $i = 0, \dots, m+n$ . For instance, when  $m = 3$  and  $n = 2$ , they are

**1.9.2.**

$$\begin{aligned} c_0 &= a_0b_0 \\ c_1 &= a_1b_0 + a_0b_1 \\ c_2 &= a_2b_0 + a_1b_1 + a_0b_2 \\ c_3 &= a_3b_0 + a_2b_1 + a_1b_2 \\ c_4 &= a_3b_1 + a_2b_2 \\ c_5 &= a_3b_2 \end{aligned}$$

Let  $J$  denote the Jacobian matrix of partial derivatives of  $c_1, \dots, c_{m+n}$  with respect to the variables  $b_1, \dots, b_n$  and  $a_1, \dots, a_m$ , treating  $a_0, b_0$  and  $c_0$  as constants. When  $m, n = 3, 2$ ,

$$(1.9.3) \quad J = \frac{\partial(c_i)}{\partial(b_j, a_k)} = \begin{pmatrix} a_0 & . & b_0 & . & . \\ a_1 & a_0 & b_1 & b_0 & . \\ a_2 & a_1 & b_2 & b_1 & b_0 \\ a_3 & a_2 & . & b_2 & b_1 \\ . & a_3 & . & . & b_2 \end{pmatrix}$$

**1.9.4. Lemma.** The Jacobian matrix  $J$  is the transpose of the resultant matrix  $\mathcal{R}$  (1.7.4). □

**1.9.5. Corollary.** Let  $F$  and  $G$  be polynomials with complex coefficients. The Jacobian matrix is singular if and only if, either  $F$  and  $G$  have a common root, or  $a_0 = b_0 = 0$ .  $\square$

This corollary has an application to polynomials with analytic coefficients. Let

$$(1.9.6) \quad P(t, x) = c_0(t)x^d + c_1(t)x^{d-1} + \cdots + c_d(t)$$

be a polynomial in  $x$  whose coefficients  $c_i$  are analytic functions, defined for small values of  $t$ , and let  $\bar{P} = P(0, x) = \bar{c}_0x^d + \bar{c}_1x^{d-1} + \cdots + \bar{c}_d$  be the evaluation of  $P$  at  $t = 0$ , so that  $\bar{c}_i = c_i(0)$ . Suppose given a factorization  $\bar{P} = \bar{F}\bar{G}$ , where  $\bar{G} = \bar{b}_0x^n + \bar{b}_1x^{n-1} + \cdots + \bar{b}_n$  is a polynomial and  $\bar{F} = x^m + \bar{a}_1x^{m-1} + \cdots + \bar{a}_m$  is a monic polynomial, both with complex coefficients. Are there polynomials  $F(t, x) = x^m + a_1x^{m-1} + \cdots + a_m$  and  $G(t, x) = b_0x^n + b_1x^{n-1} + \cdots + b_n$ , with  $F$  monic, whose coefficients  $a_i$  and  $b_i$  are analytic functions defined for small  $t$ , such that  $P = FG$ ,  $F(0, x) = \bar{F}$ , and  $G(0, x) = \bar{G}$ ?

**1.9.7. Hensel's Lemma.** With notation as above, suppose that  $\bar{F}$  and  $\bar{G}$  have no common root. Then  $P$  factors, as above.

*proof.* Since  $F$  is supposed to be monic, we set  $a_0(t) = 1$ . The first product equation tells us that  $b_0(t) = c_0(t)$ . Corollary 1.9.5 tells us that the Jacobian matrix for the remaining product equations is nonsingular at  $t = 0$ , so according to the Implicit Function Theorem, the product equations have a unique solution in analytic functions  $a_i(t), b_j(t)$  for small  $t$ .  $\square$

Note that  $P$  isn't assumed to be monic. If  $\bar{c}_0 = 0$ , the degree of  $\bar{P}$  will be less than the degree of  $P$ . In that case,  $\bar{G}$  will have lower degree than  $G$ .

figure

**1.9.8. Example.** Let  $P = c_0(t)x^2 + c_1(t)x + c_2(t)$ . The product equations for factoring  $P$  as a product  $FG = (x + a_1)(b_0x + b_1)$  of linear polynomials, with  $F$  monic, are

$$c_0 = b_0, \quad c_1 = a_1b_0 + b_1, \quad c_2 = a_1b_1$$

and the Jacobian matrix is

$$\frac{\partial(c_1, c_2)}{\partial(b_1, a_0)} = \begin{pmatrix} 1 & b_0 \\ a_1 & b_1 \end{pmatrix}$$

Suppose that  $\bar{P} = P(0, x)$  factors:  $\bar{c}_0x^2 + \bar{c}_1x + \bar{c}_2 = (x + \bar{a}_1)(\bar{b}_0x + \bar{b}_1) = \bar{F}\bar{G}$ . The determinant of the Jacobian matrix at  $t = 0$  is  $\bar{b}_1 - \bar{a}_1\bar{b}_0$ . It is nonzero if and only if the two factors are relatively prime, in which case  $P$  factors too.

On the other hand, the one-variable Jacobian criterion allows us to solve the equation  $P(t, x) = 0$  for  $x$  as function of  $t$  with  $x(0) = -\bar{a}_1$ , provided that  $\frac{\partial P}{\partial x} = 2c_0x + c_1$  isn't zero at the point  $(t, x) = (0, -\bar{a}_1)$ . In that case,  $\bar{P}$  factors. Substituting  $\bar{c}_0 = \bar{b}_0$  and  $\bar{c}_1 = \bar{a}_1\bar{b}_0 + \bar{b}_1$ , shows that  $-2\bar{c}_0\bar{a}_1 + \bar{c}_1 = \bar{b}_1 - \bar{a}_1\bar{b}_0$ . Not surprisingly, the two conditions for factoring are the same.  $\square$

### (1.9.9) general position

In algebraic geometry, the phrases *general position* and *generic* indicate an object, such as a point, has no special 'bad' properties. Typically, the object will be parametrized somehow, and the word *generic* indicates that the parameter representing that particular object avoids a proper closed subset of the parameter space that may be described explicitly or not. Proposition 1.9.13 below refers to a generic point  $q$ . In this case we require that  $q$  shall not lie on any of these lines:

(1.9.10)

- flex tangent lines and bitangent lines,
- lines that contain more than one singular point,
- special lines through singular points (see (1.8.2)),
- tangent lines that contain a singular point of  $C$ .

**1.9.11. Lemma.** *This is a list of finitely many lines that  $q$  must avoid.*

*beginning of the proof.* Proposition 1.4.18 shows that there are finitely many flex tangents. Since there are finitely many singular points, there are finitely many special lines and finitely many lines through pairs of singular points. To show that there are finitely many tangent lines that pass through singular points, we project  $C$  from a singular point  $p$  and apply Lemma 1.7.19. The discriminant isn't identically zero, so it vanishes finitely often. The proof that there are finitely many bitangents will be given later, in Corollary 1.10.15.

**(1.9.12) order of vanishing of the discriminant**

Let  $f(x, y, z)$  be a homogeneous polynomial with no multiple factors, and let  $C$  be the (possibly reducible) plane curve  $\{f = 0\}$ . Suppose that  $q = (0, 0, 1)$  is in general position, in the sense described above.

**1.9.13. Proposition. (i)** *If  $p$  is a smooth point of  $C$  with tangent line  $L_{pq}$ , the discriminant  $\text{Discr}_z(f)$  has a simple zero at  $\tilde{p}$ .*

**(ii)** *If  $p$  is a node of  $C$ ,  $\text{Discr}_z(f)$  has a double zero at  $\tilde{p}$ .*

**(iii)** *If  $p$  is a cusp,  $\text{Discr}_z(f)$  has a triple zero at  $\tilde{p}$ .*

**(iv)** *If  $p$  is an ordinary flex point of  $C$  (1.4.8) with tangent line  $L_{pq}$ ,  $\text{Discr}_z(f)$  has a double zero at  $\tilde{p}$ .*

*proof. (i)–(iii)* There are several ways to proceed, none especially simple. We'll use Hensel's Lemma. We set  $x = 1$ , to work in the standard affine open set  $\mathbb{U}$  with coordinates  $y, z$ . In affine coordinates, the projection  $\pi$  is the map  $(y, z) \rightarrow y$ . We may suppose that  $p$  is the origin in  $\mathbb{U}$ . Its image  $\tilde{p}$  will be the point  $y = 0$  of the affine  $y$ -line, and the intersection of the line  $L_{pq}$  with  $\mathbb{U}$  will be the line  $\tilde{L} : \{y = 0\}$ . We'll denote the defining polynomial of the curve  $C$ , restricted to  $\mathbb{U}$ , by  $f(y, z)$  instead of  $f(1, y, z)$ . Let  $\tilde{f}(z) = f(0, z)$ .

In each of the cases under discussion, the polynomial  $\tilde{f}(z) = f(0, z)$  will have a double zero at  $z = 0$ , so we will have  $\tilde{f}(z) = z^2 \bar{h}(z)$ , with  $\bar{h}(0) \neq 0$ . Then  $z^2$  and  $\bar{h}(z)$  have no common root, so we may apply Hensel's Lemma to write  $f(y, z) = g(y, z)h(y, z)$ , where  $g$  and  $h$  are polynomials in  $z$  whose coefficients are analytic functions of  $y$ , defined for small  $y$ ,  $g$  is monic,  $g(0, z) = z^2$ , and  $h(0, z) = \bar{h}$ . Then (1.7.18)

$$(1.9.14) \quad \text{Discr}_z(f) = \pm \text{Discr}_z(g) \text{Discr}_z(h) \text{Res}_z(g, h)^2$$

Since  $q$  is in general position,  $\bar{h}$  will have simple zeros. Then  $\text{Discr}_z(h)$  doesn't vanish at  $y = 0$ . Neither does  $\text{Res}_z(g, h)$ . So the orders of vanishing of  $\text{Discr}_z(f)$  and  $\text{Discr}_z(g)$  are equal. We replace  $f$  by  $g$ .

Since  $g$  is a monic quadratic polynomial, it will have the form

$$g(y, z) = z^2 + b(y)z + c(y)$$

The coefficients  $b$  and  $c$  are analytic functions of  $y$ , and  $g(0, z) = z^2$ . The discriminant  $\text{Discr}_z(g) = b^2 - 4c$  is unchanged when we complete the square by the substitution of  $z - \frac{1}{2}b$  for  $z$ , and if  $\tilde{p}$  is a node or a cusp, that property isn't affected by this change of coordinates (Lemma ??). So we may assume that  $g$  has the form  $z^2 + c(y)$ . The discriminant is  $D = 4c(y)$ .

We write  $c(y)$  as a series in  $y$ :

$$c(y) = c_0 + c_1y + c_2y^2 + c_3y^3 + \cdots$$

The constant coefficient  $c_0$  is zero because  $\tilde{p}$  is a point of  $C$ . If  $c_1 \neq 0$ ,  $\tilde{p}$  is a smooth point with tangent line  $\tilde{L} : \{y = 0\}$ , and  $D$  has a simple zero. If  $\tilde{p}$  is a node,  $c_0 = c_1 = 0$  and  $c_2 \neq 0$ . Then  $D$  has a double zero. If  $\tilde{p}$  is a cusp,  $c_0 = c_1 = c_2 = 0$ , and  $c_3 \neq 0$ . Then  $D$  has a triple zero at  $\tilde{p}$ .

**(iv)** In this case, the polynomial  $\tilde{f}(z) = f(0, z)$  will have a triple zero at  $z = 0$ . Proceeding as above, we may factor:  $f = gh$  where  $g$  and  $h$  are polynomials in  $z$  with analytic coefficients in  $y$ , and  $g(y, z) = z^3 + a(y)z^2 + b(y)z + c(y)$ . We eliminate the quadratic coefficient  $a$  by substituting  $z - \frac{1}{3a}$  for  $z$ . With  $g = z^3 + az + b$ , the discriminant  $\text{Discr}_z(g)$  is  $4b^3 + 27c^2$  (1.7.14). We write  $c(y) = c_0 + c_1y + \cdots$  and  $b(y) = b_0 + b_1y + \cdots$ . Since  $p$  is a point of  $C$  with tangent line  $\{y = 0\}$ ,  $c_0 = 0$  and  $c_1 \neq 0$ . Since the intersection multiplicity of  $C$  with the line  $\{y = 0\}$  at  $\tilde{p}$  is three,  $b_0 = 0$ . The discriminant has a zero of order two.  $\square$

**1.9.15. Corollary.** Let  $C : \{g = 0\}$  and  $D : \{h = 0\}$  be plane curves that intersect transversally at a point  $p = (x_0, y_0, z_0)$ . With coordinates in general position,  $\text{Res}_z(g, h)$  has a simple zero at  $(x_0, y_0)$ .

Two curves are said to intersect *transversally* at a point  $p$  if they are smooth at  $p$  and their tangent lines there are distinct.

*proof.* Proposition 1.9.13 (ii) applies to the product  $fg$ , whose zero locus is the union  $C \cup D$ . It shows that the discriminant  $\text{Discr}_z(fg)$  has a double zero at  $\tilde{p}$ . We also have the formula (1.9.14) with  $f = gh$ . Since coordinates are in general position,  $\text{Discr}_z(g)$  and  $\text{Discr}_z(h)$  will not be zero at  $\tilde{p}$ . Then  $\text{Res}_z(g, h)$  has a simple zero there.  $\square$

## 1.10 Bézout's Theorem

Bézout's Theorem counts intersections of plane curves. We state it here in a form that is ambiguous because it contains a term "multiplicity" that hasn't yet been defined.

**1.10.1. Bézout's Theorem.** Let  $C$  and  $D$  be distinct curves of degrees  $m$  and  $n$ , respectively. When intersections are counted with the appropriate multiplicity, the number of intersections is equal to  $mn$ . Moreover, the multiplicity at a point is 1 at a transversal intersection.

As before,  $C$  and  $D$  intersect transversally at  $p$  if they are smooth at  $p$  and their tangent lines there are distinct.

**1.10.2. Corollary.** Bézout's Theorem is true when one of the curves is a line.

See Corollary 1.3.9. The multiplicity of intersection of a curve and a line is the one that was defined there.  $\square$

The proof in the general case requires some algebra that we would rather defer. It will be given later (Theorem 7.9.1). It is possible to determine the intersections by counting the zeros of the resultant with respect to one of the variables. To do this, one chooses generic coordinates  $x, y, z$ . Then neither  $C$  nor  $D$  contains the point  $(0, 0, 1)$ . One writes their defining polynomials  $f$  and  $g$  as polynomials in  $z$  with coefficients in  $\mathbb{C}[x, y]$ . The resultant  $R$  with respect to  $z$  will be a homogeneous polynomial in  $x, y$ , of degree  $mn$ . It will have  $mn$  zeros in  $\mathbb{P}_{x,y}^1$ , counted with multiplicity. If  $\tilde{p} = (x_0, y_0)$  is a zero of  $R$ ,  $f(x_0, y_0, z)$  and  $g(x_0, y_0, z)$ , which are polynomials in  $z$ , have a common root  $z = z_0$ , and then  $p = (x_0, y_0, z_0)$  will be a point of  $C \cap D$ . It is a fact that the multiplicity of the zero of the resultant  $R$  at the image  $\tilde{p}$  is the (as yet undefined) intersection multiplicity of  $C$  and  $D$  at  $p$ . Unfortunately, this won't be obvious, even when multiplicity is defined. However, one can prove the next proposition using this approach.

**1.10.3. Proposition.** Let  $C$  and  $D$  be distinct plane curves of degrees  $m$  and  $n$ , respectively.

(i) The curves  $C$  and  $D$  have at least one point of intersection, and the number of intersections is at most  $mn$ .

(ii) If all intersections are transversal, the number of intersections is precisely  $mn$ .

It isn't obvious that two curves in the projective plane intersect. If two curves in the affine plane have no intersection, if they are parallel lines, for instance, their closures in the projective plane meet on the line at infinity.

**1.10.4. Lemma.** Let  $f$  and  $g$  be homogeneous polynomials in  $x, y, z$  of degrees  $m$  and  $n$ , respectively, and suppose that the point  $(0, 0, 1)$  isn't a zero of  $f$  or  $g$ . If the resultant  $\text{Res}_z(f, g)$  with respect to  $z$  is identically zero, then  $f$  and  $g$  have a common factor.

*proof.* Let the degrees of  $f$  and  $g$  be  $m$  and  $n$ , respectively, and let  $F$  denote the field of rational functions  $\mathbb{C}(x, y)$ . If the resultant is zero,  $f$  and  $g$  have a common factor in  $F[z]$  (Corollary 1.7.6). There will be polynomials  $p$  and  $q$  in  $F[z]$ , of degrees at most  $n-1$  and  $m-1$  in  $z$ , respectively, such that  $pf = qg$  (1.7.2). We may clear denominators, so we may assume that the coefficients of  $p$  and  $q$  are in  $\mathbb{C}[x, y]$ . Then  $pf = qg$  is an equation in  $\mathbb{C}[x, y, z]$ . Since  $p$  has degree at most  $n-1$  in  $z$ , it isn't divisible by  $g$ , which has degree  $n$  in  $z$ . Since  $\mathbb{C}[x, y, z]$  is a unique factorization domain,  $f$  and  $g$  have a common factor.  $\square$

*proof of Proposition 1.10.3.* (i) Let  $f$  and  $g$  be irreducible polynomials whose zero sets  $C$  and  $D$ , are distinct. Proposition 1.3.11 shows that there are finitely many intersections. We project to  $\mathbb{P}^1$  from a point  $q$  that doesn't



lie on any of the finitely many lines through pairs of intersection points. Then a line through  $q$  passes through at most one intersection, and the zeros of the resultant  $\text{Res}_z(f, g)$  that correspond to the intersection points will be distinct. Since the resultant has degree  $mn$  (1.7.8), it has at least one zero, and at most  $mn$  of them. Therefore  $C$  and  $D$  have at least one and at most  $mn$  intersections.

(ii) Every zero of the resultant will be the image of an intersection of  $C$  and  $D$ . To show that there are  $mn$  intersections if all intersections are transversal, it suffices to show that the resultant has simple zeros. This is Corollary 1.9.15.  $\square$

**1.10.5. Corollary.** *If the curve  $X$  defined by a homogeneous polynomial  $f(x, y, z)$  is smooth, then  $f$  is irreducible, and therefore  $X$  is a smooth curve.*

*proof.* Suppose that  $f = gh$ , and let  $p$  be a point of intersection of the loci  $\{g = 0\}$  and  $\{h = 0\}$ . The previous proposition shows that such a point exists. All partial derivatives of  $f$  vanish at  $p$ , so  $p$  is a singular point of  $X$ .  $\square$

**1.10.6. Corollary.** (i) *Let  $d$  be an integer  $\geq 3$ . A smooth plane curve of degree  $d$  has at least one flex point, and the number of flex points is at most  $3d(d-2)$ .*

(ii) *If all flex points are ordinary, the number of flex points is equal to  $3d(d-2)$ .*

Thus smooth curves of degrees 2, 3, 4, 5, ... have at most 0, 9, 24, 45, ... flex points, respectively.

*proof.* (i) The flex points are intersections of a smooth curve  $C$  with its Hessian divisor  $D : \{\det H = 0\}$ . (The definition of divisor is given in (1.3.12.)) Let  $C : \{f(x_0, x_1, x_2) = 0\}$  be a smooth curve of degree  $d$ . The entries of the  $3 \times 3$  Hessian matrix  $H$  are the second partial derivatives  $\frac{\partial^2 f}{\partial x_i \partial x_j}$ . They are homogeneous polynomials of degree  $d-2$ , so the Hessian determinant is homogeneous, of degree  $3(d-2)$ . Propositions 1.4.18 and 1.10.3 tell us that there are at most  $3d(d-2)$  intersections.

(ii) Recall that a flex point is ordinary if the multiplicity of intersection of the curve and its tangent line is 3. Bézout's Theorem asserts that the number of flex points is equal to  $3d(d-2)$  if the intersections of  $C$  with its Hessian divisor  $D$  are transversal, and therefore have multiplicity 1. So the next lemma completes the proof.

**1.10.7. Lemma.** *A curve  $C : \{f = 0\}$  intersects its Hessian divisor  $D$  transversally at a point  $p$  if and only if  $p$  is an ordinary flex point of  $C$ .*

*proof.* We prove this by computation. There may be a conceptual proof, but I don't know one.

Let  $L$  be the tangent line to  $C$  at the flex point  $p$ , and let  $h$  denote the restriction of the Hessian determinant to  $L$ . The Hessian divisor  $D$  will be transversal to  $C$  at  $p$  if and only if it is transversal to  $L$ , and this will be true if and only if the order of vanishing of  $h$  at  $p$  is 1.

We adjust coordinates  $x, y, z$  so that  $p = (0, 0, 1)$  and  $L$  is the line  $\{y = 0\}$ , and we write the polynomial  $f$  of degree  $d$  as

$$(1.10.8) \quad f(x, y, z) = \sum_{i+j+k=d} a_{ij} x^i y^j z^k,$$

We set  $y = 0$  and  $z = 1$ , to restrict  $f$  to  $L$ . The restricted polynomial is

$$f(x, 0, 1) = \sum_{i \leq d} a_{i0} x^i$$

Since  $p$  is a flex point with tangent line  $L$ , the coefficients  $a_{00}$ ,  $a_{10}$ , and  $a_{20}$  are zero, and  $p$  is an ordinary flex point if and only if the coefficient  $a_{30}$  is nonzero.

Let  $h$  be the restriction of  $\det H$  to  $L$ :  $h = \det H(x, 0, 1)$ . We must show that  $p$  is an ordinary flex point if and only if  $h$  has a simple zero at  $x = 0$ .

To evaluate the restriction  $f_{xx}(x, 0, 1)$  of the partial derivative to  $L$ , the relevant terms in the sum (1.10.8) have  $j = 0$ . Since  $a_{00} = a_{10} = 0$ ,

$$f_{xx}(x, 0, 1) = 6a_{30} + 12a_{40}x^2 + \cdots = 6a_{30}x + O(2)$$

Similarly,

$$\begin{aligned} f_{xz}(x, 0, 1) &= 0 + O(2) \\ f_{zz}(x, 0, 1) &= 0 + O(2) \end{aligned}$$

For the restriction of  $f_{yz}$ , the relevant terms are those with  $j = 1$ :

$$f_{yz}(x, 0, 1) = (d-1)a_{01} + (d-2)a_{11}x + O(2)$$

We don't need  $f_{xy}$  or  $f_{yy}$ .

Let  $v = 6a_{30}x$  and  $w = (d-1)a_{01} + (d-2)a_{11}x$ . The restricted Hessian matrix has the form

$$(1.10.9) \quad H(x, 0, 1) = \begin{pmatrix} v & * & 0 \\ * & * & w \\ 0 & w & 0 \end{pmatrix} + O(2)$$

where  $*$  are entries that don't affect terms of degree at most one in the determinant. The determinant is

$$h = -vw^2 + O(2) = -6(d-1)^2 a_{30}a_{01}^2 x + O(2)$$

It has a zero of order 1 at  $x = 0$  if and only if  $a_{30}$  and  $a_{01}$  aren't zero. Since  $C$  is smooth at  $p$  and  $a_{10} = 0$ , the coefficient  $a_{01}$  isn't zero. Thus the curve  $C$  and its Hessian divisor  $D$  intersect transversally, and  $C$  and  $L$  intersect with multiplicity 3, if and only if  $a_{30}$  is nonzero, which is true if and only if  $p$  is an ordinary flex.  $\square$

**1.10.10. Corollary.** *A smooth cubic curve contains exactly 9 flex points.*

*proof.* Let  $f$  be the irreducible cubic polynomial whose zero locus is a smooth cubic  $C$ . The degree of the Hessian divisor  $D$  is also 3, so Bézout predicts at most 9 intersections of  $D$  with  $C$ . To derive the corollary, we show that  $C$  intersects  $D$  transversally. According to Proposition 1.10.7, a nontransversal intersection would correspond to a point at which the curve and its tangent line intersect with multiplicity greater than 3. This is impossible when the curve is a cubic.  $\square$

#### (1.10.11) singularities of the dual curve

Let  $C$  be a plane curve. As before, an *ordinary flex point* is a smooth point  $p$  such that the intersection multiplicity of the curve and its tangent line  $L$  at  $p$  is precisely 3. A *bitangent* to  $C$  is a line  $L$  that is tangent to  $C$  at distinct smooth points  $p$  and  $q$ , and an *ordinary bitangent* is one such that neither  $p$  nor  $q$  is a flex point. A tangent line  $L$  at a smooth point  $p$  of  $C$  is an *ordinary tangent* if it isn't a flex point or a bitangent.

The line  $L$  will have other intersections with  $C$ . Most often, these other intersections will be transversal. However, it may happen that  $L$  is tangent to  $C$  at such a point, or that it is a singular point of  $C$ . Let's call such occurrences *accidents*.

**1.10.12. Proposition.** *Let  $p$  be a smooth point of a curve  $C$ , and let  $L$  be the tangent line at  $p$ . Suppose that there are no accidents.*

- (i) *If  $L$  is an ordinary tangent at  $p$ , then  $L^*$  is a smooth point of  $C^*$ .*
- (ii) *If  $L$  is an ordinary bitangent, then  $L^*$  is a node of  $C^*$ .*
- (iii) *If  $p$  is an ordinary flex point, then  $L^*$  is a cusp of  $C^*$ .*

*proof.* We refer to the map  $U \xrightarrow{t} C^*$  (1.6.3) from the set of smooth points of  $C$  to the dual curve. We set  $z = 1$  and choose affine coordinates so that  $p$  is the origin, and the tangent line  $L$  at  $p$  is the line  $\{y = 0\}$ . Let  $\tilde{f}(x, y) = f(x, y, 1)$ . We solve  $\tilde{f} = 0$  for  $y = y(x)$  as analytic function of  $x$  near zero, as before. The tangent line  $L_1$  to  $C$  at a nearby point  $p_1 = (x, y)$  has the equation (1.6.11), and  $L_1^*$  is the point  $(u, v, w) = (-y', 1, y'x - y)$  of  $\mathbb{P}^*$  (1.6.12). Since there are no accidents, this path traces out all points of  $C^*$  near to  $L^*$ .

If  $L$  is an ordinary tangent line,  $y(x)$  will have a zero of order 2 at  $x = 0$ . Then  $u = -y'$  will have a simple zero. So the path  $(-y', 1, y'x - y)$  is smooth at  $x = 0$ , and therefore  $C^*$  is smooth at the origin.

If  $L$  is an ordinary bitangent, tangent to  $C$  at two points  $p$  and  $p'$ , the reasoning given for an ordinary tangent shows that the images in  $C^*$  of small neighborhoods of  $p$  and  $p'$  in  $C$  will be smooth at  $L^*$ . Their tangent lines  $p^*$  and  $p'^*$  will be distinct, so  $p$  is a node.

The case that  $p$  is an ordinary flex point of  $C$  is trickier. Most probably, we won't know the defining equation  $f = 0$  of  $C$ . We write the analytic function  $y(x)$  that solves  $f(x, y) = 0$  as a power series. Since  $p$  is a flex point, the coefficients of  $x^i$  are zero when  $i < 3$ :  $y(x) = cx^3 + \dots$ . Since the flex is ordinary, we may assume that  $c = 1$ . In the local equation  $(u, v, w) = (-y', 1, y'x - y)$  for the dual curve,  $u = -3x^2 + \dots$  and  $w = 2x^3 + \dots$ . The locus

$$(1.10.13) \quad (u, w) = (-y', y'x - y) = (-3x^2 + \dots, 2x^3 + \dots)$$

contains the points of  $C^*$  near to  $L^*$ .

Let  $X$  and  $U$  denote the  $x$ -line and the  $u$ -line, respectively. We substitute (1.10.13) for  $u$  and  $v$ :  $u = -3x^2 + \dots$  and  $w = 2x^3 + \dots$ . This gives us a diagram of maps

$$\begin{array}{ccc} C & \xrightarrow{b} & U \\ t \downarrow & & \parallel \\ C^* & \xrightarrow{c} & U \end{array}$$

that are defined in small neighborhoods of the origins in the three spaces. The map  $t$  is locally bijective, and since the leading term of  $u(x)$  is  $3x^2$ ,  $b$  has degree 2. Therefore  $c$  also has degree 2. This implies that the origin in  $C^*$  is a point of multiplicity 2, a double point.

Let  $g(u, w) = \sum_{ij} g_{ij} u^i w^j$  be the irreducible polynomial equation for  $C^*$ . Substituting for  $u$  and  $w$ , the series in  $x$  that we obtain evaluates to zero for all small  $x$ , and this implies that it is the zero series. The orders of vanishing of the monomials  $u^i w^j$  as functions of  $x$  are as follows:

$$(1.10.14) \quad \begin{array}{cccccccccccc} 1 & u & w & u^2 & uw & w^2 & u^3 & u^2w & uw^2 & w^3 & \dots \\ 0 & 2 & 3 & 4 & 5 & 6 & 6 & 7 & 8 & 9 & \dots \end{array}$$

Looking at these orders of vanishing, one sees that the coefficients  $g_{00}, g_{10}, g_{01}, g_{20}$  and  $g_{11}$  in the series  $g = \sum g_{ij} u^i w^j$  must be zero, and that  $g_{02} + g_{30} = 0$ . Since the origin is a double point of  $C^*$ ,  $g_{02} \neq 0$ , and therefore  $g_{30} \neq 0$ . The origin of  $C^*$  is a cusp.  $\square$

figure

**1.10.15. Corollary.** *A plane curve has finitely many bitangents.*

This corollary is true whether or not the bitangents are ordinary. It follows from the fact that the dual curve  $C^*$  has finitely many singular points (1.4.7). If  $L$  is a bitangent, ordinary or not,  $L^*$  will be a singular point of  $C^*$ .  $\square$

## 1.11 The Plücker Formulas

A plane curve  $C$  is *ordinary* if it is smooth, if all of its bitangents and flex points are ordinary (see (1.10.11)), and if there are no accidents. The *Plücker formulas* compute the number of flexes and bitangents of an ordinary plane curve.

For the next theorem, we refer back to the notation of Section 1.7.20. With coordinates in general position, let  $\pi : C \rightarrow X$  be the projection of a plane curve  $C$  to the projective line  $X$  from  $q = (0, 0, 1)$ . If  $\tilde{p} = (x_0, y_0)$  is a point of  $X$ , we denote by  $L_{\tilde{p}}$  the line in  $\mathbb{P}^2$  such that the fibre of  $\pi$  over  $\tilde{p}$  is the complement of  $q$  in  $L_{\tilde{p}}$ .

The covering  $\pi$  will be branched at the points  $\tilde{p} = (x_0, y_0)$  of  $X$  such that  $L_{\tilde{p}}$  is the tangent line at some point of  $C$ , or contains a singular point of  $C$ .

**1.11.1. Theorem: Plücker Formulas.** *Let  $C$  be an ordinary curve of degree  $d$  at least two, and let  $C^*$  be its dual curve. Let  $f$  and  $b$  denote the numbers of flex points and bitangents of  $C$ , and let  $f^*$ ,  $\delta^*$  and  $\kappa^*$  denote the degree, the numbers of nodes, and the number of cusps of  $C^*$ , respectively. Then:*

(i) *The dual curve  $C^*$  has no flexes or bitangents. Its singularities are nodes and cusps.*

(ii)  $d^* = d^2 - 2$ ,  $f = \kappa^* = 3d(d - 2)$ , and  $b = \delta^* = \frac{1}{2}d(d - 2)(d^2 - 9)$ .

*proof.* (i) A bitangent or a flex on  $C^*$  would produce a singularity on the bidual  $C^{**}$ , which is the smooth curve  $C$ .

(ii) The degree  $d^*$  was computed in Corollary 1.7.30. Bézout's Theorem counts the flex points (see (1.10.6)). The facts that  $\kappa^* = f$  and  $\delta^* = b$  are dealt with in Proposition 1.10.12. Thus  $\kappa^* = f = 3d(d-2)$ .

We project  $C^*$  to  $\mathbb{P}^1$  from a generic point  $s$  of  $\mathbb{P}^*$ . The number of branch points that correspond to tangent lines through  $s$  at smooth points of  $C^*$  is the degree of  $C^{**} = C$  (1.7.30), which is  $d$ .

Next, let  $F$  be the defining polynomial for  $C^*$ . The discriminant  $\text{Discr}_z(F)$  has degree  $d^{*2} - d^*$ . Proposition 1.9.13 describes the order of vanishing of the discriminant at the images of the  $d$  tangent lines through  $s$ , the  $\delta$  nodes of  $C^*$ , and the  $\kappa$  cusps of  $C^*$ . It tells us that

$$d^{*2} - d^* = d + 2\delta^* + 3\kappa^*$$

Substituting the known values  $d^* = d^2 - d$ , and  $\kappa^* = 3d(d-2)$  into this formula gives us

$$(d^2 - d)^2 - (d^2 - d) = d + 2\delta^* + 9d(d-2) \quad \text{or} \quad 2\delta^* = d^4 - 2d^3 - 9d^2 + 18d \quad \square$$

**Note.** It isn't easy to count the number of bitangents directly.

### 1.11.2. Examples.

- (i) All curves of degree 2 and all smooth curves of degree 3 are ordinary.
- (ii) A curve of degree 2 has no flexes and no bitangents. Its dual curve has degree 2.
- (iii) A smooth curve of degree 3 has 9 flexes and no bitangents. Its dual curve has degree 6.
- (iv) An ordinary curve  $C$  of degree 4 has 24 flexes and 28 bitangents. Its dual curve has degree 12.  $\square$

We will make use of the fact that a quartic curve has 28 bitangents in Chapter ?? (see (4.9.15)). The Plücker Formulas are rarely used for curves of degree greater than four.

**1.11.3. Example.** The dual of a cubic curve  $C$  with a cusp was computed in (1.6.8). Let's do this again, using what we have learned. When we project to  $\mathbb{P}^1$  from a generic point, the discriminant will have degree 6. I will have a triple zero at the image of the cusp, and three simple zeros. Therefore the dual curve  $C^*$  has degree 3. Since  $C$  has genus zero, so does  $C^*$ . Since  $C$  has degree 3, there are no bitangents, and therefore  $C^*$  has no nodes. So  $C^*$  is another cuspidal cubic.  $\square$

## Chapter 2 AFFINE ALGEBRAIC GEOMETRY

- 2.1 Rings and Modules
- 2.2 The Zariski Topology
- 2.3 Some Affine Varieties
- 2.4 The Nullstellensatz
- 2.5 The Spectrum
- 2.6 Localization
- 2.7 Morphisms of Affine Varieties
- 2.8 Finite Group Actions

We study varieties of arbitrary dimension in the next chapters. We will use some of the basic terminology that was introduced in Chapter 1, including the concepts of discriminant and transcendence degree, but most of the results of Chapter 1 won't be used again until Chapter 8.

We begin by reviewing some basic facts about rings and modules, omitting proofs. Please look up information on the concepts that aren't familiar, as needed.

### 2.1 Rings and Modules

By the word 'ring', we mean 'commutative ring',  $ab = ba$ , unless when the contrary is stated explicitly. A *domain* is a ring that has no zero divisors and isn't the zero ring. An *algebra* is a ring that contains the field  $\mathbb{C}$  of complex numbers as subring.

A set of elements  $\alpha = \{\alpha_1, \dots, \alpha_n\}$  *generates* an algebra  $A$  if every element of  $A$  can be expressed (usually not uniquely) as a polynomial in  $\alpha_1, \dots, \alpha_n$ , with complex coefficients. Another way to state this is that  $\alpha$  generates  $A$  if the homomorphism  $\mathbb{C}[x_1, \dots, x_n] \xrightarrow{\tau} A$  that evaluates a polynomial at  $\alpha$  is surjective. If  $\alpha$  generates  $A$ , then  $A$  will be isomorphic to the quotient  $\mathbb{C}[x]/I$  of the polynomial algebra  $\mathbb{C}[x]$ , where  $I$  is the kernel of  $\tau$ . A *finite-type algebra* is one that can be generated by a finite set of elements.

If  $I$  and  $J$  are ideals of a ring  $R$ , the *product ideal*, which is denoted by  $IJ$ , is the ideal whose elements are finite sums of products  $\sum a_i b_i$ , with  $a_i \in I$  and  $b_i \in J$ . (This is not the product set, whose elements are the products  $ab$ , with  $a \in I$  and  $b \in J$ .) The power  $I^k$  of  $I$  is the product of  $k$  copies of  $I$ , the ideal spanned by products of  $k$  elements of  $I$ . The intersection  $I \cap J$  is also an ideal, and

$$(2.1.1) \quad (I \cap J)^2 \subset IJ \subset I \cap J$$

An ideal  $M$  of a ring  $R$  is *maximal* if it isn't the unit ideal  $R$ , and if there is no ideal  $I$  such that  $M < I < R$ . This is true if and only if the quotient ring  $R/M$  is a field.

An ideal  $P$  of a ring  $R$  is a *prime ideal* if the quotient  $R/P$  is a domain. A maximal ideal is a prime ideal.

**2.1.2. Lemma.** *Let  $P$  be an ideal of a ring  $R$  that isn't the unit ideal. The following conditions are equivalent.*

- (i)  $P$  is a prime ideal.
- (ii) If  $a$  and  $b$  are elements of  $R$  and if  $ab \in P$ , then  $a \in P$  or  $b \in P$ .
- (iii) If  $A$  and  $B$  are ideals of  $R$ , and if the product ideal  $AB$  is contained in  $P$ , then  $A \subset P$  or  $B \subset P$ . □

It is sometimes convenient to state (iii) this way:

- (iii') If  $A$  and  $B$  are ideals that contain  $P$ , and if the product ideal  $AB$  is contained in  $P$ , then  $A = P$  or  $B = P$ .

**2.1.3. Mapping Property of Quotient Rings.** Let  $R$  and  $S$  be rings, let  $K$  be an ideal of  $R$ , and let  $R \xrightarrow{\tau} \bar{R}$  denote the canonical map from  $R$  to the quotient ring  $\bar{R} = R/K$ . Homomorphisms  $\bar{R} \xrightarrow{\bar{\varphi}} S$  correspond bijectively to homomorphisms  $R \xrightarrow{\varphi} S$  whose kernels contain  $K$ , the correspondence being  $\varphi = \bar{\varphi} \circ \tau$ :

$$\begin{array}{ccc} R & \xrightarrow{\varphi} & S \\ \tau \downarrow & & \parallel \\ \bar{R} & \xrightarrow{\bar{\varphi}} & S \end{array}$$

If  $\ker \varphi = I$ , then  $\ker \bar{\varphi} = I/K$ . □

#### (2.1.4) commutative diagrams

In the diagram displayed above, the maps  $\bar{\varphi}\tau$  and  $\varphi$  from  $R$  to  $S$  are equal. This is referred to by saying that the diagram is *commutative*. A commutative diagram is one in which every map that can be obtained by composing its arrows depends only on the domain and range of the map. In these notes, all diagrams of maps are commutative. We won't mention commutativity most of the time. □

#### 2.1.5. Correspondence Theorem.

(i) Let  $R \xrightarrow{\varphi} S$  be a **surjective** ring homomorphism with kernel  $K$ . For instance,  $\varphi$  might be the canonical map from  $R$  to the quotient algebra  $R/K$ . There is a bijective correspondence

$$\{\text{ideals of } R \text{ that contain } K\} \longleftrightarrow \{\text{ideals of } S\}$$

This correspondence associates an ideal  $I$  of  $R$  that contains  $K$  with its image  $\varphi(I)$  in  $S$  and it associates an ideal  $J$  of  $S$  with its inverse image  $\varphi^{-1}(J)$  in  $R$ .

If an ideal  $I$  of  $R$  that contains  $K$  corresponds to an ideal  $J$  of  $S$ , then  $\varphi$  induces an isomorphism of quotient rings  $R/I \rightarrow S/J$ . If one of the ideals,  $I$  or  $J$ , is prime or maximal, they both are.

(ii) Let  $R$  be a ring, and let  $M \xrightarrow{\varphi} N$  be a surjective homomorphism of  $R$ -modules with kernel  $L$ . There is a bijective correspondence

$$\{\text{submodules of } M \text{ that contain } L\} \longleftrightarrow \{\text{submodules of } N\}$$

This correspondence associates a submodule  $S$  of  $M$  that contains  $L$  with its image  $\varphi(S)$  in  $N$  and it associates a submodule  $T$  of  $N$  with its inverse image  $\varphi^{-1}(T)$  in  $M$ . □

**2.1.6. Chinese Remainder Theorem.** Let  $I_1, \dots, I_k$  be comaximal ideals of a ring  $R$ .

(i) The product ideal  $I_1 \cdots I_k$  is equal to the intersection  $I_1 \cap \cdots \cap I_k$ .

(ii) The map  $R \rightarrow R/I_1 \times \cdots \times R/I_k$  that sends an element  $a$  of  $R$  to its vector of residues is a surjective homomorphism whose kernel is  $I_1 \cap \cdots \cap I_k$  ( $= I_1 \cdots I_k$ ).

(iii) Let  $M$  be an  $R$ -module. The canonical homomorphism  $M \rightarrow M/I_1 M \times \cdots \times M/I_k M$  is surjective. □

Ideals  $I_1, \dots, I_k$  of a ring  $R$  are said to be *comaximal* if the sum of any two of them is the unit ideal.

**2.1.7. Proposition.** Let  $R$  be a product of rings,  $R = R_1 \times \cdots \times R_k$ , let  $I$  be an ideal of  $R$ , and let  $\bar{R} = R/I$  be the quotient ring. There are ideals  $I_j$  of  $R_j$  such that  $I = I_1 \times \cdots \times I_k$  and  $\bar{R} = R_1/I_1 \times \cdots \times R_k/I_k$ . □

#### (2.1.8) Noetherian rings

A *finite module*  $M$  over a ring  $R$  is a module that is spanned, or generated, by a finite set  $\{m_1, \dots, m_k\}$  of elements. To say that the set generates means that every element of  $M$  can be obtained as a combination  $r_1 m_1 + \cdots + r_k m_k$  with coefficients  $r_i$  in  $R$ , or that the homomorphism from the free  $R$ -module  $R^k$  to  $M$  that sends a vector  $(r_1, \dots, r_k)$  to the combination  $r_1 m_1 + \cdots + r_k m_k$  is surjective. An ideal of a ring  $R$  is *finitely generated* if, when regarded as an  $R$ -module, it is a finite module.

A ring  $R$  is *noetherian* if all of its ideals are finitely generated. The ring  $\mathbb{Z}$  of integers is noetherian. Fields are noetherian. If  $I$  is an ideal of a noetherian ring  $R$ , the quotient ring  $R/I$  is noetherian.

**2.1.9. Hilbert Basis Theorem.** *If  $R$  is a noetherian ring, the ring  $R[x_1, \dots, x_n]$  of polynomials with coefficients in  $R$  is noetherian.*  $\square$

Thus  $\mathbb{Z}[x_1, \dots, x_n]$  and  $F[x_1, \dots, x_n]$ ,  $F$  a field, are noetherian rings.

**2.1.10. Corollary.** *Every finite-type algebra is noetherian.*  $\square$

**Note.** It is important not to confuse the concept of a finite-type algebra with that of a finite module. A finite  $R$ -module  $M$  is a module in which every element can be written as a (linear) combination  $r_1 m_1 + \dots + r_k m_k$  of some finite set  $\{m_1, \dots, m_k\}$  of elements of  $M$ , with coefficients in  $R$ . A finite-type algebra  $A$  is an algebra in which every element can be written as a polynomial  $f(\alpha_1, \dots, \alpha_k)$  in some finite set of elements  $\{\alpha_1, \dots, \alpha_k\}$  of  $A$ , with complex coefficients.

### (2.1.11) the ascending chain condition

The condition that a ring  $R$  be noetherian can be rewritten in several ways that we review here.

Our convention is that if  $X'$  and  $X$  are sets, the notation  $X' \subset X$  means that  $X'$  is a subset of  $X$ , while  $X' < X$  means that  $X'$  is a subset that is different from  $X$ . A *proper subset*  $X'$  of a set  $X$  is a nonempty subset different from  $X$  – a set such that  $\emptyset < X' < X$ .

A sequence  $X_1, X_2, \dots$ , finite or infinite, of subsets of a set  $Z$  forms an *increasing chain* if  $X_n \subset X_{n+1}$  for all  $n$ , equality  $X_n = X_{n+1}$  being permitted. If  $X_n < X_{n+1}$  for all  $n$ , the chain is *strictly increasing*.

Let  $\mathcal{S}$  be a set whose elements are subsets of a set  $Z$ . A member  $M$  of  $\mathcal{S}$  is a *maximal member* if there is no member  $M'$  of  $\mathcal{S}$  such that  $M < M'$ . For example, the set of proper subsets of a set of five elements contains five maximal members, the subsets of order four. The set of finite subsets of the set of integers contains no maximal member.

A maximal ideal of a ring  $R$  is a maximal member of the set of ideals of  $R$  different from the unit ideal.

**2.1.12. Proposition.** *The following conditions on a ring  $R$  are equivalent:*

- (i)  $R$  is **noetherian**: Every ideal of  $R$  is finitely generated.
- (ii) The **ascending chain condition**: Every strictly increasing chain  $I_1 < I_2 < \dots$  of ideals of  $R$  is finite.
- (iii) Every nonempty set of ideals of  $R$  contains a maximal member.  $\square$

The next corollary follows from the ascending chain condition, but the conclusions are true whether or not  $R$  is noetherian.

**2.1.13. Corollary.** *Let  $R$  be a noetherian ring.*

- (i) If  $R$  isn't the zero ring, every ideal of  $R$  except the unit ideal is contained in a maximal ideal.
- (ii) A nonzero ring  $R$  contains at least one maximal ideal.
- (iii) An element of a ring  $R$  that isn't in any maximal ideal is a unit – an invertible element of  $R$ .  $\square$

**2.1.14. Corollary.** *Let  $s_1, \dots, s_k$  be elements that generate the unit ideal of a noetherian ring  $R$ . For any positive integer  $n$ , the powers  $s_1^n, \dots, s_k^n$  generate the unit ideal.*  $\square$

**2.1.15. Proposition.** *Let  $R$  be a noetherian ring, and let  $M$  be a finite  $R$ -module.*

- (i) Every submodule of  $M$  is a finite module.
- (ii) The set of submodules of  $M$  satisfies the ascending chain condition.
- (iii) Every nonempty set of submodules of  $M$  contains a maximal member.  $\square$

This concludes our review of rings and modules.

## 2.2 The Zariski Topology

As before, the affine space  $\mathbb{A}^n$  is the space of  $n$ -tuples  $(a_1, \dots, a_n)$  of complex numbers. Algebraic geometry studies polynomial equations in terms of their solutions in affine space. If  $f_1, \dots, f_k$  are polynomials in  $x_1, \dots, x_n$ , the set of points of  $\mathbb{A}^n$  that solve the system of equations

$$(2.2.1) \quad f_1 = 0, \dots, f_k = 0$$

is a *Zariski closed* subset of  $\mathbb{A}^n$ . A *Zariski open* subset  $U$  is a subset whose complement, the set of points not in  $U$ , is Zariski closed.

When it seems unlikely to cause confusion, we may abbreviate the notation for an indexed set, using a single letter. The polynomial algebra  $\mathbb{C}[x_1, \dots, x_n]$  may be denoted by  $\mathbb{C}[x]$ , and the system of equations (2.2.1) by  $f = 0$ . The locus of solutions of the equations  $f = 0$  may be denoted by  $V(f_1, \dots, f_k)$  or by  $V(f)$ . Its points are called the *zeros* of the polynomials  $f$ .

We use analogous notation for infinite sets. If  $\mathcal{F}$  is any set of polynomials,  $V(\mathcal{F})$  denotes the set of points of affine space at which all elements of  $\mathcal{F}$  are zero. In particular, if  $I$  is an ideal of the polynomial ring,  $V(I)$  denotes the set of points at which all elements of  $I$  vanish.

The ideal  $I$  of  $\mathbb{C}[x]$  that is *generated* by the polynomials  $f_1, \dots, f_k$  is the set of combinations  $r_1 f_1 + \dots + r_k f_k$  with *polynomial* coefficients  $r_i$ . Some notations for this ideal are  $(f_1, \dots, f_k)$  and  $(f)$ . All elements of  $I$  vanish on the zero set  $V(f)$ , so  $V(f) = V(I)$ . The Zariski closed subsets of  $\mathbb{A}^n$  are the sets  $V(I)$ , where  $I$  is an ideal.

We note a few simple relations among ideals and their zero sets here. To begin with, we note that an ideal  $I$  isn't determined by its zero locus  $V(I)$ . For any  $k > 0$ , the power  $f^k$  has the same zeros as  $f$ .

The *radical* of an ideal  $I$  of a ring  $R$ , which will be denoted by  $\text{rad } I$ , is the set of elements  $\alpha$  of  $R$  such that some power  $\alpha^r$  is in  $I$ .

$$(2.2.2) \quad \text{rad } I = \{\alpha \in R \mid \alpha^r \in I \text{ for some } r > 0\}$$

The radical of  $I$  is an ideal that contains  $I$ . An ideal that is equal to its radical is a *radical ideal*. A prime ideal is a radical ideal.

The radical describes the ideals that define the same closed set.

**2.2.3. Lemma.** *If  $I$  is an ideal of the polynomial ring  $\mathbb{C}[x]$ , then  $V(I) = V(\text{rad } I)$ .* □

Consequently, if  $I$  and  $J$  are ideals and if  $\text{rad } I = \text{rad } J$ , then  $V(I) = V(J)$ . The converse of this statement is also true: If  $V(I) = V(J)$ , then  $\text{rad } I = \text{rad } J$ . This is a consequence of the *Strong Nullstellensatz* that will be proved later in this chapter. (See (2.4.9).)

Because  $(I \cap J)^2 \subset IJ \subset I \cap J$ ,

$$(2.2.4) \quad \text{rad}(IJ) = \text{rad}(I \cap J)$$

and  $\text{rad}(I \cap J) = (\text{rad } I) \cap (\text{rad } J)$ .

**2.2.5. Lemma.** *Let  $I$  and  $J$  be ideals of the polynomial ring  $\mathbb{C}[x]$ .*

- (i) *If  $I \subset J$ , then  $V(I) \supset V(J)$ .*
- (ii)  *$V(I^k) = V(I)$ .*
- (iii)  *$V(I \cap J) = V(IJ) = V(I) \cup V(J)$ .*
- (iv) *If  $I_\nu$  are ideals, then  $V(\sum I_\nu) = \bigcap V(I_\nu)$ .*

*proof.* (iii)  $V(I \cap J) = V(IJ)$  because the two ideals have the same radical, and because  $I$  and  $J$  contain  $IJ$ ,  $V(IJ) \supset V(I) \cup V(J)$ . To prove that  $V(IJ) \subset V(I) \cup V(J)$ , we note that  $V(IJ)$  is the locus of common zeros of the products  $fg$  with  $f$  in  $I$  and  $g$  in  $J$ . Suppose that a point  $p$  is a common zero:  $f(p)g(p) = 0$  for all  $f$  in  $I$  and all  $g$  in  $J$ . If  $f(p) \neq 0$  for some  $f$  in  $I$ , we must have  $g(p) = 0$  for every  $g$  in  $J$ , and then  $p$  is a point of  $V(J)$ . If  $f(p) = 0$  for all  $f$  in  $I$ , then  $p$  is a point of  $V(I)$ . In either case,  $p$  is a point of  $V(I) \cup V(J)$ . □

Zariski closed sets are the closed sets in the *Zariski topology* on  $\mathbb{A}^n$ . This topology is very useful in algebraic geometry, though it is very different from the classical topology.

To verify that the Zariski closed sets are the closed sets of a topology, one must show that

- the empty set and the whole space are Zariski closed,
- the intersection  $\bigcap C_\nu$  of an arbitrary family of Zariski closed sets is Zariski closed, and
- the union  $C \cup D$  of two Zariski closed sets is Zariski closed.

The empty set and the whole space are the zero sets of the elements 1 and 0, respectively. The other conditions follow from Lemma 2.2.5. □



**2.2.6. Example.** The proper Zariski closed subsets of the affine line, or of a plane affine curve, are finite sets. The proper Zariski closed subsets of the affine plane  $\mathbb{A}^2$  are finite unions of points and curves. Let's omit the proofs of these facts. The corresponding facts for loci in the projective line and the projective plane have been noted before (see (1.3.4) and (1.3.14)).  $\square$

figure

(Caption: A Zariski closed subset of the affine plane (real locus).)

A subset  $S$  of a topological space  $X$  becomes a topological space with its *induced topology*. The closed (or open) subsets of  $S$  in the induced topology are intersections  $S \cap Y$ , where  $Y$  is closed (or open) in  $X$ .

The induced topology on a subset  $S$  of  $\mathbb{A}^n$  will be called its *Zariski topology* too. A subset of  $S$  is closed in the Zariski topology if it has the form  $S \cap Y$  for some Zariski closed subset  $Y$  of  $\mathbb{A}^n$ . If  $S$  itself is a Zariski closed subset of  $\mathbb{A}^n$ , a closed subset of  $S$  will be a closed subset of  $\mathbb{A}^n$  that is contained in  $S$ .

Affine space also has a *classical topology*. A subset  $U$  of  $\mathbb{A}^n$  is open in the classical topology if, whenever a point  $p$  is in  $U$ , all points sufficiently near to  $p$  are in  $U$ . Since polynomial functions are continuous, their zero sets are closed in the classical topology. Therefore Zariski closed sets are closed in the classical topology too.

When two topologies  $T$  and  $T'$  on a set  $X$  are given,  $T'$  is said to be *coarser* than  $T$  if it contains fewer closed sets or fewer open sets, and *finer* than  $T$  if it contains more closed sets or more open sets. The Zariski topology is coarser than the classical topology. As the next proposition shows, it is much coarser.

**2.2.7. Proposition.** *Every nonempty Zariski open subset of  $\mathbb{A}^n$  is dense and path connected in the classical topology.*

*proof.* The (complex) line  $L$  through distinct points  $p$  and  $q$  of  $\mathbb{A}^n$  is a Zariski closed set whose points can be written as  $p + t(q - p)$ , with  $t$  in  $\mathbb{C}$ . It corresponds bijectively to the one-dimensional affine  $t$ -space  $\mathbb{A}^1$ , and the Zariski closed subsets of  $L$  correspond to Zariski closed subsets of  $\mathbb{A}^1$ . They are the finite subsets of  $L$ , and  $L$  itself.

Let  $U$  be a nonempty Zariski open set, and let  $C$  be its Zariski closed complement. To show that  $U$  is dense in the classical topology, we choose distinct points  $p$  and  $q$  of  $\mathbb{A}^n$ , with  $p$  in  $U$ . If  $L$  is the line through  $p$  and  $q$ ,  $C \cap L$  will be a Zariski closed subset of  $L$  that doesn't contain  $p$ , a finite set. In the classical topology, the closure of the complement of this finite set, which is  $U \cap L$ , will be the whole line  $L$ . Therefore the closure of  $U$  contains  $q$ , and since  $q$  was arbitrary, the closure of  $U$  is  $\mathbb{A}^n$ .

Next, let  $L$  be the line through two points  $p$  and  $q$  of  $U$ . As before,  $C \cap L$  will be a finite set. In the classical topology,  $L$  is a complex plane. The points  $p$  and  $q$  can be joined by a path in  $L$  that avoids a finite set.  $\square$

Though we will refer to the classical topology from time to time, the Zariski topology will appear more often. For this reason, we will refer to a Zariski closed subset simply as a *closed set*. Similarly, by an *open set* we mean a Zariski open set. We will mention the adjective "Zariski" only for emphasis.

## (2.2.8) irreducible closed sets

The fact that the polynomial algebra is a noetherian ring has important consequences for the Zariski topology that we discuss here.

A topological space  $X$  satisfies the *descending chain condition* on closed subsets if there is no infinite, strictly descending chain  $C_1 > C_2 > \dots$  of closed subsets of  $X$ . The descending chain condition on closed subsets is equivalent with the *ascending chain condition* on open sets.

A topological space that satisfies the descending chain condition on closed sets is called a *noetherian space*. In a noetherian space, every nonempty family  $\mathcal{S}$  of closed subsets has a minimal member, one that doesn't contain any other member of  $\mathcal{S}$ , and every nonempty family of open sets has a maximal member. (See (2.1.11).)

**2.2.9. Lemma.** *A noetherian topological space is quasicompact: Every open covering has a finite subcovering.*  $\square$

**2.2.10. Proposition.** *With its Zariski topology,  $\mathbb{A}^n$  is a noetherian space.*

*proof.* Suppose that a strictly descending chain  $C_1 > C_2 > \cdots$  of closed subsets of  $\mathbb{A}^n$  is given. Let  $I_j$  be the ideal of all elements of  $\mathbb{C}[x_1, \dots, x_n]$  that are identically zero on  $C_j$ . Then  $C_j = V(I_j)$ , and  $V(I_1) > V(I_2) > \cdots$ . Therefore  $I_1 < I_2 < \cdots$ . The ascending chain condition for ideals in  $\mathbb{C}[x_1, \dots, x_n]$  shows that the given chain is finite.  $\square$

**2.2.11. Definition.** A topological space  $X$  is *irreducible* if it isn't the union of two proper closed subsets. An *affine variety* is an irreducible closed subset of affine space  $\mathbb{A}^n$ .

Another way to say that  $X$  is irreducible is this:

*If  $C$  and  $D$  are closed subsets of  $X$ , and if  $X = C \cup D$ , then  $X = C$  or  $X = D$ .*

The concept of irreducibility is useful primarily for noetherian spaces. The only irreducible subsets of a Hausdorff space are its points. In particular, with the classical topology, the only irreducible subsets of affine space are points.

Irreducibility is somewhat analogous to connectedness. A topological space is *connected* if it isn't the union  $C \cup D$  of two proper *disjoint* closed subsets. However, the condition that a space be irreducible is much more restrictive because, in Definition 2.2.11, the closed sets  $C$  and  $D$  aren't required to be disjoint. In the Zariski topology on the affine plane, the union of two intersecting lines is connected, but not irreducible.

**2.2.12. Lemma.** *The following conditions on topological space  $X$  are equivalent.*

- $X$  is irreducible.
- The intersection  $U \cap V$  of two nonempty open subsets  $U$  and  $V$  of  $X$  is nonempty.
- Every nonempty open subset  $U$  of  $X$  is dense – its closure is  $X$ .

$\square$

**2.2.13. Theorem.** *In a noetherian topological space, every closed subset is the union of finitely many irreducible closed sets.*

*proof.* If a closed subset  $C_0$  of a topological space  $X$  isn't a union of finitely many irreducible closed sets, then it isn't irreducible, so it is a union  $C_1 \cup D_1$ , where  $C_1$  and  $D_1$  are proper closed subsets of  $C_0$ , and therefore closed subsets of  $X$ . Since  $C_0$  isn't a finite union of irreducible closed sets,  $C_1$  and  $D_1$  cannot both be finite unions of irreducible closed sets. Say that  $C_1$  isn't such a union. We have the beginning  $C_0 > C_1$  of a chain of closed subsets. We repeat the argument, replacing  $C_0$  by  $C_1$ , and we continue in this way, to construct an infinite, strictly descending chain  $C_0 > C_1 > C_2 > \cdots$ . So  $X$  isn't a noetherian space.  $\square$

Theorem 2.2.13 tells us that every closed subset of  $\mathbb{A}^n$  is a finite union of affine varieties. Since an affine variety is irreducible, it is connected in the Zariski topology. It is also connected in the classical topology, but this isn't very easy to prove. We may not get to it.

The *closure* of a subset  $S$  of a topological space  $X$  is the smallest closed subset that contains  $S$ . It is the intersection of all closed subsets that contain  $S$ .

**2.2.14. Lemma. (i)** *Let  $Z$  be a subspace of a topological space  $X$ , let  $S$  be a subset of  $Z$ , and let  $\bar{S}$  denote the closure of  $S$  in  $X$ . The closure of  $S$  in  $Z$  is the intersection  $\bar{S} \cap Z$ .*

**(ii)** *The closure  $\bar{Z}$  of a subspace  $Z$  of a topological space  $X$  is irreducible if and only if  $Z$  is irreducible.*

**(iii)** *A nonempty open subspace  $W$  of an irreducible space  $X$  is irreducible.*

*proof. (ii)* Let  $Z$  be an irreducible subset of  $X$ , and suppose that its closure  $\bar{Z}$  is the union  $\bar{C} \cup \bar{D}$  of two closed sets  $\bar{C}$  and  $\bar{D}$ . Then  $Z$  is the union of the sets  $C = \bar{C} \cap Z$  and  $D = \bar{D} \cap Z$ , and they are closed in  $Z$ . Therefore  $Z$  is one of those two sets; say  $Z = C$ . Then  $Z \subset \bar{C}$ , and since  $\bar{C}$  is closed,  $\bar{Z} \subset \bar{C}$ . Because  $\bar{C} \subset \bar{Z}$  as well,  $\bar{C} = \bar{Z}$ . Conversely, suppose that the closure  $\bar{Z}$  of a subset  $Z$  of  $X$  is irreducible, and that  $Z$  is a union  $C \cup D$  of closed subsets. Then  $\bar{Z} = \bar{C} \cup \bar{D}$ , and therefore  $\bar{Z} = \bar{C}$  or  $\bar{Z} = \bar{D}$ , say  $\bar{Z} = \bar{C}$ . So  $Z = \bar{C} \cap Z = C$ , and  $C$  is not a proper subset.

**(iii)** The closure of  $W$  is the irreducible space  $X$ .  $\square$

**(2.2.15) the coordinate algebra of a variety**

**2.2.16. Proposition.** *The affine varieties in  $\mathbb{A}^n$  are the sets  $V(P)$ , where  $P$  is a prime ideal of the polynomial algebra  $\mathbb{C}[x] = \mathbb{C}[x_1, \dots, x_n]$ . If  $P$  is a radical ideal, then  $V(P)$  is an affine variety if and only if  $P$  is a prime ideal.*

We will use this proposition in the next section, where we give a few examples of varieties, but we defer the proof to Section 2.5.

As before, an *algebra* is a ring that contains the complex numbers.

**2.2.17. Definition.** Let  $P$  be a prime ideal of the polynomial ring  $\mathbb{C}[x_1, \dots, x_n]$ , and let  $V$  be the affine variety  $V(P)$  in  $\mathbb{A}^n$ . The *coordinate algebra* of  $V$  is the quotient algebra  $A = \mathbb{C}[x]/P$ .

Geometric properties of the variety are reflected in algebraic properties of its coordinate algebra and vice versa. In a primitive sense, one can regard the geometry of an affine variety  $V$  as given by closed subsets and incidence relations – the inclusion of one closed set into another, as when a point lies on a line. A finer study of the geometry takes into account things such as tangency, but it is reasonable to begin by studying incidences  $C' \subset C$  among closed subvarieties. Such incidences translate into inclusions  $P' \supset P$  in the opposite direction among prime ideals. This is one reason that prime ideals are important.

## 2.3 Some affine varieties

This section contains a few simple examples of varieties.

**2.3.1.** A point  $p = (a_1, \dots, a_n)$  of affine space  $\mathbb{A}^n$  is the set of solutions of the  $n$  equations  $x_i - a_i = 0$ ,  $i = 1, \dots, n$ . A point is a variety because the polynomials  $x_i - a_i$  generate a maximal ideal in the polynomial algebra  $\mathbb{C}[x]$ , and a maximal ideal is a prime ideal. We denote that maximal ideal by  $\mathfrak{m}_p$ . It is the kernel of the substitution homomorphism  $\pi_p : \mathbb{C}[x] \rightarrow \mathbb{C}$  that evaluates a polynomial  $g(x_1, \dots, x_n)$  at  $p$ :  $\pi_p(g(x)) = g(a_1, \dots, a_n) = g(p)$ . As here, we denote that homomorphism by  $\pi_p$ .

The coordinate algebra of a point  $p$  is the quotient algebra  $\mathbb{C}[x]/\mathfrak{m}_p$ . It is also called the *residue field* at  $p$ , and it will be denoted by  $k(p)$ . The residue field at  $p$  is isomorphic to the image of  $\pi_p$ , which is the field  $\mathbb{C}$  of complex numbers, but  $k(p)$  is a particular quotient of the polynomial ring.

**2.3.2.** The varieties in the affine line  $\mathbb{A}^1$  are points, and the whole line  $\mathbb{A}^1$ . The varieties in the affine plane  $\mathbb{A}^2$  are points, plane affine curves, and the whole plane.

This is true because the varieties correspond to the prime ideals of the polynomial ring. The prime ideals of  $\mathbb{C}[x_1, x_2]$  are the maximal ideals, the principal ideals generated by irreducible polynomials, and the zero ideal. The proof of this is a good exercise.

**2.3.3.** The set  $X$  of solutions of a single irreducible polynomial equation  $f_1(x_1, \dots, x_n) = 0$  in  $\mathbb{A}^n$  is a variety, called an *affine hypersurface*.

For instance, the *special linear group*  $SL_2$ , the group of complex  $2 \times 2$  matrices with determinant 1, is a hypersurface in  $\mathbb{A}^4$ , the locus of zeros of the irreducible polynomial  $x_{11}x_{22} - x_{12}x_{21} - 1$ .

The reason that an affine hypersurface is a variety is that an irreducible element of a unique factorization domain is a prime element, and a prime element generates a prime ideal. The polynomial ring  $\mathbb{C}[x_1, \dots, x_n]$  is a unique factorization domain.

**2.3.4.** A hypersurface in the affine plane  $\mathbb{A}^2$  is a plane affine curve.

A *line* in the plane, the locus of a linear equation  $ax + by - c = 0$ , is a plane affine curve. Its coordinate algebra is isomorphic to a polynomial ring in one variable. Every line is isomorphic to the affine line  $\mathbb{A}^1$ .

**2.3.5.** Let  $p = (a_1, \dots, a_n)$  and  $q = (b_1, \dots, b_n)$  be distinct points of  $\mathbb{A}^n$ . The *point pair*  $(p, q)$  is the closed set defined by the system of  $n^2$  equations  $(x_i - a_i)(x_j - b_j) = 0$ ,  $1 \leq i, j \leq n$ . A point pair isn't a variety because the ideal  $I$  generated by the polynomials  $(x_i - a_i)(x_j - b_j)$  isn't a prime ideal. The next proposition, which follows from the Chinese Remainder Theorem 2.1.6, describes the ideal  $I$ .

**2.3.6. Proposition.** *The ideal of polynomials that vanish on a point pair  $p, q$  is the product  $\mathfrak{m}_p \mathfrak{m}_q$  of the maximal ideals at the points, and the quotient algebra  $\mathbb{C}[x]/I$  is isomorphic to the product algebra  $\mathbb{C} \times \mathbb{C}$ .  $\square$*

## 2.4 Hilbert's Nullstellensatz

**2.4.1. Nullstellensatz (version 1).** Let  $\mathbb{C}[x]$  be the polynomial algebra in the variables  $x_1, \dots, x_n$ . There are bijective correspondences between the following sets:

- points  $p$  of the affine space  $\mathbb{A}^n$ ,
- algebra homomorphisms  $\pi_p : \mathbb{C}[x] \rightarrow \mathbb{C}$ ,
- maximal ideals  $\mathfrak{m}_p$  of  $\mathbb{C}[x]$ .

If  $p = (a_1, \dots, a_n)$  is a point of  $\mathbb{A}^n$ , the corresponding homomorphism  $\pi_p$  evaluates a polynomial at  $p$ :  $\pi_p(g) = g(a_1, \dots, a_n) = g(p)$ , and the maximal ideal  $\mathfrak{m}_p$  is the kernel of  $\pi_p$ , which is the ideal generated by the linear polynomials  $x_1 - a_1, \dots, x_n - a_n$ .  $\square$

It is obvious that every algebra homomorphism  $\mathbb{C}[x] \rightarrow \mathbb{C}$  is surjective and that its kernel is a maximal ideal. It isn't obvious that every maximal ideal of  $\mathbb{C}[x]$  is the kernel of such a homomorphism. The proof can be found manywhere.<sup>1</sup>

The Nullstellensatz gives us a way to describe the closed set  $V(I)$  of zeros of an ideal  $I$  in affine space in terms of maximal ideals. The points of  $V(I)$  are those at which all elements of  $I$  vanish. Thus

$$(2.4.2) \quad V(I) = \{p \in \mathbb{A}^n \mid I \subset \mathfrak{m}_p\}$$

**2.4.3. Proposition.** Let  $I$  be an ideal of the polynomial ring  $\mathbb{C}[x]$ . If the zero locus  $V(I)$  is empty, then  $I$  is the unit ideal.

*proof.* Every ideal  $I$  that is not the unit ideal is contained in a maximal ideal (Corollary 2.1.13).  $\square$

**2.4.4. Nullstellensatz (version 2).** Let  $A$  be a finite-type algebra. There are bijective correspondences between the following sets:

- algebra homomorphisms  $\bar{\pi} : A \rightarrow \mathbb{C}$ ,
- maximal ideals  $\bar{\mathfrak{m}}$  of  $A$ .

The maximal ideal  $\bar{\mathfrak{m}}$  that corresponds to a homomorphism  $\bar{\pi}$  is the kernel of  $\bar{\pi}$ .

If  $A$  is presented as a quotient of a polynomial ring, say  $A \approx \mathbb{C}[x_1, \dots, x_n]/I$ , then these sets also correspond bijectively to points of the set  $V(I)$  of zeros of  $I$  in  $\mathbb{A}^n$ .

(We use the symbol  $\approx$  to indicate an isomorphism.) As before, a finite-type algebra is an algebra that can be generated by a finite set of elements.

*proof.* We choose a presentation of  $A$  as a quotient of a polynomial ring to identify  $A$  with a quotient  $\mathbb{C}[x]/I$ . The Correspondence Theorem tells us that maximal ideals of  $A$  correspond to maximal ideals of  $\mathbb{C}[x]$  that contain  $I$ . Those maximal ideals correspond to points of  $V(I)$  (see (2.4.2)).

Let  $\tau$  denote the canonical homomorphism  $\mathbb{C}[x] \rightarrow A$ . The Mapping Property 2.1.3, applied to  $\tau$ , tells us that homomorphisms  $A \xrightarrow{\bar{\pi}} \mathbb{C}$  correspond to homomorphisms  $\mathbb{C}[x] \xrightarrow{\pi} \mathbb{C}$  whose kernels contain  $I$ . Those homomorphisms also correspond to points of  $V(I)$ .

$$(2.4.5) \quad \begin{array}{ccc} \mathbb{C}[x] & \xrightarrow{\pi} & \mathbb{C} \\ \tau \downarrow & & \parallel \\ A & \xrightarrow{\bar{\pi}} & \mathbb{C} \end{array}$$

**2.4.6. Strong Nullstellensatz.** Let  $I$  be an ideal of the polynomial algebra  $\mathbb{C}[x_1, \dots, x_n]$ , and let  $V$  be the locus of zeros of  $I$  in  $\mathbb{A}^n$ :  $V = V(I)$ . If a polynomial  $g$  vanishes at every point of  $V$ , then  $I$  contains a power of  $g$ .

*proof.* This beautiful proof is due to Rainich. Let  $g(x)$  be a polynomial that is identically zero on  $V$ . We are to show that  $I$  contains a power of  $g$ . If  $g$  is the zero polynomial, it is in  $I$ , so we may assume that  $g$  isn't zero.

<sup>1</sup>While writing a paper, the mathematician Nagata decided that the English language needed this unusual word, and then he managed to find it in a dictionary.

The Hilbert Basis Theorem tells us that  $I$  is a finitely generated ideal; let  $f = f_1, \dots, f_k$  be a set of generators. In the  $n+1$ -dimensional affine space with coordinates  $(x_1, \dots, x_n, y)$ , let  $W$  be the locus of solutions of the  $k+1$  equations

$$(2.4.7) \quad f_1(x) = \dots = f_k(x) = 0 \quad \text{and} \quad g(x)y - 1 = 0$$

Suppose that we have a solution  $x$  of the equations  $f(x) = 0$ , say  $(x_1, \dots, x_n) = (a_1, \dots, a_n)$ . Then  $a$  is a point of  $V$ , and our hypothesis tells us that  $g(a) = 0$  too. So there can be no  $b$  such that  $g(a)b = 1$ . There is no point  $(a_1, \dots, a_n, b)$  that solves the equations (2.4.7): The locus  $W$  is empty. Proposition 2.4.3 tells us that the polynomials  $f_1, \dots, f_k, gy - 1$  generate the unit ideal of  $\mathbb{C}[x_1, \dots, x_n, y]$ . There are polynomials  $p_1(x, y), \dots, p_k(x, y)$  and  $q(x, y)$  such that

$$(2.4.8) \quad p_1 f_1 + \dots + p_k f_k + q(gy - 1) = 1$$

The ring  $R = \mathbb{C}[x, y]/(gy - 1)$  can be described as the one obtained by adjoining an inverse of  $g$  to the polynomial ring  $\mathbb{C}[x]$ . The residue of  $y$  is the inverse of  $g$ . Since  $g$  isn't zero,  $\mathbb{C}[x]$  is a subring of  $R$ . In  $R$ ,  $gy - 1 = 0$ , and the equation (2.4.8) becomes  $p_1 f_1 + \dots + p_k f_k = 1$ . When we multiply both sides of this equation by a large power  $g^N$  of  $g$ , we can use the equation  $gy = 1$ , which is true in  $R$ , to cancel all occurrences of  $y$  in the polynomials  $p_i(x, y)$ . Let  $h_i(x)$  denote the polynomial in  $x$  that is obtained by cancelling  $y$  in  $g^N p_i$ . Then

$$h_1(x)f_1(x) + \dots + h_k(x)f_k(x) = g^N(x)$$

is a polynomial equation that is true in  $R$  and in its subring  $\mathbb{C}[x]$ . Since  $f_1, \dots, f_k$  are in  $I$ , this equation shows that  $g^N$  is in  $I$ .  $\square$

**2.4.9. Corollary.** Let  $\mathbb{C}[x]$  denote the polynomial ring  $\mathbb{C}[x_1, \dots, x_n]$ .

- (i) Let  $P$  be a prime ideal of  $\mathbb{C}[x]$ , and let  $V = V(P)$  be the variety of zeros of  $P$  in  $\mathbb{A}^n$ . If a polynomial  $g$  vanishes at every point of  $V$ , then  $g$  is an element of  $P$ .
- (ii) Let  $f$  be an irreducible polynomial in  $\mathbb{C}[x]$ . If a polynomial  $g$  vanishes at every point of  $V(f)$ , then  $f$  divides  $g$ .
- (iii) Let  $I$  and  $J$  be ideals of  $\mathbb{C}[x]$ . Then  $V(I) \supset V(J)$  if and only if  $\text{rad } I \subset \text{rad } J$ , and  $V(I) > V(J)$  if and only if  $\text{rad } I > \text{rad } J$  (see (2.2.2)).  $\square$

#### 2.4.10. Examples.

(i) Let  $I$  be the ideal generated by  $y^5$  and  $y^2 - x^3$  in the polynomial algebra  $\mathbb{C}[x, y]$  in two variables. The origin  $y = x = 0$  is the only common zero of these polynomials in the affine plane, and the polynomial  $x$  also vanishes at the origin. The Strong Nullstellensatz predicts that  $I$  contains a power of  $x$ . This is verified by the following equation:

$$yy^5 - (y^4 + y^2x^3 + x^6)(y^2 - x^3) = x^9$$

(ii) We may regard pairs  $A, B$  of  $n \times n$  matrices as points of an affine space  $\mathbb{A}^{2n^2}$  with coordinates  $a_{ij}, b_{ij}$ ,  $1 \leq i, j \leq n$ . The pairs of commuting matrices ( $AB = BA$ ) form a closed subset of  $\mathbb{A}^{2n^2}$ , the locus of common zeros of the  $n^2$  polynomials  $c_{ij}$  that compute the entries of the matrix  $AB - BA$ :

$$(2.4.11) \quad c_{ij}(a, b) = \sum_{\nu} a_{i\nu} b_{\nu j} - b_{i\nu} a_{\nu j}$$

Let  $I$  denote the ideal of the polynomial algebra  $\mathbb{C}[a, b]$  generated by the polynomials  $c_{ij}$ . Then  $V(I)$  is the set of pairs of commuting complex matrices. The Strong Nullstellensatz asserts that if a polynomial  $g(a, b)$  vanishes on every pair of commuting matrices, some power of  $g$  is in  $I$ . Is  $g$  itself in  $I$ ? It is a famous conjecture that  $I$  is a prime ideal. If so,  $g$  would be in  $I$ . Proving the conjecture would establish your reputation as a mathematician, but I don't recommend spending very much time on it right now.  $\square$

## 2.5 The Spectrum

The Nullstellensatz allows us to associate a set of points to a finite-type domain  $A$  without reference to a presentation. We can do this because the maximal ideals of  $A$  and the homomorphisms  $A \rightarrow \mathbb{C}$  don't depend

on the presentation. When a finite-type domain  $A$  is presented as a quotient  $\mathbb{C}[x]/P$  of a polynomial ring, where  $P$  is a prime ideal, it becomes the coordinate algebra of the variety  $V(P)$  in affine space. Then the points of  $V(P)$  correspond to maximal ideals of  $A$  and also to homomorphisms  $A \rightarrow \mathbb{C}$ .

When a finite-type domain  $A$  is given without a presentation, we replace the variety  $V(P)$  by an abstract set of points, the *spectrum* of  $A$ , that we denote by  $\text{Spec } A$  and call an *affine variety*. We put one point  $p$  into the spectrum for every maximal ideal of  $A$ , and then we turn around and denote the maximal ideal that corresponds to a point  $p$  by  $\bar{m}_p$ . The Nullstellensatz tells us that  $p$  also corresponds to a homomorphism  $A \rightarrow \mathbb{C}$  whose kernel is  $\bar{m}_p$ . We denote that homomorphism by  $\bar{\pi}_p$ . In analogy with (2.2.17), the domain  $A$  is called the *coordinate algebra* of the affine variety  $\text{Spec } A$ . To work with  $\text{Spec } A$ , we may interpret its points as maximal ideals or as homomorphisms to  $\mathbb{C}$ , whichever is convenient.

When defined in this way, the variety  $\text{Spec } A$  isn't embedded into affine space, but if we present  $A$  as a quotient  $\mathbb{C}[x]/P$ , points of  $\text{Spec } A$  correspond to points of the subset  $V(P)$  in  $\mathbb{A}^n$ . Even when the coordinate ring  $A$  of an affine variety is presented as  $\mathbb{C}[x]/P$ , we may denote the variety by  $\text{Spec } A$  rather than by  $V(P)$ .

### Note about spec as prime ideals. Bijective correspondences between open or closed sets in maxspec and in spec ###

###

function determines element

###

Let  $X = \text{Spec } A$ . The elements of  $A$  define (complex-valued) functions on  $X$  as follows: A point  $p$  of  $X$  corresponds to a homomorphism  $A \xrightarrow{\bar{\pi}_p} \mathbb{C}$ . If  $\alpha$  is an element of  $A$ , the *value* of the function  $\alpha$  at  $p$  is defined to be  $\bar{\pi}_p(\alpha)$ :

$$(2.5.1) \quad \alpha(p) \stackrel{\text{def}}{=} \bar{\pi}_p(\alpha)$$

The kernel  $\bar{m}_p$  of  $\bar{\pi}_p$  is the set of elements  $\alpha$  of the coordinate algebra  $A$  such that  $\alpha(p) = 0$ :

$$\bar{m}_p = \{\alpha \in A \mid \alpha(p) = 0\}$$

The functions defined by the elements of  $A$  are called *regular functions* on  $X$ . (See Proposition 2.7.2 below.)

**2.5.2. Lemma.** *The regular function determined by distinct elements  $\alpha$  and  $\beta$  of  $A$  are distinct.*

*proof.* We replace  $\alpha$  by  $\alpha - \beta$ . Then what is to be shown is that, if the function determined by an element  $\alpha$  is the zero function, then  $\alpha$  is the zero element.

We present  $A$  as  $\mathbb{C}[x]/P$ ,  $x = x_1, \dots, x_n$ , where  $P$  is a prime ideal. Then  $X$  is the locus of zeros of  $P$  in  $\mathbb{A}^n$ , and  $P$  is also the ideal of *all* elements that are zero on  $X$  (??). Let  $g(x)$  be a polynomial that represents  $\alpha$ . If  $p$  is a point of  $X$  at which  $\alpha$  is zero, then  $g(p) = 0$ . So if  $\alpha$  is the zero function, then  $g$  is in  $P$ , and therefore  $\alpha = 0$ .  $\square$

###kill this??### For example, the spectrum  $\text{Spec } \mathbb{C}[x_1, \dots, x_n]$  of the polynomial algebra is the affine space  $\mathbb{A}^n$ . The homomorphism  $\pi_p : \mathbb{C}[x] \rightarrow \mathbb{C}$  that corresponds to a point  $p = (a_1, \dots, a_n)$  of  $\mathbb{A}^n$  is evaluation at  $p$ . So  $\pi_p(g) = g(a_1, \dots, a_n) = g(p)$ . The function defined by a polynomial  $g(x)$  is simply the polynomial function.

**Note.** In modern terminology, the word “spectrum” is usually used to denote the set of prime ideals of a ring. This becomes important when one studies rings that aren't finite-type algebras, but when working with finite-type algebras, there are enough maximal ideals. The other prime ideals aren't needed, so we have eliminated them.

**2.5.3. Lemma.** *Let  $A$  be a quotient  $\mathbb{C}[x]/P$  of the polynomial ring  $\mathbb{C}[x_1, \dots, x_n]$  modulo a prime ideal  $P$ , so that  $\text{Spec } A$  becomes the closed subset  $V(P)$  of  $\mathbb{A}^n$ . Then a point  $p$  of  $\text{Spec } A$  becomes a point  $p = (a_1, \dots, a_n)$  of  $\mathbb{A}^n$ . When an element  $\alpha$  of  $A$  is represented by a polynomial  $g(x)$ , the value of  $\alpha$  at  $p$  is  $\alpha(p) = g(a_1, \dots, a_n) = g(p)$ .*

*proof.* The point  $p$  of  $\text{Spec } A$  gives us a diagram (2.4.5), with  $\pi = \pi_p$  and  $\bar{\pi} = \bar{\pi}_p$ , and where  $\tau$  is the canonical map  $\mathbb{C}[x] \rightarrow A$ . Then  $\alpha = \tau(p)$ , and

$$(2.5.4) \quad g(p) \stackrel{\text{defn}}{=} \pi_p(g) = \bar{\pi}_p \tau(g) = \bar{\pi}_p(\alpha) \stackrel{\text{defn}}{=} \alpha(p).$$

□

Thus the value  $\alpha(p)$  at a point  $p$  of  $\text{Spec } A$  can be obtained by evaluating a polynomial  $g$  at  $p$ , though not in a unique way. The polynomial  $g$  that represents the regular function  $\alpha$  won't be unique unless  $P$  is the zero ideal.

### (2.5.5) the Zariski topology on an affine variety

Let  $X = \text{Spec } A$  be an affine variety with coordinate algebra  $A$ . An ideal  $\bar{J}$  of  $A$  defines a locus in  $X$ , a *closed subset*, that we denote by  $V_X(\bar{J})$ :

$$(2.5.6) \quad V_X(\bar{J}) = \{p \in \text{Spec } A \mid \bar{J} \subset \bar{\mathfrak{m}}_p\}$$

When a presentation  $\mathbb{C}[x]/P \approx A$ , is given, the ideal  $\bar{J}$  of  $A$  corresponds to an ideal  $J$  of  $\mathbb{C}[x]$  that contains  $P$ . Then if  $V_{\mathbb{A}^n}(J)$  denotes the zero locus of  $J$  in  $\mathbb{A}^n$ ,  $V_X(\bar{J}) = V_{\mathbb{A}^n}(J)$ .

The properties of closed sets in affine space that are given in Lemmas 2.2.3 and 2.2.5 are true for closed subsets of an affine variety. In particular,  $V_X(\bar{J}) = V_X(\text{rad } \bar{J})$ , and  $V_X(\bar{I}\bar{J}) = V_X(\bar{I} \cap \bar{J}) = V_X(\bar{I}) \cup V_X(\bar{J})$ .

**2.5.7. Proposition.** *Let  $\bar{J}$  be an ideal of a finite-type domain  $A$ , and let  $X = \text{Spec } A$ . The zero set  $V_X(\bar{J})$  is empty if and only if  $\bar{J}$  is the unit ideal of  $A$ . If  $X$  is empty, then  $A$  is the zero ring.*

*proof.* The zero ring is the only ring with no maximal ideals. □

**2.5.8. Note.** We have put bars on the symbols  $\bar{\mathfrak{m}}$  and  $\bar{\pi}$  here in order to distinguish maximal ideals of  $A$  from maximal ideals of  $\mathbb{C}[x]$  and homomorphisms  $A \rightarrow \mathbb{C}$  from homomorphisms  $\mathbb{C}[x_1, \dots, x_n] \rightarrow \mathbb{C}$ . In the future, we will put bars over the letters only when there is a danger of confusion. □

### (2.5.9) ideals whose zero sets are equal

**2.5.10. Lemma.** *An ideal  $I$  of a noetherian ring  $R$  contains a power of its radical.*

*proof.* Since  $R$  is noetherian, the ideal  $\text{rad } I$  is generated by a finite set of elements  $\alpha = \{\alpha_1, \dots, \alpha_k\}$ , and for large  $r$ ,  $\alpha_i^r$  is in  $I$ . We can use the same large integer  $r$  for every  $i$ . A monomial  $\beta = \alpha_1^{e_1} \cdots \alpha_k^{e_k}$  of sufficiently large degree  $n$  in  $\alpha$  will be divisible  $\alpha_i^r$  for at least one  $i$ , and therefore it will be in  $I$ . The monomials of degree  $n$  generate  $(\text{rad } I)^n$ , so  $(\text{rad } I)^n \subset I$ . And as has been remarked,  $I \subset \text{rad } I$ . □

**2.5.11. Corollary.** *Let  $I$  and  $J$  be ideals of a finite-type domain  $A$ , and let  $X = \text{Spec } A$ . Then  $V_X(I) \supset V_X(J)$  if and only if  $\text{rad } I \subset \text{rad } J$ .*

This follows from Corollary 2.4.9. □

For example, there is a bijective correspondence between radical ideals in the polynomial ring  $\mathbb{C}[x_1, \dots, x_n]$  and closed subsets of  $\mathbb{A}^n$ . #####

The next proposition includes Proposition 2.2.16 as a special case.

**2.5.12. Proposition.** *Let  $X = \text{Spec } A$ , where  $A$  is a finite-type domain. The closed subset  $V_X(P)$  defined by a radical ideal  $P$  is irreducible if and only if  $P$  is a prime ideal.*

*proof.* Let  $P$  be a radical ideal of  $A$ , and let  $Y = V_X(P)$ . Let  $C$  and  $D$  be closed subsets of  $X$  such that  $Y = C \cup D$ . Say  $C = V_X(I)$ ,  $D = V_X(J)$ . We may suppose that  $I$  and  $J$  are radical ideals. Then the inclusion  $C \subset Y$  implies that  $I \supset P$ . Similarly,  $J \supset P$ . Because  $Y = C \cup D$ , we also have  $Y = V_X(I \cap J) = V_X(IJ)$ . So  $IJ \subset P$  (Corollary 2.5.11). If  $P$  is a prime ideal, then  $I = P$  or  $J = P$ , and therefore  $C = Y$  or  $D = Y$ . So  $Y$  is irreducible. Conversely, suppose that  $P$  is not a prime ideal. Then there are ideals  $I, J$  strictly larger than  $P$ , such that  $IJ \subset P$  (2.1.2). Then  $Y$  will be the union of the two proper closed subsets  $V_X(I)$  and  $V_X(J)$ , and is not irreducible (2.5.11). □

### (2.5.13) the nilradical

The *nilradical* of a ring is the set of its nilpotent elements. It is the radical of the zero ideal. If a ring  $R$  is noetherian, its nilradical will be *nilpotent*: some power of it will be the zero ideal (Lemma 2.5.10). The nilradical of a domain is the zero ideal.

**2.5.14. Proposition.** *The nilradical of a noetherian ring  $R$  is the intersection of the prime ideals of  $R$ .*

*proof.* Let  $x$  be an element of the nilradical  $N$ . So some power of  $x$  is zero. Since the zero element is in every prime ideal,  $x$  is in every prime ideal. Therefore  $N$  is contained in every prime ideal. Conversely, let  $x$  be an element not in  $N$ , i.e., not nilpotent. We show that there is a prime ideal that doesn't contain any power of  $x$ . Let  $\mathcal{S}$  be the set of ideals that don't contain a power of  $x$ . The zero ideal is one such ideal, so  $\mathcal{S}$  isn't empty. Since  $R$  is noetherian,  $\mathcal{S}$  contains a maximal member  $P$  (2.1.11). We show that  $P$  is a prime ideal by showing that, if two ideals  $A$  and  $B$  are strictly larger than  $P$ , their product  $AB$  isn't contained in  $P$ . Since  $P$  is a maximal member of  $\mathcal{S}$ ,  $A$  and  $B$  aren't in  $\mathcal{S}$ . They contain powers of  $x$ , say  $x^k \in A$  and  $x^\ell \in B$ . Then  $x^{k+\ell}$  is in  $AB$  but not in  $P$ . Therefore  $AB \not\subset P$ .  $\square$

The conclusion of this proposition is true whether or not the ring  $R$  is noetherian.

### 2.5.15. Corollary.

- (i) *Let  $A$  be a finite-type algebra. An element that is in every maximal ideal of  $A$  is nilpotent.*
- (ii) *Let  $A$  be a finite-type domain. The intersection of the maximal ideals of  $A$  is the zero ideal.*

*proof.* (i) Say that  $A$  is presented as  $\mathbb{C}[x]/I$ . Let  $\alpha$  be an element of  $A$  that is in every maximal ideal, and let  $g(x)$  be a polynomial whose residue in  $A$  is  $\alpha$ . Then  $\alpha$  is in every maximal ideal of  $A$  if and only if  $g = 0$  at all points of  $V_{\mathbb{A}}(I)$ . If so, the Strong Nullstellensatz asserts that some power  $g^n$  is in  $I$ . Then  $\alpha^n = 0$ .  $\square$

**2.5.16. Corollary.** *An element  $\alpha$  of a finite-type domain  $A$  is determined by the function that it defines on  $X = \text{Spec } A$ .*

*proof.* It is enough to show that an element  $\alpha$  that defines the zero function is the zero element. Such an element is in every maximal ideal (2.5.7), so  $\alpha$  is nilpotent, and since  $A$  is a domain,  $\alpha = 0$ .  $\square$

## 2.6 Localization

Let  $s$  be a nonzero element of a domain  $A$ . The ring  $A[s^{-1}]$ , obtained by adjoining an inverse of  $s$  to  $A$  is called a *localization* of  $A$ . It is isomorphic to the quotient  $A[z]/(sz - 1)$  of the polynomial ring  $A[z]$  by the principal ideal generated by  $sz - 1$ . We will denote this localization by  $A_s$ . If  $A$  is a finite-type domain, so is  $A_s$ . In that case, if  $X$  denotes the variety  $= \text{Spec } A$ ,  $X_s$  will denote the variety  $\text{Spec } A_s$ , and  $X_s$  will be called a *localization* of  $X$  too.

**2.6.1. Proposition.** (i) *With terminology as above, points of the localization  $X_s = \text{Spec } A_s$  correspond bijectively to the open subset of  $X$  of points at which the function defined by  $s$  is nonzero.*

(ii) *When we identify a localization  $X_s$  with a subset of  $X$ , the Zariski topology on  $X_s$  is the induced topology from  $X$ . So  $X_s$  is an open subspace of  $X$ .*

*proof.* (i) Let  $p$  be a point of  $X$ , let  $A \xrightarrow{\pi_p} \mathbb{C}$  be the corresponding homomorphism. If  $s$  isn't zero at  $p$ , i.e.,  $c = s(p) \neq 0$ , then  $\pi_p$  extends uniquely to a homomorphism  $A_s \rightarrow \mathbb{C}$  that sends  $s^{-1}$  to  $c^{-1}$ . This gives us a unique point of  $X_s$  whose image in  $X$  is  $p$ . If  $c = 0$ , then  $\pi_p$  doesn't extend to  $A_s$ .

(ii) Let  $C$  be a closed subset of  $X$ , say the zero set of a set of elements  $a_1, \dots, a_k$  of  $A$ . Then  $C \cap X_s$  is the zero set in  $X_s$  of those same elements, so it is closed in  $X_s$ . Conversely, let  $D$  be a closed subset of  $X_s$ , say the zero set in  $X_s$  of some elements  $b_1 s^{-n}, \dots, b_k s^{-n}$ . We can use the same exponent  $n$  for each element. Since  $s^{-1}$  doesn't vanish on  $X_s$ , the elements  $b_i$  and  $b_i s^{-1}$  have the same zeros on  $X_s$ . If we let  $C$  be the zero set of  $b_1, \dots, b_k$  in  $X$ , we will have  $C \cap X_s = D$ .  $\square$



We usually identify  $X_s$  as an open subset of  $X$ . Then the effect of adjoining the inverse is to throw out the points of  $X$  at which  $s$  vanishes. For example, the spectrum of the Laurent polynomial ring  $\mathbb{C}[t, t^{-1}]$  becomes the complement of the origin in the affine line  $\mathbb{A}^1 = \text{Spec } \mathbb{C}[t]$ .

This illustrates the benefit of working with an affine variety without a fixing an embedding into affine space. If  $X$  is embedded into  $\mathbb{A}^n$ , the localization  $X_s$  wants to be in  $\mathbb{A}^{n+1}$ .

As is true for many open sets, the complement  $X'$  of the origin in the affine plane  $\text{Spec } \mathbb{C}[x_1, x_2]$  isn't a localization. Every polynomial that vanishes at the origin vanishes on an affine curve, which has points different from the origin. Its inverse doesn't define a function on  $X'$ . It may be hard to tell whether or not a given open set is a localization.

Localizations are important for two reasons:

- The relation between an algebra  $A$  and a localization  $A_s$  is easy to understand, and
- The localizations  $X_s$  of an affine variety  $X$  form a basis for the Zariski topology on  $X$ .

A *basis* for the topology on a topological space  $X$  is a family  $\mathcal{B}$  of open sets such that every open set is a union of open sets that are members of  $\mathcal{B}$ .

We verify the second statement marked with a bullet. We must show that if  $U$  is an open subset of  $X = \text{Spec } A$ , then  $U$  can be covered by sets of the form  $X_s$ . The complement of  $X_s$  in  $X$ , the set  $X - X_s$ , is the zero set  $V_X(s)$  of  $s$ . Let  $C$  be the closed complement  $X - U$  of  $U$  in  $X$ . Since  $C$  is closed, it is the set of common zeros of some elements  $s_1, \dots, s_k$  of  $A$ . So  $C$  is the intersection  $\bigcap V_X(s_i)$  of zero sets. Then  $U$  is the union of the sets  $X_{s_i}$ .  $\square$

**2.6.2. Lemma.** *Let  $U$  and  $V$  be affine open subsets of an affine variety  $X$ .*

- (i) *If  $V$  is a localization of  $U$  and  $U$  is a localization of  $X$ , then  $V$  is a localization of  $X$ .*
- (ii) *If  $V \subset U$  and if  $V$  is a localization of  $X$ , then  $V$  is a localization of  $U$ .*
- (iii) *Let  $p$  be a point of  $U \cap V$ . There is an open set  $Z$  containing  $p$  that is a localization of  $U$  and also a localization of  $V$ .*

*proof.* (i) Say that  $X = \text{Spec } A$ ,  $U = X_s = \text{Spec } A_s$  and  $V = U_t = \text{Spec}(A_s)_t$ . Then  $t$  is an element of  $A_s$ , say  $t = s^{-k}r$  with  $r$  in  $A$ . The localizations  $(A_s)_t$ ,  $(A_s)_r$  are equal, and  $(A_s)_r = A_{sr}$ . So  $V = X_{sr}$ .

(ii) Say that  $X = \text{Spec } A$ ,  $U = \text{Spec } B$ , and  $V = \text{Spec } A_t$ , where  $t$  is a nonzero element of  $A$ . The elements of  $B$  are the fractions  $t^{-k}a$  with  $a \in A$ , and  $t$  is an element of  $B$ . So  $B_t = A_t$ .

(iii) Since localizations form a basis for the topology,  $U \cap V$  contains a localization  $X_s$  of  $X$  that contains  $p$ . By (ii),  $X_s$  is a localization of  $U$  and of  $V$ .  $\square$

### (2.6.3) extension and contraction of ideals

Let  $A \subset B$  be the inclusion of a ring  $A$  as a subring of a ring  $B$ . The *extension* of an ideal  $I$  of  $A$  is the ideal  $IB$  of  $B$  generated by  $I$ . Its elements are finite sums  $\sum_i z_i b_i$  with  $z_i$  in  $I$  and  $b_i$  in  $B$ . The *contraction* of an ideal  $J$  of  $B$  is the intersection  $J \cap A$ . It is an ideal of  $A$ .

The next lemma explains what happens when one combines extension and contraction.

**2.6.4. Lemma.** *Let  $A \subset B$  be rings, and let  $I$  and  $J$  be ideals of  $A$  and  $B$ , respectively. Then  $I \subset (IB) \cap A$ , and  $(J \cap A)B \subset J$ .*  $\square$

If  $A_s$  is a localization of  $A$  and  $I$  is an ideal of  $A$ , the elements of the extended ideal  $IA_s$  are fractions of the form  $zs^{-k}$ , with  $z$  in  $I$ . We denote this extended ideal by  $I_s$ .

**2.6.5. Lemma.**

- (i) *Let  $A \subset B$  be rings, let  $I$  be an ideal of  $A$  and let  $J$  be an ideal of  $B$ . Then  $I \subset (IB) \cap A$  and  $(J \cap A)B \subset J$ .*
- (ii) *Let  $A_s$  be a localization of  $A$ , let  $I'$  be an ideal of  $A_s$  and let  $I = I' \cap A$ . Then  $I' = IA_s$ . Every ideal of  $A_s$  is the extension of an ideal of  $A$ .*
- (iii) *Let  $P$  be a prime ideal of  $A$ . If  $s$  is an element of  $A$  that isn't in  $P$ , the extended ideal  $P_s$  is a prime ideal of  $A_s$ . If  $s$  is in  $P$ , the extended ideal is the unit ideal.*  $\square$

## 2.7 Morphisms of Affine Varieties

Morphisms are the allowed maps between varieties. Morphisms between affine varieties are defined below. They correspond to algebra homomorphisms in the opposite direction between their coordinate algebras. Morphisms of projective varieties require more thought. They will be defined in the next chapter.

### (2.7.1) regular functions

The *function field*  $K$  of an affine variety  $X = \text{Spec } A$  is the field of fractions of  $A$ . A *rational function* on  $X$  is a nonzero element of the function field  $K$ .

As we have seen, (2.5.1) elements of the coordinate algebra  $A$  define functions on  $X$ , the rule being  $\alpha(p) = \pi_p(\alpha)$ , where  $\pi_p$  is the homomorphism  $A \rightarrow \mathbb{C}$  that corresponds to  $p$ . A rational function  $f = a/s$  with  $a$  and  $s$  in  $A$  is an element of  $A_s$ , and it defines a function on the open subset  $X_s$ . A rational function  $f$  is *regular* at a point  $p$  of  $X$  if it can be written as a fraction  $a/s$  such that  $s(p) \neq 0$ . A rational function is a *regular function* on  $X$  if it is regular at every point of  $X$ .

**2.7.2. Proposition.** *The regular functions on an affine variety  $X = \text{Spec } A$  are the elements of the coordinate algebra  $A$ .*

*proof.* Let  $f$  be a rational function that is regular on  $X$ . So for every point  $p$  of  $X$ , there is a localization  $X_s = \text{Spec } A_s$  that contains  $p$ , such that  $f$  is an element of  $A_s$ . Because  $X$  is quasicompact, a finite set of these localizations, say  $X_{s_1}, \dots, X_{s_k}$ , will cover  $X$ . Then  $s_1, \dots, s_k$  have no common zeros on  $X$ , so they generate the unit ideal of  $A$  (2.5.7). Since  $f$  is in  $A_{s_i}$ , we can write  $f = s_i^{-n} b_i$ , or  $s_i^n f = b_i$ , with  $b_i$  in  $A$ , and we can use the same exponent  $n$  for each  $i$ . Since the elements  $s_i$  generate the unit ideal of  $A$ , so do the powers  $s_i^n$ . Say that  $\sum s_i^n c_i = 1$ , with  $c_i$  in  $A$ . Then  $f = \sum s_i^n c_i f = \sum c_i b_i$  is an element of  $A$ .  $\square$

### (2.7.3) morphisms

Let  $X = \text{Spec } A$  and  $Y = \text{Spec } B$  be affine varieties, and let  $A \xrightarrow{\varphi} B$  be an algebra homomorphism. A point  $q$  of  $Y$  corresponds to an algebra homomorphism  $B \xrightarrow{\pi_q} \mathbb{C}$ . When we compose  $\pi_q \varphi$ , we obtain a homomorphism  $A \xrightarrow{\pi_q \varphi} \mathbb{C}$ . By the definition, points of  $\text{Spec } A$  correspond to homomorphisms  $A \xrightarrow{\pi_p} \mathbb{C}$ . So there is a unique point  $p$  of  $X = \text{Spec } A$  such that  $\pi_p = \pi_q \varphi$ :

$$(2.7.4) \quad \begin{array}{ccc} A & \xrightarrow{\varphi} & B \\ \pi_p \downarrow & & \downarrow \pi_q \\ \mathbb{C} & \xlongequal{\quad} & \mathbb{C} \end{array}$$

**2.7.5. Definition.** Let  $X = \text{Spec } A$  and  $Y = \text{Spec } B$ . A *morphism*  $Y \xrightarrow{u} X$  is a map defined, as above, by an algebra homomorphism  $A \xrightarrow{\varphi} B$ : If  $q$  is a point of  $Y$ , then  $uq$  is the point  $p$  of  $X$  such that  $\pi_p = \pi_q \varphi$ .

Then if  $\alpha$  is an element of  $A$  and  $\beta = \varphi(\alpha)$ ,

$$(2.7.6) \quad \beta(q) = \pi_q(\beta) = \pi_q(\varphi\alpha) = \pi_p(\alpha) = \alpha(p)$$

The morphism  $Y \xrightarrow{u} X$  is an *isomorphism* if and only if there is an inverse morphism. This will be true if and only if  $A \xrightarrow{\varphi} B$  is an isomorphism of algebras.  $\square$

The relationship between a homomorphism  $A \xrightarrow{\varphi} B$  and the associated morphism  $Y \xrightarrow{u} X$  can be summed up by the next formula. If  $q$  is a point of  $Y$  and  $\alpha$  is an element of  $A$ , then

$$(2.7.7) \quad \alpha[u(q)] = [\varphi\alpha](q)$$

Thus the homomorphism  $\varphi$  is determined by the map  $u$ . But most maps  $Y \rightarrow X$  aren't morphisms.

The description of a morphism can be confusing because the direction of the arrow is reversed. It will become clearer as we expand the discussion.

*Morphisms to the affine line.*

A morphism  $Y \xrightarrow{u} \mathbb{A}^1$  from a variety  $Y = \text{Spec } B$  to the affine line  $\text{Spec } \mathbb{C}[t]$  is defined by an algebra homomorphism  $\mathbb{C}[x] \xrightarrow{\varphi} B$ , which substitutes an element  $\beta$  of  $B$  for  $x$ . The morphism  $u$  that corresponds to  $\varphi$  sends a point  $q$  of  $Y$  to the point of the  $x$ -line at which  $x = \beta(q)$ .

For example, let  $Y$  be the space of  $2 \times 2$  matrices, so that  $B = \mathbb{C}[y_{ij}]$ ,  $1 \leq i, j \leq 2$ . The determinant defines a morphism  $Y \rightarrow \mathbb{A}^1$  that sends a matrix  $\beta = \begin{pmatrix} b_{11} & b_{12} \\ b_{21} & b_{22} \end{pmatrix}$  to its determinant  $b_{11}b_{22} - b_{12}b_{21}$ . The corresponding algebra homomorphism  $\mathbb{C}[x] \xrightarrow{\varphi} \mathbb{C}[y_{ij}]$  substitutes  $y_{11}y_{22} - y_{12}y_{21}$  for  $x$ . It sends a polynomial  $f(x)$  to  $f(y_{11}y_{22} - y_{12}y_{21})$ .

In the other direction, a morphism from  $\mathbb{A}^1$  to a variety  $Y$  is a (complex) polynomial path in  $Y$ . For example, if  $Y$  is the space of matrices, a morphism  $\mathbb{A}^1 \rightarrow Y$  corresponds to a homomorphism  $\mathbb{C}[y_{ij}] \rightarrow \mathbb{C}[x]$ , which substitutes polynomials in  $x$  for the variables  $y_{ij}$ .

*Morphisms to affine space.*

A morphism from an affine variety  $Y = \text{Spec } B$  to affine space  $\mathbb{A}^n$  will be defined by a homomorphism  $\mathbb{C}[x_1, \dots, x_n] \xrightarrow{\Phi} B$  which substitutes elements  $\beta_i$  of  $B$  for  $x_i$ :  $\Phi(f(x)) = f(\beta)$ . The corresponding morphism  $Y \xrightarrow{u} \mathbb{A}^n$  sends a point  $q$  of  $Y$  to the point  $(\beta_1(q), \dots, \beta_n(q))$  of  $\mathbb{A}^n$ .

*Morphisms to affine varieties.*

Let  $X = \text{Spec } A$  and  $Y = \text{Spec } B$  be affine varieties. Say that we have chosen a presentation  $A = \mathbb{C}[x_1, \dots, x_m]/(f_1, \dots, f_k)$  of  $A$ , so that  $X$  becomes the closed subvariety  $V(f)$  of affine space  $\mathbb{A}^m$ . There is no need to choose a presentation of  $B$ . A natural way to define a morphism from a variety  $Y$  to  $X$  is as a morphism  $Y \xrightarrow{u} \mathbb{A}^m$  to affine space, whose image is contained in  $X$ . We check that this agrees with Definition 2.7.5:

As above, a morphism  $Y \xrightarrow{u} \mathbb{A}^m$  corresponds to a homomorphism  $\mathbb{C}[x_1, \dots, x_m] \xrightarrow{\Phi} B$ , and defined by a set  $(\beta_1, \dots, \beta_m)$  of elements of  $B$ . Since  $X$  is the locus of zeros of the polynomials  $f$ , the image of  $Y$  will be contained in  $X$  if and only if  $f_i(\beta_1(q), \dots, \beta_m(q)) = 0$  for every point  $q$  of  $Y$  and every  $i$ , i.e.,  $f_i(\beta)$  is in every maximal ideal of  $B$ , in which case  $f_i(\beta) = 0$  for every  $i$  (2.5.15)(i). Another way to say this is:

The image of  $Y$  is contained in  $X$  if and only if  $\beta = (\beta_1, \dots, \beta_m)$  solves the equations  $f(x) = 0$ .

And, if  $\beta$  is a solution, the map  $\Phi$  defines a map  $A \xrightarrow{\varphi} B$ .

$$\begin{array}{ccc} \mathbb{C}[x] & \xrightarrow{\Phi} & B \\ \downarrow & & \parallel \\ A & \xrightarrow{\varphi} & B \end{array}$$

This is an elementary, but important, principle:

• Homomorphisms from an algebra  $A = \mathbb{C}[x]/(f)$  to an algebra  $B$  correspond to solutions of the equations  $f = 0$  in  $B$ .

### give examples here.  $t$ -line maps to cusp curve, maps to  $SL_2$  and of course, the blowup map###

**2.7.8. Corollary.** Let  $X = \text{Spec } A$  and let  $Y = \text{Spec } B$  be affine varieties. Suppose that  $A$  is presented as the quotient  $\mathbb{C}[x_1, \dots, x_m]/(f_1, \dots, f_k)$  of a polynomial ring. There are bijective correspondences between the following sets:

- algebra homomorphisms  $A \rightarrow B$ , or morphisms  $Y \rightarrow X$ ,
- morphisms  $Y \rightarrow \mathbb{A}^n$  whose images are contained in  $X$ ,
- solutions of the equations  $f_i(x) = 0$  in  $B$ ,

□

The second and third sets refer to an embedding of the variety  $X$  into affine space, but the first one does not. It shows that a morphism depends only on the varieties  $X$  and  $Y$ , not on the embedding of  $X$  into affine space.

The geometry of a morphism will be described more completely in Chapters ?? and ?. We note a few more facts about them here.

**2.7.9. Proposition.** Let  $X \xleftarrow{u} Y$  be the morphism of affine varieties that corresponds to a homomorphism of coordinate algebras  $A \xrightarrow{\varphi} B$ .

(i) Let  $Y \xleftarrow{v} Z$  be another morphism, that corresponds to another homomorphism  $B \xrightarrow{\psi} R$  of finite-type domains. The composition  $Z \xrightarrow{uv} X$  is the morphism that corresponds to the composed homomorphism  $A \xrightarrow{\psi\varphi} R$ .

(ii) Suppose that  $B = A/P$ , where  $P$  is a prime ideal of  $A$ , and that  $\varphi$  is the canonical homomorphism  $A \rightarrow A/P$ . Then  $u$  is the inclusion of the closed subvariety  $Y = V_X(P)$  into  $X$ .

(iii)  $\varphi$  is surjective if and only if  $u$  maps  $Y$  isomorphically to a closed subvariety of  $X$ .  $\square$

It is useful to rephrase the definition of the morphism  $Y \xrightarrow{u} X$  that corresponds to a homomorphism  $A \xrightarrow{\varphi} B$  in terms of maximal ideals. Let  $\mathfrak{m}_q$  be the maximal ideal of  $B$  at a point  $q$  of  $Y$ . The inverse image of  $\mathfrak{m}_q$  in  $A$  is the kernel of the composed homomorphism  $A \xrightarrow{\varphi} B \xrightarrow{\pi_q} \mathbb{C}$ , so it is a maximal ideal of  $A$ :  $\varphi^{-1}\mathfrak{m}_q = \mathfrak{m}_p$  for some point  $p$  of  $X$ . That point is the image of  $q$ :  $p = uq$ .

In the other direction, let  $\mathfrak{m}_p$  be the maximal ideal at a point  $p$  of  $X$ , and let  $J$  be the ideal generated by the image of  $\mathfrak{m}_p$  in  $B$ . This ideal is called the *extension* of  $\mathfrak{m}_p$  to  $B$ . Its elements are finite sums  $\sum \varphi(z_i)b_i$  with  $z_i$  in  $\mathfrak{m}_p$  and  $b_i$  in  $B$ . If  $q$  is a point of  $Y$ , then  $uq = p$  if and only if  $\mathfrak{m}_p = \varphi^{-1}\mathfrak{m}_q$ , and this will be true if and only if  $J$  is contained in  $\mathfrak{m}_q$ .

Recall that, if  $Y \xrightarrow{u} X$  is a map of sets, the *fibre* of  $Y$  over a point  $p$  of  $X$  is the set of points  $q$  of  $Y$  that map to  $p$ .

**2.7.10. Corollary.** Let  $X = \text{Spec } A$  and  $Y = \text{Spec } B$ , and let  $Y \xrightarrow{u} X$  be the morphism corresponding to a homomorphism  $A \xrightarrow{\varphi} B$ , let  $\mathfrak{m}_p$  be the maximal ideal at a point  $p$  of  $X$ , and let  $J = \mathfrak{m}_p B$  be the extended ideal.

(i) The fibre of  $Y$  over  $p$  is the set  $V_Y(J)$  of points  $q$  such that  $J \subset \mathfrak{m}_q$ .

(ii) The fibre of  $Y$  over  $p$  is empty if and only if  $J$  is the unit ideal of  $B$ .  $\square$

**2.7.11. Example.** (blowing up the plane)

Let  $Z$  and  $Y$  be the affine planes with coordinates  $x, z$  and  $x, y$ , respectively. The map  $Z \xrightarrow{\pi} Y$  defined by  $y = xz$ , the morphism that corresponds to the algebra homomorphism  $\mathbb{C}[x, y] \xrightarrow{\varphi} \mathbb{C}[x, z]$  defined by  $\varphi(x) = x$ ,  $\varphi(y) = xz$ .

The morphism  $\pi$  is bijective at points  $(x, y)$  with  $x \neq 0$ . At such a point,  $y = x^{-1}z$ . The fibre of  $Z$  over a point of  $Y$  of the form  $(0, y)$  is empty unless  $y = 0$ , and the fibre over the origin  $(0, 0)$  in  $Y$  is the  $z$ -axis  $\{(0, z)\}$  in the plane  $Z$ . Because the origin in  $Y$  is replaced by a line in  $Z$ , this morphism is called a *blowup* of the affine plane  $Y$ .  $\square$

figure

**2.7.12. Proposition.** A morphism  $Y \xrightarrow{u} X$  of affine varieties is a continuous map in the Zariski topology and also in the classical topology.

*proof.* First, the Zariski topology: Let  $X = \text{Spec } A$  and  $Y = \text{Spec } B$ , so that  $u$  corresponds to an algebra homomorphism  $A \xrightarrow{\varphi} B$ . A closed subset  $C$  of  $X$  will be the zero locus of a set  $\alpha = \{\alpha_1, \dots, \alpha_k\}$  of elements of  $A$ . Let  $\beta_i = \varphi\alpha_i$ . The inverse image  $u^{-1}C$  is the set of points  $q$  such that  $p = uq$  is in  $C$ , i.e., such that  $\alpha_i(uq) = 0$ , and  $\alpha_i(uq) = \beta_i(q)$  (2.7.5). So  $u^{-1}C$  is the zero locus in  $Y$  of the elements  $\beta_i = \varphi(\alpha_i)$  of  $B$ . It is a closed set.

Next, for the classical topology, we use the fact that polynomials are continuous functions. A morphism of affine spaces  $\mathbb{A}_Y^n \xrightarrow{\tilde{u}} \mathbb{A}_X^m$  is defined by an algebra homomorphism  $\mathbb{C}[x_1, \dots, x_m] \xrightarrow{\Phi} \mathbb{C}[y_1, \dots, y_n]$ , and this homomorphism is determined by the polynomials  $h_1(y), \dots, h_m(y)$  that are the images of the variables  $x$ . The morphism  $\tilde{u}$  sends the point  $(y_1, \dots, y_n)$  of  $\mathbb{A}^n$  to the point  $(h_1(y), \dots, h_m(y))$  of  $\mathbb{A}^m$ . It is continuous.

The morphism  $Y \xrightarrow{u} X$  is defined by a homomorphism  $A \xrightarrow{\varphi} B$ . We choose presentations  $A = \mathbb{C}[x]/I$  and  $B = \mathbb{C}[y]/J$ , and we form a diagram of homomorphisms and the associated diagram of morphisms:

$$\begin{array}{ccc} \mathbb{C}[x] & \xrightarrow{\Phi} & \mathbb{C}[y] \\ \alpha \downarrow & & \downarrow \beta \\ A & \xrightarrow{\varphi} & B \end{array} \quad \begin{array}{ccc} \mathbb{A}_x^n & \xleftarrow{\tilde{u}} & \mathbb{A}_y^m \\ \uparrow & & \uparrow \\ X & \xleftarrow{u} & Y \end{array}$$

Here  $\alpha$  and  $\beta$  are the canonical maps of a ring to a quotient ring. The map  $\alpha$  sends  $x_1, \dots, x_n$  to  $\alpha_1, \dots, \alpha_n$ . Then  $\Phi$  is obtained by choosing elements  $h_i$  whose images in  $B$  are the same as the images of  $\alpha_i$ . In the diagram of morphisms,  $\tilde{u}$  is continuous, and the vertical arrows are the embeddings of  $X$  and  $Y$  into their affine spaces. Since the topologies on  $X$  and  $Y$  are induced from their embeddings,  $u$  is continuous.  $\square$

As we see here, every morphism of affine varieties can be obtained by restricting a morphism of affine spaces. However, in the diagram above, the morphism  $\tilde{u}$  isn't unique. It depends on the choice of the polynomials  $h_i$ .

## 2.8 Finite Group Actions

Let  $G$  be a finite group of automorphisms of a finite-type domain  $B$ . An *invariant* element of  $B$  is an element that is sent to itself by every element  $\sigma$  of  $G$ . For example, the product and the sum

$$(2.8.1) \quad \prod_{\sigma \in G} \sigma b \quad , \quad \sum_{\sigma \in G} \sigma b$$

are invariant elements. The invariant elements form a subalgebra of  $B$  that is often denoted by  $B^G$ . Theorem 2.8.5 below asserts that  $B^G$  is a finite-type algebra, and that points of  $\text{Spec } B^G$  correspond bijectively to  $G$ -orbits in  $\text{Spec } B$ .

### 2.8.2. Examples.

(i) The symmetric group  $G = S_n$  operates on the polynomial ring  $R = \mathbb{C}[x_1, \dots, x_n]$  by permuting the variables, and the Symmetric Functions Theorem asserts that the elementary symmetric functions

$$s_1 = \sum_i x_i, \quad s_2 = \sum_{i < j} x_i x_j, \quad \dots, \quad s_n = x_1 x_2 \cdots x_n$$

generate the ring  $R^G$  of invariant polynomials. Moreover,  $s_1, \dots, s_n$  are algebraically independent, so  $R^G$  is the polynomial algebra  $\mathbb{C}[s_1, \dots, s_n]$ . The inclusion of  $R^G$  into  $R$  gives us a morphism from affine  $x$ -space  $\mathbb{A}_x^n$  to affine  $s$ -space  $\mathbb{A}_s^n = \text{Spec } R^G$ . If  $a = (a_1, \dots, a_n)$  is a point of  $\mathbb{A}_s^n$ , the points  $b = (b_1, \dots, b_n)$  of  $\mathbb{A}_x^n$  that map to  $a$  are those such that  $s_i(b) = a_i$ . They are the roots of the polynomial  $x^n - a_1 x^{n-1} + \cdots \pm a_n$ . Since the roots form a  $G$ -orbit, the set of  $G$ -orbits of points of  $\mathbb{A}_x^n$  maps bijectively to  $\mathbb{A}_s^n$ .

(ii) Let  $\zeta = e^{2\pi i/n}$ , let  $\sigma$  be the automorphism of the polynomial ring  $B = \mathbb{C}[y_1, y_2]$  defined by  $\sigma y_1 = \zeta y_1$  and  $\sigma y_2 = \zeta^{-1} y_2$ . Let  $G$  be the cyclic group of order  $n$  generated by  $\sigma$ , and let  $A$  denote the algebra  $B^G$  of invariant elements. A monomial  $m = y_1^i y_2^j$  is invariant if and only if  $n$  divides  $i - j$ , and an invariant polynomial is a linear combination of invariant monomials. You will be able to show that the three monomials

$$(2.8.3) \quad u_1 = y_1^n, \quad u_2 = y_2^n, \quad \text{and} \quad w = y_1 y_2$$

generate  $A$ . We'll use the same symbols  $u_1, u_2, w$  to denote variables in the polynomial ring  $\mathbb{C}[u_1, u_2, w]$ . Let  $J$  be the kernel of the canonical homomorphism  $\mathbb{C}[u_1, u_2, w] \xrightarrow{\tau} A$  that sends  $u_1, u_2, w$  to  $y_1^n, y_2^n, y_1 y_2$ .

**2.8.4. Lemma.** *With notation as above, the kernel  $J$  of  $\tau$  is the principal ideal of  $\mathbb{C}[u_1, u_2, w]$  generated by the polynomial  $f = w^n - u_1 u_2$ .*

*proof.* First,  $f$  is an element of  $J$ . Let  $g(u_1, u_2, w)$  be an element of  $J$ . So  $g(y_1^n, y_2^n, y_1 y_2) = 0$ . We divide  $g$  by  $f$ , considered as a monic polynomial in  $w$ , say  $g = fq + r$ , where the remainder  $r$  has degree  $< n$  in  $w$ . The remainder will be in  $J$  too:  $r(y_1^n, y_2^n, y_1 y_2) = 0$ . We write  $r$  as a polynomial in  $w$ :  $r = r_0(u_1, u_2) + r_1(u_1, u_2)w + \cdots + r_{n-1}(u_1, u_2)w^{n-1}$ . When we substitute  $y_1^n, y_2^n, y_1 y_2$ , the term  $r_i(u_1, u_2)w^i$  becomes  $r_i(y_1^n, y_2^n)(y_1 y_2)^i$ . The degree in  $y_1$  of every monomial that appears here will be congruent to  $i$  modulo  $n$ , and the same is true for  $y_2$ . Since  $r(y_1^n, y_2^n, y_1 y_2) = 0$ , and because the indices  $i$  are distinct,  $r_i(y_1^n, y_2^n)$  will be zero for every  $i$ . And if  $r_i(y_1^n, y_2^n)$  is zero, then  $r_i(u_1, u_2) = 0$ . So  $r = 0$ , which means that  $f$  divides  $g$ .  $\square$

We go back to the operation of the cyclic group on  $B$ . Let  $Y$  denote the affine plane  $\text{Spec } B$ , and let  $X = \text{Spec } A$ . The group  $G$  operates on  $Y$ , and except for the origin, which is a fixed point, the orbit of a point  $(y_1, y_2)$  consists of the  $n$  points  $(\zeta^i y_1, \zeta^{-i} y_2)$ ,  $i = 0, \dots, n-1$ . To show that  $G$ -orbits in  $Y$  correspond bijectively to points of  $X$ , we fix complex numbers  $u_1, u_2, w$  with  $w^n = u_1 u_2$ , and we look for solutions of the equations (2.8.3). When  $u_1 \neq 0$ , the equation  $u_1 = y_1^n$  has  $n$  solutions for  $y_1$ , and then  $y_2$  is determined by the equation  $w = y_1 y_2$ . So the fibre has order  $n$ . Similarly, there are  $n$  points in the fibre if  $u_2 \neq 0$ . If  $u_1 = u_2 = 0$ , then  $y_1 = y_2 = w = 0$ . In all cases, the fibres are the  $G$ -orbits.  $\square$

**2.8.5. Theorem.** *Let  $G$  be a finite group of automorphisms of a finite-type domain  $B$ , and let  $A$  denote the algebra  $B^G$  of invariant elements. Let  $Y = \text{Spec } B$  and  $X = \text{Spec } A$ .*

- (i)  *$A$  is a finite-type domain and  $B$  is a finite  $A$ -module.*
- (ii)  *$G$  operates by automorphisms on  $Y$ .*
- (iii) *The morphism  $Y \rightarrow X$  defined by the inclusion  $A \subset B$  is surjective, and its fibres are the  $G$ -orbits of points of  $Y$ .*

When a group  $G$  operates on a set  $Y$ , one often denotes the set of  $G$ -orbits of  $Y$  by  $Y/G$ . With that notation, the theorem asserts that there is a bijective map  $Y/G \rightarrow X$ .

*proof of 2.8.5 (i):* The invariant algebra  $A = B^G$  is a finite-type algebra, and  $B$  is a finite  $A$ -module.

This is an interesting indirect proof. To show that  $A$  is a finite-type algebra, one constructs a finite-type subalgebra  $R$  of  $A$  such that  $B$  is a finite  $R$ -module.

Let  $\{z_1, \dots, z_k\}$  be the  $G$ -orbit of an element  $z_1$  of  $B$ . The orbit is the set of roots of the polynomial

$$f(t) = (t - z_1) \cdots (t - z_k) = t^k - s_1 t^{k-1} + \cdots \pm s_k$$

whose coefficients are the elementary symmetric functions in  $\{z_1, \dots, z_k\}$ . Let  $R_1$  denote the algebra generated by those symmetric functions. Because the symmetric functions are invariant,  $R_1 \subset A$ . Using the equation  $f(z_1) = 0$ , we can write any power of  $z_1$  as a polynomial in  $z_1$  of degree less than  $k$ , with coefficients in  $R_1$ .

We choose a finite set of generators  $\{y_1, \dots, y_r\}$  for the algebra  $B$ . If the order of the orbit of  $y_j$  is  $k_j$ , then  $y_j$  will be the root of a monic polynomial  $f_j$  of degree  $k_j$  with coefficients in  $A$ . Let  $R$  denote the finite-type algebra generated by all of the coefficients of all of the polynomials  $f_1, \dots, f_r$ . We can write any power of  $y_j$  as a polynomial in  $y_j$  with coefficients in  $R$ , and of degree less than  $k_j$ . Using such expressions, we can write every monomial in  $y_1, \dots, y_r$  as a polynomial  $y_1, \dots, y_r$  with coefficients in  $R$ , whose degree in each variable  $y_j$  is less than  $k_j$ . Since  $y_1, \dots, y_r$  generate  $B$ , we can write every element of  $B$  as such a polynomial. Then the finite set of monomials  $y_1^{e_1} \cdots y_r^{e_r}$  with  $e_j < k_j$  spans  $B$  as an  $R$ -module. Therefore  $B$  is a finite  $R$ -module.

Since  $R$  is a finite-type algebra, it is noetherian. The algebra  $A$  of invariants is a subalgebra of  $B$  that contains  $R$ . So when regarded as an  $R$ -module,  $A$  is a submodule of the finite  $R$ -module  $B$ . Since  $R$  is noetherian,  $A$  is also a finite  $R$ -module. When we put a finite set of algebra generators for  $R$  together with a finite set of  $R$ -module generators for  $A$ , we obtain a finite set of algebra generators for  $A$ . So  $A$  is a finite-type algebra. And, since  $B$  is a finite  $R$ -module, it is also a finite module over the larger ring  $A$ .

*proof of 2.8.5(ii):* The group  $G$  operates on  $Y$ .

A group element  $\sigma$  is a homomorphism  $B \xrightarrow{\sigma} B$ , which defines a morphism  $Y \xleftarrow{u_\sigma} Y$ , as in Definition 2.7.5. Since  $\sigma$  is an invertible homomorphism, i.e., an automorphism,  $u_\sigma$  is also an automorphism. Thus  $G$  operates on  $Y$ . However, there is a point that should be mentioned.

Let's write the operation of  $G$  on  $B$  on the left as usual, so that a group element  $\sigma$  maps an element  $\beta$  of  $B$  to  $\sigma\beta$ . Then if  $\sigma$  and  $\tau$  are two group elements, the product  $\sigma\tau$  acts as *first do  $\tau$ , then  $\sigma$* :  $(\sigma\tau)\beta = \sigma(\tau\beta)$ .

$$(2.8.6) \quad B \xrightarrow{\tau} B \xrightarrow{\sigma} B$$

We substitute  $u = u_\sigma$  into Definition 2.7.5: If  $q$  is a point of  $Y$ , the morphism  $Y \xleftarrow{u_\sigma} Y$  sends  $q$  to the point  $p$  such that  $\pi_p = \pi_q \sigma$ . It seems permissible to drop the symbol  $u$ , and to write the morphism simply as  $Y \xleftarrow{\sigma} Y$ . But since arrows are reversed when going from homomorphisms of algebras to morphisms of their spectra, the maps displayed in (2.8.6), give us morphisms

$$(2.8.7) \quad Y \xleftarrow{\tau} Y \xleftarrow{\sigma} Y$$

On  $Y = \text{Spec } B$ , the product  $\sigma\tau$  acts as *first do  $\sigma$ , then  $\tau$* .

To get around this problem, we can put the symbol  $\sigma$  on the right when it operates on  $Y$ , so that  $\sigma$  sends a point  $q$  to  $q\sigma$ . Then if  $q$  is a point of  $Y$ , we will have  $q(\sigma\tau) = (q\sigma)\tau$ , as required of an operation.

- If  $G$  operates on the left on  $B$ , it operates on the right on  $\text{Spec } B$ .

This is important only when one wants to compose morphisms. In Definition 2.7.5, we followed custom and wrote the morphism  $u$  that corresponds to an algebra homomorphism  $\varphi$  on the left. We will continue to write morphisms on the left when possible, but not here.

Let  $\beta$  be an element of  $B$  and let  $q$  be a point of  $Y$ . The value  $[\sigma\beta](q)$  of the function  $\sigma\beta$  at  $q$  is the same as the value of  $\beta$  at  $q\sigma$ :  $[\sigma\beta](q) \stackrel{\text{defn}}{=} \pi_q(\sigma\beta) = \pi_{q\sigma}(\beta) \stackrel{\text{defn}}{=} \beta(q\sigma)$  (2.7.6):

$$(2.8.8) \quad [\sigma\beta](q) = \beta(q\sigma) \quad \square$$

*proof of 2.8.5 (iii): The fibres of the morphism  $Y \rightarrow X$  are the  $G$ -orbits in  $Y$ .*

We go back to the subalgebra  $A = B^G$ . For  $\sigma$  in  $G$ , we have a diagram of algebra homomorphisms and the corresponding diagram of morphisms

$$(2.8.9) \quad \begin{array}{ccc} B & \xrightarrow{\sigma} & B \\ \uparrow & & \uparrow \\ A & \xlongequal{\quad} & A \end{array} \quad \begin{array}{ccc} Y & \xleftarrow{\sigma} & Y \\ \downarrow & & \downarrow \\ X & \xlongequal{\quad} & X \end{array}$$

The diagram of morphisms shows that the elements of  $Y$  forming a  $G$ -orbit have the same image in  $X$ , and therefore that the set of  $G$ -orbits in  $Y$ , which we denote by  $Y/G$ , maps to  $X$ . We show that the map  $Y/G \rightarrow X$  is bijective.

**2.8.10. Lemma. (i)** Let  $p_1, \dots, p_k$  be distinct points of affine space  $\mathbb{A}^n$ , and let  $c_1, \dots, c_k$  be complex numbers. There is a polynomial  $f(x_1, \dots, x_n)$  such that  $f(p_i) = c_i$  for  $i = 1, \dots, k$ .

**(ii)** Let  $B$  be a finite-type algebra, let  $q_1, \dots, q_k$  be points of  $\text{Spec } B$ , and let  $c_1, \dots, c_k$  be complex numbers. There is an element  $\beta$  in  $B$  such that  $\beta(q_i) = c_i$  for  $i = 1, \dots, k$ .  $\square$

*injectivity of the map  $Y/G \rightarrow X$ :*

Let  $O_1$  and  $O_2$  be distinct  $G$ -orbits. Lemma 2.8.10 tells us that there is an element  $\beta$  in  $B$  whose value is 0 at every point of  $O_1$ , is 1 at every point of  $O_2$ . Since  $G$  permutes the orbits,  $\sigma\beta$  will also be 0 at points of  $O_1$  and 1 at points of  $O_2$ . Then the product  $\gamma = \prod_{\sigma} \sigma\beta$  will be 0 at points of  $O_1$  and 1 at points of  $O_2$ , and  $\gamma$  is invariant. If  $p_i$  denotes the image in  $X$  of the orbit  $O_i$ , the maximal ideal  $\mathfrak{m}_{p_i}$  of  $A$  is the intersection  $A \cap \mathfrak{m}_q$ , where  $q$  is any point in  $O_i$ . Therefore  $\gamma$  is in the maximal ideal  $\mathfrak{m}_{p_1}$ , but not in  $\mathfrak{m}_{p_2}$ . The images of the two orbits are distinct.

*surjectivity of the map  $Y/G \rightarrow X$ :*

It suffices to show that the map  $Y \rightarrow X$  is surjective.

**2.8.11. Lemma.** If  $I$  is an ideal of the invariant algebra  $A$ , and if the extended ideal  $IB$  is the unit ideal of  $B$ , then  $I$  is the unit ideal of  $A$ .

As before, the extended ideal  $IB$  is the ideal of  $B$  generated by  $I$ .

Let's assume the lemma for the moment, and use it to prove surjectivity of the map  $Y \rightarrow X$ . Let  $p$  be a point of  $X$ . The lemma tells us that the extended ideal  $\mathfrak{m}_p B$  isn't the unit ideal. So it is contained in a maximal ideal  $\mathfrak{m}_q$  of  $B$ , where  $q$  is a point of  $Y$ . Then  $\mathfrak{m}_p \subset (\mathfrak{m}_p B) \cap A \subset \mathfrak{m}_q \cap A$ .

The contraction  $\mathfrak{m}_q \cap A$  is an ideal of  $A$ , and it isn't the unit ideal because 1 isn't in  $\mathfrak{m}_q$ . Since  $\mathfrak{m}_p$  is a maximal ideal,  $\mathfrak{m}_p = \mathfrak{m}_q \cap A$ . This means that  $q$  maps to  $p$  in  $X$ .  $\square$

*proof of the lemma.* If  $IB = B$ , there will be an equation  $\sum_i z_i b_i = 1$ , with  $z_i$  in  $I$  and  $b_i$  in  $B$ . The sums  $\alpha_i = \sum_{\sigma} \sigma b_i$  are invariant, so they are elements of  $A$ , and the elements  $z_i$  are invariant. Therefore  $\sum_{\sigma} \sigma(z_i b_i) = z_i \sum_{\sigma} \sigma b_i = z_i \alpha_i$  is in  $I$ . Then

$$\sum_{\sigma} 1 = \sum_{\sigma} \sigma(1) = \sum_{\sigma, i} \sigma(z_i b_i) = \sum_i z_i \alpha_i$$

The right side is in  $I$ , and the left side is the order of the group which, because  $A$  contains the complex numbers, is an invertible element of  $A$ . So  $I$  is the unit ideal.  $\square$



## Chapter 3 PROJECTIVE ALGEBRAIC GEOMETRY

- 3.1 Projective Varieties
- 3.2 Homogeneous Ideals
- 3.3 Product Varieties
- 3.4 Morphisms and Isomorphisms
- 3.5 Affine Varieties
- 3.6 Lines in Projective Three-Space

### 3.1 Projective Varieties

The projective space  $\mathbb{P}^n$  of dimension  $n$  was defined in Chapter 1. Its points are equivalence classes of nonzero vectors  $(x_0, \dots, x_n)$ , the equivalence relation being that, for any nonzero complex number  $\lambda$ ,

$$(3.1.1) \quad (x_0, \dots, x_n) \sim (\lambda x_0, \dots, \lambda x_n).$$

A subset of  $\mathbb{P}^n$  is *Zariski closed* if it is the set of common zeros of a family of *homogeneous* polynomials  $f_1, \dots, f_k$  in the coordinate variables  $x_0, \dots, x_n$ , or if it is the set of zeros of the ideal  $\mathcal{I}$  generated by such a family. Homogeneity is required because the vectors  $(x)$  and  $(\lambda x)$  represent the same point of  $\mathbb{P}^n$ . As explained in (1.3.1),  $f(\lambda x) = 0$  for all  $\lambda$  if and only if  $f$  is homogeneous. We usually refer to the Zariski closed sets simply as closed sets. They are the closed sets in the *Zariski topology* on  $\mathbb{P}^n$ .

Because the polynomial ring  $\mathbb{C}[x_0, \dots, x_n]$  is noetherian,  $\mathbb{P}^n$  is a noetherian space: Every strictly increasing family of ideals of  $\mathbb{C}[x]$  is finite, and every strictly decreasing family of closed subsets of  $\mathbb{P}^n$  is finite. Therefore every closed subset of  $\mathbb{P}^n$  is a finite union of irreducible closed sets (2.2.13). The irreducible closed sets are the *projective varieties*, the closed subvarieties of  $\mathbb{P}^n$ ,

Thus a projective variety  $X$  is an irreducible closed subset of some projective space. We will also want to know when two projective varieties are isomorphic. This will be explained in Section 3.4, where morphisms are defined.

The Zariski topology on a projective variety  $X$  is induced from the topology on the projective space that contains it. Since a projective variety  $X$  is closed in  $\mathbb{P}^n$ , a subset of  $X$  is closed in  $X$  if it is closed in  $\mathbb{P}^n$ .

**3.1.2. Lemma.** *The one-point sets in projective space are closed.*

*proof.* This simple proof illustrates a general method. Let  $p$  be the point  $(a_0, \dots, a_n)$ . The first guess might be that the one-point set  $\{p\}$  is defined by the equations  $x_i = a_i$ , but the polynomials  $x_i - a_i$  aren't homogeneous in  $x$ . This is reflected in the fact that, for any  $\lambda \neq 0$ , the vector  $(\lambda a_0, \dots, \lambda a_n)$  represents the same point, though it won't satisfy those equations. The equations that define the set  $\{p\}$  are

$$(3.1.3) \quad a_i x_j = a_j x_i,$$

for  $i, j = 0, \dots, n$ , which show that the ratios  $a_i/a_j$  and  $x_i/x_j$  are equal. □

**3.1.4. Lemma.** *The proper closed subsets of the projective line are the nonempty finite subsets, and the proper closed subsets of the projective plane are finite unions of points and curves.* □

Though affine varieties are important, most of algebraic geometry concerns projective varieties. It isn't completely clear why this is so, but one property of projective space gives a hint of its importance: With its classical topology, projective space is *compact*.

A topological space is compact if it has these properties:

*Hausdorff property*: Distinct points  $p, q$  of  $X$  have disjoint open neighborhoods, and

*quasicompactness*: If  $X$  is covered by a family  $\{U^i\}$  of open sets, then a finite subfamily covers  $X$ .

By the way, when we say that the sets  $\{U^i\}$  cover a topological space  $X$ , we mean that  $X$  is the union  $\bigcup U^i$ . We don't allow  $U^i$  to contain elements that aren't in  $X$ , though that would be a customary English usage.

In the classical topology, affine space  $\mathbb{A}^n$  isn't quasicompact, and therefore it isn't compact. The *Heine-Borel Theorem* asserts that a subset of  $\mathbb{A}^n$  is compact in the classical topology if and only if it is closed and bounded.

We'll show that  $\mathbb{P}^n$  is compact, assuming that the Hausdorff property has been verified. The  $2n+1$ -dimensional sphere  $\mathbb{S}$  of unit length vectors in  $\mathbb{A}^{n+1}$  is a bounded set, and because it is the zero locus of the equation  $\bar{x}_0x_0 + \cdots + \bar{x}_nx_n = 1$ , it is closed. The Heine-Borel Theorem tells us that  $\mathbb{S}$  is compact. The map  $\mathbb{S} \rightarrow \mathbb{P}^n$  that sends a vector  $(x_0, \dots, x_n)$  to the point of projective space with that coordinate vector is continuous and surjective. The next lemma of topology shows that  $\mathbb{P}^n$  is compact.

**3.1.5. Lemma.** *Let  $Y \xrightarrow{f} X$  be a continuous map. Suppose that  $Y$  is compact and that  $X$  is a Hausdorff space. Then the image  $Z = f(Y)$  is a closed and compact subset of  $X$ .*  $\square$

The rest of this section contains a few examples of projective varieties.

### (3.1.6) linear subspaces

If  $W$  is a subspace of dimension  $r+1$  of the vector space  $V$ , the points of  $\mathbb{P}^n$  that are represented by the nonzero vectors in  $W$  form a *linear subspace*  $L$  of  $\mathbb{P}^n$ , of dimension  $r$ . If  $(w_0, \dots, w_r)$  is a basis of  $W$ , the linear subspace  $L$  corresponds bijectively to a projective space of dimension  $r$ , by

$$c_0w_0 + \cdots + c_rw_r \longleftrightarrow (c_0, \dots, c_r)$$

For example, the set of points  $(x_0, \dots, x_r, 0, \dots, 0)$  is a linear subspace.  $\square$

### (3.1.7) a quadric surface

A *quadric* in  $\mathbb{P}^3$  is the locus of zeros of an irreducible homogeneous quadratic equation in four variables.

We describe a bijective map from the product  $\mathbb{P}^1 \times \mathbb{P}^1$  of projective lines to a quadric. Let coordinates in the two copies of  $\mathbb{P}^1$  be  $(x_0, x_1)$  and  $(y_0, y_1)$ , respectively, and let the four coordinates in  $\mathbb{P}^3$  be  $w_{ij}$ , with  $0 \leq i, j \leq 1$ . The map is defined by  $w_{ij} = x_iy_j$ . Its image is the quadric  $Q$  whose equation is

$$(3.1.8) \quad w_{00}w_{11} = w_{01}w_{10}$$

Let's check that the map  $\mathbb{P}^1 \times \mathbb{P}^1 \rightarrow Q$  is bijective. If  $w$  is a point of  $Q$ , one of the coordinates, say  $w_{00}$ , will be nonzero. Then if  $(x, y)$  is a point of  $\mathbb{P}^1 \times \mathbb{P}^1$  whose image is  $w$ , so that  $w_{ij} = x_iy_j$ , the coordinates  $x_0$  and  $y_0$  must be nonzero. When we normalize  $w_{00}, x_0$ , and  $y_0$  to 1, there is a unique solution for  $x$  and  $y$  such that  $w_{ij} = x_iy_j$ , namely  $x_1 = w_{10}$  and  $y_1 = w_{01}$ .

The quadric with the equation (3.1.8) contains two families of *lines* (one dimensional linear subspaces), the images of the subsets  $x \times \mathbb{P}^1$  and  $\mathbb{P}^1 \times y$  of  $\mathbb{P}^1 \times \mathbb{P}^1$ .

**Note.** Equation (3.1.8) can be diagonalized by the substitution  $w_{00} = s+t$ ,  $w_{11} = s-t$ ,  $w_{01} = u+v$ ,  $w_{10} = u-v$ . This substitution changes the equation (3.1.8) to  $s^2 - t^2 = u^2 - v^2$ . When we look at the affine open set  $\{u=1\}$ , the equation becomes  $s^2 + v^2 - t^2 = 1$ . The real locus of this equation is a one-sheeted hyperboloid in  $\mathbb{R}^3$ , and the two families of complex lines in the quadric correspond to the familiar rulings of this hyperboloid by real lines.

### (3.1.9) hypersurfaces

A *hypersurface* in projective space  $\mathbb{P}^n$  is the locus of zeros of an irreducible homogeneous polynomial  $f(x_0, \dots, x_n)$ . Plane projective curves and quadric surfaces are hypersurfaces.

### (3.1.10) the Segre embedding of a product

The product  $\mathbb{P}_x^m \times \mathbb{P}_y^n$  of projective spaces can be embedded by its *Segre embedding* into a projective space  $\mathbb{P}_w^N$  that has coordinates  $w_{ij}$ , with  $i = 0, \dots, m$  and  $j = 0, \dots, n$ . So  $N = (m+1)(n+1) - 1$ . The Segre embedding is defined by

$$(3.1.11) \quad w_{ij} = x_i y_j.$$

We call the coordinates  $w_{ij}$  the *Segre variables*.

The map from  $\mathbb{P}^1 \times \mathbb{P}^1$  to  $\mathbb{P}^3$  that was described in (3.1.7) is the simplest case of a Segre embedding.

**3.1.12. Proposition.** *The Segre embedding maps the product  $\mathbb{P}^m \times \mathbb{P}^n$  bijectively to the locus  $S$  of the Segre equations*

$$(3.1.13) \quad w_{ij}w_{k\ell} - w_{i\ell}w_{kj} = 0.$$

*proof.* When one substitutes (3.1.11) into the Segre equations, one obtains equations in  $\{x_i, y_j\}$  that are true. So the image of the Segre embedding is contained in  $S$ .

Say that we have a point  $p$  of  $S$  that is the image of  $(x, y)$ . Some coordinate of  $p$ , say  $w_{00}$ , will be nonzero, and then  $x_0$  and  $y_0$  are also nonzero. We normalize  $w_{00}, x_0$ , and  $y_0$  to 1. Then  $w_{ij} = w_{i0}w_{0j}$  for all  $i, j$ . The unique solution of the Segre equations is  $x_i = w_{i0}$  and  $y_j = w_{0j}$ .  $\square$

The Segre embedding is important because it makes the product of projective spaces into a projective variety – the closed subvariety of  $\mathbb{P}^N$  defined by the Segre equations. However, to show that the product is a variety, we need to show that the locus  $S$  of the Segre equations is irreducible. This is less obvious than one might expect, so we defer the discussion to Section 3.3 (see Proposition 3.3.1).

### (3.1.14) the Veronese embedding of projective space

Let the coordinates in  $\mathbb{P}^n$  be  $x_i$ , and let those in  $\mathbb{P}^N$  be  $v_{ij}$ , with  $0 \leq i \leq j \leq n$ . Then  $N = \binom{n+2}{2} - 1$ . The Veronese embedding is the map  $\mathbb{P}^n \xrightarrow{f} \mathbb{P}^N$  defined by  $v_{ij} = x_i x_j$ . The Veronese embedding resembles the Segre embedding, but in the Segre embedding, there are distinct sets of coordinates  $x$  and  $y$ , and there is no requirement that  $i \leq j$ .

The proof of the next proposition is similar to the proof of (3.1.12), once one has untangled the inequalities.

**3.1.15. Proposition.** *The Veronese embedding  $f$  maps  $\mathbb{P}^n$  bijectively to the locus  $X$  in  $\mathbb{P}^N$  of the equations*

$$v_{ij}v_{k\ell} = v_{i\ell}v_{kj} \quad \text{for } 0 \leq i \leq k \leq j \leq \ell \leq n \quad \square$$

For example, the Veronese embedding maps  $\mathbb{P}^1$  bijectively to the conic  $v_{00}v_{11} = v_{01}^2$  in  $\mathbb{P}^2$ .

### (3.1.16) the twisted cubic

There are higher order Veronese embeddings, defined in an analogous way by the monomials of some degree  $d > 2$ . The first example is the embedding of  $\mathbb{P}^1$  by the cubic monomials in two variables, which maps  $\mathbb{P}_x^1$  to  $\mathbb{P}_v^3$ . Let the coordinates in  $\mathbb{P}^3$  be  $v_0, \dots, v_3$ . The cubic Veronese embedding is defined by

$$v_0 = x_0^3, \quad v_1 = x_0^2 x_1, \quad v_2 = x_0 x_1^2, \quad v_3 = x_1^3$$

Its image is a *twisted cubic* in  $\mathbb{P}^3$ , the locus  $(v_0, v_1, v_2, v_3) = (x_0^3, x_0^2x_1, x_0x_1^2, x_1^3)$ , which is the set of common zeros of the three polynomials

$$(3.1.17) \quad v_0v_2 - v_1^2, \quad v_1v_2 - v_0v_3, \quad v_1v_3 - v_2^2$$

These polynomials are the  $2 \times 2$  minors of the  $2 \times 3$  matrix

$$(3.1.18) \quad \begin{pmatrix} v_0 & v_1 & v_2 \\ v_1 & v_2 & v_3 \end{pmatrix}$$

A  $2 \times 3$  matrix has rank  $\leq 1$  if and only if its  $2 \times 2$  minors are zero. So a point  $(v_0, v_1, v_2, v_3)$  lies on the twisted cubic if (3.1.18) has rank one. This means that the vectors  $(v_0, v_1, v_2)$  and  $(v_1, v_2, v_3)$ , if both are nonzero, represent the same point of  $\mathbb{P}^2$ . Setting  $x_0 = v_0 = 1$  and  $x_1 = v_1 = t$ , the twisted cubic becomes the locus of points  $(1, t, t^2, t^3)$ . There is also one point on the twisted cubic at which  $v_0 = 0$ , the point  $(0, 0, 0, 1)$ .  $\square$

## 3.2 Homogeneous Ideals

We denote the polynomial algebra  $\mathbb{C}[x_0, \dots, x_n]$  by  $R$  here.

**3.2.1. Lemma.** *Let  $\mathcal{I}$  be an ideal of  $R$ . The following conditions are equivalent.*

(i)  *$\mathcal{I}$  can be generated by homogeneous polynomials.*

(ii) *A polynomial is in  $\mathcal{I}$  if and only if its homogeneous parts are in  $\mathcal{I}$ .*  $\square$

An ideal  $\mathcal{I}$  of  $R$  that satisfies these conditions is a *homogeneous ideal*.

**3.2.2. Lemma.** *The radical (2.2.2) of a homogeneous ideal is homogeneous.*

*proof.* Let  $\mathcal{I}$  be a homogeneous ideal, and let  $f$  be an element of its radical  $\text{rad } \mathcal{I}$ . So  $f^r$  is in  $\mathcal{I}$  for some  $r$ . When  $f$  is written as a sum  $f_0 + \dots + f_d$  of its homogeneous parts, the highest degree part of  $f^r$  is  $(f_d)^r$ . Since  $\mathcal{I}$  is homogeneous,  $(f_d)^r$  is in  $\mathcal{I}$  and  $f_d$  is in  $\text{rad } \mathcal{I}$ . Then  $f_0 + \dots + f_{d-1}$  is also in  $\text{rad } \mathcal{I}$ . By induction on  $d$ , all of the homogeneous parts  $f_0, \dots, f_d$  are in  $\text{rad } \mathcal{I}$ .  $\square$

If  $f$  is a set of homogeneous polynomials, the set of its zeros in  $\mathbb{P}^n$  may be denoted by  $V(f)$  or  $V_{\mathbb{P}^n}(f)$ , and the set of zeros of a homogeneous ideal  $\mathcal{I}$  by  $V(\mathcal{I})$  or  $V_{\mathbb{P}^n}(\mathcal{I})$ . We use the same notation as for closed subsets of affine space.

The complement of the origin in the affine space  $\mathbb{A}^{n+1}$  is mapped to the projective space  $\mathbb{P}^n$  by sending a vector  $(x_0, \dots, x_n)$  to the point of  $\mathbb{P}^n$  it defines. This map can be useful when one studies projective space.

A homogeneous ideal  $\mathcal{I}$  has a zero locus in projective space  $\mathbb{P}^n$  and also a zero locus in the affine space  $\mathbb{A}^{n+1}$ . We can't use the  $V(\mathcal{I})$  notation for both of them here, so let's denote these two loci by  $V$  and  $W$ , respectively. Unless  $\mathcal{I}$  is the unit ideal, the origin  $x = 0$  will be a point of  $W$ , and the complement of the origin will map surjectively to  $V$ . If a point  $x$  other than the origin is in  $W$ , then every point of the one-dimensional subspace of  $\mathbb{A}^{n+1}$  spanned by  $x$  is in  $W$ , because a homogeneous polynomial  $f$  vanishes at  $x$  if and only if it vanishes at  $\lambda x$ . An affine variety that is the union of such lines through the origin is called an *affine cone*. If the locus  $W$  contains a point  $x$  other than the origin, it is an affine cone.

The familiar locus  $x_0^2 + x_1^2 - x_2^2 = 0$  is a cone in  $\mathbb{A}^3$ . The zero locus of the polynomial  $x_0^3 + x_1^3 - x_2^3$  is also called a cone.

**Note.** The real locus  $x_0^2 + x_1^2 - x_2^2 = 0$  in  $\mathbb{R}^3$  decomposes into two parts when the origin is removed. Because of this, it is sometimes called a "double cone". However, the complex locus doesn't decompose.

### (3.2.3) the irrelevant ideal

In the polynomial algebra  $R = \mathbb{C}[x_0, \dots, x_n]$ , the maximal ideal  $\mathcal{M} = (x_0, \dots, x_n)$  generated by the variables is called the *irrelevant ideal* because its zero locus in projective space is empty.

**3.2.4. Proposition.** *The zero locus in  $\mathbb{P}^n$  of a homogeneous ideal  $\mathcal{I}$  of  $R$  is empty if and only if  $\mathcal{I}$  contains a power of the irrelevant ideal.*

Another way to say this is that the zero locus  $V(\mathcal{I})$  in projective space of a homogeneous ideal  $\mathcal{I}$  is empty if and only if either  $\mathcal{I}$  is the unit ideal  $R$ , or its radical is the irrelevant ideal.

*proof of Proposition 3.2.4.* Let  $Z$  be the zero locus of  $\mathcal{I}$  in  $\mathbb{P}^n$ . If  $\mathcal{I}$  contains a power of  $\mathcal{M}$ , it contains a power of each variable. Powers of the variables have no common zeros in projective space, so  $Z$  is empty.

Suppose that  $Z$  is empty, and let  $W$  be the locus of zeros of  $\mathcal{I}$  in the affine space  $\mathbb{A}^{n+1}$  with the same coordinates  $x_0, \dots, x_n$ , as above. Since the complement of the origin in  $W$  maps to the empty locus  $Z$ , it is empty. The origin is the only point that might be in  $W$ . If  $W$  is the one point space consisting of the origin, then  $\text{rad } \mathcal{I}$  is the irrelevant ideal  $\mathcal{M}$ . If  $W$  is empty,  $\mathcal{I}$  is the unit ideal.  $\square$

**3.2.5. Lemma.** *Let  $\mathcal{P}$  be a homogeneous ideal in the polynomial algebra  $R$ , not the unit ideal. The following conditions are equivalent:*

- (i)  $\mathcal{P}$  is a prime ideal.
- (ii) If  $f$  and  $g$  are homogeneous polynomials, and if  $fg \in \mathcal{P}$ , then  $f \in \mathcal{P}$  or  $g \in \mathcal{P}$ .
- (iii) If  $\mathcal{A}$  and  $\mathcal{B}$  are homogeneous ideals, and if  $\mathcal{A}\mathcal{B} \subset \mathcal{P}$ , then  $\mathcal{A} \subset \mathcal{P}$  or  $\mathcal{B} \subset \mathcal{P}$ .

In other words, a homogeneous ideal is a prime ideal if the usual conditions for a prime ideal are satisfied when the polynomials or ideals are homogeneous.

*proof of the lemma.* When the word homogeneous is omitted, (ii) and (iii) become the definition of a prime ideal. So (i) implies (ii) and (iii). The fact that (iii)  $\Rightarrow$  (ii) is proved by considering the principal ideals generated by  $f$  and  $g$ .

(ii)  $\Rightarrow$  (i) Suppose that a homogeneous ideal  $\mathcal{P}$  satisfies the condition (ii), and that the product  $fg$  of two polynomials, not necessarily homogeneous, is in  $\mathcal{P}$ . If  $f$  has degree  $d$  and  $g$  has degree  $e$ , the highest degree part of  $fg$  is the product  $f_d g_e$  of the homogeneous parts of  $f$  and  $g$  of maximal degree. Since  $\mathcal{P}$  is a homogeneous ideal, it contains  $f_d g_e$ . Therefore one of the factors, say  $f_d$ , is in  $\mathcal{P}$ . Let  $h = f - f_d$ . Then  $hg$  is in  $\mathcal{P}$ , and it has lower degree than  $fg$ . By induction on the degree of  $fg$ ,  $h$  or  $g$  is in  $\mathcal{P}$ , and if  $h$  is in  $\mathcal{P}$ , so is  $f$ .  $\square$

**3.2.6. Proposition.** *Let  $Y$  be the zero locus in  $\mathbb{P}^n$  of a homogeneous radical ideal  $\mathcal{I}$ , that isn't the irrelevant ideal. Then  $Y$  is a projective variety (an irreducible closed subset of  $\mathbb{P}^n$ ) if and only if  $\mathcal{I}$  is a prime ideal. Thus a subset  $Y$  of  $\mathbb{P}^n$  is a projective variety if and only if it is the zero locus of a homogeneous prime ideal that isn't the irrelevant ideal.*

*proof.* The locus  $W$  of zeros of  $\mathcal{I}$  in the affine space  $\mathbb{A}^{n+1}$  is irreducible if and only if  $Y$  is irreducible. This is easy to see. Proposition 2.2.16 tells us that  $W$  is irreducible if and only if the radical ideal  $\mathcal{I}$  is a prime ideal.  $\square$

### 3.2.7. Strong Nullstellensatz, projective version.

- (i) Let  $g$  be a nonconstant homogeneous polynomial in  $x_0, \dots, x_n$ , and let  $\mathcal{I}$  be a homogeneous ideal of  $\mathbb{C}[x]$ . If  $g$  vanishes at every point of the zero locus  $V(\mathcal{I})$  in  $\mathbb{P}^n$ , then  $\mathcal{I}$  contains a power of  $g$ .
- (ii) Let  $f$  and  $g$  be homogeneous polynomials. If  $f$  is irreducible and if  $V(f) \subset V(g)$ , then  $f$  divides  $g$ .
- (iii) Let  $\mathcal{I}$  and  $\mathcal{J}$  be homogeneous ideals, and suppose that  $\text{rad } \mathcal{I}$  isn't the irrelevant ideal or the unit ideal. Then  $V(\mathcal{I}) = V(\mathcal{J})$  if and only if  $\text{rad } \mathcal{I} = \text{rad } \mathcal{J}$ .

*proof.* (i) Let  $W$  be the locus of zeros of  $\mathcal{I}$  in the affine space  $\mathbb{A}^{n+1}$  with coordinates  $x_0, \dots, x_n$ . The homogeneous polynomial  $g$  vanishes at every point of  $W$  different from the origin, and since  $g$  isn't a constant, it vanishes at the origin too. So the affine Strong Nullstellensatz applies.  $\square$

### (3.2.8) quasiprojective varieties

### I don't like subsection quasiprojective. Too pompous. Just talk about nonempty open subsets of a projective variety. Mention quasiprojective in passing. ###

A nonempty (Zariski) open subset  $X$  of a projective variety is called a *quasiprojective variety*. For instance, a projective variety is quasiprojective. The complement of a point in a projective variety is a quasiprojective variety. An affine variety  $X = \text{Spec } A$  may be regarded as a quasiprojective variety by embedding it as a closed subvariety of the standard affine space  $\mathbb{U}^0 : \{x_0 \neq 0\}$ . It becomes an open subvariety of its closure in  $\mathbb{P}^n$ , which is a projective variety (Lemma 2.2.14 (ii)).

These days, it is customary to define varieties without reference to an embedding into projective space, as we did for affine varieties in Chapter 2 (??). However, to do this requires work. Most operations that one wants to make preserve the quasiprojective property, and though there are varieties that cannot be embedded into any projective space, they aren't very important. In fact, it is hard enough to find convincing examples of such varieties that we won't try to give one here. All varieties that we consider will be quasiprojective. In order to simplify terminology, and because the word "quasiprojective" is ugly, we will henceforth use the word "variety" to mean "quasiprojective variety".

The topology on a (quasiprojective) variety is induced from the topology on projective space.

**3.2.9. Lemma.** *The topology on the affine open subset  $\mathbb{U}^0 : x_0 \neq 0$  of  $\mathbb{P}^n$  that is induced from the Zariski topology on  $\mathbb{P}^n$  is the Zariski topology obtained by viewing  $\mathbb{U}^0$  as the affine space  $\text{Spec } \mathbb{C}[u_1, \dots, u_n]$ ,  $u_i = x_i/x_0$ .  $\square$*

Here is a description of a quasiprojective variety  $X$  in terms of equations:

**3.2.10.** Let  $\overline{X}$  be the closure of  $X$  in projective space  $\mathbb{P}^n$ , and let  $C = \overline{X} - X$  be the (closed) complement of  $X$  in  $\overline{X}$ . The closed set  $\overline{X}$  will be the zero set of a family  $f_1, \dots, f_k$  of homogeneous polynomials, and  $C$  will be the zero set of another family  $g_1, \dots, g_\ell$ . A point  $p$  of  $X$  will be a point of projective space that solves the equations  $f = 0$  but doesn't solve  $g = 0$ : All of the polynomials  $f_i$  vanish at  $p$ , and at least one polynomial  $g_j$  doesn't vanish there.

For example, suppose that an affine variety  $X$  is embedded as a closed subvariety of  $\mathbb{U}^0$  and that its closure  $\overline{X}$  in  $\mathbb{P}^n$  is the locus  $f = 0$ . Then a point of  $X$  is the set of points at which  $f = 0$ , but  $x_0 \neq 0$ .  $\square$

### 3.3 Product Varieties

The properties of products of varieties seem intuitive, but some of the proofs aren't obvious. As we will see, the (Zariski) topology on a product of varieties isn't the product topology.

The *product topology* on the product  $X \times Y$  of topological spaces is the coarsest topology such that the projection maps  $X \times Y \rightarrow X$  and  $X \times Y \rightarrow Y$  are continuous. So if  $C$  and  $D$  are closed subsets of  $X$  and  $Y$ , then  $C \times D$  is a closed subset of  $X \times Y$  in the product topology. Every closed set in the product topology is a finite union of such subsets.

So the first examples of closed subsets of  $\mathbb{P}^m \times \mathbb{P}^n$  are products of the form  $C \times D$ , where  $C$  is a closed subset of  $\mathbb{P}^m$  and  $D$  is a closed subset of  $\mathbb{P}^n$ . The product topology on  $\mathbb{P}^m \times \mathbb{P}^n$  is much coarser than the Zariski topology. For example, the proper (Zariski) closed subsets of  $\mathbb{P}^1$  are the nonempty finite subsets. In the product topology, the proper closed subsets of  $\mathbb{P}^1 \times \mathbb{P}^1$  are finite unions of points and sets of the form  $\mathbb{P}^1 \times q$  and  $p \times \mathbb{P}^1$  ('horizontal' and 'vertical' lines). Most Zariski closed subsets of  $\mathbb{P}^1 \times \mathbb{P}^1$  aren't of this form.

**3.3.1. Proposition.** *Let  $X$  and  $Y$  be irreducible topological spaces, and suppose that a topology is given on the product  $\Pi = X \times Y$ , with the following properties:*

- *The projections  $\Pi \xrightarrow{\pi_1} X$  and  $\Pi \xrightarrow{\pi_2} Y$  are continuous.*
- *For all  $x$  in  $X$  and all  $y$  in  $Y$ , the fibres  $x \times Y$  and  $X \times y$ , with topologies induced from  $\Pi$ , are homeomorphic to  $Y$  and  $X$ , respectively.*

*Then  $\Pi$  is an irreducible topological space.*

The product of varieties has these two properties.

The first condition means that the topology on  $X \times Y$  is at least as fine as the product topology, and the second one assures us that the topology isn't too fine. (We don't want the discrete topology on  $\Pi$ , for example.)

**3.3.2. Lemma.** *Let  $X, Y$ , and  $\Pi$  be as in the proposition. If  $W$  is an open subset of  $\Pi$ , its image  $U$  via the projection  $\Pi \rightarrow Y$  is an open subset of  $Y$ .*

*proof.* The intersection  ${}_xW = W \cap (x \times Y)$  is an open subset of the fibre  $x \times Y$ , and its image  ${}_xU$  in the homeomorphic space  $Y$  is also open. Since  $W$  is the union of the sets  ${}_xW$ ,  $U$  is the union of the open sets  ${}_xU$ . So  $U$  is open.  $\square$

*proof of Proposition 3.3.1.* Let  $C$  and  $C'$  be closed subsets of the product  $\Pi$ . Suppose that  $C < \Pi$  and  $C' < \Pi$ , and let  $W = \Pi - C$  and  $W' = \Pi - C'$  be the open complements of  $C$  and  $C'$  in  $\Pi$ . To show that  $\Pi$  is irreducible, we must show that  $C \cup C' < \Pi$ . We do this by showing that  $W \cap W'$  isn't the empty set.

Since  $C < \Pi$ ,  $W$  isn't empty, and similarly,  $W'$  isn't empty. The lemma tells us that the images  $U$  and  $U'$  of  $W$  and  $W'$  via projection to  $Y$  are nonempty open subsets of  $Y$ . Since  $Y$  is irreducible,  $U \cap U'$  is nonempty. Let  $y$  be a point of  $U \cap U'$ . The intersection  $W_y = W \cap (X \times y)$  is an open subset of  $X \times y$ , and since its image  $U$  contains  $y$ ,  $W_y$  contains a point of the form  $p = (x, y)$ , it is nonempty. Similarly,  $W'_y = W' \cap (X \times y)$  is a nonempty open subset of  $X \times y$ . Since  $X \times y$  is homeomorphic to the irreducible space  $X$ , it is irreducible. So  $W_y \cap W'_y$  is nonempty, and therefore  $W \cap W'$  is nonempty, as was to be shown.  $\square$

### (3.3.3) products of affine varieties

We inspect the product  $X \times Y$  of the affine varieties  $X = \text{Spec } A$  and  $Y = \text{Spec } B$ . Say that  $X$  is embedded as a closed subvariety of  $\mathbb{A}^m$ , so that  $A = \mathbb{C}[x_1, \dots, x_m]/P$  for some prime ideal  $P$ , and that  $Y$  is embedded similarly into  $\mathbb{A}^n$ , and  $B = \mathbb{C}[y_1, \dots, y_n]/Q$ . Then in affine  $x, y$ -space  $\mathbb{A}^{m+n}$ ,  $X \times Y$  is the locus of the equations  $f(x) = 0$  and  $g(y) = 0$  with  $f \in P$  and  $g \in Q$ . Proposition 3.3.1 shows that  $X \times Y$  is irreducible. Therefore it is a variety.

**3.3.4. Proposition.** *Let  $P$  and  $Q$  be prime ideals of  $\mathbb{C}[x]$  and  $\mathbb{C}[y]$ , respectively, and let  $X = V(P)$  and  $Y = V(Q)$  be their zero sets in  $\mathbb{A}^m$  and  $\mathbb{A}^n$ , respectively. The ideal  $I$  of all elements of  $\mathbb{C}[x, y]$  that vanish on the variety  $X \times Y$  is generated by the elements of  $P$  and  $Q$ . Therefore  $I$  is a prime ideal.*

The fact that  $X \times Y$  is a variety tells us only that the radical of  $I$  is a prime ideal.

*proof of Proposition 3.3.4* The ideal of  $\mathbb{C}[x, y]$  generated by  $P$  consists of combinations of elements of  $P$  with polynomials in  $x, y$  as coefficients. Let's denote that ideal by  $P'$ . Similarly, let  $Q'$  denote the ideal generated by  $Q$  in  $\mathbb{C}[x, y]$ . Then  $I = P' + Q'$ .

Let  $A = \mathbb{C}[x]/P$ ,  $B = \mathbb{C}[y]/Q$ , and  $R = \mathbb{C}[x, y]/I$ . Any polynomial in  $x, y$  can be written, in many ways, as a sum of products of polynomials in one set of variables:  $p(x, y) = \sum a_i(x)b_i(y)$ . Therefore any element  $p$  of  $R$  can be written as a finite sum

$$(3.3.5) \quad p = \sum_{i=1}^k a_i b_i$$

with  $a_i$  in  $A$  and  $b_i$  in  $B$ . We show that if  $p$  vanishes identically on  $X \times Y$ , then  $p = 0$ . To do this, we show that the same element  $p$  can also be written as a sum of  $k - 1$  products.

If  $a_k$  is zero, then  $p = \sum_{i=1}^{k-1} a_i b_i$ . Suppose that  $a_k \neq 0$ . Then the function defined by  $a_k$  isn't identically zero on  $X$ . We choose a point  $\bar{x}$  of  $X$  such that  $a_k(\bar{x}) \neq 0$ . Writing  $a_i(\bar{x}) = \bar{a}_i$  and  $\bar{p}(y) = p(\bar{x}, y)$ , gives us the equation  $\bar{p}(y) = \sum_{i=1}^k \bar{a}_i b_i$ . Since  $p$  vanishes on  $X \times Y$ ,  $\bar{p}$  vanishes on  $Y$ , and this implies that  $\bar{p} = 0$ . Since  $\bar{a}_k \neq 0$ , we can solve the equation  $\sum_{i=1}^k \bar{a}_i b_i = 0$  for  $b_k$ :  $b_k = \sum_{i=1}^{k-1} c_i b_i$ , where  $c_i = -\bar{a}_i / \bar{a}_k$ . Substituting into  $p$  gives us an expression for  $p$  as a sum of  $k - 1$  terms. Finally, when  $k = 1$ ,  $\bar{a}_1 b_1 = 0$ . Then  $b_1 = 0$ , and  $p = 0$ .  $\square$

### (3.3.6) the Zariski topology on $\mathbb{P}^m \times \mathbb{P}^n$

As mentioned above (3.1.10), the product of projective spaces  $\mathbb{P}^m \times \mathbb{P}^n$  is made into a projective variety by identifying it with its Segre image, the locus of the Segre equations  $w_{ij}w_{k\ell} = w_{i\ell}w_{kj}$ . Since  $\mathbb{P}^m \times \mathbb{P}^n$ , with its Segre embedding, is a projective variety, we don't really need a separate definition of its Zariski topology. Its closed subsets are the zero sets of families of homogeneous polynomials in the Segre variables  $w_{ij}$  including the Segre equations.

One can also describe the closed subsets of  $\mathbb{P}^m \times \mathbb{P}^n$  directly, in terms of bihomogeneous polynomials. A polynomial  $f(x, y)$  is *bihomogeneous* if it is homogeneous in the variables  $x$  and also in the variables  $y$ . For example, the polynomial  $x_0^2 y_0 + x_0 x_1 y_1$  is bihomogeneous, of degree 2 in  $x$  and degree 1 in  $y$ .

Because  $(x, y)$  and  $(\lambda x, \mu y)$  represent the same point of  $\mathbb{P}^m \times \mathbb{P}^n$  for all nonzero  $\lambda$  and  $\mu$ , we want to know that  $f(x, y) = 0$  if and only if  $f(\lambda x, \mu y) = 0$ , and this is true for all nonzero  $\lambda$  and  $\mu$  if and only if  $f$  is bihomogeneous.

**3.3.7. Lemma.** *The (Zariski) topology on  $\mathbb{P}^m \times \mathbb{P}^n$  has the properties listed in Proposition 3.3.1.*

- *The projections  $\mathbb{P}^m \times \mathbb{P}^n \rightarrow \mathbb{P}^m$  and  $\mathbb{P}^m \times \mathbb{P}^n \rightarrow \mathbb{P}^n$  are continuous maps.*
- *For all  $y$  in  $\mathbb{P}^n$ , the fibre  $\mathbb{P}^m \times y$ , with its topology induced from  $\mathbb{P}^m \times \mathbb{P}^n$ , is homeomorphic to  $\mathbb{P}^m$ .*

*proof.* We look at the projection  $\mathbb{P}^m \times \mathbb{P}^n \rightarrow \mathbb{P}^m$ . If  $X$  is the closed subset of  $\mathbb{P}^m$  defined by a system of homogeneous polynomials  $f_i(x)$ , its inverse image in  $\mathbb{P}^m \times \mathbb{P}^n$  is the zero set of the same system, considered as a family of bihomogeneous polynomials of degree zero in  $y$ . So the inverse image is closed.

Since the projections are continuous, it suffices, for the second property, to show that the inclusion map  $\mathbb{P}^m \rightarrow \mathbb{P}^m \times \mathbb{P}^n$  that sends  $\mathbb{P}^m$  to  $\mathbb{P}^m \times y$  is continuous. If  $f(x, y)$  is a bihomogeneous polynomial and  $y^0$  is a point of  $Y$ , the zero set of  $f$  in  $\mathbb{P}^m \times y^0$  is the zero set of  $f(x, y^0)$ . This polynomial defines a closed subset of  $\mathbb{P}^m$ .  $\square$

**3.3.8. Proposition.**

- (i) *A subset of  $\mathbb{P}^m \times \mathbb{P}^n$  is closed if and only if it is the locus of zeros of a family of bihomogeneous polynomials.*
- (ii) *If  $X$  and  $Y$  are closed subsets of  $\mathbb{P}^m$  and  $\mathbb{P}^n$ , respectively, then  $X \times Y$  is a closed subset of  $\mathbb{P}^m \times \mathbb{P}^n$ .*

*proof.* (i) For the proof, we denote the Segre image of  $\mathbb{P}^m \times \mathbb{P}^n$  by  $\Pi$ . Let  $f(w)$  be a homogeneous polynomial in the Segre variables  $w_{ij}$ . When we substitute  $w_{ij} = x_i y_j$  into  $f$ , we obtain a polynomial  $f(x_i y_j)$  that is bihomogeneous and that has the same degree as  $f$  in  $x$  and in  $y$ . Let's denote that bihomogeneous polynomial by  $\tilde{f}(x, y)$ . The inverse image of the zero set of  $f$  in  $\Pi$  is the zero set of  $\tilde{f}$  in  $\mathbb{P}^m \times \mathbb{P}^n$ . Therefore the inverse image of a closed subset of  $\Pi$  is the zero set of a family of bihomogeneous polynomials in  $\mathbb{P}^m \times \mathbb{P}^n$ .

Conversely, let  $g(x, y)$  be a bihomogeneous polynomial, say of degree  $r$  and  $s$  in  $y$ . If  $r = s$ , we may collect variables that appear in  $g$  in pairs  $x_i y_j$  and replace each pair  $x_i y_j$  by  $w_{ij}$ . We will obtain a homogeneous polynomial  $G$  in  $w$  such that  $G(w) = g(x, y)$  when  $w_{ij} = x_i y_j$ . The zero set of  $G$  in  $\Pi$  is the image of the zero set of  $g$  in  $\mathbb{P}^m \times \mathbb{P}^n$ .

Suppose that  $r \geq s$ , and let  $k = r - s$ . Because the variables  $y$  cannot all be zero at any point of  $\mathbb{P}^n$ , the equation  $g = 0$  on  $\mathbb{P}^m \times \mathbb{P}^n$  is equivalent with the system of equations  $gy_0^k = gy_1^k = \cdots = gy_n^k = 0$ . The polynomials  $gy_i^k$  are bihomogeneous, of same degree in  $x$  and in  $y$ .

- (ii) A polynomial  $f(x)$  can be viewed as a bihomogeneous polynomial of degree zero in  $y$ , and a polynomial  $g(y)$  is bihomogeneous of degree zero in  $x$ . So  $X \times Y$ , which is the locus  $f = g = 0$  in  $\mathbb{A}^{m+n}$ , is a closed subset of  $\mathbb{P}^m \times \mathbb{P}^n$ .  $\square$

**3.3.9. Corollary.** *Let  $X$  and  $Y$  be projective varieties, and let  $\Pi$  denote the product  $X \times Y$ , a closed subset of  $\mathbb{P}^m \times \mathbb{P}^n$ .*

- (i) *The projections  $\Pi \rightarrow X$  and  $\Pi \rightarrow Y$  are continuous.*
- (ii) *For all  $x$  in  $X$  and all  $y$  in  $Y$ , the fibres  $x \times Y$  and  $X \times y$ , with topologies induced from  $\Pi$ , are homeomorphic to  $Y$  and  $X$ , respectively.*  $\square$

The next corollary follows from Proposition 3.3.1 and Corollary 3.3.9.

**3.3.10. Corollary.** *If  $X$  and  $Y$  are projective varieties, so is  $X \times Y$ .*  $\square$

We will come back to products in Chapter ??.

## 3.4 Morphisms and Isomorphisms

### (3.4.1) the function field



Let  $X$  be a projective variety, and let  $X^i$  be its intersection with the standard affine open subset  $\mathbb{U}^i$  of projective space with coordinates  $x_0, \dots, x_n$ . If nonempty,  $X^i$  will be an affine variety – an irreducible closed subset of  $\mathbb{U}^i$ . Let's omit the indices for which  $X^i$  is empty. Then the intersection  $X^{ij} = X^i \cap X^j$  will be a localization of  $X^i$  and also a localization of  $X^j$ . If  $X^i = \text{Spec } A_i$  and  $u_{ij} = x_j/x_i$ , then  $X^{ij} = \text{Spec } A_{ij}$ , and  $A_{ij} = A_i[u_{ij}^{-1}] = A_j[u_{ji}^{-1}]$ . Therefore the fraction fields of the coordinate algebras  $A_i$  are equal for all  $i$  such that  $X^i$  isn't empty.

**3.4.2. Definition.** The *function field*  $K_X$  of a projective variety  $X$  is the field of fractions of the coordinate algebra  $A_i$  of any one of its nonempty affine open subsets  $X^i = X \cap \mathbb{U}^i$ . If  $X'$  is an open subvariety of a projective variety  $X$ , the *function field* of  $X'$  is the function field of  $X$ .  $\square$

Thus all open subvarieties of a variety have the same function field. In particular, suppose that we regard an affine variety  $X = \text{Spec } A$  as a closed subvariety of  $\mathbb{U}^0$ . The function field of  $X$  will be the field of fractions of  $A$ .

A *rational function* on a variety  $X'$  is an element of its function field of  $K$ .

A rational function can be evaluated at some points of  $X'$ , but probably not all of them. Suppose that  $X'$  is an open subvariety of a projective variety  $X$ , and that  $p$  is a point of the affine open set  $X^i = \text{Spec } A_i$ , as above. A rational function  $\alpha$  on  $X$  is *regular* at  $p$  if it is a regular function at  $p$  on one  $X^i$ . This means that one can write  $\alpha$  as a fraction  $a/b$  of elements of  $A_i$ , with  $b(p) \neq 0$ . Then the value of  $\alpha$  at  $p$  is  $\alpha(p) = a(p)/b(p)$ .

**3.4.3. Lemma.** *a rational function  $\alpha$  that is regular on an open subset  $U$  of  $X$  is determined by the function it defines on  $U$ .*

*proof.* We show that if the function is identically zero, then  $\alpha = 0$ . For some  $i$ , the intersection  $U \cap X^i$  will be nonempty. We may replace  $U$  by a localization  $X_s^i$  of  $X^i$ , so we may assume that  $U$  is affine, say  $U = \text{Spec } A$ . Then what is to be proved is that if an element  $\alpha$  of  $A$  defines the zero function, it is zero.

**3.4.4. Lemma.** *Let  $A$  be a finite-type algebra. The intersection of the maximal ideals of  $A$  is the zero ideal.*

*proof.* We present  $A$  as  $\mathbb{C}[x_1, \dots, x_n]/P$  where  $P$  is a prime ideal. Then  $U$  becomes a closed subset of  $\mathbb{A}^n$ . Let  $g(x)$  be a polynomial that represents  $\alpha$ . The function defined by  $\alpha$  is restriction of the polynomial function  $g$  to  $U$ . The Strong Nullstellensatz tells us that if  $g$  vanishes on  $U$ , it is in the prime ideal  $P$ , and then  $\alpha = 0$ .  $\square$

### (3.4.5) points with values in a field

Let  $K$  be a field that contains the complex numbers. A *point* of projective space  $\mathbb{P}^n$  with values in  $K$  is an equivalence class of nonzero vectors  $\alpha = (\alpha_0, \dots, \alpha_n)$  with  $\alpha_i$  in  $K$ , the equivalence relation being analogous to the one for ordinary points:  $\alpha \sim \alpha'$  if  $\alpha' = \lambda\alpha$  for some  $\lambda$  in  $K$ .

Let  $\bar{Y}$  be a projective variety, defined by a system of polynomial equations  $f(y) = 0$  in projective space  $\mathbb{P}^n$ . A point  $\beta$  of  $\bar{Y}$  with values in  $K$  is a point of  $\mathbb{P}^n$  with values in  $K$  such that  $f(\beta) = 0$ . If  $Y$  is an open subvariety with closure  $\bar{Y}$ , and  $C = \bar{Y} - Y$  is the complement of  $Y$  in  $\bar{Y}$ , then a point  $\beta$  of  $Y$  with values in  $K$  is a point of  $\bar{Y}$  that isn't also a point of  $C$ .

In particular, if  $K$  is the function field of a variety  $X$  the embedding of  $X$  into projective space  $\mathbb{P}^m$  defines a point  $(\alpha_0, \dots, \alpha_m)$  of with values in  $K$ .

### (3.4.6) morphisms to projective space

For the rest of this section, it will be helpful to have a separate notation for the point with values in  $K$  determined by a nonzero vector  $\alpha$ . We'll denote that point by  $\underline{\alpha}$ . Thus  $\underline{\alpha} = \underline{\alpha'}$  if  $\alpha' = \lambda\alpha$  for some nonzero  $\lambda$  in  $K$ . We'll drop this notation later.

We will define a morphism from a variety  $Y$  to projective space using a point of  $\mathbb{P}^n$  with values in the function field  $K$  of  $Y$ . When doing this, we must keep in mind that the points of projective space are equivalence classes of vectors, not the vectors themselves. This complication turns out to be useful.

We begin with a simple example.

**3.4.7. Example.** Let  $C$  be the conic in the projective plane  $\mathbb{P}^2$  defined by the polynomial  $f(x_0, x_1, x_2) = x_0x_1 + x_0x_2 + x_1x_2$ . We project  $C$  to the projective line  $L_0 : \{x_0 = 0\}$ , defining  $C \xrightarrow{\pi} \mathbb{P}^1$  by  $\pi(x_0, x_1, x_2) = (x_1, x_2)$ . This formula is undefined at the point  $q = (1, 0, 0)$ , though the map extends to the whole conic  $C$ . The image of  $q$  is the point at which the tangent line  $L_q$  to  $C$  at  $q$  intersects  $L_0$ .

Let's write this projection using a point with values in the function field  $K$  of  $C$ . The affine open set  $\{x_0 \neq 0\}$  of  $\mathbb{P}^2$  is the polynomial algebra  $\mathbb{C}[u_1, u_2]$ , with  $u_1 = x_1/x_0$  and  $u_2 = x_2/x_0$ . We also denote by  $u_i$  the restriction of the function  $u_i$  to  $C^0 = C \cap \mathbb{U}^0$ . The restricted functions are related by the equation  $u_1 + u_2 + u_1u_2 = 0$  obtained by dehomogenizing  $f$ . We solve for  $u_2$ :  $u_2 = -u_1/(1 + u_1)$ .

The projection is given by  $\pi(x_0, x_1, x_2) = \pi(1, u_1, u_2) = (u_1, -u_1/(1 + u_1)) \approx (1 + u_1, -1)$ . The formula  $(1 + u_1, -1)$  is defined at all points at which  $x_0 \neq 0$ , including  $q$ . Thus  $\pi(q) = (1, -1)$ .

To define  $\pi$  at the remaining points, we can look on another standard affine open set. Let  $v_i = x_i/x_1$ ,  $i = 0, 2$ , and  $w_i = x_i/x_2$ ,  $j = 0, 1$ . Then  $(x_0, x_1, x_2) = (v_0, 1, v_2) = (w_0, w_1, 1)$ . The projection can also be written as  $\pi(x) = (1, v_2)$ , which is valid provided that  $x_1 \neq 0$ , or as  $\pi(x) = (w_1, 1)$ , which is valid provided that  $x_2 \neq 0$ .  $\square$

Let  $\alpha = (\alpha_0, \dots, \alpha_n)$  be a nonzero vector with entries in the function field  $K$  of  $Y$ . We try to define a morphism  $Y \xrightarrow{u_\alpha} \mathbb{P}^n$  from  $Y$  to projective space  $\mathbb{P}^n$  using the point  $\underline{\alpha}$  with values in  $K$  that is defined by  $\alpha$ . To define the image  $\underline{\alpha}(q)$  of a point  $q$  of  $Y$  (an ordinary point), we look for a vector  $\alpha' = (\alpha'_0, \dots, \alpha'_n)$ , with  $\underline{\alpha}' = \underline{\alpha}$ , i.e.,  $\alpha' = \lambda\alpha$ , such that the rational functions  $\alpha'_i$  are all regular and not all zero at  $q$ . Such a vector may exist or not. If it exists, we define

$$(3.4.8) \quad \underline{\alpha}(q) = (\alpha'_0(q), \dots, \alpha'_n(q)) \quad (= \underline{\alpha}'(q))$$

If such a vector  $\alpha'$  exists for every point  $q$  of  $Y$ , we call  $\underline{\alpha}$  a *good point*.

**3.4.9. Lemma.** A point  $\underline{\alpha}$  of  $\mathbb{P}^n$  with values in the function field  $K_Y$  of  $Y$  is a good point if either one of the two following conditions holds for every point  $q$  of  $Y$ :

- There is an element  $\lambda$  in  $K_Y$  such that the rational functions  $\alpha'_i = \lambda\alpha_i$ ,  $i = 0, \dots, n$ , are regular and not all zero at  $q$ .
- There is an index  $j$ ,  $0 \leq j \leq n$ , such that the rational functions  $\alpha_i/\alpha_j$ ,  $j = 0, \dots, n$ , are regular at  $q$ .

*proof.* The first condition simply restates the definition. We show that it is equivalent with the second one. Suppose that  $\alpha_i/\alpha_j$  are regular at  $q$  for all  $i$ . Let  $\lambda = \alpha_j^{-1}$ , and let  $\alpha'_i = \lambda\alpha_i = \alpha_i/\alpha_j$ . The rational functions  $\alpha'_i$  are regular at  $q$ , and they aren't all zero there because  $\alpha'_j = 1$ .

Conversely, suppose that  $\alpha'_i = \lambda\alpha_i$  are all regular at  $q$  and that  $\alpha'_j$  isn't zero there. Then  $\alpha_j'^{-1}$  is a regular function at  $q$ , so the rational functions  $\alpha'_i/\alpha'_j$ , which are equal to  $\alpha_i/\alpha_j$ , are regular at  $q$  for all  $i$ .  $\square$

**3.4.10. Lemma.** With notation as in (3.4.8), the image  $\underline{\alpha}(q)$  of a point  $q$  is independent of the choice of the vector  $\alpha'$ .

*proof.* Let  $\alpha' = \lambda\alpha$  and  $\alpha'' = \mu\alpha$ . Suppose that  $\alpha'_i$  are all regular and not all zero at  $q$ , and that  $\alpha''_i$  are also regular and not all zero there. We need to show that  $\underline{\alpha}' = \underline{\alpha}''$ , and since we can replace  $\alpha$  by  $\mu\alpha$  and  $\alpha'$  by  $\mu^{-1}\lambda\alpha$ , we may assume that  $\alpha'' = \alpha$ , so that  $\alpha' = \lambda\alpha$ . The rational functions  $\alpha_i$  and  $\alpha'_i$  are all regular at  $q$ , and there are indices  $j, k$  such that  $\alpha_j(q)$  and  $\alpha'_k(q)$  are nonzero. Then  $\lambda = \alpha'_j/\alpha_j$  and  $\lambda^{-1} = \alpha_k/\alpha'_k$  are both regular at  $q$ . So  $\lambda(q) \neq 0$ ,  $\alpha'(q) = \lambda(q)\alpha(q)$ , and  $\underline{\alpha}'(q) = \underline{\alpha}(q)$ .  $\square$

**3.4.11. Definition.** Let  $Y$  be a variety with function field  $K_Y$ . A *morphism* from  $Y$  to projective space  $\mathbb{P}^n$  is a map that is defined, as in (3.4.8)

, by a good point  $\underline{\alpha}$  with values in  $K_Y$ . We denote that morphism by  $\underline{\alpha}$ .

**3.4.12. Example.** The identity map  $\mathbb{P}^1 \rightarrow \mathbb{P}^1$ .

Let  $X = \mathbb{P}^1$ , and let  $(x_0, x_1)$  be coordinates in  $X$ . The function field of  $X$  is the field  $K = \mathbb{C}(t)$  of rational functions in the variable  $t = x_1/x_0$ . The identity map  $X \rightarrow X$  is the map  $\underline{\alpha}$  defined by the point  $\alpha = (1, t)$  with values in  $K$ . For every point  $p$  of  $X$  except the point  $(1, 0)$ ,  $\alpha(p)$  is defined and not zero, so  $\underline{\alpha}(p) = \alpha(p)$ . At the point  $q = (0, 1)$ ,  $\alpha' = (t^{-1}, 1) = t^{-1}\alpha$  defines  $\underline{\alpha}$ :  $\underline{\alpha}(q) = (0, 1)$ .  $\square$

### (3.4.13) morphisms to quasiprojective varieties

**3.4.14. Definition.** Let  $Y$  be a variety, and let  $X$  be a subvariety of a projective space  $\mathbb{P}^n$ . A morphism of varieties  $Y \xrightarrow{\alpha} X$  is the restriction of a morphism  $Y \xrightarrow{\alpha} \mathbb{P}^n$  whose image is contained in  $X$ .

Thus if a projective variety  $X$  is the locus of zeros of a family  $f$  of homogeneous polynomials, a morphism  $Y \xrightarrow{\alpha} \mathbb{P}^n$  defines a morphism  $Y \rightarrow X$  if  $f(\alpha) = 0$ . This follows from Lemma 3.4.3. The image of  $Y$  will be contained in  $X$  if and only if  $f(\alpha(q)) = 0$  for every point  $q$  of  $Y$ . If so, the lemma tells us that  $f(\alpha) = 0$ .

A word of caution: A morphism  $Y \xrightarrow{\alpha} X$  won't define a map of function fields  $K_X \rightarrow K_Y$  unless the image of  $Y$  is dense in  $X$ .

**3.4.15. Proposition.** A morphism of varieties  $Y \xrightarrow{\alpha} X$  is a continuous map in the Zariski topology, and also in the classical topology.

*proof.* This proposition is trivial, once one has unraveled the notation. Let  $\mathbb{U}^i$  be the standard affine open subset of  $\mathbb{P}^m$ , and let  $Y^i$  be an affine open subset of the inverse image of  $\mathbb{U}^i$ . If  $X = \mathbb{P}^m$ , the restriction  $Y^i \rightarrow \mathbb{U}^i$  of  $\alpha$  is continuous in both topologies because it is a morphism of affine varieties. Since  $Y$  can be covered by affine open sets such as  $Y^i$ ,  $\alpha$  is continuous. Continuity for a morphism to a subvariety  $X$  of  $\mathbb{P}^m$  follows because the topology on  $X$  is the induced topology.  $\square$

**3.4.16. Proposition.** Let  $X, Y$ , and  $Z$  be varieties and let  $Z \xrightarrow{\beta} Y$  and  $Y \xrightarrow{\alpha} X$  be morphisms. The composed map  $Z \xrightarrow{\alpha\beta} X$  is a morphism.

*proof.* Say that  $X$  is a subvariety of  $\mathbb{P}^m$ . The morphism  $\alpha$  is the restriction of a morphism  $Y \rightarrow \mathbb{P}^m$  whose image is in  $X$ , and that is defined by a good point  $\underline{\alpha}$ ,  $\alpha = (\alpha_0, \dots, \alpha_m)$  of  $\mathbb{P}^m$  with values in the function field  $K_Y$  of  $Y$ . Similarly, if  $Y$  is a subvariety of  $\mathbb{P}^n$ , the morphism  $\beta$  is the restriction of a morphism  $Z \rightarrow \mathbb{P}^n$  whose image is contained in  $Y$ , and that is defined by a good point  $\underline{\beta}$ ,  $\beta = (\beta_0, \dots, \beta_n)$  of  $\mathbb{P}^n$  with values in the function field  $K_Z$  of  $Z$ .

Let  $z$  be a point (an ordinary point) of  $Z$ . Since  $\underline{\beta}$  is a good point, we may adjust  $\beta$  by a factor in  $K_Z$  so that the rational functions  $\beta_i$  are regular and not all zero at  $z$ . Then  $\beta(z)$  is the point  $(\beta_0(z), \dots, \beta_n(z))$  of  $Y$ . Let's denote that point by  $q = (q_0, \dots, q_m)$ . So  $q_i = \beta_i(z)$ . The elements  $\alpha_j$  are rational functions on  $Y$ . We may adjust  $\alpha$  by a factor in  $K_Y$ , so that they are regular and not all zero at  $q$ . Then  $[\alpha\beta](z) = \underline{\alpha}(q) = (\alpha_0(q), \dots, \alpha_m(q))$ , and  $\alpha_j(q) = \alpha_j(\beta_0(z), \dots, \beta_n(z)) = \alpha_j(\beta(z))$  are not all zero. When these adjustments have been made, the point of  $\mathbb{P}^m$  with values in  $K_Z$  that defines  $\underline{\alpha\beta}$  is  $(\alpha_0(\beta(z)), \dots, \alpha_m(\beta(z)))$ .  $\square$

This next is a lemma of topology.

**3.4.17. Lemma.** Let  $\{X^i\}$  be a covering of a topological space  $X$  by open sets. A subset  $Y$  of  $X$  is open (or closed) if and only if  $Y \cap X^i$  is open (or closed) in  $X^i$  for every  $i$ . In particular, if  $\{\mathbb{U}^i\}$  is the standard affine cover of  $\mathbb{P}^n$ , a subset  $Y$  of  $\mathbb{P}^n$  is open (or closed) if and only if  $Y \cap \mathbb{U}^i$  is open (or closed) in  $\mathbb{U}^i$  for every  $i$ .  $\square$

**3.4.18. Lemma.**

(i) The inclusion of an open or a closed subvariety  $Y$  into a variety  $X$  is a morphism.

(ii) Let  $Y \xrightarrow{f} X$  be a map whose image lies in an open or a closed subvariety  $Z$  of  $X$ . Then  $f$  is a morphism if and only if its restriction  $Y \rightarrow Z$  is a morphism.

(iii) Let  $\{Y^i\}$  be an open covering of a variety  $Y$ , and let  $Y^i \xrightarrow{f^i} X$  be morphisms. If the restrictions of  $f^i$  and  $f^j$  to the intersections  $Y^i \cap Y^j$  are equal for all  $i, j$ , there is a unique morphism  $f$  whose restriction to  $Y^i$  is  $f^i$ .

Part (iii) is trivial because the points with vals in  $K$  are the same. We omit the proofs of (i) and (ii).  $\square$

### (3.4.19) isomorphisms

A bijective morphism  $Y \xrightarrow{u} X$  of quasiprojective varieties is an *isomorphism* if its inverse function is also a morphism. Isomorphisms are important because they allow us to identify different incarnations of the

“same variety”, i.e., to describe an isomorphism class of varieties. The projective line  $\mathbb{P}^1$ , a conic in  $\mathbb{P}^2$ , and the twisted cubic in  $\mathbb{P}^3$  are isomorphic varieties, for example.

**3.4.20. Example.** Let  $Y$  denote the projective line, with coordinates  $y_0, y_1$ . As before, the function field of  $Y$  is the field  $K = \mathbb{C}(t)$  of rational functions in  $t = y_1/y_0$ . The degree 3 Veronese map  $Y \rightarrow \mathbb{P}^3$  (3.1.16) defines an isomorphism of  $Y$  to its image, a twisted cubic  $X$ . The Veronese map is defined by the point  $\alpha = (1, t, t^2, t^3)$  of  $\mathbb{P}^3$  with values in  $K$ . On the open set  $\{y_0 \neq 0\}$  of  $Y$ , the rational functions  $1, t, t^2, t^3$  are regular and not all zero. Let  $\lambda = t^{-3}$  and  $\alpha' = \lambda\alpha = (t^{-3}, t^{-2}, t^{-1}, 1)$ . The functions  $t^{-k}$  are regular on the open set  $\{y_1 \neq 0\}$ . So  $\alpha$  is a good point that defines a morphism  $Y \xrightarrow{\alpha} X$ .

The twisted cubic  $X$  is the locus of zeros of the equations (3.1.17).

$$v_0v_2 = v_1^2, \quad v_2v_1 = v_0v_3, \quad v_1v_3 = v_2^2$$

To identify the function field  $F$  of  $X$ , we put  $v_0 = 1$ , obtaining relations  $v_2 = v_1^2, v_3 = v_1^3$ . Then  $F$  is the field  $\mathbb{C}(v_1)$ . The point of  $Y = \mathbb{P}^1$  with values in  $F$  that defines the inverse of the morphism  $\underline{\alpha}$  is  $\beta = (1, v_1)$ .  $\square$

**3.4.21. Lemma.** Let  $Y \xrightarrow{f} X$  be a morphism of varieties, let  $\{X^i\}$  be an open covering of  $X$ , and let  $Y^i = f^{-1}X^i$ . If the restrictions  $Y^i \xrightarrow{f^i} X^i$  of  $f$  are isomorphisms, then  $f$  is an isomorphism.

*proof.* Let  $g^i$  denote the inverse of the morphism  $f^i$ . Then  $g^i = g^j$  on  $X^i \cap X^j$  because  $f^i = f^j$  on  $Y^i \cap Y^j$ . By (3.4.18) (iii), there is a unique morphism  $X \xrightarrow{g} Y$  whose restriction to  $Y^i$  is  $g^i$ . That morphism is the inverse of  $f$ .  $\square$

### (3.4.22) the diagonal

Let  $X$  be a variety. The *diagonal*  $X_\Delta$  is the set of points  $(p, p)$  in  $X \times X$ . It is a subset of  $X \times X$  that is closed in the Zariski topology, but not in the product topology.

**3.4.23. Proposition.** Let  $X$  be a variety. The diagonal  $X_\Delta$  is a closed subvariety of the product variety  $X \times X$ .

*proof.* Let  $\mathbb{P}$  denote the projective space  $\mathbb{P}^n$  that contains  $X$ , and let  $x_0, \dots, x_n$  and  $y_0, \dots, y_n$  be coordinates in the two factors of  $\mathbb{P} \times \mathbb{P}$ . The diagonal  $\mathbb{P}_\Delta$  in  $\mathbb{P} \times \mathbb{P}$  is the closed subvariety defined by the bilinear equations  $x_i y_j = x_j y_i$ , or in the Segre variables, by the equations  $w_{ij} = w_{ji}$ , which show that the ratios  $x_i/x_j$  and  $y_i/y_j$  are equal.

Next, suppose that  $X$  is the closed subvariety of  $\mathbb{P}$  defined by a system of homogeneous equations  $f(x) = 0$ . The diagonal  $X_\Delta$  can be identified as the intersection of the product  $X \times X$  with the diagonal  $\mathbb{P}_\Delta$  in  $\mathbb{P} \times \mathbb{P}$ , so it is a closed subvariety of  $X \times X$ . As a closed subvariety of  $\mathbb{P} \times \mathbb{P}$ , the diagonal  $X_\Delta$  is defined by the equations

$$(3.4.24) \quad x_i y_j = x_j y_i \quad \text{and} \quad f(x) = 0$$

The equations  $f(y) = 0$  are redundant. Finally,  $X_\Delta$  is irreducible because it is homeomorphic to  $X$ .  $\square$

It is interesting to compare Proposition 3.4.23 with the Hausdorff condition for a topological space. The proof of the next lemma is often assigned as an exercise in topology.

**3.4.25. Lemma.** A topological space  $X$  is a Hausdorff space if and only if, when  $X \times X$  is given the product topology, the diagonal  $X_\Delta$  is a closed subset of  $X \times X$ .  $\square$

Though a variety  $X$  with its Zariski topology isn't a Hausdorff space unless it is a point, Lemma 3.4.25 doesn't contradict Proposition 3.4.23 because the Zariski topology on  $X \times X$  is finer than the product topology.

### (3.4.26) the graph of a morphism

Let  $Y \xrightarrow{f} X$  be a morphism of varieties. The *graph*  $\Gamma$  of  $f$  is the subset of  $Y \times X$  of pairs  $(q, p)$  such that  $p = f(q)$ .

**3.4.27. Proposition.** The graph  $\Gamma_f$  of a morphism  $Y \xrightarrow{f} X$  is a closed subvariety of  $Y \times X$ , and it is isomorphic to  $Y$ .

*proof.* We form a diagram of morphisms

$$(3.4.28) \quad \begin{array}{ccc} \Gamma_f & \longrightarrow & Y \times X \\ v \downarrow & & \downarrow f \times \text{id} \\ X_\Delta & \longrightarrow & X \times X \end{array}$$

where  $v$  sends a point  $(q, p)$  of  $\Gamma_f$  with  $f(q) = p$  to  $(p, p)$ . The graph  $\Gamma_f$  is the inverse image in  $Y \times X$  of the diagonal  $X_\Delta$ . Since the diagonal is closed in  $X \times X$ ,  $\Gamma_f$  is closed in  $Y \times X$ .

Let  $\pi_1$  denote the projection from  $X \times Y$  to  $Y$ . The composition of the morphisms  $Y \xrightarrow{(id, f)} Y \times X \xrightarrow{\pi_1} Y$  is the identity map on  $Y$ , and the image of the map  $(id, f)$  is the graph  $\Gamma_f$ . Therefore  $Y$  maps bijectively to  $\Gamma_f$ . The two maps  $Y \rightarrow \Gamma_f$  and  $\Gamma_f \rightarrow Y$  are inverses, so  $\Gamma_f$  is isomorphic to  $Y$ .  $\square$

### (3.4.29) projection

The map

$$(3.4.30) \quad \mathbb{P}^n \xrightarrow{\pi} \mathbb{P}^{n-1}$$

that drops the last coordinate of a point:  $\pi(x_0, \dots, x_n) = (x_0, \dots, x_{n-1})$  is called a *projection*. It is defined at all points of  $\mathbb{P}^n$  except at the point  $q = (0, \dots, 0, 1)$ . This point is called the *center of projection*. So  $\pi$  is a morphism from the complement  $U = \mathbb{P}^n - \{q\}$  to  $\mathbb{P}^{n-1}$ .

Let the coordinates in  $\mathbb{P}^n$  and  $\mathbb{P}^{n-1}$  be  $x = x_0, \dots, x_n$  and  $y = y_0, \dots, y_{n-1}$ , respectively. The fibre  $\pi^{-1}(y)$  over a point  $(y_0, \dots, y_{n-1})$  is the set of points  $(x_0, \dots, x_n)$  such that  $(x_0, \dots, x_{n-1}) = (\lambda y_0, \dots, \lambda y_{n-1})$ , while  $x_n$  is arbitrary. It is the line in  $\mathbb{P}^n$  through the points  $(y_1, \dots, y_{n-1}, 0)$  and  $q = (0, \dots, 0, 1)$ , with the center of projection  $q$  omitted.

The graph  $\Gamma$  of  $\pi$  in  $U \times \mathbb{P}^{n-1}$  is the locus of solutions of the equations  $w_{ij} = w_{ji}$  for  $0 \leq i, j \leq n-1$ , which imply that the vectors  $(x_0, \dots, x_{n-1})$  and  $(y_0, \dots, y_{n-1})$  are proportional.

**3.4.31. Proposition.** In  $\mathbb{P}_x^n \times \mathbb{P}_y^{n-1}$ , the locus  $\bar{\Gamma}$  of the equations  $x_i y_j = x_j y_i$ , or  $w_{ij} = w_{ji}$ , with  $0 \leq i, j \leq n-1$ , is the closure of the graph  $\Gamma$  of  $\pi$ .

*proof.* The equations are true at points  $(x, y)$  of  $\Gamma$  at which  $x \neq q$ , and also at all points  $(q, y)$ . So the locus  $\bar{\Gamma}$ , a closed set, is the union of the graph  $\Gamma$  and the set  $q \times \mathbb{P}^{n-1}$ . We must show that a homogeneous polynomial  $g(w)$  that vanishes on  $\Gamma$  vanishes at all points of  $q \times \mathbb{P}^{n-1}$ . Given  $y$ , let  $x = (ty_0, \dots, ty_{n-1}, 1)$ . For all  $t \neq 0$ , the point  $(x, y)$  is in  $\Gamma$  and therefore  $g(x, y) = 0$ . Since  $g$  is a continuous function,  $g(x, y)$  approaches  $g(q, y)$  as  $t \rightarrow 0$ . So  $g(q, y) = 0$ .  $\square$

The projection  $\bar{\Gamma} \rightarrow \mathbb{P}_x^n$  that sends a point  $(x, y)$  to  $x$  is bijective except when  $x = q$ . The fibre over  $q$ , which is  $q \times \mathbb{P}^{n-1}$ , is a projective space of dimension  $n-1$ . Because the point  $q$  of  $\mathbb{P}^n$  is replaced by a projective space in  $\bar{\Gamma}$ , the map  $\bar{\Gamma} \rightarrow \mathbb{P}_x^n$  is called a *blowup* of the point  $q$ .

figure: projection with closure of graph??

**3.4.32. Proposition.** Let  $Y \xrightarrow{\alpha} X$  and  $Z \xrightarrow{\beta} W$  be morphisms of varieties. The product map  $Y \times Z \xrightarrow{\alpha \times \beta} X \times W$  that sends  $(y, z)$  to  $(\alpha(y), \beta(z))$  is a morphism

*proof.* Let  $p$  and  $q$  be points of  $X$  and  $Y$ , respectively. We may assume that  $\alpha_i$  are regular and not all zero at  $p$  and that  $\beta_j$  are regular and not all zero at  $q$ . Then, in the Segre coordinates  $w_{ij}$ ,  $[\underline{\alpha} \times \underline{\beta}](p, q)$  is the point  $w_{ij} = \alpha_i(p)\beta_j(q)$ . The fact that  $\alpha_i\beta_j$  are all regular at  $(p, q)$  and are not all zero there follows from the analogous properties of  $\alpha_i$  and  $\beta_j$ .  $\square$

### (3.4.33) digression: the function field of a product

To define the function field of a product  $X \times Y$  of projective varieties, one can use the Segre embedding  $\mathbb{P}_x^m \times \mathbb{P}_y^n \rightarrow \mathbb{P}^N$ . We use notation as in (3.1.10), and let's denote the product  $X \times Y$  by  $\Pi$ . So  $x_i, y_j$ , and  $w_{ij}$  are coordinates in the three projective spaces, and the Segre map is defined by  $w_{ij} = x_i y_j$ . Let  $\mathbb{U}^i, \mathbb{V}^j$ , and  $\mathbb{W}^{ij}$  be the standard affine open sets  $x_i \neq 0, y_j \neq 0$  and  $w_{ij} \neq 0$ . The function field will be the field of fractions of the nonempty intersections  $\Pi \cap \mathbb{W}^{ij} = \Pi^{ij}$ , and  $\Pi^{ij} \approx X^i \times Y^j$ , where  $X^i = X \cap \mathbb{U}^i$  and  $Y^j = Y \cap \mathbb{V}^j$ . Since  $\Pi^{ij}, X^i$ , and  $Y^j$  are affine varieties, the function field of the product  $\Pi = X \times Y$  is the field of fractions of any one of the nonempty affine open sets  $\Pi^{ij}$ .

Since  $\Pi^{ij} = X^i \cap Y^j$ , all that remains to do is to describe the field of fractions of a product of affine varieties  $\Pi = X \times Y$ , when  $X = \text{Spec } A$  and  $Y = \text{Spec } B$ . If  $A = \mathbb{C}[x]/P$  and  $B = \mathbb{C}[y]/Q$ , the coordinate algebra of  $\Pi$  is the algebra  $\mathbb{C}[x, y]/(P, Q)$ . This is the tensor product algebra  $A \otimes B$ . We don't need to know much about the tensor product algebra here, but let's use the tensor notation.

The function field  $K_X$  of  $X$  is the field of fractions of the coordinate algebra  $A$ . Similarly,  $K_Y$  is the field of fractions of  $B$  and  $K_{X \times Y}$  is the field of fractions of  $A \otimes B$ . The one important fact to note is that  $K_{X \times Y}$  isn't generated by  $K_X$  and  $K_Y$ . For example, if  $A = \mathbb{C}[x]$  and  $B = \mathbb{C}[y]$  (one  $x$  and one  $y$ ), then  $K_{X \times Y}$  is the field of rational functions in two variables  $\mathbb{C}(x, y)$ . The algebra generated by the fraction fields  $\mathbb{C}(x)$  and  $\mathbb{C}(y)$  consists of the rational functions  $p(x, y)/q(x, y)$  in which  $q(x, y)$  is a product  $f(x)g(y)$  of a polynomial in  $x$  and a polynomial in  $y$ . Most rational functions,  $1/(x + y)$  for example, aren't of this type.

But,  $K_{X \times Y}$  is the fraction field of  $A \otimes B$ .

### (3.4.34) digression: rational functions on projective space

Let  $R$  denote the polynomial ring  $\mathbb{C}[x_0, \dots, x_n]$ . If  $f$  is a homogeneous polynomial of positive degree  $d$ , it makes sense to say that  $f$  vanishes at a point of  $\mathbb{P}^n$ , because  $f(\lambda x) = \lambda^d f(x)$ . But  $f$  doesn't define a function on  $\mathbb{P}^n$ . On the other hand, a fraction  $g/h$  of homogeneous polynomials of the same degree  $d$  does define a function wherever  $h$  isn't zero, because

$$g(\lambda x)/h(\lambda x) = \lambda^d g(x)/\lambda^d h(x) = g(x)/h(x)$$

A *homogeneous fraction*  $f$  is a fraction of homogeneous polynomials. The *degree* of a homogeneous fraction  $f = g/h$  is the difference of degrees:  $\deg f = \deg g - \deg h$ .

**3.4.35. Definition.** A homogeneous fraction  $f$  is *regular* at a point  $p$  of  $\mathbb{P}^n$  if, when it is written as a fraction  $g/h$  of relatively prime homogeneous polynomials, the denominator  $h$  isn't zero at  $p$ , and  $f$  is *regular* on a subset  $U$  if it is regular at every point of  $U$ .  $\square$

This definition agrees with the one given above, in Definition 3.4.2.

**3.4.36. Lemma. (i)** Let  $h$  be a homogeneous polynomial of positive degree  $d$ , and let  $V$  be the open subset of  $\mathbb{P}^n$ , of points at which  $h$  isn't zero. The nonzero rational functions that are regular on  $V$  are those of the form  $g/h^k$ , where  $k \geq 0$  and  $g$  is a homogeneous polynomial of degree  $dk$ .

(ii) The only rational functions that are regular at every point of  $\mathbb{P}^n$  are the constant functions.

For example, the homogeneous polynomials that are nonzero at every point of the standard affine open set  $\mathbb{U}^0$  are the scalar multiples of powers of  $x_0$ . So the rational functions that are regular on  $\mathbb{U}^0$  are those of the form  $g/x_0^k$ ,  $g$  homogeneous of degree  $k$ . This agrees with the fact that the coordinate algebra of  $\mathbb{U}^0$  is the polynomial ring  $\mathbb{C}[u_1, \dots, u_n]$ , with  $u_i = x_i/x_0$ :  $g(x_0, \dots, x_n)/x_0^k = g(u_0, \dots, u_n)$  (with  $u_0 = 1$ ).

*proof of Lemma 3.4.36 (i)* Let  $\alpha$  be a regular function on the open set  $U$ , say  $g_1/h_1$ , where  $g_1$  and  $h_1$  are relatively prime homogeneous polynomials. Then  $h_1$  doesn't vanish on  $U$ , so its zero locus in  $\mathbb{P}^n$  is contained in the zero locus of  $h$ . According to the Strong Nullstellensatz 3.2.7,  $h_1$  divides a power of  $h$ , say  $h^k = f h_1$ . Then  $g_1/h_1 = f g_1/f h_1 = f g_1/h^k$ .

(ii) If a rational function  $f$  is regular at every point of  $\mathbb{P}^n$ , then it is regular on  $\mathbb{U}^0$ . It will have the form  $g/x_0^k$ ,  $g$  has degree  $k$  and isn't divisible by  $x_0$ . And since  $f$  is also regular on  $\mathbb{U}^1$ , it will have the form  $h/x_1^\ell$ , where  $x_1$  doesn't divide  $h$ . Then  $g x_1^\ell = h x_0^k$ . Since  $x_0$  doesn't divide  $g$ ,  $k = 0$ . Then  $g$  is a constant, and  $f = g$ .  $\square$

It is also true that the only rational functions on a projective variety  $X$  that are regular at every point of  $X$  are the constant functions. The proof of this will be given later (see Corollary 8.4.8). The constant functions are useless, so one has to look at regular functions on open subsets. One way that affine varieties appear in projective algebraic geometry is as open subsets on which there are enough regular functions.

### 3.5 Affine Varieties

We have used the term 'affine variety' in several contexts:

A closed subset of affine space  $\mathbb{A}_x^n$  is an affine variety, the set of zeros of a prime ideal  $P$  of  $\mathbb{C}[x]$ . Its coordinate algebra is  $A = \mathbb{C}[x]/P$ .

The spectrum  $\text{Spec } A$  of a finite type domain  $A$  is an affine variety that becomes a closed subvariety of affine space when one chooses a presentation  $A = \mathbb{C}[x]/P$ .

An affine variety becomes a quasiprojective variety by identifying the ambient affine space  $\mathbb{A}^n$  with the open subset  $\mathbb{U}^0$  of projective space.

We combine these definitions now: An *affine variety*  $X$  is a variety that is isomorphic to a variety of the form  $\text{Spec } A$ .

If  $X = \text{Spec } A$  is an affine variety with function field  $K$ , its coordinate algebra  $A$  will be the subalgebra of  $K$  of regular functions on  $X$ . So  $A$  and therefore  $\text{Spec } A$ , are determined uniquely by  $X$ , and the isomorphism  $\text{Spec } A \rightarrow X$  is determined uniquely too. When  $X$  is affine, it seems permissible to identify  $X$  with  $\text{Spec } A$ .

#### (3.5.1) regular functions on affine varieties

Let  $X = \text{Spec } A$  be an affine variety. Its function field  $K$  is the field of fractions of  $A$ . A rational function  $\alpha$  is regular at a point  $p$  of  $X$  if it can be written as a fraction  $a/s$  where  $a, s$  are in  $A$  and  $s(p) \neq 0$ , and  $\alpha$  is regular on  $X$  if it is regular at every point of  $X$ . On the other hand, in Chapter 2 (2.7.1),  $\alpha$  is defined to be a regular function on  $X$  if and only if it is an element of the coordinate algebra  $A$ . The next lemma shows that the two conditions are equivalent.

**3.5.2. Lemma.** *The regular functions on an affine variety  $X = \text{Spec } A$ , as defined in (3.4.2), are the elements of its coordinate algebra  $A$ .*

*proof.* Let  $\alpha$  be a regular function on  $X$ , as defined above. So for every point  $p$  of  $X$ , there is a localization  $X_s = \text{Spec } A_s$  that contains  $p$ , such that  $\alpha$  is an element of  $A_s$ . Because  $X$  is quasicompact, a finite set of these localizations, say  $X_{s_1}, \dots, X_{s_k}$ , will cover  $X$ . Then  $s_1, \dots, s_k$  have no common zeros on  $X$ , so they generate the unit ideal of  $A$ . Since  $\alpha$  is in  $A_{s_i}$ , we can write  $\alpha = s_i^{-n} b_i$  with  $b_i$  in  $A$ , and we can use the same exponent  $n$  for all  $i$ . Since the elements  $s_i$  generate the unit ideal of  $A$ , so do the powers  $s_i^n$ . Say that  $\sum s_i^n a_i = 1$ , with  $a_i$  in  $A$ . Then  $\alpha = \sum s_i^n a_i \alpha = \sum a_i b_i$  is in  $A$ .  $\square$

#### 3.5.3. Proposition.

(i) *Let  $R$  be the algebra of regular functions on a variety  $Y$ , and let  $A$  be a finite-type domain. A homomorphism  $A \rightarrow R$  defines a morphism  $Y \xrightarrow{f} \text{Spec } A$ .*

(ii) *When  $X$  and  $Y$  are affine varieties, say  $X = \text{Spec } A$  and  $Y = \text{Spec } B$ , morphisms  $Y \rightarrow X$ , as defined in Definition 3.4.14 correspond bijectively to algebra homomorphisms  $A \rightarrow B$ , as in Definition 2.7.5.*

**Note.** Since  $Y$  isn't affine, all that we know about the algebra  $R$  is that its elements are rational functions that are regular on  $Y$ .

*proof of Proposition 3.5.3. (i)* Let  $\{Y^i\}$  be an affine open covering of  $Y$ , and let  $R_i$  be the coordinate algebra of  $Y^i$ . The inclusions  $A \subset R \subset R_i$  define morphisms  $Y^i = \text{Spec } R_i \xrightarrow{f^i} \text{Spec } A$ . It is true that  $f^i = f^j$  on  $Y^i \cap Y^j$ , so Lemma 3.4.18 (iii) applies.  $\square$

**3.5.4. Lemma.** *Let  $X$  and  $Y$  be affine varieties, say  $X = \text{Spec } A$  and  $Y = \text{Spec } B$ . Morphisms  $Y \xrightarrow{\alpha} X$ , as defined in (3.4.11) and (3.4.14), correspond bijectively to algebra homomorphisms  $A \xrightarrow{\varphi} B$ .*

*proof.* We choose a presentation of  $A$ , to embed  $X$  as a closed subvariety of affine space, and we identify that affine space with the standard affine open set  $\mathbb{U}^0$  of  $\mathbb{P}^n$ . Let  $K$  be the function field of  $Y$  – the field of fractions of  $B$ . A morphism  $Y \xrightarrow{u} X$  is determined by a good point  $\alpha$  with values in  $K$ , and since  $\alpha_0 \neq 0$ , we may suppose that this point has the form  $\alpha = (\alpha_0, \dots, \alpha_n)$ . Then the rational functions  $\alpha_i/\alpha_0 = \alpha_i$  will be regular at every point of  $Y$ . So they are elements of  $B$ . The coordinate algebra  $A$  of  $X$  is generated by the residues of the coordinate variables  $x_1, \dots, x_n$ , with  $x_0 = 1$ . Sending  $x_i \rightarrow \alpha_i$  defines a homomorphism

$A \xrightarrow{\varphi} B$ . Conversely, if  $\varphi$  is such a homomorphism, the good point that defines the morphism  $Y \xrightarrow{u} X$  is  $(1, \varphi(x_1), \dots, \varphi(x_n))$ .  $\square$

### (3.5.5) affine open subsets

An *affine open subset* of a variety  $X$  is an open subset that is an affine variety. If  $V$  is a nonempty open subset of  $X$  and  $R$  is the algebra of rational functions that are regular on  $V$ , then  $V$  is an affine open subset if and only if  $R$  is a finite-type domain, and  $V$  is isomorphic to  $\text{Spec } R$ .

**3.5.6. Proposition.** *The complement of a hypersurface is an affine open subvariety of  $\mathbb{P}^n$ .*

*proof.* Let  $V$  be the complement of the hypersurface  $\{f = 0\}$ , where  $f$  is an irreducible homogeneous polynomial of degree  $d$ , let  $R$  be the algebra of regular functions on  $V$ , and let  $K$  be its fraction de field.

The regular functions on  $V$  are the homogeneous fractions of degree zero of the form  $g/f^k$  (3.4.34), and the fractions  $m/f$ , where  $m$  is a monomial of degree  $d$ , generate  $R$ . Since there are finitely many monomials of degree  $d$ ,  $R$  is a finite-type domain. Let  $w$  be an arbitrary monomial of degree  $d-1$ , and let  $s_i = x_i w/f$ . The point  $(x_0, \dots, x_n)$  of  $V$  can also be written as  $(s_0, \dots, s_n)$ , and the fractions  $s_i$  are among the generators for  $R$ . So  $(s_0, \dots, s_n)$  is a point of  $W = \text{Spec } R$  with values in  $K$  that defines a morphism  $W \xrightarrow{z} V$ . We show that  $z$  is an isomorphism.

**3.5.7. Lemma.** *Let  $U^i$  be the standard affine open subset of  $\mathbb{P}^n$ . With  $s_i$  as above, the intersection  $V^i = V \cap U^i$  is isomorphic to the localization  $W_{s_i}$  of  $W$ .*

*proof.* Say that  $i = 0$ , and let  $s = s_0 = x_0^d/f$  and  $t = s^{-1} = f/x_0^d$ . Let  $A$  be the coordinate algebra of  $U^0$ . Then  $V^0 = V \cap U^0$  is the set of points of  $U^0$  at which  $t$  isn't zero. Its coordinate algebra is the localization  $A_t$ , and  $V^0$  is the affine variety  $\text{Spec } A_t$ .

It suffices to show that  $A_t = R_s$ . With coordinates  $u_j = x_j/x_0$  for  $U^0$ , a fraction  $m/f$ , where  $m = x_{j_1} \cdots x_{j_d}$ , can be written as  $(u_{j_1} \cdots u_{j_d})/t$ . These fractions generate  $R$ , so  $R \subset A_t$ , and since  $s^{-1} = t$  is in  $A_t$ ,  $R_s \subset A_t$ . For the other inclusion, we write  $u_j = (x_j x_0^{d-1}/f) s^{-1}$ . This is an element of  $R_s$  because  $x_j x_0^{d-1}/f$  is in  $R$ . Therefore  $A \subset R_s$  and  $A_t \subset R_s$ . So  $A_t = R_s$ , as claimed.  $\square$

We go back to the proof of Proposition 3.5.6. The sets  $V^i = V \cap U^i$  for  $i = 0, \dots, n$  cover  $V$ , and the morphism  $z$  restricts to an isomorphism  $V^i \rightarrow \text{Spec } R_{s_i}$ . So the morphism  $z$  defined above is an isomorphism (3.4.21).  $\square$

**3.5.8. Lemma.** *The affine open subsets of a variety  $X$  form a basis for the topology on  $X$ .*  $\square$

**3.5.9. Theorem** *Let  $U$  and  $V$  be affine open subvarieties of a variety  $X$ , say  $U \approx \text{Spec } A$  and  $V \approx \text{Spec } B$ . The intersection  $U \cap V$  is an affine open subvariety. Its coordinate algebra is generated by the two algebras  $A$  and  $B$ .*

*proof.* We will denote the algebra generated by two subalgebras  $A$  and  $B$  of the function field  $K$  of  $X$  by  $[A, B]$ . The elements of  $[A, B]$  are finite sums of products of elements of  $A$  and  $B$ . If  $A = \mathbb{C}[a]$ ,  $a = a_1, \dots, a_r$ , and  $B = \mathbb{C}[b]$ ,  $b = b_1, \dots, b_s$ , then  $[A, B]$  is the finite-type subalgebra of  $K$  generated by the set  $a \cup b$ .

Let  $R = [A, B]$  and let  $W = \text{Spec } R$ . We are to show that  $W$  is isomorphic to  $U \cap V$ . The inclusions of coordinate algebras  $A \rightarrow R$  and  $B \rightarrow R$  give us morphisms  $W \rightarrow U$  and  $W \rightarrow V$ . We also have inclusions  $U \subset X$  and  $V \subset X$ , and  $X$  is a subvariety of a projective space  $\mathbb{P}^n$ .

Let  $\alpha$  be the point of  $\mathbb{P}^n$  with values in  $K$  that defines the projective embedding  $X \xrightarrow{\varphi} \mathbb{P}^n$ . The varieties  $U, V, W, X$  have the same function field  $K$ , and  $\alpha$  also defines the morphisms  $U \rightarrow \mathbb{P}^n$  and  $V \rightarrow \mathbb{P}^n$  obtained by restricting the embedding  $\varphi$  of  $X$ .

The maps from  $U$  and  $V$  to  $\mathbb{P}^n$  defined by  $\alpha$  are restrictions of  $\varphi$ . The variety  $W$  also has function field  $K$ , and  $\alpha$  defines a morphism  $W \xrightarrow{\psi} \mathbb{P}^n$  that can be obtained either as the composition  $W \rightarrow U \rightarrow X$ , or as  $W \rightarrow V \rightarrow X$ . So its image is in  $U \cap V$ . This gives us a morphism  $W \xrightarrow{\epsilon} U \cap V$ . We show that  $\epsilon$  is an isomorphism.

Let  $p$  be a point of  $U \cap V$ . We choose an affine open subset  $Z$  of  $U \cap V$  that is a localization of  $U$  and of  $V$ , and that contains  $p$  (2.6.2)(ii). Let  $S$  be the coordinate ring of  $Z$ . So  $S = A_s$  for some nonzero  $s$  in  $A$  and also  $S = B_t$  for some nonzero  $t$  in  $B$ . Then

$$R_s = [A, B]_s = [A_s, B] = [B_t, B] = B_t = S$$



So  $\epsilon$  maps the localization  $W_s = \text{Spec } R_s$  of  $W$  isomorphically to the open subset  $Z$  of  $U \cap V$ . Since we can cover  $U \cap V$  by open sets such as  $Z$ , Lemma 3.4.18 (ii) shows that  $\epsilon$  is an isomorphism.  $\square$

### 3.6 Lines in Projective Three-Space

The *Grassmanian*  $G(m, n)$  is a variety whose points correspond to subspaces of dimension  $m$  of the vector space  $\mathbb{C}^n$ , and to linear subspaces of dimension  $m - 1$  of  $\mathbb{P}^{n-1}$ . One says that  $G(m, n)$  *parametrizes* those subspaces. For example, the Grassmanian  $G(1, n+1)$  is the projective space  $\mathbb{P}^n$ . Points of  $\mathbb{P}^n$  parametrize one-dimensional subspaces of  $\mathbb{C}^{n+1}$ .

The Grassmanian  $G(2, 4)$  parametrizes two-dimensional subspaces of  $\mathbb{C}^4$ , and lines in  $\mathbb{P}^3$ . In this section we describe this Grassmanian, which we denote by  $\mathbb{G}$ . The point of  $\mathbb{G}$  that corresponds to a line  $\ell$  in  $\mathbb{P}^3$  will be denoted by  $[\ell]$ .

One can get some insight into the structure of  $\mathbb{G}$  using row reduction. Let  $V = \mathbb{C}^4$ , let  $u_1, u_2$  be a basis of a two-dimensional subspace  $U$  of  $V$  and let  $M$  be the  $2 \times 4$  matrix whose rows are  $u_1, u_2$ . The rows of the matrix  $M'$  obtained from  $M$  by row reduction span the same space  $U$ , and the row-reduced matrix  $M'$  is uniquely determined by  $U$ . Provided that the left hand  $2 \times 2$  submatrix of  $M$  is invertible,  $M'$  will have the form

$$(3.6.1) \quad M' = \begin{pmatrix} 1 & 0 & * & * \\ 0 & 1 & * & * \end{pmatrix}$$

So the Grassmanian  $\mathbb{G}$  contains, as an open subset, a four-dimensional affine space whose coordinates are the variable entries of  $M'$ .

In any  $2 \times 4$  matrix  $M$  with independent rows, some pair of columns will be independent. Those columns can be used in place of the first two in a row reduction. So  $\mathbb{G}$  is covered by six four-dimensional affine spaces that we denote by  $\mathbb{W}^{ij}$ ,  $1 \leq i < j \leq 4$ ,  $\mathbb{W}^{ij}$  being the space of  $2 \times 4$  matrices such that  $\text{column}_i = (1, 0)^t$  and  $\text{column}_j = (0, 1)^t$ . Since  $\mathbb{P}^4$  and the Grassmanian are both covered by affine spaces of dimension four, they may seem similar, but they aren't the same.

#### (3.6.2) the exterior algebra

Let  $V$  be a complex vector space. The *exterior algebra*  $\bigwedge V$  (read 'wedge  $V$ ') is a noncommutative ring that contains the complex numbers and is generated by the elements of  $V$ , with the relations

$$(3.6.3) \quad vw = -wv \quad \text{for all } v, w \text{ in } V.$$

**3.6.4. Lemma.** *The condition (3.6.3) is equivalent with:  $vv = 0$  for all  $v$  in  $V$ .*

*proof.* To get  $vv = 0$  from (3.6.3), one sets  $w = v$ . Suppose that  $vv = 0$  for all  $v$  in  $V$ . Then  $(v+w)(v+w) = vv + ww = 0$ , and since  $(v+w)(v+w) = vv + vw + wv + ww$ ,  $vw + wv = 0$ .  $\square$

To familiarize yourself with computation in  $\bigwedge V$ , verify that  $v_2 v_3 v_1 v_4 = v_1 v_2 v_3 v_4$  and that  $v_2 v_3 v_4 v_1 = -v_1 v_2 v_3 v_4$ .

Let  $\bigwedge^r V$  denote the subspace of  $\bigwedge V$  spanned by products of length  $r$  of elements of  $V$ . The exterior algebra  $\bigwedge V$  is the direct sum of the subspaces  $\bigwedge^r V$ . An algebra  $A$  that is a direct sum of subspaces  $A^i$ , and such that multiplication maps  $A^i \times A^j$  to  $A^{i+j}$  is called a *graded algebra*. Since its multiplication law isn't commutative, the exterior algebra is a noncommutative graded algebra.

**3.6.5. Proposition.** *If  $(v_1, \dots, v_n)$  is a basis for  $V$ , the products  $v_{i_1} \cdots v_{i_r}$  of length  $r$  with increasing indices  $i_1 < i_2 < \cdots < i_r$  form a basis for  $\bigwedge^r V$ .*

The proof is at the end of the section.

**3.6.6. Corollary.** *Let  $v_1, \dots, v_r$  be elements of  $V$ . The product  $v_1 \cdots v_r$  in  $\bigwedge^r V$  is zero if and only if the set  $(v_1, \dots, v_r)$  is dependent.*  $\square$

For the rest of the section, we let  $V$  be a vector space of dimension four with basis  $(v_1, \dots, v_4)$ . Proposition 3.6.5 tells us that

(3.6.7)

$\bigwedge^0 V = \mathbb{C}$  is a space of dimension 1, with basis  $\{1\}$

$\bigwedge^1 V = V$  is a space of dimension 4, with basis  $\{v_1, v_2, v_3, v_4\}$

$\bigwedge^2 V$  is a space of dimension 6, with basis  $\{v_i v_j \mid i < j\} = \{v_1 v_2, v_1 v_3, v_1 v_4, v_2 v_3, v_2 v_4, v_3 v_4\}$

$\bigwedge^3 V$  is a space of dimension 4, with basis  $\{v_i v_j v_k \mid i < j < k\} = \{v_1 v_2 v_3, v_1 v_2 v_4, v_1 v_3 v_4, v_2 v_3 v_4\}$

$\bigwedge^4 V$  is a space of dimension 1, with basis  $\{v_1 v_2 v_3 v_4\}$

$\bigwedge^q V = 0$  when  $q > 4$

The elements of  $\bigwedge^2 V$  are combinations

$$(3.6.8) \quad w = \sum_{i < j} a_{ij} v_i v_j$$

We regard  $\bigwedge^2 V$  as an affine space of dimension 6, identifying the combination  $w$  with the vector whose coordinates are the six coefficients  $a_{ij}$  ( $i < j$ ). We use the same symbol  $w$  to denote the point of the projective space  $\mathbb{P}^5$  with those coordinates:  $w = (a_{12}, a_{13}, a_{14}, a_{23}, a_{24}, a_{34})$ .

**3.6.9. Definition.** An element  $w$  of  $\bigwedge^2 V$  is *decomposable* if it is a product of two elements of  $V$ .

**3.6.10. Proposition.** The decomposable elements  $w = \sum_{i < j} a_{ij} v_i v_j$  of  $\bigwedge^2 V$  are those such that  $ww = 0$ , and the relation  $ww = 0$  is given by the following equation in the coefficients  $a_{ij}$ :

$$(3.6.11) \quad a_{12}a_{34} - a_{13}a_{24} + a_{14}a_{23} = 0$$

*proof.* If  $w$  is decomposable, say  $w = u_1 u_2$ , then  $w^2 = u_1 u_2 u_1 u_2 = -u_1^2 u_2^2$  is zero because  $u_1^2 = 0$ . For the converse, we compute  $w^2$  when  $w = \sum_{i < j} a_{ij} v_i v_j$ . The answer is

$$ww = 2(a_{12}a_{34} - a_{13}a_{24} + a_{14}a_{23})v_1 v_2 v_3 v_4$$

To show that  $w$  is decomposable if  $w^2 = 0$ , it seems simplest to factor  $w$  explicitly. Since the assertion is trivial when  $w = 0$ , we may suppose that some coefficient of  $w$ , say  $a_{12}$ , is nonzero. Then if  $w^2 = 0$ ,  $w$  is the product

$$(3.6.12) \quad w = \frac{1}{a_{12}}(a_{12}v_2 + a_{13}v_3 + a_{14}v_4)(-a_{12}v_1 + a_{23}v_3 + a_{24}v_4) \quad \square$$

**3.6.13. Corollary. (i)** Let  $w$  be a nonzero decomposable element of  $\bigwedge^2 V$ , say  $w = u_1 u_2$ , with  $u_i$  in  $V$ . Then  $(u_1, u_2)$  is a basis for a two-dimensional subspace of  $V$ .

**(ii)** If  $(u_1, u_2)$  and  $(u'_1, u'_2)$  are bases for the same subspace  $U$  of  $V$ , then  $w = u_1 u_2$  and  $w' = u'_1 u'_2$  differ by a scalar factor. Their coefficients represent the same point of  $\mathbb{P}^5$ .

**(iii)** Let  $u_1, u_2$  be a basis for a two-dimensional subspace  $U$  of  $V$ , and let  $w = u_1 u_2$ . The rule  $\epsilon(U) = w$  defines a bijection  $\epsilon$  from  $\mathbb{G}$  to the quadric  $Q$  in  $\mathbb{P}^5$  whose equation is (3.6.11).

Thus  $\mathbb{G}$  can be represented as the quadric (3.6.11).

*proof. (i)* If an element  $w$  of  $\bigwedge^2 V$  is decomposable, say  $w = u_1 u_2$ , and if  $w$  is nonzero, then  $u_1$  and  $u_2$  must be independent (3.6.6). They span a two-dimensional subspace.

**(ii)** When we write the second basis in terms of the first one, say  $(u'_1, u'_2) = (au_1 + bu_2, cu_1 + du_2)$ , the product  $u'_1 u'_2$  becomes  $(ad - bc)u_1 u_2$ , and  $ad - bc \neq 0$ .

**(iii)** In view of **(i)** and **(ii)**, all that remains to show is that, if  $(u_1, u_2)$  and  $(u'_1, u'_2)$  are bases for distinct two-dimensional subspaces  $U$  and  $U'$ , then  $u_1 u_2 \neq u'_1 u'_2$  in  $\bigwedge^2 V$ .

Since  $U \neq U'$ , the intersection  $W = U \cap U'$  has dimension at most 1, so at least three of the vectors  $u_1, u_2, u'_1, u'_2$  will be independent. Therefore  $u_1 u_2 \neq u'_1 u'_2$ .  $\square$

For the rest of this section, we use the algebraic dimension of a variety, a concept that will be studied in the next chapter. We refer to the algebraic dimension simply as the *dimension*. The *dimension* of a variety  $X$  can be defined as the length  $d$  of the longest chain  $C_0 > C_1 > \cdots > C_d$  of closed subvarieties of  $X$ .

As was mentioned in Chapter 1, the topological dimension of  $X$  its dimension in the classical topology, is always twice the algebraic dimension. Because the Grassmanian  $\mathbb{G}$  is covered by affine spaces of dimension 4, its algebraic dimension is 4 and its topological dimension is 8.

**3.6.14. Proposition.** *Let  $\mathbb{P}^3$  be the projective space associated to a four dimensional vector space  $V$ . In the product  $\mathbb{P}^3 \times \mathbb{G}$ , the locus  $\Gamma$  of pairs  $p, [\ell]$  such that the point  $p$  of  $\mathbb{P}^3$  lies on the line  $\ell$  is a closed subset of dimension 5.*

*proof.* Let  $\ell$  be the line in  $\mathbb{P}^3$  that corresponds to the subspace  $U$  with basis  $(u_1, u_2)$ , and say that  $p$  represented by the vector  $x$  in  $V$ . Let  $w = u_1 u_2$ . Then  $p \in \ell$  means  $x \in U$ , which is true if and only if  $(x, u_1, u_2)$  is a dependent set, and this happens if and only if  $xw = 0$  (3.6.5). So  $\Gamma$  is the closed subset of points  $(x, w)$  of  $\mathbb{P}^3 \times \mathbb{P}^5$  defined by the bihomogeneous equations  $w^2 = 0$  and  $xw = 0$ .

When we project  $\Gamma$  to  $\mathbb{G}$ , The fibre over a point  $[\ell]$  of  $\mathbb{G}$  is the set of points  $p, [\ell]$  such that  $p$  is a point of the line  $\ell$ . The fibre over the point  $[\ell]$  of  $\mathbb{G}$  is the line  $\ell$ . Thus  $\Gamma$  can be viewed as a family of lines, parametrized by the four-dimensional variety  $\mathbb{G}$ . Its dimension is  $\dim \ell + \dim \mathbb{G} = 1 + 4 = 5$ .  $\square$

### (3.6.15) lines on a surface

One may ask whether or not a given surface in  $\mathbb{P}^3$  contains a line. One surface that contains lines is the quadric  $Q$  in  $\mathbb{P}^3$  with equation  $w_{01}w_{10} = w_{00}w_{11}$ , the image of the Segre embedding  $\mathbb{P}^1 \times \mathbb{P}^1 \rightarrow \mathbb{P}_w^3$  (3.1.7). It contains two families of lines, corresponding to the two “rulings”  $p \times \mathbb{P}^1$  and  $\mathbb{P}^1 \times q$  of  $\mathbb{P}^1 \times \mathbb{P}^1$ . There are surfaces of arbitrary degree that contain lines, but, that a generic surface of degree four or more doesn’t contain any line.

We use coordinates  $x_i$  with  $i = 1, 2, 3, 4$  for  $\mathbb{P}^3$  here. There are  $N = \binom{d+3}{3}$  monomials of degree  $d$  in four variables, so homogeneous polynomials of degree  $d$  are parametrized by an affine space of dimension  $N$ , and surfaces of degree  $d$  in  $\mathbb{P}^3$  by a projective space of dimension  $N - 1$ . Let  $\mathbb{S}$  denote that projective space, and let  $[S]$  denote the point of  $\mathbb{S}$  that corresponds to a surface  $S$ . The coordinates of  $[S]$  are the coefficients of the monomials in the defining polynomial  $f$  of  $S$ . Speaking informally, we say that a point of  $\mathbb{S}$  “is” a surface of degree  $d$  in  $\mathbb{P}^3$ . (When  $f$  is reducible, its zero locus isn’t a variety. Let’s not worry about this.)

Consider the line  $\ell_0$  defined by  $x_3 = x_4 = 0$ . Its points are those of the form  $(x_1, x_2, 0, 0)$ , so a surface  $S : \{f = 0\}$  will contain  $\ell_0$  if and only if  $f(x_1, x_2, 0, 0) = 0$  for all  $x_1, x_2$ . Substituting  $x_3 = x_4 = 0$  into  $f$  leaves us with a polynomial in two variables:

$$(3.6.16) \quad f(x_1, x_2, 0, 0) = c_0 x_1^d + c_1 x_1^{d-1} x_2 + \cdots + c_d x_2^d,$$

where the coefficients  $c_i$  are among the coefficients of the polynomial  $f$ . If  $f(x_1, x_2, 0, 0)$  is identically zero, all of its coefficients must be zero. So the surfaces that contain  $\ell_0$  correspond to the points of the linear subspace  $\mathbb{L}_0$  of  $\mathbb{S}$  defined by the equations  $c_0 = \cdots = c_d = 0$ . Its dimension is  $(N - 1) - (d + 1) = N - d - 2$ . This is a satisfactory answer to the question of which surfaces contain  $\ell_0$ , and we can use it to make a guess about lines in a generic surface of degree  $d$ .

**3.6.17. Lemma.** *In the product variety  $\mathbb{G} \times \mathbb{S}$ , the set  $\Gamma$  of pairs  $[\ell], [S]$  such that  $\ell \subset S$  is a closed subset.*

*proof.* Let  $\mathbb{W}^{ij}$ ,  $1 \leq i < j \leq 4$  denote the six affine spaces that cover the Grassmanian, as at the beginning of this section. It suffices to show that the intersection  $\Gamma^{ij} = \Gamma \cap (\mathbb{W}^{ij} \times \mathbb{S})$  is closed in  $\mathbb{W}^{ij} \times \mathbb{S}$  (3.4.17). We inspect the case  $i, j = 1, 2$ .

A line  $\ell$  such that  $[\ell]$  is in  $\mathbb{W}^{12}$  corresponds to a subspace of  $\mathbb{C}^2$  with basis of the form  $u_1 = (1, 0, a_2, a_3)$ ,  $u_2 = (0, 1, b_2, b_3)$  and  $\ell$  is the line  $\{ru_1 + su_2\}$ . Let  $f(x_1, x_2, x_3, x_4)$  be the polynomial that defines a surface  $S$ . The line  $\ell$  is contained in  $S$  if and only if  $f(r, s, ra_2 + sb_2, ra_3 + sb_3)$  is zero for all  $r$  and  $s$ . This is a homogeneous polynomial of degree  $d$  in  $r, s$ . Let’s call it  $\tilde{f}(r, s)$ . If we write  $\tilde{f}(r, s) = z_0 r^d + z_1 r^{d-1} s + \cdots + z_d s^d$ , the coefficients  $z_\nu$  will be polynomials in  $a_i, b_i$  and in the coefficients of  $f$ . The locus  $z_0 = \cdots = z_d = 0$  is the closed set  $\Gamma^{12}$  of  $\mathbb{W}^{12} \times \mathbb{S}$ .  $\square$

The set of surfaces that contain our special line  $\ell_0$  corresponds to the linear space  $\mathbb{L}_0$  of  $\mathbb{S}$  of dimension  $N - d - 2$ , and  $\ell_0$  can be carried to any other line  $\ell$  by a linear map  $\mathbb{P}^3 \rightarrow \mathbb{P}^3$ . So the surfaces that contain another line  $\ell$  also form a linear subspace of  $\mathbb{S}$  of dimension  $N - d - 2$ . They are the fibres of  $\Gamma$  over  $\mathbb{G}$ . The dimension of the Grassmanian  $\mathbb{G}$  is 4. Therefore the dimension of  $\Gamma$  is  $\dim \Gamma = \dim \mathbb{L}_0 + \dim \mathbb{G} = (N - d - 2) + 4$ . Since  $\mathbb{S}$  has dimension  $N - 1$ ,

$$(3.6.18) \quad \dim \Gamma = \dim \mathbb{S} - d + 3.$$

We project the product  $\mathbb{G} \times \mathbb{S}$  and its subvariety  $\Gamma$  to  $\mathbb{S}$ . The fibre of  $\Gamma$  over a point  $[S]$  is the set of pairs  $[\ell], [S]$  such that  $\ell$  is contained in  $S$  – the set of lines in  $S$ .

When the degree  $d$  of the surfaces we are studying is 1,  $\dim \Gamma = \dim \mathbb{S} + 2$ . Every fibre of  $\Gamma$  over  $\mathbb{S}$  will have dimension at least 2. In fact, every fibre has dimension equal to 2. Surfaces of degree 1 are planes, and the lines in a plane form a two-dimensional family.

When  $d = 2$ ,  $\dim \Gamma = \dim \mathbb{S} + 1$ . We can expect that most fibres of  $\Gamma$  over  $\mathbb{S}$  will have dimension 1. This is true: A smooth quadric contains two one-dimensional families of lines. (All smooth quadrics are equivalent with the quadric (3.1.8).) But if a quadratic polynomial  $f(x_1, x_2, x_3, x_4)$  is the product of linear polynomials, its locus of zeros will be a union of planes. It will contain two-dimensional families of lines. Some fibres have dimension 2.

When  $d \geq 4$ ,  $\dim \Gamma < \dim \mathbb{S}$ . The projection  $\Gamma \rightarrow \mathbb{S}$  cannot be surjective. Most surfaces of degree 4 or more contain no lines.

The most interesting case is that  $d = 3$ . In this case,  $\dim \Gamma = \dim \mathbb{S}$ . Most fibres will have dimension zero. They will be finite sets. In fact, a generic cubic surface contains 27 lines. We have to wait to see why the number is precisely 27 (see Theorem 4.9.17).

Our conclusions are intuitively plausible, but to be sure about them, we need to study dimension carefully. We do this in the next chapters.

*proof of Proposition 3.6.5.* Let  $v = (v_1, \dots, v_n)$  be a basis of the vector space  $V$ . The proposition asserts that the products  $v_{i_1} \cdots v_{i_r}$  of length  $r$  with increasing indices  $i_1 < i_2 < \cdots < i_r$  form a basis for  $\bigwedge^r V$ .

To prove this, we need to be more precise about the definition of the exterior algebra  $\bigwedge V$ . We start with the algebra  $T(V)$  of noncommutative polynomials in the basis  $v$ , which is also called the *tensor algebra* on  $V$ . The part  $T^r(V)$  of  $T(V)$  of degree  $r$  has as basis the  $n^r$  noncommutative monomials of degree  $r$ , products  $v_{i_1} \cdots v_{i_r}$  of length  $r$  of elements of the basis  $v$ . Its dimension is  $n^r$ . When  $n = r = 2$ ,  $T^2(V)$  the basis is  $(x_1^2, x_1x_2, x_2x_1, x_2^2)$ .

The exterior algebra  $\bigwedge V$  is the quotient of  $T(V)$  obtained by forcing the relations  $vw + wv = 0$  (3.6.3). Using the distributive law, one sees that the relations  $v_i v_j + v_j v_i = 0$ ,  $1 \leq i, j \leq n$ , are sufficient to define this quotient. The relations  $v_i v_i = 0$  are included when  $i = j$ .

To obtain  $\bigwedge^r V$ , we multiply the relations  $v_i v_j + v_j v_i$  on left and right by arbitrary noncommutative monomials  $p$  and  $q$  in  $v_1, \dots, v_n$  whose degrees add to  $r - 2$ . The noncommutative polynomials

$$(3.6.19) \quad p(v_i v_j + v_j v_i)q$$

span the kernel of the linear map  $T^r(V) \rightarrow \bigwedge^r V$ . So in  $\bigwedge^r V$ ,  $p(v_i v_j)q = -p(v_j v_i)q$ . Using these relations, any product  $v_{i_1} \cdots v_{i_r}$  in  $\bigwedge^r V$  is, up to sign, equal to a product in which the elements  $v_{i_\nu}$  are listed in increasing order. Thus the products with indices in increasing order span  $\bigwedge^r V$ , and because  $v_i v_i = 0$ , such a product will be zero unless the indices are strictly increasing.

We go to the proof now. Let  $v = (v_1, \dots, v_n)$  be a basis for  $V$ . We show first that the product  $w = v_1 \cdots v_n$  in increasing order of the basis elements of  $V$  is a basis of  $\bigwedge^n V$ . We have shown that this product spans  $\bigwedge^n V$ , and it remains to show that  $w \neq 0$ , or that  $\bigwedge^n V \neq 0$ .

Let's use multi-index notation:  $(i) = (i_1, \dots, i_r)$ , and  $v_{(i)} = v_{i_1} \cdots v_{i_r}$ . We define a surjective linear map  $T^n(V) \xrightarrow{\varphi} \mathbb{C}$  on the basis of  $T^n(V)$  of products  $v_{(i)} = (v_{i_1} \cdots v_{i_n})$  of length  $n$ . If there is no repetition among the indices  $i_1, \dots, i_n$ , then  $(i)$  will be a permutation of the indices  $1, \dots, n$ . In that case, we set  $\varphi(v_{(i)}) = \varphi(v_{i_1} \cdots v_{i_n}) = \text{sign}(i)$ . If there is a repetition, we set  $\varphi(v_{(i)}) = 0$ .

Let  $p$  and  $q$  be noncommutative monomials whose degrees add to  $n - 2$ . If the product  $p(v_i v_j)q$  has no repeated index, the indices in  $p(v_i v_j)q$  and  $p(v_j v_i)q$  will be permutations of  $1, \dots, n$ , and those permutations will have opposite signs. Then  $p(v_i v_j + v_j v_i)q$  will be in the kernel of  $\varphi$ . Since these elements span the space of relations,  $\varphi$  defines a surjective linear map  $\bigwedge^n V \rightarrow \mathbb{C}$ . Therefore  $\bigwedge^n V \neq 0$ .

To prove (3.6.5), we must show that for  $r \leq n$ , the products  $v_{i_1} \cdots v_{i_r}$  with  $i_1 < i_2 < \cdots < i_r$  form a basis for  $\bigwedge^r V$ , and we know that those products span  $\bigwedge^r V$ . We must show that they are independent. Suppose that a combination  $z = \sum c_{(i)} v_{(i)}$  is zero, the sum being over sets of strictly increasing indices. We choose a

set  $(j_1, \dots, j_r)$  of strictly increasing indices, and we let  $(k) = (k_1, \dots, k_{n-r})$  be the set of indices not occurring in  $(j)$ , listed in arbitrary order. Then all terms in the sum  $zv_{(k)} = \sum c_{(i)}v_{(i)}v_{(k)}$  will be zero except the term with  $(i) = (j)$ . On the other hand, since  $z = 0$ ,  $zv_{(k)} = 0$ . Therefore  $c_{(j)}v_{(j)}v_{(k)} = 0$ , and since  $v_{(j)}v_{(k)}$  differs by sign from  $v_1 \cdots v_n$ , it isn't zero. It follows that  $c_{(j)} = 0$ . This is true for all  $(j)$ , so  $z = 0$ .  $\square$



## Chapter 4 INTEGRAL MORPHISMS OF AFFINE VARIETIES

- 4.1 The Nakayama Lemma
- 4.2 Integral Extensions
- 4.3 Normalization
- 4.4 Geometry of Integral Morphisms
- 4.5 Finite Group Actions II
- 4.6 Chevalley's Finiteness Theorem
- 4.7 Dimension
- 4.8 Krull's Theorem
- 4.9 Double Planes

The concept of an algebraic integer was one of the most important ideas contributing to the development of algebraic number theory in the 19th century. Then, largely through the work of Noether and Zariski, an analog was seen to be essential in algebraic geometry. We study this analog here.

### 4.1 The Nakayama Lemma

#### (4.1.1) eigenvectors

It won't surprise you that eigenvectors are important, but the way that they are used to study modules may be unfamiliar.

Let  $P$  be an  $n \times n$  matrix with entries in a ring  $A$ . The concept of an eigenvector for  $P$  makes sense when the entries of a vector are in an  $A$ -module. A column vector  $v = (v_1, \dots, v_n)^t$  with entries in a module  $M$  is an *eigenvector* of  $P$  with *eigenvalue*  $\lambda$  if  $Pv = \lambda v$ .

When the entries of a vector are in a module, it becomes hard to adapt the usual requirement that an eigenvector must be nonzero, so we drop it, though the zero eigenvector tells us nothing.

**4.1.2. Lemma.** *Let  $p(t)$  be the characteristic polynomial  $\det(tI - P)$  of a square matrix  $P$ . If  $v$  is an eigenvector of  $P$  with eigenvalue  $\lambda$ , then  $p(\lambda)v = 0$ .*

The usual proof, in which one multiplies the equation  $(\lambda I - P)v = 0$  by the cofactor matrix of  $(\lambda I - P)$ , carries over.  $\square$

This lemma is a cornerstone of the theory of modules:

**4.1.3. Nakayama Lemma.** Let  $M$  be a finite module over a ring  $A$ , and let  $J$  be an ideal of  $A$  such that  $M = JM$ . There is an element  $z$  in  $J$  such that  $m = zm$  for all  $m$  in  $M$ , or such that  $(1 - z)M = 0$ .

It is always true that  $M \supset JM$ , so the hypothesis  $M = JM$  can be replaced by  $M \subset JM$ .

*proof.* By definition,  $JM$  denotes the set of (finite) sums  $\sum a_i m_i$  with  $a_i$  in  $J$  and  $m_i$  in  $M$ .

Let  $v_1, \dots, v_n$  be generators for the finite  $A$ -module  $M$ , and let  $v$  be the vector  $(v_1, \dots, v_n)^t$ . The equation  $M = JM$  tells us that there are elements  $p_{ij}$  in  $J$  such that  $v_i = \sum p_{ij} v_j$ . In matrix notation,  $v = Pv$ . So

$v$  is an eigenvector of  $P$  with eigenvalue 1, and if  $p(t)$  is the characteristic polynomial of  $P$ , then  $p(1)v = 0$ . Since the entries of  $P$  are in  $J$ , inspection of the determinant of  $I - P$  shows that  $p(1)$  has the form  $1 - z$ , with  $z$  in  $J$ . Then  $(1 - z)v_i = 0$  for all  $i$ . Since  $v_1, \dots, v_n$  generate  $M$ ,  $(1 - z)M = 0$ .  $\square$

**4.1.4. Corollary.** *With notation as in the Nakayama Lemma, let  $s = 1 - z$ , so that  $sM = 0$ . The localized module  $M_s$  is the zero module.*

**4.1.5. Corollary.** (i) *Let  $I$  and  $J$  be ideals of a noetherian domain  $A$ . If  $I = JI$ , then either  $I$  is the zero ideal or  $J$  is the unit ideal.*

(ii) *Let  $A \subset B$  be rings, and suppose that  $B$  is a finite  $A$ -module. If  $J$  is an ideal of  $A$ , and if the extended ideal  $JB$  is the unit ideal of  $B$ , then  $J$  is the unit ideal of  $A$ .*

(iii) *Let  $x$  be an element of a noetherian domain  $A$ , not a unit and let  $J$  be the ideal  $xA$ . The intersection  $\bigcap J^n$  is the zero ideal. Therefore, if  $y$  is a nonzero element of  $A$ , the integers  $k$  such that  $x^k$  divides  $y$  in  $A$  are bounded.*

*proof.* (i) Since  $A$  is noetherian,  $I$  is a finite  $A$ -module. If  $I = JI$ , the Nakayama Lemma tells us that there is an element  $z$  of  $J$  such that  $zx = x$  for all  $x$  in  $I$ . Suppose that  $I$  isn't the zero ideal. We choose a nonzero element  $x$  of  $I$ . Because  $A$  is a domain, we can cancel  $x$  from the equation  $zx = x$ , obtaining  $z = 1$ . Then 1 is in  $J$ , and  $J$  is the unit ideal.

(ii) Suppose that  $B = JB$ . The Nakayama Lemma tells us that there is an element  $z$  in  $J$  such that  $zb = b$  for all  $b$  in  $B$ . Setting  $b = 1$  shows that  $z = 1$ . So  $J$  is the unit ideal.

(iii) Let  $I = \bigcap J^n$ . The elements of  $I$  are the elements of  $A$  that are divisible by  $x^n$  for every  $n$ . Let  $y$  be an element of  $I$ . So for every  $n$ , there is an element  $a_n$  in  $A$  such that  $y = a_n x^n$ . Then  $y/x = a_n x^{n-1}$ , which is an element of  $J^{n-1}$ . This is true for every  $n$ , so  $y/x$  is in  $I$ , and  $y$  is in  $JI$ . Since  $y$  can be any element of  $I$ ,  $I = JI$ . But since  $x$  isn't a unit,  $J$  isn't the unit ideal. So (i) tells us that  $I = 0$ .  $\square$

**4.1.6. Corollary.** *Let  $I$  be a nonzero ideal of a noetherian domain  $A$ , and let  $B$  be a domain that contains  $A$  as subring. If  $\beta$  is an element of  $B$  and if  $\beta I \subset I$ , then  $\beta$  is integral over  $A$ .*

*proof.* Because  $A$  is noetherian,  $I$  is finitely generated. Let  $v = (v_1, \dots, v_n)^t$  be a vector whose entries generate  $I$ . The hypothesis  $\beta I \subset I$  allows us to write  $\beta v_i = \sum p_{ij} v_j$  with  $p_{ij}$  in  $A$ , or in matrix notation,  $Pv = \beta v$ . So  $v$  is an eigenvector, and  $p(\beta)v = 0$ . Since at least one  $v_i$  is nonzero and since  $A$  is a domain,  $p(\beta) = 0$ . The characteristic polynomial  $p(t)$  is a monic polynomial with coefficients in  $A$ , so  $\beta$  is integral over  $A$ .  $\square$

## 4.2 Integral Extensions

Let  $A$  be a domain. An *extension*  $B$  of  $A$  is a ring that contains  $A$  as a subring. An element  $\beta$  of an extension  $B$  is *integral over*  $A$  if it is a root of a monic polynomial

$$(4.2.1) \quad f(x) = x^n + a_{n-1}x^{n-1} + \dots + a_0,$$

with coefficients  $a_i$  in  $A$ , and an extension  $B$  is an *integral extension* if all of its elements are integral over  $A$ .

**4.2.2. Lemma.** *Let  $A \subset B$  be an extension of domains.*

(i) *An element  $b$  of  $B$  is integral over  $A$  if and only if the subring  $A[b]$  of  $B$  generated by  $b$  is a finite  $A$ -module.*

(ii) *The set of elements of  $B$  that are integral over  $A$  is a subring of  $B$ .*

(iii) *If  $B$  is generated as  $A$ -algebra by finitely many integral elements, it is a finite  $A$ -module.*

(iv) *Let  $R \subset A \subset B$  be rings, and suppose that  $A$  is an integral extension of  $R$ . An element of  $B$  is integral over  $A$  if and only if it is integral over  $R$ .*  $\square$

**4.2.3. Corollary.** *An extension  $A \subset B$  of finite-type domains is an integral extension if and only if  $B$  is a finite  $A$ -module.*  $\square$

**4.2.4. Definition.** Let  $Y \xrightarrow{u} X$  be a morphism of affine varieties  $Y = \text{Spec } B$  and  $X = \text{Spec } A$ , and let  $A \xrightarrow{\varphi} B$  be the corresponding homomorphism of finite-type domains. If  $\varphi$  makes  $B$  into a finite  $A$ -module, we call  $u$  a *finite morphism* of affine varieties. If  $A$  is a subring of  $B$ , and  $B$  is an integral extension of  $A$ , we call  $u$  an *integral morphism* of affine varieties.



An integral morphism is a finite morphism whose associated algebra homomorphism  $A \xrightarrow{\varphi} B$  is injective.

**4.2.5. Example.** If  $G$  is a finite group of automorphisms of a finite-type domain  $B$  and  $A = B^G$  is the algebra of invariants, then  $B$  is an integral extension of  $A$ . (See Theorem 2.8.5.)

The next example is helpful for an intuitive understanding of the geometric meaning of integrality.

**4.2.6. Example.** Let  $f(x, y)$  be an irreducible polynomial in  $\mathbb{C}[x, y]$  (one  $x$  and one  $y$ ), let  $B = \mathbb{C}[x, y]/(f)$ , and let  $A = \mathbb{C}[x]$ . So  $X = \text{Spec } A$  is the affine line  $\mathbb{A}_x^1$ , and  $Y = \text{Spec } B$  is an affine plane curve. The canonical map  $A \rightarrow B$  defines a morphism  $Y \xrightarrow{u} X$ , which can be described as the restriction to  $Y$  of the projection  $\mathbb{A}_{x,y}^2 \rightarrow \mathbb{A}_x^1$ .

We write  $f$  as a polynomial in  $y$ , whose coefficients  $a_i(x)$  are polynomials in  $x$ :

$$(4.2.7) \quad f(x, y) = a_0(x)y^n + a_1(x)y^{n-1} + \cdots + a_n(x)$$

The fibre of  $Y$  over a point  $x = x_0$  of  $X$  is the set of points  $(x_0, y_0)$  such that  $y_0$  is a root of the one-variable polynomial  $f(x_0, y) = \tilde{f}(y)$ . Because  $f$  is irreducible, its discriminant with respect to the variable  $y$  isn't identically zero (1.7.19). For all but finitely many values of  $x$ ,  $\tilde{f}(y)$  will have degree  $n$  and its discriminant will be nonzero. Then  $\tilde{f}(y)$  will have  $n$  distinct roots.

When  $f(x, y)$  is a monic polynomial in  $y$ ,  $u$  will be an integral morphism. If so, the leading term  $y^n$  of  $f$  will be the dominant term, when  $y$  is large. Near a point  $x_0$  of  $X$ , there will be a positive real number  $B$  such that

$$|y^n| > |a_1(x)y^{n-1} + \cdots + a_n(x)|$$

when  $|y| > B$ , and then  $\tilde{f}(y) \neq 0$ . Therefore the roots  $y$  of  $f(x, y)$  are bounded for all  $x$  near to  $x_0$ .

On the other hand, when the leading coefficient  $a_0(x)$  isn't a constant,  $B$  won't be integral over  $A$ , and when  $x_0$  is a root of  $a_0$ ,  $f(x_0, y)$  will have degree less than  $n$ . In this case, as a point  $x_1$  of  $X$  approaches  $x_0$ , at least one root of  $f(x_1, y)$  tends to infinity. In calculus, one says that the locus  $f(x, y) = 0$  has a vertical asymptote at  $x_0$ .

To see this, we divide  $f$  by its leading coefficient. Let  $g(x, y) = f(x, y)/a_0 = y^n + c_1y^{n-1} + \cdots + c_n$  with  $c_i(x) = a_i(x)/a_0(x)$ . For any  $x$  at which  $a_0(x)$  isn't zero, the roots of  $g$  are the same as those of  $f$ . However, let  $x_0$  be a root of  $a_0$ . Because  $f$  is irreducible, there is at least one coefficient  $a_j(x)$  doesn't have  $x_0$  as a root. Then  $c_j(x)$  is unbounded near  $x_0$ , and because the coefficient  $c_j$  is an elementary symmetric function in the roots, the roots aren't all bounded.

This is the general picture: The roots of a polynomial remain bounded where the leading coefficient isn't zero. If the leading coefficient vanishes at a point, some roots are unbounded near that point.  $\square$

*figure : nonmonic polynomial, but compare with figure for Hensel's Lemma*

**4.2.8. Noether Normalization Theorem.** Let  $A$  be a finite-type algebra over an infinite field  $k$ . There exist elements  $y_1, \dots, y_n$  in  $A$  that are algebraically independent over  $k$ , such that  $A$  is a finite module over its polynomial subalgebra  $k[y_1, \dots, y_n]$ .

The Noether Normalization Theorem is also true when  $k$  is a finite field, though the proof given below needs to be modified. When  $K = \mathbb{C}$ , the theorem can be stated by saying that every affine variety  $X$  admits an integral morphism to an affine space.

**4.2.9. Lemma.** Let  $k$  be an infinite field, and let  $f(x)$  be a nonzero polynomial of degree  $d$  in  $x_1, \dots, x_n$ , with coefficients in  $k$ . After a suitable linear change of variable, the coefficient of  $x_n^d$  in  $f$  will be nonzero.

*proof.* Let  $f_d$  be the homogeneous part of  $f$  of maximal degree  $d$ . We regard  $f_d$  as a polynomial function. Since  $k$  is infinite, this function isn't identically zero. We choose coordinates  $x_1, \dots, x_n$  so that the point  $q = (0, \dots, 0, 1)$  isn't a zero of  $f_d$ . Then  $f_d(0, \dots, 0, x_n) = cx_n^d$ , and the coefficient  $c$ , which is  $f_d(0, \dots, 0, 1)$ , will be nonzero. By scaling  $x_n$ , we can make  $c = 1$ .  $\square$

*proof of the Noether Normalization Theorem.* Say that the finite-type algebra  $A$  is generated by elements  $x_1, \dots, x_n$ . If those elements are algebraically independent over  $k$ ,  $A$  will be isomorphic to the polynomial

algebra  $\mathbb{C}[x]$ , and we will be done. If not, they will satisfy a polynomial relation  $f(x) = 0$  of some degree  $d$ , with coefficients in  $k$ . The lemma tells us that, after a suitable change of variable, the coefficient of  $x_n^d$  in  $f$  will be 1. Then  $f$  will be a monic polynomial in  $x_n$  with coefficients in the subalgebra  $R$  generated by  $x_1, \dots, x_{n-1}$ . So  $x_n$  will be integral over  $R$ , and  $A$  will be a finite  $R$ -module. By induction on  $n$ , we may assume that  $R$  is a finite module over a polynomial subalgebra  $P$ . Then  $A$  will be a finite module over  $P$  too.  $\square$

The next proposition is an example of a general principle: A construction involving finitely many operations can be done in a simple localization.

**4.2.10. Proposition.** *Let  $A \subset B$  be finite-type domains. There is a nonzero element  $s$  in  $A$  such that  $B_s$  is a finite module over a polynomial subring  $A_s[y_1, \dots, y_r]$ .*

*proof.* Let  $S$  be the set of nonzero elements of  $A$ , so that  $K = AS^{-1}$  is the fraction field of  $A$ , and let  $B_K = BS^{-1}$  be the ring obtained from  $B$  by inverting all elements of  $S$ . Also, let  $\beta = (\beta_1, \dots, \beta_k)$  be a set of elements of the finite-type algebra  $B$  that generates  $B$  as algebra. Then  $B_K$  is a finite-type  $K$ -algebra, generated as  $K$ -algebra by  $\beta$ . (A  $K$ -algebra is a ring that contains  $K$  as subring. The Noether Normalization Theorem tells us that  $B_K$  is a finite module over a polynomial subring  $P = K[y_1, \dots, y_r]$ . So  $B_K$  is an integral extension of  $P$ . Any element of  $B$  will be in  $B_K$ , and therefore it will be the root of a monic polynomial, say

$$f(x) = x^n + c_{n-1}(y)x^{n-1} + \dots + c_0(y) = 0$$

where the coefficients  $c_j(y)$  are elements of  $P$ . Each coefficient is a combination of finitely many monomials in  $y$ , with coefficients in  $K$ . If  $d \in A$  is a common denominator for those coefficients,  $c_j(x)$  will have coefficients in  $A_d[y]$ . Since the generators  $\beta$  of  $B$  are integral over  $P$ , we may choose a denominator  $s$  so that all of the generators  $\beta_1, \dots, \beta_k$  are integral over  $A_s[y]$ . The algebra  $B_s$  is generated over  $A_s$  by  $\beta$ , so it will be an integral extension of  $A_s[y]$ .  $\square$

### 4.3 Normalization

Let  $A$  be a domain with fraction field  $K$ . The *normalization*  $A^\#$  of  $A$  is the set of elements of  $K$  that are integral over  $A$ . It follows from Lemma 4.2.2 (ii) that the normalization is a domain that contains  $A$ .

A domain  $A$  is *normal* if it is equal to its normalization, and a *normal variety*  $X$  is a variety that has an affine covering  $\{X^i = \text{Spec } A_i\}$  in which the algebras  $A_i$  are normal domains.

To justify the definition of normal variety, we need to show that if an affine variety  $X = \text{Spec } A$  has an affine covering  $X^i = \text{Spec } A_i$ , in which  $A_i$  are normal domains, then  $A$  is normal. This follows from Lemma 4.3.3 (iii) below.

Our goal here is the next theorem, whose proof is at the end of the section.

**4.3.1. Theorem.** *Let  $A$  be a finite-type domain with fraction field  $K$  of characteristic zero. The normalization of  $A$  is a finite  $A$ -module and a finite-type domain.*

Thus, if  $A^\#$  is the normalization of  $A$ , there will be an integral morphism  $\text{Spec } A^\# \rightarrow \text{Spec } A$ .

The proof given here makes use of the characteristic zero hypothesis, though the theorem is true for a finite-type  $k$ -algebra when  $k$  is a field of characteristic  $p$ .

**4.3.2. Example.** (*normalization of a nodal cubic curve*) The algebra  $A = \mathbb{C}[u, v]/(v^2 - u^3 - u^2)$  can be embedded into the one-variable polynomial algebra  $B = \mathbb{C}[x]$ , by  $u = x^2 - 1$  and  $v = x^3 - x$ . The fraction fields of  $A$  and  $B$  are equal because  $x = v/u$ , and the equation  $x^2 - (u+1) = 0$  shows that  $x$  is integral over  $A$ . Since  $B$  is normal, it is the normalization of  $A$  (see Lemma 4.3.3 (i)).

In this example,  $\text{Spec } B$  is the affine line  $\mathbb{A}_x^1$ , and the plane curve  $C = \text{Spec } A$  has a node at the origin  $p = (0, 0)$ . The inclusion  $A \subset B$  defines an integral morphism  $\mathbb{A}_x^1 \rightarrow C$  whose fibre over  $p$  is the point pair  $x = \pm 1$ . The morphism is bijective at all other points. I think of  $C$  as the variety obtained by gluing the points  $x = \pm 1$  of the affine line together.

figure: curve, not quite glued

In this example, the effect of normalization can be visualized geometrically. This isn't always so. Normalization is an algebraic process, whose effect on geometry may be subtle.  $\square$

**4.3.3. Lemma.** (i) A unique factorization domain is normal. In particular, a polynomial algebra over a field is normal.

(ii) If  $s$  is a nonzero element of a normal domain  $A$ . The localization  $A_s$  is normal.

(iii) Let  $s_1, \dots, s_k$  be nonzero elements of a domain  $A$  that generate the unit ideal. If the localizations  $A_{s_i}$  are normal for all  $i$ , then  $A$  is normal.

*proof.* (i) Let  $A$  be a unique factorization domain, and let  $\beta$  be an element of its fraction field that is integral over  $A$ . Say that

$$(4.3.4) \quad \beta^n + a_1\beta^{n-1} + \dots + a_{n-1}\beta + a_n = 0$$

with  $a_i$  in  $A$ . We write  $\beta = r/s$ , where  $r$  and  $s$  are relatively prime elements of  $A$ . Multiplying by  $s^n$  gives us the equation

$$r^n = -s(a_1r^{n-1} + \dots + a_{n-1}r + a_ns^{n-1})$$

This equation shows that if a prime element of  $A$  divides  $s$ , it also divides  $r$ . Since  $r$  and  $s$  are relatively prime, there is no such element. So  $s$  is a unit, and  $\beta$  is in  $A$ .

(ii) Let  $\beta$  be an element of the fraction field of  $A$  that is integral over  $A_s$ . There will be a polynomial relation of the form (4.3.4), except that the coefficients  $a_i$  will be elements of  $A_s$ . The element  $\gamma = s^k\beta$  satisfies the polynomial equation

$$\gamma^n + (s^k a_1)\gamma^{n-1} + \dots + (s^{(n-1)k} a_{n-1})\gamma + (s^{nk} a_n) = 0$$

Since  $a_i$  are in  $A_s$ , all coefficients of this polynomial will be in  $A$  when  $k$  is sufficiently large, and then  $\gamma$  will be integral over  $A$ . Since  $A$  is normal,  $\gamma$  will be in  $A$ , and  $\beta = s^{-k}\gamma$  will be in  $A_s$ .

(iii) This proof follows a common pattern. Suppose that  $A_{s_i}$  is normal for every  $i$ . If an element  $\beta$  of  $K$  is integral over  $A$ , it will be in  $A_{s_i}$  for all  $i$ , and  $s_i^n\beta$  will be an element of  $A$  if  $n$  is large. We can use the same exponent  $n$  for all  $i$ . Since  $s_1, \dots, s_k$  generate the unit ideal, so do their powers  $s_1^n, \dots, s_k^n$ . Say that  $\sum r_i s_i^n = 1$ , with  $r_i$  in  $A$ . Then  $\beta = \sum r_i s_i^n \beta$  is in  $A$ .  $\square$

It is convenient to state Theorem 4.3.1 more generally, especially for the proof. The more general statement is essentially the same.

Let  $A$  be a finite type domain with fraction field  $K$ , and let  $L$  be a finite field extension of  $K$ . The *integral closure* of  $A$  in  $L$  is the set of elements of  $L$  that are integral over  $A$ . As Lemma 4.2.2 (ii) shows, the integral closure is a domain that contains  $A$ .

**4.3.5. Theorem.** Let  $A$  be a finite type domain with fraction field  $K$ , and let  $L$  be a finite field extension of  $K$ . The integral closure  $B$  of  $A$  in  $L$  is a finite  $A$ -module.

**4.3.6. Lemma.** Let  $A$  be a normal noetherian domain with fraction field  $K$  of characteristic zero, and let  $L$  be an algebraic field extension of  $K$ . An element  $\beta$  of  $L$  is integral over  $A$  if and only if the coefficients of the monic irreducible polynomial  $f$  for  $\beta$  over  $K$  are in  $A$ .

*proof.* If the monic polynomial  $f$  has coefficients in  $A$ , then  $\beta$  is integral over  $A$ . Suppose that  $\beta$  is integral over  $A$ . Since we may replace  $L$  by any field extension that contains  $\beta$ , we may assume that  $L$  is a finite extension of  $K$ . A finite extension embeds into a Galois extension, so we may assume that  $L/K$  is a Galois extension. Let  $G$  be its Galois group, and let  $\{\beta_1, \dots, \beta_r\}$  be the  $G$ -orbit of  $\beta$ , with  $\beta = \beta_1$ . The irreducible polynomial for  $\beta$  over  $K$  is

$$(4.3.7) \quad f(x) = (x - \beta_1) \cdots (x - \beta_r)$$

Its coefficients are symmetric functions of the roots. If  $\beta$  is integral over  $A$ , then all elements of the orbit are integral over  $A$ . Therefore the symmetric functions are integral over  $A$  (4.2.2) (iii), and since  $A$  is normal, they are in  $A$ . So  $f$  has coefficients in  $A$ .  $\square$

**4.3.8. Example.** A polynomial in  $A = \mathbb{C}[x, y]$  is *square-free* if it has no nonconstant square factors and isn't a constant. Let  $f(x, y)$  be a square-free polynomial, and let  $B$  denote the integral extension  $\mathbb{C}[x, y, w]/(w^2 - f)$  of  $A$ . Let  $K$  and  $L$  be the fraction fields of  $A$  and  $B$ , respectively. Then  $L = K[w]/(w^2 - f)$  is a Galois extension of  $K$ . Its Galois group is generated by the automorphism  $\sigma$  of order 2 defined by  $\sigma(w) = -w$ . The elements of  $L$  have the form  $\beta = a + bw$  with  $a, b \in K$ , and  $\sigma(\beta) = \beta' = a - bw$ .

We show that  $B$  is the integral closure of  $A$  in  $L$ . Suppose that  $\beta = a + bw$  is integral over  $A$ . If  $b = 0$ , then  $\beta = a$ . This is an element of  $A$  and therefore it is in  $B$ . If  $b \neq 0$ , the irreducible polynomial for  $\beta = a + bw$  will be

$$(x - \beta)(x - \beta') = x^2 - 2ax + (a^2 - b^2f)$$

Because  $\beta$  is integral over  $A$ ,  $2a$  and  $a^2 - b^2f$  are in  $A$ . Because the characteristic isn't 2, this is true if and only if  $a$  and  $b^2f$  are in  $A$ . We write  $b = u/v$ , with  $u, v$  relatively prime elements of  $A$ , so  $b^2f = u^2f/v^2$ . If  $v$  weren't a unit, then since  $f$  is square-free, it couldn't cancel  $v^2$ . So from  $b^2f$  in  $A$  we can conclude that  $b$  is in  $A$ . Summing up,  $\beta$  is integral if and only if  $a$  and  $b$  are in  $A$ , which means that  $\beta$  is in  $B$ .  $\square$

#### (4.3.9) trace

We will use the trace in the proof of Theorem 4.3.5.

Let  $L$  be a finite field extension of a field  $K$  and let  $\beta$  be an element of  $K$ . When  $L$  is viewed as a  $K$ -vector space, multiplication by  $\beta$  becomes a linear operator  $L \xrightarrow{\beta} L$ . The *trace* of this operator will be denoted by  $\text{tr}(\beta)$ . The trace is a  $K$ -linear map  $L \rightarrow K$ .

**4.3.10. Lemma.** Let  $L/K$  be a finite field extension, let  $\beta$  be an element of  $L$  of degree  $r$  over  $K$ , let  $f(x) = x^r + a_1x^{r-1} + \cdots + a_r$  be its irreducible polynomial over  $K$ , and let  $K(\beta)$  be the extension of  $K$  generated by  $\beta$ . Say that  $[L : K(\beta)] = d$  and  $[L : K] = n (= rd)$ . Then  $\text{tr}(\beta) = -da_1$ . If  $\beta$  is an element of  $K$ , then  $\text{tr}(\beta) = n\beta$ .

*proof.* With  $r = [K(\beta) : K]$ , the set  $(1, \beta, \dots, \beta^{r-1})$  is a  $K$ -basis for  $K(\beta)$ . On this basis, the matrix  $M$  of multiplication by  $\beta$  has the form illustrated below for the case  $r = 3$ . Its trace is  $-a_1$ .

$$M = \begin{pmatrix} 0 & 0 & -a_3 \\ 1 & 0 & -a_2 \\ 0 & 1 & -a_1 \end{pmatrix}.$$

Next, let  $(u_1, \dots, u_d)$  be a basis for  $L$  over  $K(\beta)$ . Then  $\{\beta^i u_j\}$ , with  $i = 0, \dots, r-1$  and  $j = 1, \dots, d$ , will be a basis for  $L$  over  $K$ . When this basis is listed in the order

$$(u_1, u_1\beta, \dots, u_1\beta^{r-1}; u_2, u_2\beta, \dots, u_2\beta^{r-1}; \dots; u_d, u_d\beta, \dots, u_d\beta^{r-1}),$$

the matrix of multiplication by  $\beta$  will be made up of  $d$  blocks of the matrix  $M$ .  $\square$

**4.3.11. Corollary.** Let  $A$  be a normal domain with fraction field  $K$  and let  $L$  be a finite field extension of  $K$ . If an element  $\beta$  is integral over  $A$ , its trace is in  $A$ .

This follows from Lemmas 4.3.6 and 4.3.10.  $\square$

**4.3.12. Lemma.** Let  $A$  be a normal noetherian domain with fraction field  $K$  of characteristic zero, and let  $L$  be a finite field extension of  $K$ . The form  $L \times L \rightarrow K$  defined by  $\langle \alpha, \beta \rangle = \text{tr}(\alpha\beta)$  is  $K$ -bilinear, symmetric, and nondegenerate. If  $\alpha$  and  $\beta$  are integral over  $A$ , then  $\langle \alpha, \beta \rangle$  is an element of  $A$ .

*proof.* The form is obviously symmetric, and it is  $K$ -bilinear because multiplication is  $K$ -bilinear and trace is  $K$ -linear. A form is nondegenerate if its nullspace is zero, which means that when  $\alpha$  is a nonzero element, there is an element  $\beta$  such that  $\langle \alpha, \beta \rangle \neq 0$ . We let  $\beta = \alpha^{-1}$ . Then  $\langle \alpha, \beta \rangle = \text{tr}(1)$ , which, according to (4.3.10), is the degree  $[L : K]$  of the field extension. It is here that the hypothesis on the characteristic of  $K$  enters: The degree is a nonzero element of  $K$ .

If  $\alpha$  and  $\beta$  are integral over  $A$ , so is their product  $\alpha\beta$  (4.2.2) (ii). Corollary 4.3.11 shows that  $\langle \alpha, \beta \rangle$  is an element of  $A$ .  $\square$

**4.3.13. Lemma.** *Let  $A$  be a domain with fraction field  $K$ , let  $L$  be a field extension of  $K$ , and let  $\beta$  be an element of  $L$  that is algebraic over  $K$ . If  $\beta$  is a root of a polynomial  $f = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_0$  with  $a_i$  in  $A$ , then  $\gamma = a_n \beta$  is integral over  $A$ .*

One finds a monic polynomial with root  $\gamma$  by substituting  $x = y/a_n$  into  $f$  and multiplying by  $a_n^{n-1}$ .  $\square$

*proof of Theorem 4.3.1.* Let  $A$  be a finite-type domain with fraction field  $K$  of characteristic zero, and let  $L$  be a finite field extension of  $K$ . We are to show that the integral closure of  $A$  in  $L$  is a finite  $A$ -module.

*Step 1. We may assume that  $A$  is normal.*

We use the Noether Normalization Theorem to write  $A$  as a finite module over a polynomial subalgebra  $R = \mathbb{C}[y_1, \dots, y_d]$ . Let  $F$  be the fraction field of  $R$ . Then  $K$  and  $L$  are finite extensions of  $F$ . An element of  $L$  will be integral over  $A$  if and only if it is integral over  $R$  ((4.2.2) (iv)). So the integral closure of  $A$  in  $L$  is the same as the integral closure of  $R$  in  $L$ . We replace  $A$  by the normal algebra  $R$  and  $K$  by  $F$ .

*Step 2. Bounding the integral extension.*

We assume that  $A$  is normal. Let  $(v_1, \dots, v_n)$  be a  $K$ -basis for  $L$  whose elements are integral over  $A$ . Such a basis exists because we can multiply any element of  $L$  by a nonzero element of  $K$  to make it integral (Lemma 4.3.13). Let

$$(4.3.14) \quad T : L \rightarrow K^n$$

be the map  $T(\beta) = (\langle v_1, \beta \rangle, \dots, \langle v_n, \beta \rangle)$ , where  $\langle \cdot, \cdot \rangle$  is the form defined in Lemma 4.3.12. This map is  $K$ -linear. If  $\langle v_i, \beta \rangle = 0$  for all  $i$ , then because  $(v_1, \dots, v_n)$  is a basis for  $L$ ,  $\langle \gamma, \beta \rangle = 0$  for all  $\gamma$  in  $L$ , and since the form is nondegenerate,  $\beta = 0$ . Therefore  $T$  is injective.

Let  $B$  be the integral closure of  $A$  in  $L$ . The basis elements  $v_i$  are in  $B$ , and if  $\beta$  is in  $B$ ,  $v_i \beta$  will be in  $B$  too. Then  $\langle v_i, \beta \rangle$  will be in  $A$ , and  $T(\beta)$  will be in  $A^n$  (4.3.12). When we restrict  $T$  to  $B$ , we obtain an injective map  $B \rightarrow A^n$  that we denote by  $T_0$ . Since  $T$  is  $K$ -linear,  $T_0$  is a  $A$ -linear. It is an injective homomorphism of  $A$ -modules. It maps  $B$  isomorphically to its image, a submodule of  $A^n$ . Since  $A$  is noetherian, every submodule of the finite  $A$ -module  $A^n$  is finitely generated. Therefore the image of  $T_0$  is a finite  $A$ -module, and so is the isomorphic  $A$ -module  $B$ .  $\square$

## 4.4 Geometry of Integral Morphisms

The main geometric properties of an integral morphism of affine varieties are summarized in the two theorems below which show that the geometry is as nice as could be expected.

Let  $Y \xrightarrow{u} X$  be an integral morphism. We say that a closed subvariety  $D$  of  $Y$  *lies over* a closed subvariety  $C$  of  $X$  if  $C$  is the image of  $D$ .

**4.4.1. Theorem.** *Let  $Y \xrightarrow{u} X$  be an integral morphism of affine varieties.*

- (i) *The morphism  $u$  is surjective, and its fibres have bounded cardinality.*
- (ii) *The image of a closed subvariety of  $Y$  is a closed subvariety of  $X$ , and the image of a closed subset of  $Y$  is a closed subset of  $X$ .*
- (iii) *The set of closed subvarieties of  $Y$  that lie over a closed subvariety of  $X$  is finite and nonempty.*

The second theorem concerns inclusions among closed subvarieties. It refers to the diagram below:

$$\begin{array}{ccc} D' \subset D & & Y \\ C' \subset C & & X \end{array}$$

**4.4.2. Theorem.** *Let  $Y \xrightarrow{u} X$  be an integral morphism of affine varieties, and let  $C' \subset C$  be closed subvarieties of  $X$ .*

- (i) *Every closed subvariety  $D$  of  $Y$  that lies over  $C$  contains a closed subvariety  $D'$  that lies over  $C'$ .*
- (ii) *Suppose that  $X$  normal. Every closed subvariety  $D'$  of  $Y$  that lies over  $C'$  is contained in a closed subvariety  $D$  that lies over  $C$ .*

Property (ii) is more subtle than (i), as is indicated by the fact that  $X$  is assumed normal. There is an example at the end of the section showing that the hypothesis of normality cannot be dropped.

**4.4.3. Corollary.** *Let  $Y \xrightarrow{u} X$  be an integral morphism of affine varieties with  $X$  normal, and let  $C$  be a closed subvariety of  $X$ . The subvarieties of  $Y$  that lie over  $C$  are the irreducible components of the inverse image of  $C$ .*

*proof.* Let  $Z$  be the inverse image of  $C$ , let  $D'$  be a component of  $Z$ , and let  $C'$  be its image in  $X$ . Then  $C' \subset C$ , so by part (ii) of Theorem 4.4.2,  $D'$  is contained in a subvariety  $D$  of  $Y$  that lies over  $C$ . Because  $D$  is contained in  $Z$  and  $D'$  is a component of  $Z$ ,  $D' = D$ .  $\square$

We will go to algebra to prove Theorems 4.4.1 and 4.4.2.

Let  $A \subset B$  be an extension of finite-type domains. As before, if  $I$  is an ideal of  $A$ , the *extended ideal*  $IB$  is the ideal of  $B$  generated by  $I$ . Its elements are finite sums  $\sum u_i b_i$  with  $u_i$  in  $I$  and  $b_i$  in  $B$ . The *contraction* of an ideal  $J$  of  $B$  is the ideal  $J \cap A$  of  $A$ . The contraction of a prime ideal is a prime ideal.

Closed subvarieties of the affine variety  $X = \text{Spec } A$  correspond bijectively to prime ideals of  $A$ . In analogy with the terminology for closed subvarieties, we say that a prime ideal  $Q$  of  $B$  *lies over* a prime ideal  $P$  of  $A$  if its contraction is  $P$ . For example, if a point  $y$  of  $Y = \text{Spec } B$  has image  $x$  in  $X$ , the maximal ideal  $\mathfrak{m}_y$  lies over the maximal ideal  $\mathfrak{m}_x$ .

In commutative algebra books, Theorem 4.4.2 is stated in terms of prime ideals. Suppose given prime ideals  $P \subset P'$  of  $A$ : Do there exist prime ideals  $Q \subset Q'$  of  $B$  whose contractions are  $P$  and  $P'$ , respectively?

$$\begin{array}{ccc} Q \subset Q' & & B \\ P \subset P' & & A \end{array}$$

The translation of Theorem 4.4.2 to prime ideals reads as follows:

**4.4.4. Theorem.** *Let  $A \subset B$  be an integral extension of finite-type domains, and let  $P \subset P'$  be prime ideals of  $A$ .*

- (i) *Every prime ideal  $Q$  that lies over  $P$  is contained in a prime ideal  $Q'$  that lies over  $P'$ .*
- (ii) *Suppose that  $A$  is normal. Then every prime ideal  $Q'$  that lies over  $P'$  contains a prime ideal  $Q$  that lies over  $P$ .*

The statements (i) and (ii) of this theorem are often called “going up”, and “going down”, respectively. Since inclusions are reversed when one passes to closed subvarieties, those terms aren’t appropriate in Theorem 4.4.2.  $\square$

*proof of Theorem 4.4.1.* Let  $Y \xrightarrow{u} X$  be an integral morphism, with  $Y = \text{Spec } B$  and  $X = \text{Spec } A$ .

(i) (*bounding the fibres*) Let  $\mathfrak{m}_x$  be the maximal ideal at point  $x$  of  $X$ . Corollary 4.1.5 (ii) shows that the extended ideal  $\mathfrak{m}_x B$  isn’t the unit ideal of  $B$ , so it is contained in a maximal ideal of  $B$ , say  $\mathfrak{m}_y$ , where  $y$  is a point of  $Y$ , and then  $x$  is the image of  $y$ . Therefore  $u$  is surjective.

Let  $y_1, \dots, y_r$  be the points of  $Y$  in the fibre over a point  $x$  of  $X$ . Then for each  $i$ , the maximal ideal  $\mathfrak{m}_x$  of  $A$  is the contraction of the maximal ideal  $\mathfrak{m}_{y_i}$  of  $B$ . To bound the number  $r$ , we use the Chinese Remainder Theorem to show that  $B$  cannot be spanned as  $A$ -module by fewer than  $r$  elements.

Let  $k_i$  and  $k$  denote the residue fields  $B/\mathfrak{m}_{y_i}$ , and  $A/\mathfrak{m}_x$ , respectively, all of these fields being abstractly isomorphic to  $\mathbb{C}$ , and let  $\overline{B} = k_1 \times \dots \times k_r$ . We form a diagram of algebra homomorphisms

$$\begin{array}{ccc} B & \xrightarrow{\varphi} & \overline{B} \\ \uparrow & & \uparrow \\ A & \longrightarrow & k \end{array}$$

which we interpret as a diagram of  $A$ -modules. The minimal number of generators of the  $A$ -module  $\overline{B}$  is equal to its dimension as  $k$ -module, which is  $r$ . The Chinese Remainder Theorem asserts that  $\varphi$  is surjective, so  $B$  cannot be spanned by fewer than  $r$  elements.

(ii) *(the image of a closed set is closed)* It is obvious that the image of an irreducible set via continuous map is irreducible, so it suffices to show that the image of a closed subvariety is closed. Let  $D$  be the closed subvariety of  $Y$  that corresponds to a prime ideal  $Q$  of  $B$ , and let  $P$  be the contraction of  $Q$ . Then  $P$  is a prime ideal of  $A$ . Let  $C$  be its variety of zeros. The coordinate rings of the affine varieties  $D$  and  $C$  are  $\overline{B} = B/Q$  and  $\overline{A} = A/P$ , respectively, and because  $B$  is an integral extension of  $A$ ,  $\overline{B}$  is an integral extension of  $\overline{A}$ . By (i), the map  $D \rightarrow C$  is surjective. Therefore the closed set  $C$  is the image of  $D$ .

We use the next lemma for the proof of part (iii) of the theorem.

**4.4.5. Lemma.** *Let  $Y \xrightarrow{u} X$  be an integral morphism of affine varieties, and let  $D$  and  $D'$  be closed subvarieties of  $Y$  that lie over closed subvarieties  $C$  and  $C'$  of  $X$ , respectively. Then  $D' < D$  if and only if  $C' < C$ .*

*proof.* Since  $C$  and  $C'$  are images, it is clear that  $D' < D$  if  $C' < C$ . For the other implication, we may replace  $X$  and  $Y$  by  $C$  and  $D$ , respectively. Then what has to be shown is that  $D' < Y$  if and only if  $C' < X$ . We go over to ideals. Say that  $Q'$  and  $P'$  are the prime ideals of  $B$  and  $A$  corresponding to  $D'$  and  $C'$ , respectively. So  $P'$  is the contraction  $Q' \cap A$  of  $Q'$ . What has to be shown is that if  $Q'$  is nonzero, then  $P'$  is nonzero.

Let  $\beta$  be a nonzero element of  $Q'$ . Then  $\beta$  is integral over  $A$ , say  $\beta^n + a_{n-1}\beta^{n-1} + \cdots + a_0 = 0$ , with  $a_i \in A$ . If  $a_0 = 0$ , then because  $B$  is a domain, we can cancel  $\beta$  from the equation. So we may assume  $a_0 \neq 0$ . The equation shows that  $a_0$  is in  $Q'$ , and since it is also in  $A$ , it is in  $P'$ .  $\square$

*proof of Theorem 4.4.1. (iii) (subvarieties that lie over a closed subvariety)* The inverse image  $Z = u^{-1}C$  of a closed subvariety  $C$  is closed in  $Y$ . It is the union of finitely many irreducible closed subsets, say  $Z = D'_1 \cup \cdots \cup D'_k$ . Part (i) tells us that the image  $C'_i$  of  $D'_i$  is a closed subvariety of  $X$ . Since  $u$  is surjective,  $C = \bigcup C'_i$ , and since  $C$  is irreducible, it is equal to at least one  $C'_i$ . The components  $D'_i$  such that  $C'_i = C$  are the subvarieties that lie over  $C$ . Moreover, a subvariety  $D$  that lies over  $C$  will be contained in the inverse image  $Z = \bigcup D'_i$  of  $C$ . According to Lemma 4.4.5, there are no inclusions among subvarieties that lie over  $C$ . Therefore  $D$  must be one of the  $D'_i$ . So it is an element of a finite set.  $\square$

*proof of Theorem 4.4.2 (i).* We are given  $C' \subset C$  in  $X$  and  $D$  in  $Y$  that lies over  $C$ , and we must find  $D'$ . This follows from what has been done. We replace  $Y$  and  $X$  by  $D$  and  $C$ , respectively. Then what is to be proved is that there is a closed subvariety  $D'$  of  $Y$  that lies over a given closed subvariety  $C'$ . This is part (ii) of Theorem 4.4.1.

The proof of part (ii) of Theorem 4.4.2 will be given at the end of Section 4.5.

**4.4.6. Example.** In this example,  $B$  is the normalization of a finite-type domain  $A$ ,  $A$  isn't normal, and the conclusion of Theorem 4.4.2 (ii) fails.

We add a variable to the rings  $R = \mathbb{C}[u, v]/(v^2 - u^3 - u^2)$  and  $S = \mathbb{C}[x]$  of Example 4.3.2, where  $u = x^2 - 1$  and  $v = x^3 - x$ . Let  $A = R[y]$ , and  $B = S[y]$ . Then  $A \subset B$  and  $B$  is the normalization of  $A$ . Let  $Y$  be the affine  $x, y$ -plane, and let  $X$  be the locus  $v^2 - u^3 - u^2 = 0$  in the affine 3-space with coordinates  $u, v, y$ . So  $X$  is the product  $V \times \mathbb{A}_y^1$ , where  $V = \text{Spec } R$ . The integral morphism  $Y \rightarrow X$  that corresponds to the inclusion  $A \subset B$  sends a point  $(x, y)$  of  $Y$  to  $(x^2 - 1, x^3 - x, y)$  in  $X$ . We noted before that  $V$  is the nodal curve obtained by identifying the points  $x = \pm 1$  of the affine  $x$ -line. So  $X$  can be obtained geometrically by identifying the two lines  $\ell_1 : \{x = 1\}$  and  $\ell_2 : \{x = -1\}$  in the affine plane  $Y$ . The image of these lines in  $X$  is the line  $L : \{u = v = 0\}$ .

Let  $C'$  be the point  $(0, 0, 1)$  of  $Y$ , let  $D'$  be the point  $(-1, 1)$  of  $X$ , and let  $C$  be the image of the diagonal line  $Z : \{y = x\}$  in  $Y$ , a closed subvariety of  $X$ . The inverse image of  $C$  is  $Z \cup (-1, 1) \cup (1, -1)$ . Then  $C' \subset C$ , and  $D'$  lies over  $C'$ , but there is no subvariety  $D$  that lies over  $C$  and contains  $D'$ . So the assertion of Theorem 4.4.2 (ii) is false.  $\square$

##figure##

$\square$

## 4.5 Finite group actions II

Let  $G$  be a finite group of automorphisms of a normal, finite-type domain  $B$ , let  $A$  be the algebra of invariant elements of  $B$ . According to Theorem 2.8.5,  $A$  is a finite-type domain, and  $B$  is a finite  $A$ -module. Let

$Y = \text{Spec } B$  and  $X = \text{Spec } A$ . Points of  $X$  correspond to  $G$ -orbits of points of  $Y$ . Theorem 4.5.2 below extends these facts to closed subvarieties.

**4.5.1. Lemma.** Let  $G$  be a finite group of automorphisms of a normal, finite-type domain  $B$  and let  $A$  be the subalgebra of invariant elements. Let  $L$  and  $K$  be the fraction fields of  $B$  and  $A$ , respectively.

- (i) The algebra  $A$  is normal.
- (ii) Every element of  $L$  can be written as a fraction  $b/s$ , with  $b$  in  $B$ , and  $s$  in  $A$ .
- (iii) The operation of  $G$  on  $B$  induces an operation on  $L$ , and the ring of invariant elements of  $L$  is  $K$ .  $\square$

Since  $B$  is a finite  $A$ -module, the results of Section 4.4 apply to the integral morphism  $Y \xrightarrow{u} X$ . Recall that, since  $G$  operates on the left on  $B$ , it operates on the right on  $Y$  (2.8.9).

**4.5.2. Theorem.** Let  $G$  be a finite group of automorphisms of a normal, finite-type domain  $B$ , let  $A$  be the algebra of invariant elements of  $B$ , and let  $Y \xrightarrow{u} X$  be the integral morphism of varieties corresponding to the inclusion  $A \subset B$ .

- (i) There is a bijective correspondence between orbits of closed subvarieties of  $Y$  and closed subvarieties of  $X$ :

$$\{\text{closed subvarieties of } Y\}/G \longleftrightarrow \{\text{closed subvarieties of } X\}$$

in which the orbit that corresponds to a closed subvariety  $C$  of  $X$  is the set of closed subvarieties of  $Y$  that lie over  $C$ .

- (ii) Let  $\{D_1, \dots, D_r\}$  and  $\{D'_1, \dots, D'_s\}$  be orbits of closed subvarieties  $Y$  that lie over  $C$  and  $C'$ , respectively, in  $X$ . If  $C \supset C'$ , then every  $D_i$  contains some  $D'_j$ . Conversely, if  $D_i \supset D'_j$  for some  $i$  and  $j$ , then  $C \supset C'$ .

*proof.* (i) Let  $D$  be a closed subvariety of  $Y$  with image  $C$  in  $X$ . Because points of  $X$  correspond to  $\sigma$ -orbits in  $Y$ , the subvarieties  $D\sigma$  in the orbit of  $D$  have the same image  $C$ . The fact that distinct orbits of closed subvarieties of  $Y$  lie over distinct closed subvarieties of  $X$  will follow from (ii).

- (ii) It is clear that if  $D_i \supset D'_j$ , then  $C \supset C'$ . We suppose that  $D_i \not\supset D'_j$  for all  $i$  and  $j$ , and we show that  $C \not\supset C'$ . The lemma below, whose proof is an exercise, shows that there is an element  $\beta$  that is identically zero on every  $D_i$  and isn't identically zero on any  $D'_j$ . Then for all  $\sigma$  in  $G$ ,  $\sigma\beta$  has the same property. So  $\alpha = \prod \sigma\beta$  is an element of  $A$  that is identically zero on every  $D_i$  but not on any  $D'_j$ . Then  $\alpha$  is identically zero on  $C$  but not on  $C'$ . So  $C$  doesn't contain  $C'$ .  $\square$

**4.5.3. Lemma.** Let  $Y = \text{Spec } B$  be an affine variety, let  $D_1, \dots, D_n$  be distinct closed subvarieties of  $Y$  and let  $V$  be a closed subset of  $Y$ . Assume that  $V$  doesn't contain any of the sets  $D_j$ . There is an element  $\beta$  of  $B$  that vanishes on  $V$ , but isn't identically zero on any  $D_j$ .

We now complete the proof of Theorem 4.4.2.

*proof of Theorem 4.4.2 (ii).* We are given an integral morphism  $\text{Spec } B = Y \xrightarrow{u} X = \text{Spec } A$  with  $X$  normal, and we are given closed subvarieties  $C \supset C'$  of  $X$  and a closed subvariety  $D'$  of  $Y$  that lies over  $C'$ . We are to find a closed subvariety  $D$  that lies over  $C$  and that contains  $D'$ . Let  $K$  and  $L$  denote the fraction fields of  $A$  and  $B$ , respectively. Since  $B$  is a finite  $A$ -module,  $L$  is a finite extension of  $K$ .

*Case 1:*  $L$  is a Galois extension of  $K$  and  $B$  is normal. Then  $A$  and  $B^G$  have the same fields of fractions,  $A \subset B^G$ , and  $A$  is normal. So  $A = B^G$ . This case follows from Theorem 4.5.2.

*Case 2:* the general case. We put  $L$  into a Galois extension  $F$ , and we let  $R$  be the integral closure of  $B$  in  $F$ . Then  $R$  is a finite  $B$ -module and a finite  $A$ -module. Let  $Z = \text{Spec } R$  and let  $E$  be a closed subvariety of  $Z$  that lies over  $D$ . Then  $E$  also lies over  $C$ . By Case 1, there is a closed subvariety  $E'$  of  $Z$  that lies over  $C'$  and is contained in  $E$ . The image  $D'$  of  $E'$  in  $Y$  is the required closed subvariety of  $Y$ .

$$\begin{array}{ccc} E' \subset E & & Z \\ & D & Y \\ C' \subset C & & X \end{array}$$

$\square$ .



## 4.6 Chevalley's Finiteness Theorem

### (4.6.1) finite morphisms

The concepts of a finite morphism and an integral morphism of affine varieties were defined in Section 4.2. A morphism  $Y \xrightarrow{u} X$  of affine varieties  $X = \text{Spec } A$  and  $Y = \text{Spec } B$  is a finite morphism if the homomorphism  $A \xrightarrow{\varphi} B$  that corresponds to  $u$  makes  $B$  into a finite  $A$ -module. As was explained, the difference between a finite morphism and an integral morphism of affine varieties is that for a finite morphism, the homomorphism  $\varphi$  needn't be injective. If  $\varphi$  is injective,  $B$  will be an integral extension of  $A$ , and  $u$  will be an integral morphism. We extend the definitions to varieties that aren't necessarily affine here.

By the *restriction* of a morphism  $Y \xrightarrow{u} X$  to an open subset  $X'$  of  $X$ , we mean the induced morphism  $Y' \rightarrow X'$ , where  $Y'$  is the inverse image of  $X'$ .

**4.6.2. Definition.** A morphism of varieties  $Y \xrightarrow{u} X$  is a *finite morphism* if  $X$  can be covered by affine open subsets to which the restriction of  $u$  is a finite morphism of affine varieties, as defined in (4.2.4). A morphism  $u$  is an *integral morphism* if there is a covering of  $X$  by affine open sets to which the restriction of  $u$  is an integral morphism of affine varieties.

**4.6.3. Corollary.** *An integral morphism is a finite morphism. The composition of finite morphisms is a finite morphism. The inclusion of a closed subvariety into a variety is a finite morphism.*  $\square$

When  $X$  is affine, Definition 4.2.4 and Definition 4.6.2 both apply. The next proposition shows that the two definitions are equivalent.

**4.6.4. Proposition.** *Let  $Y \xrightarrow{u} X$  be a finite or an integral morphism, as in (4.6.2), and let  $X'$  be an affine open subset of  $X$ . The restriction of  $u$  to  $X'$  is a finite or an integral morphism of affine varieties, as defined in (4.2.4).*

**4.6.5. Lemma.** (i) *Let  $A \xrightarrow{\varphi} B$  be a homomorphism of finite-type domains that makes  $B$  into a finite  $A$ -module, and let  $s$  be a nonzero element of  $A$ . Then  $B_s$  is a finite  $A_s$ -module.*

(ii) *The restriction of a finite (or an integral) morphism  $Y \xrightarrow{u} X$  to an open subset of  $X$  is a finite (or an integral) morphism, as in Definition 4.6.2.*

*proof.* (i) In the statement,  $B_s$  denotes the localization of  $B$  as  $A$ -module. This localization can also be obtained by localizing the algebra  $B$  with respect to the image  $s' = \varphi(s)$ , provided that it isn't zero. If  $s'$  is zero, then  $s$  annihilates  $B$ , so  $B_s = 0$ . In either case, a set of elements that spans  $B$  as  $A$ -module will span  $B_s$  as  $A_s$ -module, so  $B_s$  is a finite  $A_s$ -module.

(ii) Say that  $X$  is covered by affine open sets to which the restriction of  $u$  is a finite morphism. The localizations of these open sets form a basis for the Zariski topology on  $X$ . So  $X'$  can be covered by such localizations. Part (i) shows that the restriction of  $u$  to  $X'$  is a finite morphism.  $\square$

*proof of Proposition 4.6.4.* We'll do the case of a finite morphism. The proof isn't difficult, but there are several things to check, and this makes the proof longer than one would like.

*Step 1. Preliminaries.*

We are given a morphism  $Y \xrightarrow{u} X$ ,  $X$  is covered by affine open sets  $X^i$ , and the restrictions of  $u$  to these open sets are finite morphisms of affine varieties. We are to show that the restriction to any affine open set  $X'$  is a finite morphism of affine varieties.

The affine open set  $X'$  is covered by the affine open sets  $X'^i = X' \cap X^i$ , and the restrictions to  $X'^i$  are finite morphisms ((4.6.5) (ii)). So we may replace  $X$  by  $X'$ . Since the localizations of an affine variety form a basis for its Zariski topology, we see that what is to be proved is this:

A morphism  $Y \xrightarrow{u} X$  is given in which  $X = \text{Spec } A$  is affine. There are elements  $s_1, \dots, s_k$  that generate the unit ideal of  $A$ , such that for every  $i$ , the inverse image  $Y^i$  of  $X^i = X_{s_i}$  if nonempty, is affine, and its coordinate algebra  $B_i$  is a finite module over the localized algebra  $A_i = A_{s_i}$ . We must show that  $Y$  is affine, and that its coordinate algebra  $B$  is a finite  $A$ -module.

*Step 2. The algebra  $B$  of regular functions on  $Y$ .*

If  $Y$  is affine, its coordinate algebra  $B$  will be a finite-type domain, and  $Y$  will be its spectrum. Since  $Y$  isn't assumed to be affine, we don't know very much about  $B$  other than that it is a subalgebra of the function field  $L$  of  $Y$ . On the other hand, the inverse image  $Y^i$  of  $X^i$ , if nonempty, is affine, the spectrum of a finite-type domain  $B_i$ . Since the localizations  $X^i$  cover  $X$ , the affine varieties the  $Y^i$  cover  $Y$ . We throw out the indices  $i$  such that  $Y^i$  is empty. Then a function is regular on  $Y$  if and only if it is regular on each  $Y^i$ , and

$$B = \bigcap B_i$$

the intersection being in the function field  $L$ .

Let's denote the images in  $B$  of the elements  $s_i$  of  $A$  by the same symbols.

*Step 3. For any index  $j$ ,  $B_j$  is the localization  $B[s_j^{-1}]$  of  $B$ .*

The intersection  $Y^j \cap Y^i$  is an affine variety. It is the set of points of  $Y^j$  at which  $s_i$  isn't zero, and its coordinate algebra is the localization  $B_j[s_i^{-1}]$ . Then

$$B[s_j^{-1}] \stackrel{(1)}{=} \bigcap (B_i[s_j^{-1}]) \stackrel{(2)}{=} \bigcap B_j[s_i^{-1}] \stackrel{(3)}{=} B_j[s_j^{-1}] \stackrel{(3)}{=} B_j$$

where the explanation of the numbered equalities is as follows:

- (1) A rational function  $\beta$  is in  $B_i[s_j^{-1}]$  if  $s_j^n \beta$  is in  $B_i$  for large  $n$ , and we can use the same exponent  $n$  for all  $i = 1, \dots, r$ . So  $\beta$  is in  $\bigcap (B_i[s_j^{-1}])$  if and only if  $s_j^n \beta$  is in  $\bigcap B_i = B$ , i.e., if and only if  $\beta$  is in  $B[s_j^{-1}]$ .
- (2) This is true because  $Y^j \cap Y^i = Y^i \cap Y^j$ .
- (3) For all  $i$ ,  $B_j \subset B_j[s_i^{-1}]$ . Moreover,  $s_j$  doesn't vanish on  $Y^j$ . It is a unit in  $B_j$ , and therefore  $B_j[s_j^{-1}] = B_j$ :  $B_j \subset \bigcap B_j[s_i^{-1}] \subset B_j[s_j^{-1}] \subset B_j$ .

*Step 4.  $B$  is a finite  $A$ -module.*

We choose a finite set  $b = (b_1, \dots, b_n)$  of elements of  $B$  that generates the  $A_i$ -module  $B_i$  for every  $i$ . We can do this because we can span the finite  $A_i$ -module  $B_i$  by finitely many elements of  $B$ , and there are finitely many algebras  $B_i$ . We show that the set  $b$  generates the  $A$ -module  $B$ .

Let  $x$  be an element of  $B$ . Since  $x$  is in  $B_i$ , it is a combination of the elements  $b$  with coefficients in  $A_i$ . Then for large  $k$ ,  $s_i^k x$  will be a combination of  $b$  with coefficients in  $A$ , say

$$s_i^k x = \sum_{\nu} a_{i,\nu} b_{\nu}$$

with  $a_{i,\nu} \in A$ . We can use the same exponent  $k$  for all  $i$ . Then with  $\sum r_i s_i^k = 1$ ,

$$x = \sum_i r_i s_i^k x = \sum_i r_i \sum_{\nu} a_{i,\nu} b_{\nu}$$

The right side is a combination of  $b$  with coefficients in  $A$ .

*Step 5.  $Y$  is affine.*

The algebra  $B$  of regular functions on  $Y$  is a finite-type domain because it is a finite module over the finite-type domain  $A$ . Let  $\tilde{Y} = \text{Spec } B$ . The fact that  $B$  is the algebra of regular functions on  $Y$  gives us a morphism  $Y \xrightarrow{\epsilon} \tilde{Y}$  (Corollary 3.5.3). Restricting to the open subset  $X^j$  of  $X$  gives us a morphism  $Y^j \xrightarrow{\epsilon^j} \tilde{Y}^j$  in which  $Y^j$  and  $\tilde{Y}^j$  are both equal to  $\text{Spec } B_j$ . Therefore  $\epsilon^j$  is an isomorphism. Corollary 3.4.18 (ii) shows that  $\epsilon$  is an isomorphism. So  $Y$  is affine and by Step 4, its coordinate algebra  $B$  is a finite  $A$ -module.  $\square$

We come to Chevalley's theorem now. Let  $\mathbb{P}$  denote the projective space  $\mathbb{P}^n$  with coordinates  $y_0, \dots, y_n$ .

**4.6.6. Chevalley's Finiteness Theorem.** *Let  $X$  be a variety, let  $Y$  be a closed subvariety of the product  $\mathbb{P} \times X$ , and let  $\pi$  denote the projection  $Y \rightarrow X$ . If all fibres of  $\pi$  are finite sets, then  $\pi$  is a finite morphism.*

**4.6.7. Corollary.** *Let  $Y$  be a projective variety and let  $Y \xrightarrow{u} X$  be a morphism whose fibres are finite sets. Then  $u$  is a finite morphism. In particular, if  $Y$  is a projective curve, any nonconstant morphism  $Y \xrightarrow{u} X$  is a finite morphism.*

This corollary follows from the theorem when one replaces  $Y$  by the graph of  $u$  in  $Y \times X$ . If  $Y$  is embedded as a closed subvariety of  $\mathbb{P}$ , the graph will be a closed subvariety of  $\mathbb{P} \times X$  (Proposition 3.4.27).  $\square$

In the next lemma,  $A$  denotes a finite-type domain,  $B$  denotes a quotient of the algebra  $A[u]$  of polynomials in  $n$  variables  $u_1, \dots, u_n$  with coefficients in  $A$ , and  $A \xrightarrow{\varphi} B$  denotes the canonical homomorphism. We'll use capital letters for nonhomogeneous polynomials here. If  $G(u)$  is a polynomial in  $A[u]$ , we denote its image in  $B$  by  $G(u)$ , too.

**4.6.8. Lemma.** *Let  $k$  be a positive integer. Suppose that, for each  $i = 1, \dots, n$ , there is a polynomial  $G_i(u_1, \dots, u_n)$  of degree at most  $k-1$  in  $n$  variables with coefficients in  $A$ , such that  $u_i^k = G_i(u)$  in  $B$ . Then  $B$  is a finite  $A$ -module.*

*proof.* Any monomial in  $u_1, \dots, u_n$  of degree at least  $nk$  will be divisible by  $u_i^k$  for at least one  $i$ . So if  $m$  is a monomial of degree  $d \geq nk$ , the relation  $u_i^k = G_i(u)$  shows that, in  $B$ ,  $m$  is equal to a polynomial in  $u_1, \dots, u_n$  of degree less than  $d$ , with coefficients in  $A$ . By induction, it follows that the monomials of degree at most  $nk-1$  span  $B$ .  $\square$

Let  $y_0, \dots, y_n$  be coordinates in  $\mathbb{P}^n$ , and let  $A[y_0, \dots, y_n]$  be the algebra of polynomials in  $y$  with coefficients in  $A$ . A *homogeneous element* of  $A[y]$  is an element that is a homogeneous polynomial in  $y$  with coefficients in  $A$ . A *homogeneous ideal* of  $A[y]$  is an ideal that can be generated by homogeneous polynomials.

**4.6.9. Lemma.** *Let  $Y$  be a closed subset of  $\mathbb{P} \times X$ , where  $X = \text{Spec } A$  is affine,*

- (i) *The ideal  $\mathcal{I}$  of elements of  $A[y]$  that vanish at every point of  $Y$  is a homogeneous ideal of  $A[y]$ .*
- (ii) *If the zero locus of a homogeneous ideal  $\mathcal{I}$  of  $A[y]$  is empty, then  $\mathcal{I}$  contains a power of the irrelevant ideal  $\mathcal{M} = (y_0, \dots, y_n)$  of  $A[y]$ .*

*proof.* (i) Let's write a point of  $\mathbb{P} \times X$  as  $q = (y_0, \dots, y_n, x)$ , with  $x$  representing a point of  $X$ . So  $(y, x) = (\lambda y, x)$ . Then the proof for the case  $A = \mathbb{C}$  that is given in (1.3.2) carries over.

(ii) Let  $V$  be the complement of the origin in the affine  $n+1$ -space with coordinates  $y$ . The complement of the origin in  $V \times X$  maps to  $\mathbb{P} \times X$  (see 3.2.4). If the locus of zeros of  $\mathcal{I}$  in  $\mathbb{P} \times X$  is empty, its locus of zeros in  $V \times X$  will be contained in  $o \times X$ ,  $o$  being the origin in  $\mathbb{P}$ . Then the ideal of  $o \times X$ , which is the radical ideal generated by the elements  $y_0, \dots, y_n$ , will contain  $\mathcal{I}$ .  $\square$

*proof of Chevalley's Finiteness Theorem.* This proof is adapted from a proof by Schelter.

We abbreviate the notation for a product  $Z \times X$  of a variety  $Z$  with  $X$ , denoting  $Z \times X$  by  $\tilde{Z}$ .

We are given a closed subvariety  $Y$  of  $\tilde{\mathbb{P}} = \mathbb{P} \times X$ , and the fibres over  $X$  are finite sets. We are to prove that the projection  $Y \rightarrow X$  is a finite morphism. We may assume that  $X$  is affine, say  $X = \text{Spec } A$ , and by induction on  $n$ , we may assume that the theorem is true when  $\mathbb{P}$  is a projective space of dimension  $n-1$ .

*Case 1.* *There is a hyperplane  $H$  in  $\mathbb{P}$  such that  $Y$  is disjoint from  $\tilde{H} = H \times X$  in  $\tilde{\mathbb{P}} = \mathbb{P} \times X$ .*

This is the main case. We adjust coordinates  $y_0, \dots, y_n$  in  $\mathbb{P}$  so that  $H$  is the hyperplane  $\{y_0 = 0\}$ . Because  $Y$  is a closed subvariety of  $\tilde{\mathbb{P}}$  disjoint from  $\tilde{H}$ ,  $Y$  is also a closed subvariety of  $\tilde{\mathbb{U}}^0 = \mathbb{U}^0 \times X$ ,  $\mathbb{U}^0$  being the standard affine  $\{y_0 \neq 0\}$ . So  $Y$  is affine.

Let  $\mathcal{P}$  and  $\mathcal{Q}$  be the (homogeneous) prime ideals in  $A[y]$  that define  $Y$  and  $\tilde{H}$ , respectively, and let  $\mathcal{I} = \mathcal{P} + \mathcal{Q}$ . So  $\tilde{q}$  is the principal ideal of  $A[y]$  generated by  $y_0$ . A homogeneous element of  $\mathcal{I}$  of degree  $k$  has the form  $f(y) + y_0 g(y)$ , where  $f$  is a homogeneous polynomial in  $A[y]$  of some degree  $k$ , and  $g$  is a homogeneous polynomial of degree  $k-1$ .

The closed subsets  $Y$  and  $\tilde{H}$  are disjoint. Since  $Y \cap \tilde{H}$  is empty, the sum  $\mathcal{I} = \mathcal{P} + \mathcal{Q}$  contains a power of the irrelevant ideal  $\mathcal{M} = (y_0, \dots, y_n)$ , say  $\mathcal{M}^k \subset \mathcal{I}$ . Then  $y_i^k$  is in  $\mathcal{I}$  for  $i = 0, \dots, n$ . So we may write

$$(4.6.10) \quad y_i^k = f_i(y) + y_0 g_i(y)$$

with  $f_i$  of degree  $k$  in  $\mathcal{P}$  and  $g_i$  of degree  $k-1$  in  $A[y]$ . We omit the index  $i = 0$ . For that index, one can take  $f_0 = 0$  and  $g_0 = y_0^{k-1}$ .

We dehomogenize these equations with respect to the variables  $y$ , substituting  $u_i = y_i/y_0$  for  $y_i$ ,  $i = 1, \dots, n$  with  $u_0 = 1$ . Writing dehomogenizations with capital letters, the dehomogenized equations that

correspond to the equations  $y_i^k = f_i(y) + y_0 g(y)$  have the form

$$(4.6.11) \quad u_i^k = F_i(u) + G_i(u)$$

The important point is that the degree of  $G_i$  is at most  $k-1$ .

Recall that  $Y$  is also a closed subset of  $\mathbb{U}^0$ . Let  $P$  be its (nonhomogeneous) ideal in  $A[u]$ , which contains the polynomials  $F_1, \dots, F_n$ . The coordinate algebra of  $Y$  is  $B = A[u]/P$ . In the quotient algebra  $B$ , the terms  $F_i$  drop out, leaving us with equations  $u_i^k = G_i(u)$ . These equations are true in  $B$ . Since  $G_i$  has degree at most  $k-1$ , Lemma 4.6.8 tells us that  $B$  is a finite  $A$ -algebra, as was to be shown. This completes the proof of Case 1.

*Case 2. the general case.*

We have taken care of the case in which there exists a hyperplane  $H$  such that  $Y$  is disjoint from  $\tilde{H}$ . The next lemma shows that we can cover the given variety  $X$  by open subsets to which this special case applies. Then Lemma 4.6.4 and Proposition 4.6.4 will complete the proof.

**4.6.12. Lemma.** *Let  $Y$  be a closed subvariety of  $\tilde{\mathbb{P}} = \mathbb{P}^n \times X$ , and suppose that the projection  $Y \xrightarrow{\pi} X$  has finite fibres. Suppose also that Chevalley's Theorem has been proved for closed subvarieties of  $\mathbb{P}^{n-1} \times X$ . For every point  $p$  of  $X$ , there is an open neighborhood  $X'$  of  $p$  in  $X$ , and there is a hyperplane  $H$  in  $\mathbb{P}$ , such that the inverse image  $Y' = \pi^{-1}X'$  is disjoint from  $\tilde{H}$ .*

*proof.* Let  $p$  be a point of  $X$ , and let  $\tilde{q} = (\tilde{q}_1, \dots, \tilde{q}_r)$  be the finite set of points of  $Y$  making up the fibre over  $p$ . We project  $\tilde{q}$  from  $\mathbb{P} \times X$  to  $\mathbb{P}$ , obtaining a finite set  $q = (q_1, \dots, q_r)$  of points of  $\mathbb{P}$ , and we choose a hyperplane  $H$  in  $\mathbb{P}$  that avoids this finite set. Then  $\tilde{H}$  avoids the fibre  $\tilde{q}$ . Let  $W$  denote the closed set  $Y \cap \tilde{H}$ . Because the fibres of  $Y$  over  $X$  are finite, so are the fibres of  $W$  over  $X$ . By hypothesis, Chevalley's Theorem is true for subvarieties of  $\mathbb{P}^{n-1} \times X$ , and  $\tilde{H}$  is isomorphic to  $\mathbb{P}^{n-1} \times X$ . It follows that, for every component  $W'$  of  $W$ , the morphism  $W' \rightarrow X$  is a finite morphism, and therefore its image is closed in  $X$  (Theorem 4.4.1). Thus the image  $Z$  of  $W$  is a closed subset of  $X$ , and it doesn't contain  $p$ . Then  $X' = X - Z$  is the required neighborhood of  $p$ .  $\square$

figure: ??I'm not sure

## 4.7 Dimension

Every variety has a dimension, and though it is a very coarse measure, the dimension is important, as is true for the dimension of a vector space.

A *chain* of closed subvarieties of a variety  $X$  is a strictly decreasing sequence of closed subvarieties – irreducible closed subsets

$$(4.7.1) \quad C_0 > C_1 > C_2 > \dots > C_k$$

The *length* of this chain is defined to be  $k$ . The chain is *maximal* if it cannot be lengthened by inserting another closed subvariety, which means that  $C_0 = X$ , that there is no closed subvariety  $\tilde{C}$  with  $C_i > \tilde{C} > C_{i+1}$  for  $i < k$ , and that  $C_k$  is a point. Theorem 4.7.3 below shows that all maximal chains have the same length, and the *dimension* of  $X$ , often denoted by  $\dim X$ , is the length of a maximal chain.

When  $X$  is an affine variety, say  $X = \text{Spec } A$ , the decreasing chain (4.7.1) corresponds to an increasing chain

$$(4.7.2) \quad P_0 < P_1 < P_2 < \dots < P_k,$$

of prime ideals of  $A$  of length  $k$ , a *prime chain*. This prime chain is *maximal* if it cannot be lengthened by inserting another prime ideal, which means that  $P_0$  is the zero ideal, that there is no prime ideal  $\tilde{P}$  with  $P_i < \tilde{P} < P_{i+1}$  for  $i < k$ , and that  $P_k$  is a maximal ideal. The *dimension*  $\dim A$  of a finite-type domain  $A$  is the length  $k$  of a maximal chain (4.7.2) of prime ideals. Thus if  $X = \text{Spec } A$ , then  $\dim X = \dim A$ .

**4.7.3. Theorem.** (i) *Let  $A$  be a finite-type domain whose fraction field  $K$  has transcendence degree  $n$ . All prime chains in  $A$  have length at most  $n$ , and all maximal prime chains have length equal to  $n$ . Therefore the dimension of  $A$  is the transcendence degree of  $A$ .*

(ii) Let  $X$  be a variety whose function field  $K$  has transcendence degree  $n$ . All chains of closed subvarieties of  $X$  have length at most  $n$ , and all maximal chains have length  $n$ . Therefore the dimension of  $X$  is equal to the transcendence degree of  $K$ .

The proof is below.

For example, the transcendence degree of the polynomial algebra  $\mathbb{C}[x_1, \dots, x_n]$  in  $n$  variables is  $n$ , so the polynomial algebra has dimension  $n$ . The chain of prime ideals

$$(4.7.4) \quad 0 < (x_1) < (x_1, x_2) < \dots < (x_1, \dots, x_n)$$

is a maximal prime chain. The corresponding chain

$$\mathbb{P}^n > \mathbb{P}^{n-1} > \dots > \mathbb{P}^0$$

is a maximal chain of closed subvarieties of projective space  $\mathbb{P}^n$ .

The maximal chains of closed subvarieties of  $\mathbb{P}^2$  have the form  $\mathbb{P}^2 > C > p$ , where  $C$  is a plane curve and  $p$  is a point.

If (4.7.1) is a maximal chain in  $X$ , then

$$(4.7.5) \quad C_i > C_{i+1} > \dots > C_k$$

will be a maximal chain in the variety  $C_i$ . So when  $X$  has dimension  $k$ , the dimension of  $C_i$  is  $k - i - 1$ . Similarly, let (4.7.2) be a maximal chain of prime ideals in a finite-type domain  $A$ , let  $\bar{A} = A/P_i$  and let  $\bar{P}_j$  denote the image  $P_j/P_i$  of  $P_j$  in  $\bar{A}$ , for  $j \geq i$ . The Correspondence Theorem implies that

$$\bar{0} = \bar{P}_i < \bar{P}_{i+1} < \dots < \bar{P}_k$$

will be a maximal prime chain in  $\bar{A}$ , and therefore that the dimension of the domain  $\bar{A}$  is  $k - i - 1$ . There is a bijective correspondence between maximal prime chains in  $\bar{A}$  and maximal prime chains in  $A$  whose first terms are  $P_0 < P_1 < \dots < P_{i-1}$ .

One more term: A closed subvariety  $C$  of a variety  $X$  has *codimension*  $k$  if  $\dim X - \dim C = k$ . In particular,  $C$  has codimension 1 if  $C < X$  and if there is no closed set  $\tilde{C}$  with  $C < \tilde{C} < X$ . A prime ideal  $P$  of a noetherian domain has *codimension* 1 if it is not the zero ideal, and if there is no prime ideal  $\tilde{P}$  with  $(0) < \tilde{P} < P$ . In the polynomial algebra  $\mathbb{C}[x_1, \dots, x_n]$ , the prime ideals of codimension 1 are the principal ideals generated by irreducible polynomials.

For the proof of the next proposition, see Lemma 4.4.5, and Theorem 4.4.2(i).

**4.7.6. Proposition.** Let  $Y \xrightarrow{u} X$  be an integral morphism of varieties. Every chain of closed subvarieties of  $Y$ , lies over a chain in  $X$ , and every chain of closed subvarieties of  $X$  has chain in  $Y$  lying over it.  $\square$

**4.7.7. Lemma.** Let  $A = \mathbb{C}[x_1, \dots, x_n]$  be a polynomial algebra, let  $f$  be an irreducible element of  $A$ , and let  $\bar{A} = A/(f)$ . The transcendence degree of  $\bar{A}$  is  $n - 1$ .

*proof.* We may choose coordinates so that  $f$  becomes a monic polynomial in  $x_n$  with coefficients in  $\mathbb{C}[x_1, \dots, x_{n-1}]$ , say  $f = x_n^k + c_{k-1}x_n^{k-1} + \dots + c_0$  (Lemma 4.2.9). Then  $\bar{A}$  will be integral over  $\mathbb{C}[x_1, \dots, x_{n-1}]$ , so it will have the same transcendence degree.  $\square$

*proof of theorem 4.7.3.* (i) Induction allows us to assume the theorem true for a finite-type domain whose transcendence degree is less than  $n$ . Let  $A$  be a finite-type domain of transcendence degree  $n$ .

*Case I:* The case that  $A$  is a polynomial algebra  $\mathbb{C}[x_1, \dots, x_n]$ .

Let  $P_0 < P_1 < \dots < P_k$  be a prime chain in  $A$ . We are to show that  $k \leq n$ , and that  $k = n$  if the chain is maximal. We may assume that  $P_0 = 0$  and that  $P_1$  is a codimension 1 prime, generated by an irreducible polynomial  $f$ . (If not, we can insert a prime ideal into the chain.) Let  $\bar{A} = A/P_1$ , and for  $i \geq 1$ , let  $\bar{P}_i = P_i/P_1$ . Then  $\bar{P}_1 < \bar{P}_2 < \dots < \bar{P}_k$  is a prime chain in  $\bar{A}$  of length  $k - 1$ , and if the chain  $\{P_i\}$  is maximal, the chain  $\{\bar{P}_i\}$  will be a maximal chain too. Lemma 4.7.7 shows that  $\bar{A}$  has transcendence degree  $n - 1$ . So by induction, the length of the chain  $\{\bar{P}_i\}$  is at most  $n - 1$  and is equal to  $n - 1$  if the chain  $\{P_i\}$  is maximal. Therefore the chain  $\{P_i\}$  has length at most  $n$  and has length  $n$  if it is maximal.

*Case 2: The general case.*

Let  $B$  be a finite-type domain of transcendence degree  $n$ , and let  $Q_0 < Q_1 < \cdots < Q_k$  be a prime chain in  $B$ . Again, we are to show that  $k \leq n$  and that if the chain is maximal, then  $k = n$ . We apply the Noether Normalization Theorem:  $B$  is a finite module over a polynomial subring  $A$ . The transcendence degree of  $A$  is  $n$ , and the contractions  $P_i = Q_i \cap A$  form a prime chain in  $A$ . This is the translation of Proposition 4.7.6 to prime ideals. Therefore  $k \leq n$ .

Next, suppose that the chain  $\{Q_i\}$  is maximal. Then  $Q_0 = 0$  and therefore  $P_0 = 0$ . If  $P_1$  were not a codimension 1 prime, we could choose a nonzero prime ideal  $\tilde{P}$  contained in  $P_1$ . Since  $A$  is normal, we could apply Theorem 4.4.2: there would be a nonzero prime ideal  $\tilde{Q}$  of  $B$  that lies over  $\tilde{P}$  and is contained in  $Q_1$ . This would imply that the chain  $\{Q_i\}$  wasn't maximal, contrary to hypothesis. So  $P_1$  is a codimension 1 prime, and by Lemma 4.7.7,  $\bar{A} = A/P_1$  has transcendence degree  $n - 1$ . Since  $\bar{B} = B/Q_1$  is a finite  $\bar{A}$ -module, it also has transcendence degree  $n - 1$ . Let  $\bar{Q}_i = Q_i/Q_1$  for  $i \geq 1$ . By induction, the length of the maximal chain  $\bar{Q}_1 < \cdots < \bar{Q}_k$  in  $\bar{B}$  is  $n - 1$ , and therefore  $k = n$ .

Part (ii) of Theorem 4.7.3 follows from the next lemma.

**4.7.8. Lemma.** *Let  $X'$  be an open subvariety of a variety  $X$ . There is a bijective correspondence between chains  $C_0 > \cdots > C_k$  of closed subvarieties of a variety  $X$  such that  $C_k \cap X' \neq \emptyset$  and chains  $C'_0 > \cdots > C'_k$  of closed subvarieties of  $X'$ , defined by  $C'_i = C_i \cap X'$ . Given a chain  $C'_i$  in  $X'$ , the corresponding chain in  $X$  consists of the closures  $C_i$  in  $X$  of the varieties  $C'_i$ .*

*proof.* Suppose given a chain  $C_i$  and that  $C_k \cap X' \neq \emptyset$ . Then the intersections  $C'_i = C_i \cap X'$  are dense open subsets of the irreducible closed sets  $C_i$  (2.2.12). So the closure of  $C'_i$  is  $C_i$ , and since  $C_i > C_{i+1}$ , it is also true that  $C'_i > C'_{i+1}$ . Therefore  $C'_0 > \cdots > C'_k$  is a chain of closed subsets of  $X'$ . Conversely, if  $C'_0 > \cdots > C'_k$  is a chain in  $X'$ , the closures in  $X$  will form a chain in  $X$ .  $\square$

**4.7.9. Corollary.** (i) *If  $X_s$  is a localization of an affine variety  $X$ , then  $\dim X_s = \dim X$ .*

(ii) *If  $Y$  is a proper closed subvariety of a variety  $X$ , then  $\dim Y < \dim X$ .*

(iii) *If  $Y \rightarrow X$  is an integral morphism of varieties, then  $\dim Y = \dim X$ .*  $\square$

## 4.8 Krull's Theorem

Krull's Principal Ideal Theorem completes our discussion of dimension. It asserts that the zero set of a principal ideal can't have a low dimension. Though the statement is natural, the proof isn't very easy.

**4.8.1. Krull's Theorem.** Let  $X$  be an affine variety of dimension  $d$ , and let  $\alpha$  be a nonzero element of its coordinate ring  $A$ . Every irreducible component of the zero locus  $V_X(\alpha)$  of  $\alpha$  in  $X$  has dimension  $d - 1$ .

*proof.* Let  $C$  be an irreducible component of  $V_X(\alpha)$ . Since  $\alpha$  isn't zero,  $C$  is a proper subset of  $X$ , and its dimension is less than  $d$ . We must show that the dimension is  $d - 1$ , and we prove this by contradiction. So we assume that  $\dim C < d - 1$ .

*Step 1: Reduction to the case that  $A$  is normal.*

Let  $B$  be the normalization of  $A$  and let  $Y = \text{Spec } B$ . The dimension of  $Y$  is  $d$ . The integral morphism  $Y \rightarrow X$  is surjective, and it sends closed sets to closed sets (4.4.1). So the zero locus of  $\alpha$  in  $Y$  maps surjectively to the zero locus in  $X$ , and at least one irreducible component of  $V_Y(\alpha)$ , call it  $D$ , will map surjectively to  $C$ . The map  $D \rightarrow C$  is also an integral morphism, so the dimension of  $D$  is the same as that of  $C$ . We may therefore replace  $X$  by  $Y$  and  $C$  by  $D$ . Hence we may assume that  $A$  is normal.

*Step 2: Reduction to the case that the zero locus of  $\alpha$  is irreducible.*

We do this by localizing. Say that the zero locus is  $C \cup V$ , where  $C$  is a closed subvariety of codimension at least two, and  $V$  is the union of the other irreducible components. We choose an element  $s$  of  $A$  that is identically zero on  $V$  but not identically zero on  $C$ . Inverting  $s$  eliminates the points of  $V$ , but  $X_s \cap C = C_s$  will be nonempty. If  $X$  is normal, so is  $X_s$ . Since localization doesn't change dimensions, we may replace  $X$  and  $C$  by  $X_s$  and  $C_s$ .

*Step 3: Completion of the proof.*

This is the main step: We assume that  $X$  is a normal affine variety,  $X = \text{Spec } A$ , and that the zero locus of  $\alpha$  in  $X$  is a closed subvariety  $C$  of codimension at least two. Then  $C$  is the zero locus of a prime ideal  $P$ , and also the zero locus of  $\alpha$ . So  $P$  is the radical of the principal ideal  $\alpha A$ , and  $P^n \subset \alpha A$  if  $n$  is large (see 2.5.10).

By what we know about dimension,  $C$  will be contained in a closed subvariety  $Z$  of codimension one. Let  $Q$  be the prime ideal whose locus is  $Z$ . Then  $P \supset Q$  because  $C \subset Z$ . On the other hand,  $\alpha \notin Q$  because  $\alpha$  vanishes only on  $C$ , and it follows that the principal ideal  $\alpha A$  isn't contained in  $Q$ . Since  $P$  is the radical of  $\alpha A$ , these two facts are in stress.

**4.8.2. Lemma.** *With notation as above, There is an element  $\gamma$  in  $A$  such that  $\gamma \notin \alpha A$  but  $P\gamma \subset \alpha A$ .*

*proof.* Let  $\beta$  be an element of  $Q$ . Corollary 4.1.5 (iii) tells us that the powers of  $\alpha$  that divide  $\beta$  are bounded. Let  $\alpha^k$  be the largest such power, and let  $\gamma = \beta/\alpha^k$ . Then  $\gamma$  vanishes on the dense open subset  $Z' = Z - C$  of  $Z$ , so it vanishes on  $Z$ . There is an element  $\gamma$  in  $Q$  that isn't in the principal ideal  $\alpha A$ .

Next, since  $P^n \subset \alpha A$  for large  $n$ , it is also true that  $P^n Q \subset \alpha A$  for large  $n$ . Let  $r$  be the largest integer  $\geq 0$  such that  $P^r Q \not\subset \alpha A$ . We replace  $\gamma$  by an element of  $P^r Q$  that isn't in  $\alpha A$ . Then  $\gamma \notin \alpha A$ , but  $P\gamma \subset \alpha A$ .  $\square$

Krull's Theorem follows easily from this lemma. Let  $\delta = \gamma/\alpha$ , where  $\gamma$  is as in the lemma. Then  $\delta \notin A$ , but  $P\delta \subset P$ . Since  $\gamma$  vanishes on  $Z$  while  $\alpha$  vanishes only on  $C$ , every element  $\epsilon$  of  $P\delta$  vanishes on the dense complement  $Z'$  of  $C$  in  $Z$ , and therefore on  $Z$ . So  $P\delta \subset Q \subset P$ . Corollary 5.2.7 shows that  $\delta$  is integral over  $A$ , and since  $A$  is assumed normal,  $\delta$  is in  $A$ . This is a contradiction that proves the theorem.  $\square$

**4.8.3. Corollary.** *Let  $Z$  be a proper closed subset of a variety  $X$  of dimension  $d$ , and let  $p$  be a point of  $Z$ . There is a maximal chain of closed subvarieties  $X = X_0 > X_1 > \cdots > X_d = \{p\}$  such that  $X_{d-1}$  isn't contained in  $Z$ , and therefore  $X_i$  isn't contained in  $Z$  for any  $i < d$ .*

*proof.* Lemma 4.7.8 shows that we may assume that  $X$  is affine,  $X = \text{Spec } A$ . Lemma 4.5.3 asserts that  $A$  contains an element  $\alpha$  that vanishes at  $p$  but doesn't vanish identically on any component of  $Z$ . Then at least one component of  $V_X(\alpha)$  contains  $p$ . Let  $X_1$  be such a component. Krull's Theorem tells us that  $X_1$  has dimension  $d - 1$ . Since  $\alpha$  doesn't vanish identically on any component of  $Z$ ,  $X_1$  doesn't contain any of those components. Therefore  $Z_1 = Z \cap X_1$  is a proper closed subvariety of  $X_1$ . We replace  $X$  by  $X_1$ . Then the corollary follows by induction on the dimension  $d$ .  $\square$

## 4.9 Double Planes

### (4.9.1) affine double planes

Let  $A$  be the polynomial algebra  $\mathbb{C}[x, y]$ , and let  $X$  be the affine plane  $\text{Spec } A$ . An *affine double plane* is a locus of the form  $w^2 = f(x, y)$  in affine 3-space with coordinates  $w, x, y$ , where  $f$  is a square-free polynomial in  $x, y$ , as in Example 4.3.8. Let  $B = \mathbb{C}[w, x, y]/(w^2 - f)$ . So the affine double plane is  $Y = \text{Spec } B$ .

We'll denote by  $w, x, y$  both the variables and their residues in  $B$ .

**4.9.2. Lemma** *The algebra  $B$  is a normal domain of dimension two, and a free  $A$ -module with basis  $(1, w)$ . It has an automorphism  $\sigma$  of order 2, defined by  $\sigma(a + bw) = a - bw$ .  $\square$*

The fibres of  $Y$  over  $X$  are the  $\sigma$ -orbits in  $Y$ . If  $f(x_0, y_0) \neq 0$ , the fibre consists of two points, and if  $f(x_0, y_0) = 0$ , it consists of one point. The reason that  $Y$  is called a *double plane* is that most points of the plane  $X$  are covered by two points of  $Y$ . The *branch locus* of the covering, which will be denoted by  $\Delta$ , is the (possibly reducible) curve  $\{f = 0\}$  in  $X$ . The fibres over the *branch points*, points of  $\Delta$ , are single points.

We study the closed subvarieties  $D$  of  $Y$  that lie over a curve  $C$  in  $X$ . These subvarieties will have dimension one, and we call them curves too. If  $D$  lies over  $C$ , and if  $D = D\sigma$ , then  $D$  is the only curve lying over  $C$ . Otherwise, there will be the two curves that lie over  $C$ , namely  $D$  and  $D\sigma$ . In that case we say that  $C$  *splits* in  $Y$ .

A curve  $C$  in  $X$  will be the zero set of a principal prime ideal  $P$  of  $A$ , and if  $D$  lies over  $C$ , it will be the zero set of a prime ideal  $Q$  of  $B$  that lies over  $P$  (??). The prime ideal  $Q$  isn't always a principal ideal.

**4.9.3. Example.** Let  $f(x, y) = x^2 + y^2 - 1$ . The double plane  $Y = \{w^2 = x^2 + y^2 - 1\}$  is an *affine quadric* in  $\mathbb{A}^3$ . In the affine plane, its branch locus  $\Delta$  is the curve  $\{x^2 + y^2 = 1\}$ .

The line  $C_1 : \{y = 0\}$  in  $X$  meets the branch locus  $\Delta$  transversally at the points  $(x, y) = (\pm 1, 0)$ , and  $y$  generates a prime ideal of  $B$ . When we set  $y = 0$  in the equation for  $Y$ , we obtain the irreducible polynomial  $w^2 - x^2 + 1$ . On the other hand, the line  $C_2 : \{y = 1\}$  is tangent to  $\Delta$  at the point  $(0, 1)$ , and it splits. When we set  $y = 1$  in the equation for  $Y$ , we obtain  $w^2 = x^2$ . The locus  $\{w^2 = x^2\}$  is the union of the two lines  $\{w = x\}$  and  $\{w = -x\}$  that lie over  $C_1$ . The prime ideals of  $B$  that correspond to these lines aren't principal ideals.

figure circle with two lines

□

This example illustrates a general principle: If a curve intersects the branch locus transversally, it doesn't split. We explain this now.

#### (4.9.4) local analysis

Suppose that a plane curve  $C : \{g = 0\}$  and the branch locus  $\Delta : \{f = 0\}$  of a double plane  $w^2 = f$  meet at a point  $p$ . We adjust coordinates so that  $p$  becomes the origin  $(0, 0)$ , and we write

$$f(x, y) = \sum a_{ij} x^i y^j = a_{10}x + a_{01}y + a_{20}x^2 + \cdots$$

Since  $p$  is a point of  $\Delta$ , the constant coefficient of  $f$  is zero. If the two linear coefficients aren't both zero,  $p$  will be a smooth point of  $\Delta$ , and the tangent line to  $\Delta$  at  $p$  will be the line  $\{a_{10}x + a_{01}y = 0\}$ . Similarly, writing  $g(x, y) = \sum b_{ij} x^i y^j$ , the tangent line to  $C$ , if defined, is the line  $\{b_{10}x + b_{01}y = 0\}$ .

Let's suppose that the two tangent lines are defined and distinct – that  $\Delta$  and  $C$  intersect transversally at  $p$ . We change coordinates once more, to make the tangent lines the coordinate axes. After adjusting by scalar factors, the polynomials  $f$  and  $g$  will have the form

$$f(x, y) = x + u(x, y) \quad \text{and} \quad g(x, y) = y + v(x, y),$$

where  $u$  and  $v$  are polynomials all of whose terms have degree at least 2.

Let  $X_1 = \text{Spec } \mathbb{C}[x_1, y_1]$  be another affine plane. We consider the map  $X_1 \rightarrow X$  defined by the substitution  $x_1 = x + u$ ,  $y_1 = y + v$ . In the classical topology, this map is invertible analytically near the origin, because the Jacobian matrix

$$(4.9.5) \quad \left( \frac{\partial(x_1, y_1)}{\partial(x, y)} \right)_{(0,0)}$$

at  $p$  is the identity matrix. When we make this substitution,  $\Delta$  becomes the locus  $\{x_1 = 0\}$  and  $C$  becomes the locus  $\{y_1 = 0\}$ . In this local analytic coordinate system, the equation  $w^2 = f$  that defines the double plane becomes  $w^2 = x_1$ . When we restrict it to  $C$  by setting  $y_1 = 0$ ,  $x_1$  becomes a local coordinate function on  $C$ . The restriction of the equation remains  $w^2 = x_1$ . So the inverse image  $Z$  of  $C$  doesn't split analytically near  $p$ . Therefore it doesn't split globally either.

**4.9.6. Corollary.** *A curve that meets the branch locus transversally at some point doesn't split.* □

This isn't a complete analysis. When  $C$  and  $\Delta$  are tangent at every point of intersection,  $C$  may split or not, and which possibility occurs cannot be decided locally in most cases. However, one case in which a local analysis suffices to decide splitting is that  $C$  is a line. Let  $t$  be a coordinate in a line  $C$ , so that  $C \approx \text{Spec } \mathbb{C}[t]$ . Let's assume that  $C$  doesn't intersect  $\Delta$  at  $t = \infty$ . The restriction of the polynomial  $f$  to  $C$  will give us a polynomial  $\bar{f}(t)$  in  $t$ . A root of  $\bar{f}$  corresponds to an intersection of  $C$  with  $\Delta$ , and a multiple root corresponds to an intersection at which  $C$  and  $\Delta$  are tangent, or at which  $\Delta$  is singular. The line  $C$  will split if and only if  $\bar{f}$  is a square in  $\mathbb{C}[t]$ , and this will be true if and only if the multiplicity of every root of  $\bar{f}$  is even.

A *rational curve* is a curve whose function field is a rational function field  $\mathbb{C}(t)$  in one variable. One can make a similar analysis for any rational plane curve, a conic for example, but one needs to inspect its points at infinity and its singular points as well as the smooth points at finite distance.

#### (4.9.7) projective double planes



Let  $X$  be the projective plane  $\mathbb{P}^2$ , with coordinates  $x_0, x_1, x_2$ . A *projective double plane* is a locus of the form

$$(4.9.8) \quad y^2 = f(x_0, x_1, x_2),$$

where  $f$  is a square-free, homogeneous polynomial of even degree  $2d$ . To regard this as a homogeneous equation, we must assign weight  $d$  to the variable  $y$  (see 1.7.7). Then, since we have weighted variables, we must work in a *weighted projective space*  $\mathbb{WP}$  with coordinates  $x_0, x_1, x_2, y$ , where  $x_i$  have weight 1 and  $y$  has weight  $d$ . A point of this weighted space  $\mathbb{WP}$  is represented by a nonzero vector  $(x_0, x_1, x_2, y)$  with the relation that, for all  $\lambda \neq 0$ ,  $(x_0, x_1, x_2, y) \sim (\lambda x_0, \lambda x_1, \lambda x_2, \lambda^d y)$ . The points of the projective double plane  $Y$  are the points of  $\mathbb{WP}$  that solve the equation (4.9.8).

The projection  $\mathbb{WP} \rightarrow X$  that sends  $(x, y)$  to  $x$  is defined at all points except at  $(0, 0, 0, 1)$ . If  $(x, y)$  solves (4.9.8) and if  $x = 0$ , then  $y = 0$  too. So  $(0, 0, 0, 1)$  isn't a point of  $Y$ . The projection is defined at all points of  $Y$ . The fibre of the morphism  $Y \rightarrow X$  over a point  $x$  consists of points  $(x, y)$  and  $(x, -y)$ , which will be equal if and only if  $x$  lies on the *branch locus* of the double plane, the (possibly reducible) plane curve  $\Delta : \{f = 0\}$  in  $X$ . The map  $\sigma : (x, y) \rightsquigarrow (x, -y)$  is an automorphism of  $Y$ , and points of  $X$  correspond bijectively to  $\sigma$ -orbits in  $Y$ .

Since the double plane  $Y$  is embedded into a weighted projective space, it isn't presented to us as a projective variety in the usual sense. However, it can be embedded into a projective space in the following way: The projective plane  $X$  can be embedded by a Veronese embedding of higher order, using as coordinates the monomials  $m = (m_1, m_2, \dots)$  of degree  $d$  in the variables  $x$ . This embeds  $X$  into a projective space  $\mathbb{P}^N$  where  $N = \binom{d+2}{2} - 1$ . When we add a coordinate  $y$  of weight  $d$ , we obtain an embedding of the weighted projective space  $\mathbb{WP}$  into  $\mathbb{P}^{N+1}$  that sends the point  $(x, y)$  to  $(m, y)$ . The double plane can be realized as a projective variety by this embedding.

If  $Y \rightarrow X$  is a projective double plane, then, as happens with affine double planes, a curve  $C$  in  $X$  may split in  $Y$  or not. If  $C$  has a transversal intersection with the branch locus  $\Delta$ , it will not split. On the other hand, if  $C$  is a line, and if  $C$  intersects the branch locus  $\Delta$  with multiplicity 2 at every intersection point, it will split. For example, when the branch locus  $\Delta$  is a generic quartic curve, the lines that split will be the bitangent lines (see Section 1.11).

#### (4.9.9) homogenizing an affine double plane

To construct a projective double plane from an affine double plane, we write the affine double plane as

$$(4.9.10) \quad w^2 = F(u_1, u_2)$$

for some nonhomogeneous polynomial  $F$ . We suppose that  $F$  has even degree  $2d$ , and we homogenize  $F$ , setting  $u_i = x_i/x_0$ . We multiply both sides of this equation by  $x_0^{2d}$  and set  $y = x_0^d w$ . This produces an equation of the form (4.9.8), where  $f$  is the homogenization of  $F$ .

If  $F$  has odd degree  $2d - 1$ , one needs to multiply  $F$  by  $x_0$  in order to make the substitution  $y = x_0^d w$  permissible. When we do this, the line at infinity  $\{x_0 = 0\}$  becomes a part of the branch locus.

#### (4.9.11) cubic surfaces and quartic double planes

We use coordinates  $x_0, x_1, x_2, z$  for the (unweighted) projective 3-space  $\mathbb{P}^3$  here, and  $X$  will denote the projective  $x$ -plane  $\mathbb{P}^2$ . Let  $\mathbb{P}^3 \xrightarrow{\pi} X$  denote the projection that sends  $(x, z)$  to  $x$ . It is defined at all points except at the center of projection  $q = (0, 0, 0, 1)$ , and its fibres are the lines through  $q$ , with  $q$  omitted.

Let  $S$  be a cubic surface in  $\mathbb{P}^3$ , the locus of zeros of an irreducible homogeneous cubic polynomial  $g(x, z)$ . We'll denote the restriction of  $\pi$  to  $S$  by the same symbol  $\pi$ .

Let's suppose that  $q$  is a point of  $S$ . Then the coefficient of  $z^3$  in  $g$  will be zero, and  $g$  will be quadratic in  $z$ :  $g(x, z) = az^2 + bz + c$ , where the coefficients  $a, b, c$  are homogeneous polynomials in  $x$ , of degrees 1, 2, 3, respectively. The equation for  $S$  becomes

$$(4.9.12) \quad az^2 + bz + c = 0$$

The discriminant  $f = b^2 - 4ac$  of  $g$  is a homogeneous polynomial of degree 4 in  $x$ . Let  $Y$  be the projective double plane

$$(4.9.13) \quad y^2 = b^2 - 4ac$$

We denote by  $V$  the affine space of polynomials  $a, b, c$  of degrees 1, 2, 3 in  $x$ , and by  $W$  the affine space of homogeneous quartic polynomials in  $x$ . Sending  $g$  to its discriminant  $f$  defines a morphism  $V \xrightarrow{u} W$  (4.9.12).

**4.9.14. Lemma.** *The image of the morphism  $u$  contains all quartic polynomials  $f$  such that the divisor  $D : f = 0$  has at least one bitangent line. Therefore the image of  $u$  is dense in  $W$ .*

*proof.* Given such a quartic polynomial  $f$ , let  $a$  be a linear polynomial such that the line  $\ell_1 : \{a = 0\}$  is a bitangent to  $D : \{f = 0\}$ . Then, as noted above,  $\ell_1$  splits in the double plane  $y^2 = f$ . So  $f$  is congruent to a square, modulo  $a$ . Let  $b$  be a quadratic polynomial such that  $f \equiv b^2$  modulo  $a$ . When we take this polynomial as  $b$ , we will have  $f = b^2 - 4ac$  for some cubic polynomial  $c$ .

Conversely, if  $g(x, y) = az^2 + bz + c$ , the line  $\ell_1 : \{a = 0\}$  will be a bitangent to  $D$  provides that the locus  $b = 0$  meets  $\ell_1$  in two distinct points.  $\square$

It follows from the lemma that, if  $g(x, z) = az^2 + bz + c$  is a polynomial in which  $a, b, c$  are generic homogeneous polynomials in  $x$ , of degrees 1, 2, 3, respectively, the discriminant  $b^2 - 4ac$  will be a generic homogeneous quartic polynomial in  $x$ .

We go back to the generic cubic surface  $S : az^2 + bz + c = 0$  and the generic double plane  $Y : y^2 = b^2 - 4ac$ .

**4.9.15. Theorem.** *A generic cubic surface  $S$  in  $\mathbb{P}^3$  contains precisely 27 lines.*

This theorem follows from next lemma, which relates the 27 lines in  $S$  to the 28 bitangents of the generic quartic curve  $\Delta : \{b^2 - 4ac = 0\}$  in the plane  $X$  (1.11.2).

As noted above, the line  $\ell_1$  defined by the linear equation  $a = 0$  is a bitangent to the quartic curve  $\Delta$ .

**4.9.16. Lemma.** *Let  $S$  be a generic cubic surface. The 27 bitangent lines in  $X$  that are distinct from  $\ell_1$  are the images of the lines in  $S$ , and distinct lines in  $S$  have distinct images.*

*proof.* Because the cubic surface  $S$  is generic, it contains finitely many lines (3.6). When we project to  $X$  from a generic point  $q$  of  $S$ ,  $q$  won't lie on any of those lines. The fibres of the projection  $\mathbb{P}^3 \rightarrow X$  are lines through  $q$ , and they aren't contained in  $S$ . So a line in  $S$  projects bijectively to a line in  $X$ .

A line in  $X$  is defined by a homogeneous linear equation in the variables  $x$ . The same linear equation defines a plane  $H$  in  $\mathbb{P}^3$  that contains  $q$ , and the intersection  $C = S \cap H$  will be a cubic curve in  $H$ . At least one of the irreducible components of  $C$  contains  $q$ , and that component isn't a line. So if  $C$  is reducible, it will be a union  $Q \cup L$ , where  $Q$  is a conic that contains  $q$  and  $L$  is a line in  $S$ . Thus lines  $L$  in  $S$  correspond bijectively to lines in  $X$  such that the corresponding cubic  $C$  is reducible.

Referring to (4.9.12) and (4.9.13), the quadratic formula solves for  $z$  in terms of  $y$  whenever  $a \neq 0$ :

$$(4.9.17) \quad z = \frac{-b + y}{2a} \quad \text{or} \quad y = 2az + b$$

These equations define a bijection  $S' \longleftrightarrow Y'$  between the open subsets  $S'$  and  $Y'$  of points of  $S$  and  $Y$  at which  $a \neq 0$ .

If  $\ell$  is a line in  $X$ , not the line  $\ell_1$ , the intersection  $\ell \cap \ell_1$  will be a point  $p$ . The bijection  $S' \longleftrightarrow Y'$  will be defined at all points that lie over  $\ell$  except those whose images are  $p$ . If  $\ell$  is the image of a line in  $S$ , the cubic curve  $C = S \cap H$  is reducible. Because  $\ell$  splits in  $Y$ , it is a bitangent to the quartic curve  $\Delta$ . Conversely, if  $\ell$  splits in  $Y$ , then  $C$  will be reducible. It will be the union of a line and a conic. So every bitangent line distinct from  $\ell_1$  is the image of a unique line in  $S$ .

The line  $\ell_1 : \{a = 0\}$  is special. Its inverse image  $C$  in  $S$  is the locus of zeros of the two polynomials  $a$  and  $az^2 + bz + c$ , or equivalently, the locus  $a = 0$  and  $bz + c = 0$ .

Let's adjust coordinates so that  $a$  becomes the polynomial  $x_0$ . The locus  $\{x_0 = 0\}$  in  $\mathbb{P}^3$  is the projective plane  $P$  with coordinates  $x_1, x_2, z$ , and in  $P$   $C$  is the locus  $\bar{g} = 0$  in that plane, where  $\bar{g} = \bar{b}z + \bar{c}$ ,  $\bar{b}, \bar{c}$  being

the polynomials obtained from  $b, c$  by substituting  $x_0 = 0$ . In  $P$ , the point  $q$  becomes  $(0, 0, 1)$ , and  $C$  becomes the cubic curve  $\bar{g} = 0$ . The cubic curve  $C$  is singular at  $q$  because  $\bar{g}$  has no term of degree  $> 1$  in  $z$ . As we have noted,  $C$  cannot be the union  $Q \cup L$ , of a conic and a line that meet at  $q$ . Therefore  $C$  is irreducible. It doesn't contain a line, so  $\ell_1$  doesn't split.  $\square$

Summing up: The 27 bitangents distinct from  $\ell_1$  are images of lines in  $S$ , but  $\ell_1$  is not the image of a line in  $S$ .



## Chapter 5 STRUCTURE OF VARIETIES IN THE ZARISKI TOPOLOGY

- 5.1 Modules (a review)
- 5.2 Valuations
- 5.3 Smooth Curves
- 5.4 Constructible sets
- 5.5 Closed Sets
- 5.6 Fibred Products
- 5.7 Projective Varieties are Proper
- 5.8 Fibre Dimension

The goal of this chapter is to show how algebraic curves control the geometry of higher dimensional varieties. We do this, beginning in Section 5.5.

### 5.1 Modules (a review)

We start with a brief review of modules over a ring, omitting proofs.

#### (5.1.1) exact sequences

A sequence

$$\dots \rightarrow V^{n-1} \xrightarrow{d^{n-1}} V^n \xrightarrow{d^n} V^{n+1} \xrightarrow{d^{n+1}} \dots$$

of homomorphisms of  $R$ -modules is *exact* if the image of  $d^{k-1}$  is equal to the kernel of  $d^k$ . For example, a sequence  $0 \rightarrow V \xrightarrow{d} V'$  is exact if and only if the map  $d$  is injective, and a sequence  $V \xrightarrow{d} V' \rightarrow 0$  is exact if and only if  $d$  is surjective.

Any homomorphism  $V \xrightarrow{d} V'$  can be embedded into an exact sequence

$$0 \rightarrow K \rightarrow V \xrightarrow{d} V' \rightarrow C \rightarrow 0,$$

where  $K$  and  $C$  are the kernel and cokernel of  $d$ , respectively.

A *short exact sequence* is an exact sequence of the form

$$0 \rightarrow V \xrightarrow{a} V' \xrightarrow{b} V'' \rightarrow 0.$$

The statement that this sequence is exact asserts that the map  $a$  is injective, and that  $V''$  is isomorphic to the quotient group  $V'/aV$ .

**5.1.2. Proposition.** (*functorial property of the kernel and cokernel*) Suppose given a (commutative) diagram of  $R$ -modules

$$\begin{array}{ccccccc} V & \xrightarrow{u} & V' & \longrightarrow & V'' & \longrightarrow & 0 \\ f \downarrow & & f' \downarrow & & f'' \downarrow & & \\ 0 \longrightarrow & W & \longrightarrow & W' & \xrightarrow{v} & W'' & \end{array}$$

whose rows are exact sequences. Let  $K, K', K''$  and  $C, C', C''$  denote the kernels and cokernels of  $f, f'$ , and  $f''$ , respectively.

(i) **(kernel is left exact)** The kernels form an exact sequence  $K \rightarrow K' \rightarrow K''$ . If  $u$  is injective, the sequence  $0 \rightarrow K \rightarrow K' \rightarrow K''$  is exact.

(ii) **(cokernel is right exact)** The cokernels form an exact sequence  $C \rightarrow C' \rightarrow C''$ . If  $v$  is surjective, the sequence  $C \rightarrow C' \rightarrow C'' \rightarrow 0$  is exact.

(iii) **(Snake Lemma)** There is a canonical homomorphism  $K'' \xrightarrow{d} C$  that combines with the above sequences to form an exact sequence

$$K \rightarrow K' \rightarrow K'' \xrightarrow{d} C \rightarrow C' \rightarrow C''.$$

If  $u$  is injective and/or  $v$  is surjective, the sequence remains exact with zeros at the appropriate ends.  $\square$

### (5.1.3) tensor products

Let  $U$  and  $V$  be modules over a ring  $R$ . The *tensor product*  $U \otimes_R V$  is an  $R$ -module that is generated by elements  $u \otimes v$  called tensors, one for each  $u$  in  $U$  and  $v$  in  $V$ . Its elements are combinations of tensors with coefficients in  $R$ .

The defining relations among the tensors are the *bilinear relations*:

$$(5.1.4) \quad (u_1 + u_2) \otimes v = u_1 \otimes v + u_2 \otimes v, \quad u \otimes (v_1 + v_2) = u \otimes v_1 + u \otimes v_2$$

$$\text{and} \quad r(u \otimes v) = (ru) \otimes v = u \otimes (rv)$$

for all  $u$  in  $U$ ,  $v$  in  $V$ , and  $r$  in  $R$ . The tensor symbol  $\otimes$  is used as a reminder that the elements  $u \otimes v$  are to be manipulated using these relations.

One can absorb a coefficient from  $R$  into either one of the factors of a tensor, so every element of  $U \otimes_R V$  can be written as a finite sum  $\sum u_i \otimes v_i$  with  $u_i$  in  $U$  and  $v_i$  in  $V$ .

**5.1.5. Example.** Let  $U$  be the space of  $m$  dimensional (complex) column vectors, and let  $V$  be the space of  $n$ -dimensional row vectors. Then  $U \otimes_{\mathbb{C}} V$  identifies naturally with the space of  $m \times n$ -matrices. If  $U$  and  $V$  are free  $R$ -modules with bases  $\{u_i\}$  and  $\{v_j\}$ , respectively, then  $U \otimes_R V$  is a free  $R$ -module with basis  $\{u_i \otimes v_j\}$ .

There is an obvious map of sets  $U \times V \xrightarrow{\beta} U \otimes_R V$  from the product set to the tensor product, that sends  $(u, v)$  to  $u \otimes v$ . This map isn't a module homomorphism. The defining relations (5.1.4) show that it is  $R$ -bilinear, not linear. It is a universal bilinear map.

**5.1.6. Corollary.** Let  $U, V$ , and  $W$  be  $R$ -modules. Homomorphisms of  $R$ -modules  $U \otimes_R V \rightarrow W$  correspond bijectively to  $R$ -bilinear maps  $U \times V \rightarrow W$ .  $\square$

Any  $R$ -bilinear map  $U \times V \xrightarrow{f} W$  to a module  $W$  can be obtained from a module homomorphism  $U \otimes_R V \xrightarrow{\tilde{f}} W$  by composition with the bilinear map  $\beta$  defined above:  $U \times V \xrightarrow{\beta} U \otimes_R V \xrightarrow{\tilde{f}} W$ .

This follows from the defining relations.  $\square$

**5.1.7. Proposition.** There are canonical isomorphisms

- $U \otimes_R R \approx U$ , defined by  $u \otimes r \rightsquigarrow ur$
- $(U \oplus U') \otimes_R V \approx (U \otimes_R V) \oplus (U' \otimes_R V)$ , defined by  $(u_1 + u_2) \otimes v \rightsquigarrow u_1 \otimes v + u_2 \otimes v$
- $U \otimes_R V \approx V \otimes_R U$ , defined by  $u \otimes v \rightsquigarrow v \otimes u$
- $(U \otimes_R V) \otimes_R W \approx U \otimes_R (V \otimes_R W)$ , defined by  $(u \otimes v) \otimes w \rightsquigarrow u \otimes (v \otimes w)$   $\square$

**5.1.8. Proposition.** Tensor product is right exact: Let  $U \xrightarrow{f} U' \xrightarrow{g} U'' \rightarrow 0$  be an exact sequence of  $R$ -modules. For any  $R$ -module  $V$ , the sequence below is exact:

$$U \otimes_R V \xrightarrow{f \otimes id} U' \otimes_R V \xrightarrow{g \otimes id} U'' \otimes_R V \rightarrow 0$$

$\square$

Tensor product isn't left exact. For example, Let  $R = \mathbb{C}[x]$ . Then  $R/xR \approx \mathbb{C}$ , so there is an exact sequence  $0 \rightarrow R \xrightarrow{x} R \rightarrow \mathbb{C} \rightarrow 0$ . When we tensor with  $\mathbb{C}$  we get the sequence  $0 \rightarrow \mathbb{C} \xrightarrow{0} \mathbb{C} \rightarrow \mathbb{C} \rightarrow 0$ , in which the zero map isn't injective.

**5.1.9. Corollary.** *Let  $U$  and  $V$  be modules over a domain  $R$  and let  $s$  be a nonzero element of  $R$ . Let  $R_s, U_s, V_s$  be the (simple) localizations of  $R, U, V$ , respectively.*

(i) *There is a canonical isomorphism  $U_s \approx U \otimes_R (R_s)$ .*

(ii) *Tensor product is compatible with localization:  $U_s \otimes_{R_s} V_s \approx (U \otimes_R V)_s$*  □

We note that the product module  $U \times V$  and the tensor product module  $U \otimes_R V$  are very different. For instance, when  $U$  and  $V$  are free modules of ranks  $r$  and  $s$ ,  $U \times V$  is free of rank  $r+s$ , while  $U \otimes_R V$  is free of rank  $rs$ .

### (5.1.10) extension of scalars in a module

Let  $R \xrightarrow{\rho} R'$  be a ring homomorphism. Extension of scalars constructs an  $R'$ -module from an  $R$ -module. Let's write scalar multiplication on the right. So  $M$  will be a right  $R$ -module. Then  $M \otimes_R R'$  becomes an  $R'$ -module, multiplication by  $s \in R'$  being  $(m \otimes a)r' = m \otimes (ar')$ . This gives the functor

$$R\text{-modules} \xrightarrow{\otimes R'} R'\text{-modules}$$

that is called *extension of scalars*.

### (5.1.11) localization, again

If  $s$  is a nonzero element of a domain  $A$ , the *simple localization*  $A_s$ , which is often referred to simply as a *localization*, is the ring obtained by adjoining an inverse of  $s$ , and to work with the inverses of finitely many nonzero elements, one may simply adjoin the inverse of their product. For working with an infinite set of inverses, the concept of a multiplicative system is useful.

A *multiplicative system*  $S$  in a domain  $A$  is a subset that consists of nonzero elements, is closed under multiplication, and contains 1. If  $S$  is a multiplicative system, the ring of  *$S$ -fractions*  $AS^{-1}$  is the ring obtained by adjoining inverses of all elements of  $S$ . Its elements are equivalence classes of fractions  $as^{-1}$  with  $a$  in  $A$  and  $s$  in  $S$ , the equivalence relation and the laws of composition being the usual ones for fractions. The ring  $AS^{-1}$  is called a *localization* too..

**5.1.12. Examples.** (i) The set consisting of the powers of a nonzero element  $s$  of a domain  $A$  is a multiplicative system. Its ring of fractions is the simple localization  $A_s = A[s^{-1}]$ .

(ii) The set  $S$  of all nonzero elements of a domain  $A$  is a multiplicative system. Its ring of fractions is the field of fractions of  $A$ .

(iii) An ideal  $P$  of a domain  $A$  is a prime ideal if and only if its complement, the set of elements of  $A$  **not** in  $P$ , is a multiplicative system. □

Let  $A \subset B$  be a ring extension, and let  $I$  and  $J$  be ideals of  $A$  and  $B$ , respectively. Recall that the *extension* of  $I$  is the ideal  $IB$  of  $B$  generated by  $I$ , whose elements are finite sums  $\sum_i z_i b_i$  with  $z_i$  in  $I$  and  $b_i$  in  $B$ . The *contraction* of  $J$  is the intersection  $J \cap A$ , which is an ideal of  $A$ .

**5.1.13. Proposition.** *Let  $S$  be a multiplicative system in a domain  $A$ , and let  $A'$  be the localization  $AS^{-1}$ .*

(i) *Let  $I$  be an ideal of  $A$ . The extended ideal  $IA'$  is the set  $IS^{-1}$  whose elements are classes of fractions  $xs^{-1}$ , with  $x$  in  $I$  and  $s$  in  $S$ . The extended ideal is the unit ideal if and only if  $I$  contains an element of  $S$ .*

(ii) *Let  $J$  be an ideal of the localization  $A'$  and let  $I$  denote its contraction  $J \cap A$ . The extended ideal  $IA'$  is equal to  $J$ :  $J = (J \cap A)A'$ .*

(iii) *If  $Q$  is a prime ideal of  $A$  and if  $Q \cap S$  is empty, the extended ideal  $Q' = QA'$  is a prime ideal of  $A'$ , and the contraction  $Q' \cap A$  is equal to  $Q$ . If  $Q \cap S$  isn't empty, the extended ideal is the unit ideal. Thus prime ideals of  $AS^{-1}$  correspond bijectively to prime ideals of  $A$  that don't meet  $S$ .* □

**5.1.14. Corollary.** *Every localization  $AS^{-1}$  of a noetherian domain  $A$  is noetherian.*  $\square$

**(5.1.15) a general principle**

An important, though elementary, principle for working with fractions is that any finite sequence of computations in a localization  $AS^{-1}$  will involve only finitely many denominators, and can therefore be done in a simple localization  $A_s$ , where  $s$  is a common denominator for the fractions that occur. This principle has been mentioned before, in Proposition 4.2.10.

For example, let  $A \subset B$  be finite-type domains, and let  $S$  be the multiplicative system of nonzero elements of  $A$ . Then  $AS^{-1} = K$  is the field of fractions of  $A$ , and  $B_K = BS^{-1}$  is a finite-type  $K$ -algebra. The Noether Normalization Theorem tells us that  $B_K$  is a finite module over a polynomial subring  $K[y_1, \dots, y_n]$ . Therefore there is a nonzero element  $s$  in  $A$  such that  $B_s$  is a finite module over the polynomial ring  $A_s[y_1, \dots, y_n]$ .

**(5.1.16) module homomorphisms**

Beginning in Chapter 6, we will work with modules over various rings. Let  $R \xrightarrow{\rho} R'$  be a ring homomorphism, let  $M$  be an  $R$ -module, and let  $M'$  be an  $R'$ -module. A *homomorphism*  $M \xrightarrow{\varphi} M'$  associated to the ring homomorphism  $\rho$  is an additive group homomorphism  $M \xrightarrow{\varphi} M'$  compatible with scalar multiplication. If  $m$  in  $M$  and  $a$  in  $R$ , then

$$(5.1.17) \quad \varphi(am) = \rho(a)\varphi(m)$$

For example, if  $M$  is a module over a domain  $R$  and  $s$  is a nonzero element of  $R$ , the localization  $M_s$  is an  $R_s$ -module. The canonical homomorphism  $M \rightarrow M_s$  is compatible with the localization map  $R \rightarrow R_s$ .

If  $A \xrightarrow{\rho} B$  is a ring homomorphism, a  $B$ -module  $N$  can be made into an  $A$ -module by *restriction of scalars*, scalar multiplication by an element  $a$  of  $A$  being defined by the formula

$$(5.1.18) \quad an = \rho(a)n$$

If it seems necessary in order to avoid confusion, we may denote a  $B$ -module  $N$  and the  $A$ -module obtained from it by restriction of scalars by  ${}_B N$  and  ${}_A N$ , respectively.

Let  $M \xrightarrow{\varphi} M'$  be a homomorphism compatible with a ring homomorphism  $R \xrightarrow{\rho} R'$ . When  $M'$  is made into an  $R$ -module by restriction of scalars,  $\varphi$  becomes a homomorphism of  $R$ -modules.

**(5.1.19) localizing a module**

Let  $S$  be a multiplicative system in a domain  $A$ . The *localization*  $MS^{-1}$  of an  $A$ -module  $M$  is defined in a natural way, as the  $AS^{-1}$ -module whose elements are equivalence classes of fractions  $ms^{-1}$  with  $m$  in  $M$  and  $s$  in  $S$ , and there will be a homomorphism  $M \rightarrow MS^{-1}$  that sends an element  $m$  to the fraction  $m/1$ . The only complication comes from the fact that  $M$  may have  $S$ -torsion elements – nonzero elements  $m$  such that  $ms = 0$  for some  $s$  in  $S$ . If  $ms = 0$  and  $s$  is in  $S$ , then  $m$  must map to zero in  $MS^{-1}$ , because in  $MS^{-1}$ , we will have  $m = mss^{-1}$ .

To define  $MS^{-1}$ , it suffices to modify the equivalence relation. Two fractions  $m_1s_1^{-1}$  and  $m_2s_2^{-1}$  are defined to be equal if there is an element  $\tilde{s} \in S$  such that  $m_1s_2\tilde{s} = m_2s_1\tilde{s}$ . Then  $ms_1^{-1} = 0$  if and only if  $m\tilde{s} = 0$  for some  $\tilde{s}$  in  $S$ . This takes care of torsion, and  $MS^{-1}$  becomes an  $AS^{-1}$ -module.

This is also how one localizes a ring that isn't a domain.

When  $S$  is the set of powers of an element  $s$ , the localized  $A_s$ -module will be denoted by  $M_s$ . Its elements have the form  $ms^{-k}$ , for some  $k \geq 0$ , and  $m_1s^{-k_1} = m_2s^{-k_2}$  if for sufficiently large  $n$ ,  $m_1s^{k_2+n} = m_2s^{k_1+n}$ .

**5.1.20. Proposition.** *Let  $S$  be a multiplicative system in a domain  $A$ .*



- (i) *Localization is an exact functor:* A homomorphism  $M \xrightarrow{\varphi} N$  of  $A$ -modules induces a homomorphism  $MS^{-1} \xrightarrow{\varphi'} NS^{-1}$  of  $AS^{-1}$ -modules, and if  $M \xrightarrow{\varphi} N \xrightarrow{\psi} P$  is an exact sequence of  $A$ -modules, the localized sequence  $MS^{-1} \xrightarrow{\varphi'} NS^{-1} \xrightarrow{\psi'} PS^{-1}$  is exact.
- (ii) Let  $M$  be an  $A$ -module. and let  $N$  be an  $AS^{-1}$ -module. Homomorphisms of  $AS^{-1}$ -modules  $MS^{-1} \rightarrow N$  correspond bijectively to homomorphisms of  $A$ -modules  $M \rightarrow N$ .
- (iii) If multiplication by  $s$  is an injective map  $M \rightarrow M$  for every  $s$  in  $S$ , then  $M \subset S^{-1}M$ . If multiplication by every  $s$  is a bijective map  $M \rightarrow M$ , then  $M \approx S^{-1}M$ .  $\square$

### (5.1.21) local rings

A *local ring* is a noetherian ring that contains just one maximal ideal.

An element of a local ring  $R$  that isn't in its maximal ideal  $M$  isn't in any maximal ideal, so it is a unit. A local ring  $R$  will have a *residue field*  $R/M$ . The case that is most important for us is that the residue field is the field of complex numbers.

We make a few general comments about local rings here though we will be interested mainly in some special local rings, discrete valuation rings that are discussed below.

The Nakayama Lemma 4.1.3 has a useful version for local rings:

**5.1.22. Local Nakayama Lemma.** Let  $R$  be a local ring with maximal ideal  $M$ , let  $V$  be a finite  $R$ -module, and let  $\bar{V}$  be the quotient  $V/MV$ , which is a vector space over the residue field  $k$  of  $R$  as well as an  $R$ -module.

- (i) If  $\bar{V} = 0$ , then  $V = 0$ .
- (ii) Let  $v = (v_1, \dots, v_r)$  be a set of elements of  $V$ , and let  $\bar{v} = (\bar{v}_1, \dots, \bar{v}_r)$  be the residues of  $v$  in  $\bar{V}$ . If  $\bar{v}$  spans  $\bar{V}$ , then  $v$  spans  $V$ .

*proof.* (i) If  $\bar{V} = 0$ , then  $V = MV$ . The Nakayama Lemma tells us that  $M$  contains an element  $z$  such that  $1 - z$  annihilates  $V$ . Then  $1 - z$  isn't in  $M$ , so it is a unit. A unit annihilates  $V$ , and therefore  $V = 0$ .

(ii) Let  $V'$  be the submodule of  $V$  spanned by  $v$ , let  $W = V/V'$ , and let  $\bar{W} = W/MW$ . To show that  $v$  spans  $V$ , we show that  $W = 0$ , and according to (i), it suffices to show that  $\bar{W} = 0$ .

We inspect the iagram

$$\begin{array}{ccccccccc} 0 & \longrightarrow & MV & \longrightarrow & V & \xrightarrow{a} & \bar{V} & \longrightarrow & 0 \\ & & \downarrow & & \downarrow b & & \downarrow c & & \\ 0 & \longrightarrow & MW & \longrightarrow & W & \xrightarrow{d} & \bar{W} & \longrightarrow & 0 \end{array}$$

Its rows are exact, the maps labelled  $a, b, c, d$  are surjective, and the kernel of  $b$  is  $V'$ . If  $\bar{v}$  spans  $\bar{V}$ , i.e., if  $aV' = \bar{V}$ , then  $caV' = \bar{W} = dbV'$ . Since  $bV' = 0$ , it follows that  $\bar{W} = 0$ , as required.  $\square$

**5.1.23. Corollary.** Let  $R$  be a local ring with maximal ideal  $M$  and residue field  $k$ , and let  $m = \{m_1, \dots, m_r\}$  be a set of elements of  $M$ . If the residues of  $m$  span the  $k$ -vector space  $M/M^2$ , then  $m$  spans  $M$ .  $\square$

## 5.2 Valuations

A local domain  $R$  with maximal ideal  $M$  has *dimension one* if  $(0)$  and  $M$  are the only prime ideals of  $R$ , and  $M \neq (0)$ . In this section, we describe the *normal* local domains of dimension one. They are the discrete valuation rings that are defined below.

Let  $K$  be a field. A *discrete valuation*  $v$  on  $K$  is a surjective homomorphism

$$(5.2.1) \quad K^\times \xrightarrow{v} \mathbb{Z}^+$$

from the multiplicative group of nonzero elements of  $K$  to the additive group of integers such that, if  $a, b$  are elements of  $K$  and if  $a, b$  and  $a+b$  aren't zero, then

- $v(a+b) \geq \min\{v(a), v(b)\}$ .

The word “discrete” refers to the fact that  $\mathbb{Z}^+$  has the discrete topology. Other valuations exist. They are interesting, but less important, and we won’t use them. To simplify terminology, we refer to a discrete valuation simply as a *valuation*.

Let  $k$  be a positive integer. If  $v$  is a valuation and if  $v(a) = k$ , then  $k$  is the *order of zero of  $a$* , and if  $v(a) = -k$ , then  $k$  is the *order of pole of  $a$* , with respect to the valuation.

**5.2.2. Lemma.** *If  $v$  is a valuation on a field  $K$  that contains the complex numbers, every nonzero complex number has value zero.*

*proof.* This is true because  $\mathbb{C}$  contains  $n$ th roots. If  $\gamma$  is an  $n$ th root of a nonzero complex number  $c$ , then because  $v$  is a homomorphism,  $v(\gamma) = v(c)/n$ . The only integer that is divisible by every positive integer  $n$  is zero.  $\square$

The *valuation ring*  $R$  associated to a valuation  $v$  on a field  $K$  is the subring of elements of  $K$  with non-negative values, together with zero:

$$(5.2.3) \quad R = \{a \in K^\times \mid v(a) \geq 0\} \cup \{0\}.$$

Valuation rings are usually called “discrete valuation rings”, but since we have dropped the word discrete from the valuation, we drop it from the valuation ring too.

**5.2.4. Proposition.** *Valuations of the field  $\mathbb{C}(t)$  of rational functions in one variable correspond bijectively to points of the projective line  $\mathbb{P}_t^1$ . The valuation ring that corresponds to a point  $p \neq \infty$  is the local ring of the polynomial ring  $\mathbb{C}[t]$  at  $p$ .*

*beginning of the proof.* Let  $K$  denote the field  $\mathbb{C}(t)$ , and let  $a$  be a complex number. To define the valuation  $v$  that corresponds to the point  $p : t = a$  of  $\mathbb{P}^1$ , we write a nonzero polynomial  $f$  as  $(t - a)^k h$ , where  $t - a$  doesn’t divide  $h$ , and we define,  $v(f) = k$ . Then we define  $v(f/g) = v(f) - v(g)$ . You will be able to check that with this definition,  $v$  becomes a valuation whose valuation ring is the local ring at  $p$ . The valuation that corresponds to the point at infinity of  $\mathbb{P}^1$  is obtained by working with  $t^{-1}$  in place of  $t$ .

The proof that these are all of the valuations of  $\mathbb{C}(t)$  will be given at the end of the section.

**5.2.5. Proposition.** *Let  $v$  be a valuation on a field  $K$ , let  $R$  be its valuation ring, and let  $x$  be an element of the multiplicative group  $K^\times$  with value  $v(x) = 1$ .*

(i) *The ring  $R$  is a normal local domain of dimension one. Its maximal ideal  $M$  is the principal ideal  $xR$ . The elements of  $M$  are those that have positive value:*

$$M = \{a \in K^\times \mid v(a) > 0\} \cup \{0\}$$

(ii) *The units of  $R$  are the elements of  $K^\times$  with value zero. Every element  $z$  of  $K^\times$  has the form  $z = x^k u$ , where  $u$  is a unit and  $k = v(z)$  is an integer.*

(iii) *The proper  $R$ -submodules of  $K$  are the sets  $x^k R$ , where  $k$  is a positive or negative integer. The set  $x^k R$  consists of zero and the elements of  $K^\times$  with value  $\geq k$ . The nonzero ideals of  $R$  are the principal ideals  $x^k R$  with  $k \geq 0$ , the powers of the maximal ideal.*

(iv) *There is no ring properly between  $R$  and  $K$ : If  $R'$  is a ring and if  $R \subset R' \subset K$ , then either  $R = R'$  or  $R' = K$ .*

*proof.* We prove (i) last.

(ii) Since  $v$  is a homomorphism,  $v(u^{-1}) = -v(u)$ . So  $u$  and  $u^{-1}$  are both in  $R$ , i.e.,  $u$  is a unit, if and only if  $v(u) = 0$ . If  $z$  is a nonzero element of  $K$  with  $v(z) = k$ , then  $u = x^{-k}z$  has value zero, so it is a unit, and  $z = ux^k$ .

(iii) The  $R$ -module  $x^k R$  consists of the elements of  $K$  of value at least  $k$ . Suppose that an  $R$ -submodule  $N$  of  $K$  contains an element  $z$  with value  $k$ . Then  $z = ux^k$ , where  $u$  is a unit, and therefore  $N$  contains  $x^k$  and

$x^k R$ . If  $k$  is the smallest integer such that  $N$  contains an element  $z$  with value  $k$ , then  $N = x^k R$ . If there is no minimum value of the elements of  $N$ , then  $N$  contains  $x^k R$  for every  $k$ , and  $N = K$ .

(iv) This follows from (iii). The ring  $R'$  will be an  $R$ -submodule of  $K$ . If  $R' \neq K$ , then  $R' = x^k R$  for some  $k$ , and since  $R'$  contains  $R$ ,  $k \leq 0$ . If  $k < 0$  then  $x^k R$  isn't closed under multiplication. So  $k = 0$  and  $R' = R$ .

(i) First,  $R$  is noetherian because (iii) tells us that it is a principal ideal domain, and it follows from (ii) that the only prime ideals of  $R$  are  $\{0\}$  and  $M = xR$ . So  $R$  is a local ring of dimension 1. If the normalization of  $R$  were larger than  $R$ , then according to (iv), it would be equal to  $K$ . Then  $x^{-1}$  would be integral over  $R$ . It would satisfy a polynomial relation  $x^{-r} + a_1 x^{-(r-1)} + \cdots + a_r = 0$  with  $a_i$  in  $R$ . When one multiplies this relation by  $x^r$ , one sees that 1 would be a multiple of  $x$ . Then  $x$  would be a unit, which it is not.  $\square$

### 5.2.6. Theorem.

(i) A local domain whose maximal ideal is a nonzero principal ideal is a valuation ring.

(ii) Every normal local domain of dimension 1 is a valuation ring.

*proof.* (i) Let  $R$  be a local domain whose maximal ideal  $M$  is a nonzero principal ideal, say  $M = xR$ , with  $x \neq 0$ , and let  $y$  be a nonzero element of  $R$ . The integers  $k$  such that  $x^k$  divides  $y$  are bounded (4.1.5). Let  $x^k$  be the largest power that divides  $y$ . Then  $y = ux^k$ , where  $k \geq 0$  and  $u$  isn't in  $M$ . It is a unit. Then any nonzero element  $z$  of the fraction field  $K$  of  $R$  will have the form  $z = ux^r$  where  $u$  is a unit and  $r$  is an integer, possibly negative. This is shown by writing the numerator and denominator of a fraction in such a form and dividing.

The valuation whose valuation ring is  $R$  is defined by  $v(z) = r$  when  $z = ux^r$  as above. If  $z_i = u_i x^{r_i}$ ,  $i = 1, 2$ , where  $u_i$  are units and  $r_1 \leq r_2$ , then  $z_1 + z_2 = \alpha x^{r_1}$ , where  $\alpha = u_1 + u_2 x^{r_2 - r_1}$  is an element of  $R$ . Therefore  $v(z_1 + z_2) \geq r_1 = \min\{v(z_1), v(z_2)\}$ . We also have  $v(z_1 z_2) = v(z_1) + v(z_2)$ . Thus  $v$  is a surjective homomorphism. The requirements for a valuation are satisfied.

(ii) The fact that a valuation ring is a normal, one-dimensional local ring is Proposition 5.2.5 (i). We show that a normal local domain  $R$  of dimension 1 is a valuation ring by showing that its maximal ideal is a principal ideal. The proof is a bit tricky.

Let  $z$  be a nonzero element of  $M$ . Because  $R$  is a local ring of dimension 1,  $M$  is the only prime ideal that contains  $z$ , so  $M$  is the radical of the principal ideal  $zR$ , and  $M^r \subset zR$  if  $r$  is large. Let  $r$  be the smallest integer such that  $M^r \subset zR$ . Then there is an element  $y$  in  $M^{r-1}$  that isn't in  $zR$ , but such that  $yM \subset zR$ . We restate this by saying that  $w = y/z$  isn't in  $R$ , but  $wM \subset R$ . Since  $M$  is an ideal, multiplication by an element of  $R$  carries  $wM$  to  $wM$ . So  $wM$  is an ideal. Since  $M$  is the maximal ideal of the local ring  $R$ , either  $wM \subset M$ , or  $wM = R$ . If  $wM \subset M$ , the lemma below shows that  $w$  is integral over  $R$ . This can't happen because  $R$  is normal and  $w$  isn't in  $R$ . Therefore  $wM = R$  and  $M = w^{-1}R$ . This implies that  $w^{-1}$  is in  $R$  and that  $M$  is a principal ideal.  $\square$

**5.2.7. Lemma.** Let  $I$  be a nonzero ideal of a noetherian domain  $A$ , and let  $B$  be a domain that contains  $A$ . An element  $w$  of  $B$  such that  $wI \subset I$  is integral over  $A$ .

*proof.* This is the Nakayama Lemma once more. Because  $A$  is noetherian,  $I$  is finitely generated. Let  $v = (v_1, \dots, v_n)^t$  be a vector whose entries generate  $I$ . The hypothesis  $wI \subset I$  allows us to write  $wv_i = \sum p_{ij}v_j$  with  $p_{ij}$  in  $A$ , or in matrix notation,  $wv = Pv$ . So  $w$  is an eigenvalue of  $P$ . If  $p(t)$  denotes the characteristic polynomial of  $P$ ,  $p(w)v = 0$ . Since  $I \neq 0$ , at least one  $v_i$  is nonzero. Since  $A$  is a domain,  $p(w)v_i = 0$  implies that  $p(w) = 0$ . The characteristic polynomial is a monic polynomial with coefficients in  $A$ , so  $w$  is integral over  $A$ .  $\square$

### (5.2.8) the local ring at a point

Let  $\mathfrak{m}$  be the maximal ideal at a point  $p$  of an affine variety  $X = \text{Spec } A$ . The complement  $S$  of  $\mathfrak{m}$  is a multiplicative system (5.1.12)(iii), and the prime ideals  $P$  of the localization  $AS^{-1}$  (the ring obtained by inverting the elements of  $S$ ) are extensions of the prime ideals  $Q$  of  $A$  that are contained in  $\mathfrak{m}$ :  $P = QS^{-1}$  (5.1.13). Thus  $AS^{-1}$  is a local ring whose maximal ideal is  $\mathfrak{m}S^{-1}$ . This ring is called the *local ring of  $A$  at  $p$* , and is often denoted by  $A_p$ .

For example, let  $X = \text{Spec } A$  be the affine line,  $A = \mathbb{C}[t]$ , and let  $p$  be the point  $t = 0$ . The local ring  $A_p$  is the ring whose elements are fractions  $f(t)/g(t)$  with  $g(0) \neq 0$ .

Any finite set  $\alpha_1, \dots, \alpha_k$  of elements of the local ring  $A_p$  at  $p$  will be contained in a simple localization  $A_s$ , for some  $s$  in  $S$ . It will be in the coordinate algebra of the affine open neighborhood  $X_s$  of  $p$ .

**5.2.9. Lemma.** *A rational function  $\alpha$  on a variety  $X$  is regular on  $X$  if it is in the local ring of  $X$  at every point  $p$ .*

This is true because a function  $\alpha$  is in the local ring at  $p$  if and only if it is in the coordinate algebra of some affine neighborhood of  $p$ .  $\square$

**5.2.10. Corollary.** *Let  $X = \text{Spec } A$  be an affine variety.*

(i) *The coordinate algebra  $A$  is the intersection of the local rings  $A_p$  at points of  $X$ .*

$$A = \bigcap_{p \in X} A_p$$

(ii) *The coordinate algebra  $A$  is normal if and only if all of its local rings  $A_p$  are normal.*

See Lemma 4.3.3 for (ii).  $\square$

**5.2.11. Note.** (about the overused word *local*) A property is true *locally* on a topological space  $X$  if every point  $p$  of  $X$  has an open neighborhood  $U$  such that the property is true on  $U$ .

The words *localize* and *localization* refer to the process of adjoining inverses. The localizations  $X_s$  of an affine variety  $X = \text{Spec } A$  form a basis for the topology on  $X$ . So if some property is true locally on  $X$ , one can cover  $X$  by localizations on which the property is true. There will be elements  $s_1, \dots, s_k$  of  $A$  that generate the unit ideal, such that the property is true on each of the localizations  $X_{s_i}$ .

An  $A$ -module  $M$  is *locally free* if there are elements  $s_1, \dots, s_k$  that generate the unit ideal of  $A$ , such that  $M_{s_i}$  is a free  $A_{s_i}$ -module for each  $i$ . If a locally free  $A$ -module  $U$  that is locally isomorphic to  $A^k$ , then  $U$  has rank  $k$ .

An ideal  $I$  of  $A$  is *locally principal* if there are elements  $s_i$  that generate the unit ideal, such that  $I_{s_i}$  is a principal ideal of  $A_{s_i}$ .  $\square$

**5.2.12. Corollary.** *Let  $M$  be a finite module over a finite-type domain  $A$ . If for some point  $p$  of  $X = \text{Spec } A$  the localized module  $M_p$  (5.1) is a free module, there is an element  $s$  not in  $\mathfrak{m}_p$  such that  $M_s$  is free.*

*proof.* See the general principle (5.1.15).  $\square$

We finish the proof of Proposition 5.2.4 now, by showing that every valuation  $v$  of the function field  $K = \mathbb{C}(t)$  of  $\mathbb{P}^1$  corresponds to a point of  $\mathbb{P}^1$ .

Let  $R$  be the valuation ring of  $v$ . If  $v(t) < 0$ , we replace  $t$  by  $t^{-1}$ . So we may assume that  $v(t) \geq 0$ . Then  $t$  is an element of  $R$ , and therefore  $\mathbb{C}[t] \subset R$ . The maximal ideal  $M$  of  $R$  isn't zero. It contains a nonzero element of  $K$ , a fraction  $\alpha = f/g$  of polynomials in  $t$ . The denominator  $g$  is in  $R$ , so  $M$  also contains the nonzero polynomial  $f = g\alpha$ . Since  $M$  is a prime ideal, it contains an irreducible factor of  $f$ . The irreducible polynomials in  $t$  are linear, so  $M$  contains  $t - a$  for some complex number  $a$ . Then  $t - c$  isn't in  $M$  when  $c \neq a$ , because the scalar  $c - a$  cannot be in  $M$ . Since  $R$  is a local ring,  $t - c$  is a unit of  $R$  for all  $c \neq a$ . The localization  $R_0$  of  $\mathbb{C}[t]$  at the point  $t = a$  is a valuation ring that is contained in the valuation ring  $R$  (5.2.4). There is no ring properly containing  $R_0$  except  $K$ , so  $R_0 = R$ .  $\square$

## 5.3 Smooth Curves

A *curve* is a variety of dimension 1. The proper closed subsets of a curve are its nonempty finite subsets.

**5.3.1. Definition.** A point  $p$  of a curve  $X$  is a *smooth point* if the local ring at  $p$  is a valuation ring. Otherwise,  $p$  is a *singular point*. A curve  $X$  is *smooth* if all of its points are smooth.

Let  $p$  be a smooth point of a curve  $X$ , and let  $v_p$  be the corresponding valuation. As with any valuation, we say that a rational function  $\alpha$  on  $X$  has a *zero of order*  $k > 0$  at  $p$  if  $v_p(\alpha) = k$ , and that it has a *pole of order*  $k$  at  $p$  if  $v_p(\alpha) = -k$ .

**5.3.2. Lemma.** (i) An affine curve  $X$  is smooth if and only if its coordinate algebra is a normal domain.

(ii) A curve has finitely many singular points.

(iii) The normalization  $\tilde{X}$  of a curve  $X$  is a smooth curve, and the canonical morphism  $\tilde{X} \rightarrow X$  becomes an isomorphism when the finite set of singular points of  $X$  and their inverse images are deleted.

*proof.* (i) This follows from Theorem 5.2.6 and Proposition 4.3.3.

(ii),(iii) Any nonempty open subset of a curve  $X$  will be the complement of a finite set, so we may replace  $X$  by an affine open subset, say  $\text{Spec } A$ . The normalization  $\tilde{A}$  of  $A$  will be a finite  $A$ -module, and therefore a finite-type algebra with the same fraction field as  $A$ , and  $\text{Spec } \tilde{A}$  will be a smooth curve. It follows from the principle 5.1.15 that  $\tilde{A}$  and  $A$  have a common localization, say  $A_s$ . The open subset  $X_s = \text{Spec } A_s$  of  $X$  will be smooth.  $\square$

**5.3.3. Proposition.** Let  $X$  be a smooth curve with function field  $K$ . Every point of  $\mathbb{P}^n$  with values in  $K$  defines a morphism  $X \rightarrow \mathbb{P}^n$ .

*proof.* A point  $(\alpha_0, \dots, \alpha_n)$  of  $\mathbb{P}^n$  with values in  $K$  determines a morphism  $X \rightarrow \mathbb{P}^n$  if and only if, for every point  $p$  of  $X$ , there is an index  $j$  such that the functions  $\alpha_i/\alpha_j$  are regular at  $p$  for all  $i = 0, \dots, n$  (3.4.11). This will be true when  $j$  is chosen so that the order of zero  $v_p(\alpha_j)$  of  $\alpha_j$  at  $p$  is the minimal integer among the  $v_p(\alpha_i)$ ,  $i = 0, \dots, n$ .  $\square$

As the next example shows, this proposition can't be extended to varieties  $X$  of dimension greater than one.

**5.3.4. Example.** Let  $Y$  be the complement of the origin in the affine plane  $X = \text{Spec } \mathbb{C}[x, y]$ , and let  $K = \mathbb{C}(x, y)$  be the function field of  $X$ . The vector  $(x, y)$  defines a point of  $\mathbb{P}_{x,y}^1$  with values in  $K$ . This point can be written as  $(1, y/x)$  and also as  $(x/y, 1)$ . So  $(x, y)$  defines a morphism to  $\mathbb{P}^1$  wherever at least one of the functions  $x/y$  or  $y/x$  is regular, which is true at all points of  $Y$ . However, there is no way to extend the morphism to  $X$ .  $\square$

**5.3.5. Proposition.** Let  $X = \text{Spec } A$  be a smooth affine curve with function field  $K$ . The local rings of  $X$  are the valuation rings of  $K$  that contain  $A$ . Therefore the maximal ideals of  $A$  are locally principal.

*proof.* Since  $A$  is a normal domain of dimension one, its local rings are valuation rings that contain  $A$  (Theorem 5.2.6). Let  $R$  be a valuation ring of  $K$  that contains  $A$ , let  $v$  be the associated valuation, and let  $M$  be the maximal ideal of  $R$ . The intersection  $M \cap A$  is a prime ideal of  $A$ . Since  $A$  has dimension 1, the zero ideal is the only prime ideal of  $A$  that isn't a maximal ideal. We can clear the denominator of an element of  $M$ , multiplying by an element of  $R$ , to obtain an element of  $A$  while staying in  $M$ . So  $M \cap A$  isn't the zero ideal. It is the maximal ideal  $\mathfrak{m}_p$  of  $A$  at a point  $p$  of  $X$ . The elements of  $A$  that aren't in  $\mathfrak{m}_p$  aren't in  $M$  either, so they are invertible in  $R$ . Therefore the local ring  $A_p$ , at  $p$ , a valuation ring, is contained in  $R$ . This implies that  $A_p = R$  (5.2.5) (iii).  $\square$

**5.3.6. Proposition.** Let  $X'$  and  $X$  be smooth curves with the same function field  $K$ .

(i) Any morphism  $X' \xrightarrow{f} X$  that is the identity on the function field  $K$  maps  $X'$  isomorphically to an open subvariety of  $X$ .

(ii) If  $X$  is projective, every smooth curve  $X'$  with function field  $K$  is isomorphic to an open subvariety of  $X$ .

(iii) If  $X'$  and  $X$  are both projective, they are isomorphic.

(iv) If  $X$  is projective, every valuation ring of  $K$  is the local ring at a point of  $X$ .

*proof.* (i) Let  $q$  be a point of  $X'$ , let  $U$  be an affine open neighborhood of  $p = fq$ , and let  $V$  be an affine open neighborhood of  $q$  in  $X'$  that is contained in the inverse image of  $U$ . Say  $U = \text{Spec } A$  and  $V = \text{Spec } B$ . The morphism  $f$  gives us a homomorphism  $A \rightarrow B$ , and since  $q$  maps to  $p$ , this homomorphism extends to an inclusion of local rings  $A_p \subset B_q$ . These rings are valuation rings with the same field of fractions, so they are equal. Since  $B$  is a finite-type algebra, there is an element  $s$  in  $A$ , with  $s(q) \neq 0$ , such that  $A_s = B_s$ . Then the open subsets  $\text{Spec } A_s$  of  $X$  and  $\text{Spec } B_s$  of  $X'$  are the same. Since  $q$  is an arbitrary point of  $X'$ ,  $X'$  is covered by open subvarieties of  $X$ . So it is an open subvariety of  $X$  too.

(ii) The projective embedding  $X \subset \mathbb{P}^n$  is defined by a point  $(\alpha_0, \dots, \alpha_n)$  with values in  $K$ , and that same point defines a morphism  $X' \rightarrow \mathbb{P}^n$ . If  $f(x_0, \dots, x_n) = 0$  is a set of defining equations of  $X$  in  $\mathbb{P}^n$ , then  $f(\alpha) = 0$

in  $K$ , and therefore  $f$  vanishes on  $X'$  too (??). So the image of  $X'$  is contained in the zero locus of  $f$ , which is  $X$ . Then (i) shows that  $X'$  is an open subvariety of  $X$ .

(iii) This follows from (ii).

(iv) The local rings of  $X$  are normal and of dimension one. They are valuation rings. We prove the converse. Let  $\beta = (\beta_0, \dots, \beta_n)$  be the point with values in  $K$  that defines the projective embedding of  $X$ , let  $R$  be a valuation ring of  $K$ , and let  $v$  be the corresponding valuation. We order the coordinates so that  $v(\beta_0)$  is minimal. Then the ratios  $\gamma_j = \beta_j/\beta_0$  will be in  $R$ . The coordinate algebra  $A_0$  of the affine variety  $X^0 = X \cap \mathbb{U}^0$  is generated by the coordinate functions  $\gamma_j$ . So  $A_0 \subset R$ , and  $R$  is the local ring of  $X^0$  at some point 5.3.5.  $\square$

**5.3.7. Proposition.** *Let  $X = \text{Spec } A$  be an affine curve, and let  $\mathfrak{m}$  and  $v$  be the maximal ideal and valuation, respectively, at a smooth point  $p$ . Let  $R$  be the valuation ring of  $v$  and let  $M$  be its maximal ideal.*

(i) *The power  $\mathfrak{m}^k$  of  $\mathfrak{m}$  consists of the elements of  $A$  whose values are at least  $k$ . If  $I$  is an ideal of  $A$  whose radical is  $\mathfrak{m}$ , then  $I = \mathfrak{m}^k$  for some  $k > 0$ .*

(ii) *The algebras  $A/\mathfrak{m}^{n+1}$  and  $R/M^{n+1}$  are isomorphic to the truncated polynomial ring  $\mathbb{C}[t]/(t^{n+1})$ .*

(iii) *If  $X$  is a smooth affine curve, every nonzero ideal  $I$  of  $A$  is a product  $\mathfrak{m}_1^{e_1} \cdots \mathfrak{m}_k^{e_k}$  of powers of maximal ideals.*

*proof.* (i) The nonzero ideals of  $R$  are powers of  $M$ . Let  $I$  be an ideal of  $A$  whose radical is  $\mathfrak{m}$ , and let  $k$  be the minimal value  $v(x)$  of the nonzero elements  $x$  of  $I$ . We will show that  $I$  is the set of all elements of  $A$  with value  $\geq k$ , i.e., that  $I = M^k \cap A$ . Since we can apply the same reasoning to  $\mathfrak{m}^k$ , it will follow that  $I = \mathfrak{m}^k$ .

Let  $x$  be an element of  $I$  with value  $k$ , and let  $y$  be an element with value at least  $k$ . Then  $x$  divides  $y$  in  $R$ , say  $y/x = u$ , with  $u$  in  $R$ . The element  $u$  will be a fraction  $a/s$  with  $s, a$  in  $A$  and  $s$  not in  $\mathfrak{m}$ , and  $sy = ax$ . The element  $s$  will vanish at a finite set of points  $q_1, \dots, q_r$ , but not at  $p$ . We choose an element  $z$  of  $A$  that vanishes at  $p$  but not at any of the points  $q_1, \dots, q_r$ . Then  $z$  is in  $\mathfrak{m}$ , and since the radical of  $I$  is  $\mathfrak{m}$ , some power of  $z$  is in  $I$ . We replace  $z$  by that power. Then  $z$  is in  $I$ . By our choice,  $z$  and  $s$  have no common zeros in  $X$ . They generate the unit ideal of  $A$ . We write  $1 = cs + dz$  with  $c$  and  $d$  in  $A$ . Then  $y = csy + dzy = cax + dzy$ . Since  $x$  and  $z$  are in  $I$ , so is  $y$ .

(ii) Since  $p$  is a smooth point, the local ring of  $A$  at  $p$  is the valuation ring  $R$ , and  $A$  contains an element  $t$  with value  $v(t) = 1$ . Let  $P$  be the subring  $\mathbb{C}[t]$  of  $A$ , and let  $\bar{P}_k = P/(t)^k$ ,  $\bar{A}_k = A/\mathfrak{m}^k$ , and  $\bar{R}_k = R/M^k$ . Since  $\mathfrak{m}$  isn't the zero ideal,  $\mathfrak{m}^{k-1} < \mathfrak{m}^k$  (Corollary 4.1.5(ii)). It follows from (i) that  $t\mathfrak{m}^{k-1} = \mathfrak{m}^k$ . Therefore  $\mathfrak{m}^{k-1}/\mathfrak{m}^k$  has  $\mathbb{C}$ -dimension 1. The map labelled  $g_{k-1}$  in the diagram below is bijective.

$$\begin{array}{ccccccc} 0 & \longrightarrow & (t^{k-1})/(t^k) & \longrightarrow & \bar{P}_k & \longrightarrow & \bar{P}_{k-1} \longrightarrow 0 \\ & & \downarrow g_{k-1} & & \downarrow f_k & & \downarrow f_{k-1} \\ 0 & \longrightarrow & \mathfrak{m}^{k-1}/\mathfrak{m}^k & \longrightarrow & \bar{A}_k & \longrightarrow & \bar{A}_{k-1} \longrightarrow 0 \end{array}$$

Induction on  $k$  shows that the map labelled  $f_{k-1}$  is bijective, so  $f_k$  is bijective. A similar argument shows that  $\bar{P}_k$  and  $\bar{R}_k$  are isomorphic.

(iii) Let  $I$  be a nonzero ideal of  $A$ . Because  $X$  has dimension one, the locus of zeros of  $I$  is a finite set  $\{p_1, \dots, p_k\}$ . Therefore the radical of  $I$  is the intersection  $\mathfrak{m}_1 \cap \cdots \cap \mathfrak{m}_k$  of the maximal ideals  $\mathfrak{m}_j$  at  $p_j$ , which, by the Chinese Remainder Theorem, is the product ideal  $\mathfrak{m}_1 \cdots \mathfrak{m}_k$ . Moreover,  $I$  contains a power of that product, say  $I \supset \mathfrak{m}_1^N \cdots \mathfrak{m}_k^N$ . Let  $J = \mathfrak{m}_1^N \cdots \mathfrak{m}_k^N$ . The quotient algebra  $A/J$  is the product  $B_1 \times \cdots \times B_k$ , with  $B_j = A/\mathfrak{m}_j^N$ , and  $A/I$  is a quotient of  $A/J$ . Proposition 2.1.7 tells us that  $A/I$  is a product  $\bar{A}_1 \times \cdots \times \bar{A}_k$ , where  $\bar{A}_j$  is a quotient of  $B_j$ . By part (ii), each  $B_j$  is a truncated polynomial ring. Then the quotients  $\bar{A}_j$  must also be truncated polynomial rings. So the kernel  $I$  of the map  $A \rightarrow \bar{A}_1 \times \cdots \times \bar{A}_k$  is a product of powers of the maximal ideals  $\mathfrak{m}_j$ .  $\square$

### (5.3.8) isolated points

**5.3.9. Proposition.** *A curve, smooth or not, contains no point that is isolated in the classical topology.*

This was proved before for plane curves (Proposition 1.3.18).

### 5.3.10. Lemma.

(i) Let  $Y'$  be an open subvariety of a variety  $Y$ . Then  $q$  is an isolated point of  $Y$  if and only if it is an isolated point of  $Y'$ .

(ii) Let  $Y' \xrightarrow{u'} Y$  be a nonconstant morphism of curves and let  $q'$  be a point of  $Y'$ . If the image of  $q'$  is an isolated point of  $Y$ , then  $q'$  is an isolated point of  $Y'$ .

*proof.* (i) A point  $q$  of  $Y$  is isolated if  $\{q\}$  is an open subset of  $Y$ . If  $\{q\}$  is open in  $Y'$  and  $Y'$  is open in  $Y$ , then  $\{q\}$  is open in  $Y$ . If  $\{q\}$  is open in  $Y$ , it is open in  $Y'$ .

(ii) Because  $Y'$  has dimension one, the fibre over  $q$  will be a finite set, say  $\{q'\} \cup F$ , where  $F$  is finite. Let  $Y''$  denote the (open) complement  $Y' - F$  of  $F$  in  $Y'$ , and let  $u''$  be the restriction of  $u'$  to  $Y''$ . The fibre of  $Y''$  over  $q$  is the point  $q'$ . If  $\{q\}$  is open in  $Y$ , then because  $u''$  is continuous,  $\{q'\}$  will be open in  $Y''$ , and therefore open in  $Y'$ .  $\square$

*proof of Proposition 5.3.9.* Let  $q$  be a point of a curve  $Y$ . Part (i) of Lemma 5.3.10 allows us to replace  $Y$  by an affine neighborhood of  $q$ . Let  $Y'$  be the normalization of  $Y$ . Part (ii) of the lemma allows us to replace  $Y$  by  $Y'$ . So we may assume that  $Y$  is a smooth affine curve, say  $Y = \text{Spec } B$ . We can still replace  $Y$  by an open neighborhood of  $q$ , so we may assume that the maximal ideal  $\mathfrak{m}_q$  is a principal ideal.

Say that  $B = \mathbb{C}[x_1, \dots, x_n]/(f_1, \dots, f_k)$ ,  $q$  is the origin  $(0, \dots, 0)$  in  $\mathbb{A}_x^n$ , and that the maximal ideal  $\mathfrak{m}_q$  is generated by the residue of a polynomial  $f_0$  in  $B$ . Then  $f_0, \dots, f_k$  generate the maximal ideal  $(x_1, \dots, x_n)$  in  $\mathbb{C}[x_1, \dots, x_n]$ . Let's write  $f_i = \sum_1^n c_{ij}x_j + O(2)$ , where  $O(2)$  denotes an undetermined polynomial, all of whose terms have degree  $\geq 2$  in  $x$ . The coefficient  $c_{ij}$  is the partial derivative  $\frac{\partial f_i}{\partial x_j}$ , evaluated at  $q$ . If  $J$  denotes  $(k+1) \times n$  Jacobian matrix  $(\frac{\partial f_i}{\partial x_j})$  at  $q$ , we have  $(f_0, \dots, f_k)^t = J(x_1, \dots, x_n)^t + O(2)$ . Since  $f_0, \dots, f_k$  generate the maximal ideal, there is a matrix  $P$  with polynomial entries such that  $Pf^t = x^t$ . Then  $x^t = PJx^t + O(2)$ . If  $P_0$  is the constant term of  $P$ ,  $P_0J$  will be the identity matrix. So  $J$  has rank  $n$ .

Let  $J^1$  be the matrix obtained by deleting the column with index 0 from  $J$ . This matrix has rank at least  $n-1$ , and we may assume that the submatrix with indices  $1 \leq i, j \leq n-1$  is invertible. The Implicit Function Theorem says that the equations  $f_1, \dots, f_{n-1}$  can be solved for the variables  $x_1, \dots, x_{n-1}$  as analytic functions of  $x_n$ , for small  $x_n$ . The locus  $Z$  of zeros of  $f_1, \dots, f_{n-1}$  has dimension at most 1, it is locally homeomorphic to the affine line (1.4.19), and it contains  $Y$ . Since  $Y$  has dimension 1, the component of  $Z$  that contains  $q$  must be equal to  $Y$ . So  $Y$  is locally homeomorphic to  $\mathbb{A}^1$ , which has no isolated point. Therefore  $q$  isn't an isolated point of  $Y$ .

## 5.4 Constructible Sets

In this section,  $X$  will denote a noetherian topological space. Every closed subset of  $X$  is a finite union irreducible closed sets (2.2.13).

The intersection  $L = C \cap U$  of a closed set  $C$  and an open set  $U$  is a *locally closed* set. Open sets and closed sets are examples of locally closed sets.

A *constructible set* is a set that is the union of finitely many locally closed sets.

**5.4.1. Lemma.** *The following conditions on a subset  $L$  of  $A$  are equivalent.*

- $L$  is locally closed.
- $L$  is a closed subset of an open subset  $U$  of  $X$ .
- $L$  is an open subset of a closed subset  $C$  of  $X$ .

$\square$

### 5.4.2. Examples.

(i) A subset  $S$  of a curve  $X$  is constructible if and only if it is either a finite set or the complement of a finite set. Thus  $S$  is constructible if and only if it is either closed or open.

(ii) Let  $C$  be the line  $\{y = 0\}$  in the affine plane  $X = \text{Spec } \mathbb{C}[x, y]$ , let  $U = X - C$  be its open complement, and let  $p = (0, 0)$ . The union  $U \cup \{p\}$  is constructible, but not locally closed.  $\square$

We will use the following notation:  $L$  is a locally closed set,  $C$  is a closed set, and  $U$  is an open set.

**5.4.3. Theorem.** *The family of constructible subsets of a noetherian topological space  $X$ , which we denote by  $\mathbb{S}$ , is the smallest family of subsets that contains the open sets and is closed under the three operations of finite unions, finite intersections, and complementation.*

*proof.* Let  $\mathbb{S}_1$  denote the family of subsets obtained from the open sets by the three operations mentioned in the statement. Open sets are constructible, and with those three operations, one can produce any constructible set from the open sets. So  $\mathbb{S} \subset \mathbb{S}_1$ . To show that  $\mathbb{S} = \mathbb{S}_1$ , we show that the family of constructible sets is closed under the three operations.

It is obvious that a finite union of constructible sets is constructible. The intersection of two locally closed sets  $L_1 = C_1 \cap U_1$  and  $L_2 = C_2 \cap U_2$  is locally closed because  $L_1 \cap L_2 = (C_1 \cap C_2) \cap (U_1 \cap U_2)$ . If  $S = L_1 \cup \dots \cup L_k$  and  $S' = L'_1 \cup \dots \cup L'_r$  are constructible sets, the intersection  $S \cap S'$  is the union of the locally closed intersections  $(L_i \cap L'_j)$ , so it is constructible.

Let  $S$  be the constructible set  $L_1 \cup \dots \cup L_k$ . Its complement is the intersection of the complements of  $L_i$ :  $S^c = L_1^c \cap \dots \cap L_k^c$ . We have shown that intersections of constructible sets are constructible. So to show that the complement  $S^c$  is constructible, it suffices to show that the complement of a locally closed set is constructible. Let  $L$  be the locally closed set  $C \cap U$ , and let  $C^c$  and  $U^c$  be the complements of  $C$  and  $U$ , respectively. Then  $C^c$  is open and  $U^c$  is closed. The complement  $L^c$  of  $L$  is the union  $C^c \cup U^c$  of constructible sets, so it is constructible.  $\square$

**5.4.4. Proposition.** *Let  $X$  be a noetherian topological space. Every constructible subset  $S$  is a union  $L_1 \cup \dots \cup L_k$  of locally closed sets  $L_i = C_i \cap U_i$ , in which the closed sets  $C_i$  are irreducible and distinct.*

*proof.* Suppose that  $L = C \cap U$  is a locally closed set, and let  $C = C_1 \cup \dots \cup C_r$  be the decomposition of  $C$  into irreducible components. Then  $L = (C_1 \cap U) \cup \dots \cup (C_r \cap U)$ , which is constructible. So every constructible set  $S$  is a union of locally closed sets  $L_i = C_i \cap U_i$  in which the  $C_i$  are irreducible. Next, suppose that two of the irreducible closed sets are equal, say  $C_1 = C_2$ . Then  $L_1 \cup L_2 = (C_1 \cap U_1) \cup (C_1 \cap U_2) = C_1 \cap (U_1 \cup U_2)$  is locally closed. So we can find an expression in which the closed sets are distinct as well.  $\square$

**5.4.5. Lemma.**

(i) *Let  $X_1$  be a closed subset of a variety  $X$ , and let  $X_2$  be its open complement. A subset  $S$  of  $X$  is constructible if and only if  $S \cap X_1$  and  $S \cap X_2$  are constructible.*

(ii) *Let  $X'$  be an open or a closed subvariety of a variety  $X$ .*

a) *If  $S$  is a constructible subset of  $X$ , then  $S' = S \cap X'$  is a constructible subset of  $X'$ .*

b) *If  $S'$  is a constructible subset of  $X'$ , then it is a constructible subset of  $X$ .*

*proof.* (i) This follows from Theorem 5.4.3.

(iia) It suffices to prove that the intersection  $L' = L \cap X'$  of a locally closed subset  $L$  of  $X$  is a locally closed subset of  $X'$ . If  $L = C \cap U$ , then  $C' = C \cap X'$  is closed in  $X'$ , and  $U' = U \cap X'$  is open in  $X'$ . So  $L' = C' \cap U'$  is locally closed.

(iib) It suffices to show that a locally closed subset  $L' = C' \cap U'$  of  $X'$  is locally closed in  $X$ . If  $X'$  is closed in  $X$ , then  $C'$  is closed in  $X$ , and  $U' = X \cap U$  for some open subset  $U$  of  $X$ . If  $X'$  is open in  $X$ , then  $U'$  is open in  $X$ , and if  $C$  is the closure of  $C'$  in  $X$ , then  $C \cap U' = C' \cap U'$ . So  $L' = C \cap U'$  is locally closed in  $X$ .  $\square$

The next theorem illustrates a general fact, that sets arising in algebraic geometry tend to be constructible.

**5.4.6. Theorem.** *Let  $Y \xrightarrow{f} X$  be a morphism of varieties. The inverse image of a constructible subset of  $X$  is a constructible subset of  $Y$ . The image of a constructible subset of  $Y$  is a constructible subset of  $X$ .*

*proof.* The fact that a morphism is continuous implies that the inverse image of a constructible set is constructible. To prove that the image of a constructible set is constructible, one keeps reducing the problem until there is nothing left to do.

Let  $S$  be a constructible subset of  $Y$ . Noetherian induction allows us to assume that the theorem is true when  $S$  is contained in a proper closed subvariety of  $Y$ , and also when its image  $f(S)$  is contained in a proper closed subvariety of  $X$ .



Suppose that  $Y$  is the union of a proper closed subvariety  $Y_1$  and its open complement  $Y_2$ , and let  $S_i = S \cap Y_i$ . It suffices to show that  $S_i$  is a constructible subset of  $Y_i$ ,  $i = 1, 2$ , and induction applies to  $Y_1$ . So we may replace  $Y$  by any nonempty open subvariety.

Let  $X_1$  be a proper closed subvariety of  $X$  and let  $X_2$  be its open complement. The inverse image  $Y_1 = f^{-1}(X_1)$  will be closed in  $Y$ , and its open complement will be the inverse image  $Y_2 = f^{-1}(X_2)$ . A constructible subset  $S$  of  $Y$  is the union of the constructible sets  $S_1 = S \cap Y_1$  and  $S_2 = S \cap Y_2$ . So it suffices to show that  $f(S_i)$  is constructible. To show this, it suffices to show that  $f(S_i)$  is a constructible subset of  $X_i$  for  $i = 1, 2$  (5.4.5) (iib). Moreover, induction applies to  $X_1$ . So we need only show that  $f(S_2)$  is a constructible subset of  $X_2$ . This means that we can replace  $X$  and  $Y$  by nonempty open subsets finitely many times.

Since  $S$  is a finite union of locally closed sets, it suffices to treat the case that  $S$  is locally closed. Moreover, we may suppose that  $S = C \cap U$ , where  $C$  is irreducible. Then  $Y$  is the union of the closed subset  $C = Y_1$  and its complement  $Y_2$ . Since  $S \cap Y_2 = \emptyset$ , it suffices to treat  $Y_1$ . We may replace  $Y$  by  $C$ . So we may assume that  $S = Y \cap U = U$ , and we may replace  $Y$  by  $U$ . We are thus reduced to the case that  $S = Y$ .

At this point, we may still replace  $X$  and  $Y$  by nonempty open subsets, so we may assume that they are affine, say  $Y = \text{Spec } B$  and  $X = \text{Spec } A$ . Then the morphism  $Y \rightarrow X$  corresponds to an algebra homomorphism  $A \xrightarrow{\varphi} B$ . If the kernel of  $\varphi$  were nonzero, the image of  $Y$  would be contained in a proper closed subset of  $X$  to which induction would apply. So we may assume that  $\varphi$  is injective.

Proposition 4.2.10 tells us that, for suitable nonzero  $s$  in  $A$ ,  $B_s$  will be a finite module over a polynomial subring  $A_s[y_1, \dots, y_k]$ . Then the maps  $Y_s \rightarrow \text{Spec } A_s[y]$  and  $\text{Spec } A_s[y] \rightarrow X_s$  are both surjective, so  $Y_s$  maps surjectively to  $X_s$ . When we replace  $X$  and  $Y$  by  $X_s$  and  $Y_s$ , the map  $Y \rightarrow X$  becomes surjective, and we are done.  $\square$

## 5.5 Closed Sets

Limits of sequences are often used to analyze subsets of a topological space. In the classical topology, a subset  $Y$  of  $\mathbb{C}^n$  is closed if, whenever a sequence of points in  $Y$  has a limit in  $\mathbb{C}^n$ , the limit is in  $Y$ . In algebraic geometry one uses curves as substitutes.

We use the following notation:

(5.5.1)  $C$  is a smooth affine curve,  $q$  is a point of  $C$ , and  $C'$  is the complement of  $q$  in  $C$ .

The closure of  $C'$  will be  $C$ , and we think of  $q$  as a limit point. Theorem 5.5.3, which is below, asserts that a constructible subset of a variety is closed if it contains all such limit points.

The next theorem tells us that there are enough curves to do the job.

**5.5.2. Theorem. (enough curves)** *Let  $Y$  be a constructible subset of a variety  $X$ , and let  $p$  be a point of its closure  $\overline{Y}$ . There exists a morphism  $C \xrightarrow{f} X$  from a smooth affine curve to  $X$ , and a point  $q$  of  $C$  with  $f(q) = p$ , such that the image of  $C' = C - \{q\}$  is contained in  $Y$ .*

*proof.* We use Krull's Theorem to slice  $Y$  down to dimension 1. If  $X = p$ , then  $Y = p$  too. In this case, we may take for  $f$  the constant morphism from any curve  $C$  to  $p$ . So we may assume that  $X$  has dimension at least one. Next, we may replace  $X$  by an affine open subset  $X'$  that contains  $p$ , and  $Y$  by  $Y' = Y \cap X'$ . The closure  $\overline{Y'}$  of  $Y'$  in  $X'$  will be the intersection  $\overline{Y} \cap X'$ , and it will contain  $p$ . So we may assume that  $X$  is affine, say  $X = \text{Spec } A$ .

Since  $Y$  is constructible, it is a union  $L_1 \cup \dots \cup L_k$  of locally closed sets, say  $L_i = Z_i \cap U_i$  where  $Z_i$  are irreducible closed sets and  $U_i$  are open sets. (We use  $Z_i$  in place of  $C_i$  here to avoid confusion with a curve.) The closure of  $Y$  is the union  $Z_1 \cup \dots \cup Z_k$ , and  $p$  is in one of the closed sets  $Z_i$ . We may replace  $X$  by  $Z_i$  and  $Y$  by  $L_i$ , so we may assume that  $Y$  is a nonempty open subset of  $X$ .

Suppose that the dimension  $n$  of  $X$  is at least two. Let  $D = X - Y$  be the (closed) complement of the open set  $Y$ . The components of  $D$  have dimension at most  $n - 1$ . We choose an element  $\alpha$  of the coordinate algebra  $A$  of  $X$  that is zero at  $p$  and isn't identically zero on any component of  $D$  except  $p$  itself, if  $p$  happens to be a component. Krull's Theorem tells us that every component of the zero locus of  $\alpha$  has dimension  $n - 1$ ,

and at least one of those components, call it  $V$ , contains  $p$ . If  $V$  were contained in  $D$ , it would be a component of  $D$  because  $\dim V = n - 1$  and  $\dim D \leq n - 1$ . By our choice of  $\alpha$ , this isn't the case. So  $V \not\subset D$ , and therefore  $V \cap Y \neq \emptyset$ . Because  $V$  is irreducible and  $Y$  is open,  $V \cap Y$  is an open dense subset of  $V$ , and  $p$  is a point of its closure  $V$ . We replace  $X$  by  $V$  and  $Y$  by  $V \cap Y$ . The dimension of  $X$  is thereby reduced to  $n - 1$ .

Thus it suffices to treat the case that  $X$  has dimension one. Then  $X$  will be a curve that contains  $p$  and  $Y$  will be a nonempty open subset of  $X$ . The normalization of  $X$  will be a smooth curve  $x_1$  that comes with an integral and therefore surjective morphism to  $Y$ . Finitely many points of  $X_1$  will map to  $p$ . We choose for  $C$  an affine open subvariety of  $X_1$  that contains just one of those points, and we call that point  $q$ .  $\square$

**5.5.3. Theorem** (curve criterion for a closed set) *Let  $Y$  be a constructible subset of a variety  $X$ . The following conditions are equivalent:*

(a)  $Y$  is closed.

(b) For every morphism  $C \xrightarrow{f} X$  from a smooth affine curve to  $X$ , the inverse image  $f^{-1}Y$  is closed in  $C$ .

(c) Let  $q$  be a point of a smooth affine curve  $C$ , let  $C' = C - \{q\}$ , and let  $C \xrightarrow{f} X$  be a morphism. If  $f(C') \subset Y$ , then  $f(C) \subset Y$ .

The hypothesis that  $Y$  be constructible is necessary. For example, let  $X$  be the affine line  $\mathbb{A}^1$ . The set  $Z$  of points of  $X$  with integer coordinates isn't constructible, but it satisfies the curve criterion. Any morphism  $C' \rightarrow X$  whose image is in  $Z$  will map  $C'$  to a single point, and therefore it will extend to  $C$ .

*proof.* The implications (a)  $\Rightarrow$  (b)  $\Rightarrow$  (c) are obvious. We prove the contrapositive of the implication (c)  $\Rightarrow$  (a). Suppose that  $Y$  isn't closed. We choose a point  $p$  of the closure  $\overline{Y}$  that isn't in  $Y$ , and we apply Theorem 5.5.2. There exists a morphism  $C \xrightarrow{f} X$  from a smooth curve to  $X$  and a point  $q$  of  $C$  such that  $f(q) = p$  and  $f(C') \subset Y$ . Since  $q \notin Y$ , this morphism shows that (c) doesn't hold either.  $\square$

**5.5.4. Theorem.** *A constructible subset  $Y$  of a variety  $X$  is closed in the Zariski topology if and only if it is closed in the classical topology.*

*proof.* A Zariski closed set is closed in the classical topology because the classical topology is finer than the Zariski topology.

Suppose that  $Y$  is closed in the classical topology. Let  $q$  be a point of the Zariski closure  $\overline{Y}$  of  $Y$ , and let  $C \xrightarrow{f} X$  be a morphism from a smooth affine curve to  $X$  that maps the complement  $C'$  of  $q$  to  $Y$ . Let  $Y' = f^{-1}Y$ . Then  $Y'$  contains  $C'$ , so it is either  $C'$  or  $C$ . A morphism is a continuous map in the classical topology. Since  $Y$  is closed in the classical topology,  $Y'$  is closed in  $C$ . If  $Y'$  were equal to  $C'$ , then  $\{q\}$  would be open as well as closed. It would be an isolated point of  $C$ . Since a curve contains no isolated point, the closure is  $C$ . Therefore the curve criterion (5.5.3) is satisfied, and  $Y$  is closed in the Zariski topology.  $\square$

## 5.6 Fibred Products

### (5.6.1) the mapping property of a product

The product  $X \times Y$  of two sets  $X$  and  $Y$  has a mapping property that is easy to verify: Maps from a set  $T$  to the product set  $X \times Y$ , correspond bijectively to pairs of maps  $T \xrightarrow{f} X$  and  $T \xrightarrow{g} Y$ . The map  $T \xrightarrow{(f,g)} X \times Y$  defined by the pair of maps  $f, g$  sends a point  $t$  to the point pair  $(f(t), g(t))$ .

Let  $X \times Y \xrightarrow{\pi_1} X$  and  $X \times Y \xrightarrow{\pi_2} Y$  denote the projection maps. If  $T \xrightarrow{h} X \times Y$  is a map to the product, the corresponding maps to  $X$  and  $Y$  are the compositions with the projections:  $T \xrightarrow{\pi_1 \circ h} X$  and  $T \xrightarrow{\pi_2 \circ h} Y$ :

The analogous statements are true for morphisms of varieties.

**5.6.2. Proposition.** *Let  $X$  and  $Y$  be varieties, and let  $X \times Y$  be the product variety.*

(i) *The projections  $X \times Y \xrightarrow{\pi_1} X$  and  $X \times Y \xrightarrow{\pi_2} Y$  are morphisms.*

(ii) *Morphisms from a variety  $T$  to the product variety  $X \times Y$  correspond bijectively to pairs of morphisms  $T \rightarrow X$  and  $T \rightarrow Y$ , the correspondence being the same as for maps of sets.*  $\square$

It was proved in Proposition 3.4.32 that if  $X \xrightarrow{f} Z$  and  $Y \xrightarrow{g} W$  are morphisms of varieties, the product map  $X \times Y \xrightarrow{f \times g} Z \times W$  defined by  $[f \times g](x, y) = (f(x), g(y))$  is a morphism.

### (5.6.3) fibred products of sets

If  $X \xrightarrow{f} Z$  and  $Y \xrightarrow{g} Z$  are maps of sets, the *fibred product*  $X \times_Z Y$  is the subset of the product  $X \times Y$  consisting of pairs of points  $x, y$  such that  $f(x) = g(y)$ . It fits into a diagram

$$(5.6.4) \quad \begin{array}{ccc} X \times_Z Y & \xrightarrow{\pi_2} & Y \\ \pi_1 \downarrow & & \downarrow g \\ X & \xrightarrow{f} & Z \end{array}$$

in which  $\pi_1$  and  $\pi_2$  are the projections. Many important subsets of a product can be described as fibred products. If a map  $Y \rightarrow Z$  is given, and if  $p \rightarrow Z$  is the inclusion of a point into  $Z$ , then  $p \times_Z Y$  is the fibre of  $Y$  over  $p$ . The diagonal in  $X \times X$  is the fibred product  $X \times_X X$ .

The reason for the term “fibred product” is that the fibre of  $X \times_Z Y$  over a point  $x$  of  $X$  maps bijectively to the fibre of  $Y$  over the image  $z = f(x)$ , and the analogous statement is true for fibres over points of  $Y$ .

### (5.6.5) fibred products of varieties

Since we are working with varieties, not schemes, we have a small problem: The fibred product of varieties will be a scheme, but it needn't be a variety.

**5.6.6. Example.** Let  $X = \text{Spec } \mathbb{C}[x]$ ,  $Y = \text{Spec } \mathbb{C}[y]$  and  $Z = \text{Spec } \mathbb{C}[z]$  be affine lines, let  $X \xrightarrow{f} Z$  and  $Y \xrightarrow{g} Z$  be the maps defined by  $z = x^2$  and  $z = y^2$ , respectively. The fibred product  $X \times_Z Y$  is the closed subset of the affine  $x, y$ -plane consisting of points  $(x, y)$  such that  $x^2 = y^2$ . It is the union of the two lines  $x = y$  and  $x = -y$ .  $\square$

The next proposition will be enough for our purposes.

**5.6.7. Proposition.** Let  $X \xrightarrow{f} Z$  and  $Y \xrightarrow{g} Z$  be morphisms of varieties. The fibred product  $X \times_Z Y$  is a closed subset of the product variety  $X \times Y$ .

*proof.* The graph  $\Gamma_f$  of a morphism  $X \xrightarrow{f} Z$  of varieties is a closed subvariety of  $X \times Z$  isomorphic to  $X$  (Proposition 3.4.27). Next, let  $u$  and  $v$  be two morphisms from a variety  $X$  to another variety:  $Z$ . We show that the set  $W$  consisting of points  $x$  in  $X$  such that  $u(x) = v(x)$  is a closed subset of  $X$ . In  $X \times Z$ , let  $W'$  be the intersection of the graphs of  $u$  and  $v$ :  $W' = \Gamma_u \cap \Gamma_v$ . A point  $(x, z)$  is in  $W'$  if  $z = ux = vx$ . This is an intersection of closed sets, so it is closed in  $\Gamma_u$  (and in  $\Gamma_v$ ). The projection  $\Gamma_u \rightarrow X$ , which is an isomorphism, carries  $W'$  to  $W$ , so  $W$  is closed in  $X$ .

With reference to Diagram 5.6.4,  $X \times_Z Y$  is the subset of the product  $X \times Y$  of points at which the maps  $f\pi_X$  and  $g\pi_Y$  to  $Z$  are equal, so it is closed in  $X \times Y$ .  $\square$

## 5.7 Projective Varieties are Proper

As has been noted (3.1), an important property of projective space with the classical topology is that it is a compact space. A variety isn't compact in the Zariski topology unless it is a single point. However, in the Zariski topology, projective varieties have a property closely related to compactness: They are *proper*.

Before defining the concept of a proper variety, we explain an analogous property of compact spaces.

**5.7.1. Proposition.** Let  $X$  be a compact space, let  $Z$  be a Hausdorff space, and let  $C$  be a closed subset of  $Z \times X$ . The image of  $C$  in  $Z$  is a closed subset of  $Z$ .

*proof.* Let  $D$  be the image of  $C$ . We show that if a sequence of points  $z_i$  of  $D$  has a limit  $\underline{z}$  in  $Z$ , then  $\underline{z}$  is in  $D$ . For each  $i$ , we choose a point  $p_i$  of  $C$  that lies over  $z_i$ . So  $p_i$  is a pair  $(z_i, x_i)$ ,  $x_i$  being a point of  $X$ . Since  $X$  is compact, there is a subsequence of the sequence  $x_i$  that has a limit  $\underline{x}$  in  $X$ . Passing to subsequences, we may suppose that  $x_i$  has limit  $\underline{x}$ . Then  $p_i$  will have the limit  $\underline{p} = (\underline{z}, \underline{x})$ . Since  $C$  is closed,  $\underline{p}$  is in  $C$ , and therefore  $\underline{z}$  is in  $D$ .  $\square$

**5.7.2. Definition.** A variety  $X$  is *proper* if has the following property: Let  $Z \times X$  be the product with another variety  $Z$ , let  $\pi$  denote the projection  $Z \times X \rightarrow Z$ , and let  $C$  be a closed subvariety of  $Z \times X$ . The image  $D = \pi(C)$  of  $C$  is a closed subvariety of  $Z$ .

$$(5.7.3) \quad \begin{array}{ccc} C & \xrightarrow{\subset} & Z \times X \\ \downarrow & & \downarrow \pi \\ D & \xrightarrow{\subset} & Z \end{array}$$

If  $X$  is proper, then because every closed set is a finite union of closed subvarieties, the image of any closed subset of  $Z \times X$  will be closed in  $Z$ .

**5.7.4. Theorem.** *Projective varieties are proper.*

This is the most important application of the use of curves to characterize closed sets.

*proof.* Let  $X$  be a projective variety. With notation as in Definition 5.7.2, suppose we are given a closed subvariety  $C$  of the product  $Z \times X$ . We must show that its image  $D$  is a closed subvariety of  $Z$ . If the image is a closed set, it will be irreducible. So it suffices to show that  $D$  is closed, and to do this, it suffices to show that  $D$  is closed in the classical topology (Theorem 5.5.4). Theorem 5.4.6 tells us that  $D$  is a constructible set, and since  $X$  is closed in projective space, it is compact in the classical topology. Proposition 5.7.1 tells us that  $D$  is closed in the classical topology.  $\square$

The next examples show how the theorem can be used.

**5.7.5. Example.** (*singular curves*) We parametrize the plane curves of a given degree  $d$ . The number of distinct monomials  $x_0^i x_1^j x_2^k$  of degree  $d = i+j+k$  is the binomial coefficient  $\binom{d+2}{2}$ . We order those monomials arbitrarily, and label them as  $m_0, \dots, m_r$ , with  $r = \binom{d+2}{2} - 1$ . A homogeneous polynomial of degree  $d$  will be a combination  $\sum z_i m_i$  of monomials with complex coefficients  $z_i$ , so the homogeneous polynomials  $f$  of degree  $d$  in  $x$ , taken up to scalar factors, are parametrized by the projective space of dimension  $r$  with coordinates  $z$ . Let's denote that projective space by  $Z$ . Points of  $Z$  correspond bijectively to divisors of degree  $d$  in the projective plane.

The product variety  $Z \times \mathbb{P}^2$  represents pairs  $(D, p)$ , where  $D$  is a divisor of degree  $d$  and  $p$  is a point of  $\mathbb{P}^2$ . A variable homogeneous polynomial of degree  $d$  in  $x$  will be a bihomogeneous polynomial  $f(z, x)$  of degree 1 in  $z$  and degree  $d$  in  $x$ . So the locus  $\Gamma: \{f(z, x) = 0\}$  in  $Z \times \mathbb{P}^2$  is a closed set. Its points are pairs  $(D, p)$  such that  $D$  is the divisor of  $f$  and  $p$  is a point of  $D$ .

Let  $\Sigma$  be the set of pairs  $(D, p)$  such that  $p$  is a singular point of  $D$ . This is also a closed set. It is defined by the system of equations  $f_0(z, x) = f_1(z, x) = f_2(z, x) = 0$ , where  $f_i$  are the partial derivatives  $\frac{\partial f}{\partial x_i}$ . Euler's Formula shows that then  $f(x, z) = 0$ . The partial derivatives  $f_i$  are bihomogeneous, of degree 1 in  $z$  and degree  $d-1$  in  $x$ .

The next proposition isn't very easy to prove directly, but the proof becomes easy when one uses the fact that projective space is proper.

**5.7.6. Proposition** *The singular divisors of degree  $d$ , the divisors containing at least one singular point, form a closed subset  $S$  of the projective space  $Z$  of all divisors of degree  $d$ .*

*proof.* The points of  $S$  are the images of points of the set  $\Sigma$  via projection to  $Z$ . Theorem 5.7.4 tells us that the image of  $\Sigma$  is closed.  $\square$

**5.7.7. Example.** (*surfaces that contain a line*) We go back to the discussion of lines in a surface, as in (3.6). Let  $\mathbb{S}$  denote the projective space that parametrizes surfaces of degree  $d$  in  $\mathbb{P}^3$ , as before.

**5.7.8. Proposition** In  $\mathbb{P}^3$ , the surfaces of degree  $d$  that contain a line form a closed subset of the space  $\mathbb{S}$ .

*proof.* Let  $\mathbb{G}$  be the Grassmanian  $G(2, 4)$  of lines in  $\mathbb{P}^3$ , and let  $\Xi$  be the subset of  $\mathbb{G} \times \mathbb{S}$  of pairs of pairs  $[\ell], [S]$  such that  $\ell \subset S$ . Lemma 3.6.17 tells us that  $\Xi$  is a closed subset of  $\mathbb{G} \times \mathbb{S}$ . Therefore its image in  $\mathbb{S}$  is closed.  $\square$

## 5.8 Fibre Dimension

A function  $Y \xrightarrow{\delta} \mathbb{Z}$  from a variety to the integers is *constructible* if, for every integer  $n$ , the set of points of  $Y$  such that  $\delta(p) = n$  is constructible, and  $\delta$  is *upper semicontinuous* if for every  $n$ , the set of points such that  $\delta(p) \geq n$  is closed. For brevity, we refer to an upper semicontinuous function as *semicontinuous*, though the term is ambiguous, since a function might be lower semicontinuous.

A function  $\delta$  on a curve  $C$  is semicontinuous if and only if for every integer  $n$ , there is a nonempty open subset  $C'$  of  $C$  such that  $\delta(p) = n$  for all points  $p$  of  $C'$  and  $\delta(p) \geq n$  for all points not in  $C'$ .

The next curve criterion for semicontinuous functions follows from the criterion for closed sets.

**5.8.1. Proposition.** (*curve criterion for semicontinuity*) Let  $Y$  be a variety. A function  $Y \xrightarrow{\delta} \mathbb{Z}$  is semicontinuous if and only if it is a constructible function, and for every morphism  $C \xrightarrow{f} Y$  from a smooth curve  $C$  to  $Y$ , the composition  $\delta \circ f$  is a semicontinuous function on  $C$ .  $\square$

Let  $Y \xrightarrow{f} X$  be a morphism of varieties, let  $q$  be a point of  $Y$ , and let  $Y_p$  be the fibre of  $f$  over  $p = f(q)$ . The *fibre dimension*  $\delta(q)$  of  $f$  at  $q$  is the maximum among the dimensions of the components of the fibre that contain  $q$ .

**5.8.2. Theorem.** (*semicontinuity of fibre dimension*) Let  $Y \xrightarrow{u} X$  be a morphism of varieties, and let  $\delta(q)$  denote the fibre dimension at a point  $q$  of  $Y$ .

(i) Suppose that  $X$  is a smooth curve, that  $Y$  has dimension  $n$ , and that the image of  $u$  is a point. Then  $\delta$  is constant: Every nonempty fibre has constant dimension  $n - 1$ .

(ii) Suppose that the image of  $Y$  contains a nonempty open subset of  $X$ , and let the dimensions of  $X$  and  $Y$  be  $m$  and  $n$ , respectively. There is a nonempty open subset  $X'$  of  $X$  such that  $\delta(q) = n - m$  for every point  $q$  in the inverse image of  $X'$ .

(iii)  $\delta$  is a semicontinuous function on  $Y$ .

The proof of this theorem is left as a long exercise. When you have done it, you will have understood the chapter.



## Chapter 6 MODULES

- 6.1 The Structure Sheaf
- 6.2  $\mathcal{O}$ -Modules
- 6.3 The Sheaf Property
- 6.4 Some  $\mathcal{O}$ -Modules
- 6.5 Direct Image
- 6.6 Twisting
- 6.7 Proof of Theorem 6.3.2

This chapter explains how modules on a variety are defined. variety.

We will need few facts about localization. Recall that, if  $s$  is a nonzero element of a domain  $A$ , the symbol  $A_s$  stands for the localization  $A[s^{-1}]$ , and if  $\text{Spec } A = X$ , then  $\text{Spec } A_s = X_s$ .

- Let  $U = \text{Spec } A$  be an affine variety. The intersection of two localizations  $U_s = \text{Spec } A_s$  and  $U_t = \text{Spec } A_t$  is the localization  $U_{st} = \text{Spec } A_{st}$ .
- Let  $W \subset V \subset U$  be affine open subsets of a variety  $X$ . If  $V$  is a localization of  $U$  and  $W$  is a localization of  $V$ , then  $W$  is a localization of  $U$  (2.6.2).
- The affine open subsets of a variety  $X$  form a basis for the topology on a variety  $X$ . The localizations of an affine variety form a basis for its topology (??).
- If  $U$  and  $V$  are affine open subsets of  $X$ , the open sets  $W$  that are localizations, both of  $U$  and of  $V$ , form a basis for the topology on  $U \cap V$ . (2.6.2).

### 6.1 The Structure Sheaf.

We introduce two categories associated to a variety  $X$ . The first is the category (*opens*), whose objects are the open subsets of  $X$ , and whose morphisms are inclusions: If  $U$  and  $V$  are open sets and if  $V \subset U$ , there is a unique morphism  $V \rightarrow U$  in (*opens*), and if  $V \not\subset U$  there is no morphism  $V \rightarrow U$ .

We also introduce a subcategory (*affines*) of the category (*opens*). Its objects are the affine open subsets of  $X$ , and its morphisms are localizations. A morphism  $V \rightarrow U$  in (*opens*) is a morphism in (*affines*) if  $U$  is affine and  $V$  is a localization of  $U$  – if  $V$  is an open subset of the form  $U_s$ , where  $s$  is a nonzero element of the coordinate algebra of  $U$ .

The *structure sheaf*  $\mathcal{O}_X$  on a variety  $X$  is the functor from affine open sets to algebras,

$$(6.1.1) \quad (\text{affines})^\circ \xrightarrow{\mathcal{O}_X} (\text{algebras})$$

that sends an affine open set  $U = \text{Spec } A$  to its coordinate algebra. The coordinate algebra  $A$  is then denoted by  $\mathcal{O}_X(U)$ .

As has been noted before, inclusions  $V \rightarrow U$  of affine open subsets needn't be localizations. We focus attention on localizations because the relation between the coordinate algebras of an affine variety and a localization is easy to understand. However, the structure sheaf can be extended with little difficulty to the category (*opens*), (See Corollary 6.1.3 below.)

A brief review about regular functions: The *function field* of a variety  $X$  is the field of fractions of the coordinate algebra of any one of its affine open subsets, and a *rational function* on  $X$  is a nonzero element of

the function field. A rational function  $f$  is *regular* on an affine open set  $U = \text{Spec } A$  if it is an element of  $A$ , and  $f$  is *regular* on a nonempty open set  $U$  that can be covered by affine open sets on which it is regular. Thus the function field of a variety  $X$  contains the regular functions on every nonempty open subset, and the regular functions are governed by the regular functions on the affine open subsets.

An affine variety is determined by its regular functions, but the regular functions don't suffice to determine a variety that isn't affine. For instance, the only rational functions that are regular everywhere on the projective line  $\mathbb{P}^1$  are the constant functions, which are useless. We will be interested in regular functions on non-affine open sets, especially in functions that are regular on the whole variety, but one should always work with the affine open sets, where the definition of a regular function is clear.

**6.1.2. Lemma.** *Let  $U$  and  $V$  be open subsets of a variety  $X$ , with  $V \subset U$ . If a rational function is regular on  $U$ , it is also regular on  $V$ .*  $\square$

Thus if  $U \subset V$  is an inclusion of affine open subsets, say  $U = \text{Spec } A$  and  $V = \text{Spec } B$ , then  $A \subset B$ . However, it won't be clear how to construct  $B$  from  $A$  unless  $B$  is a localization. If  $V = U_s$ , then  $B = A[s^{-1}]$ . If  $B$  isn't a localization, the exact relationship between  $A$  and  $B$  remains obscure.

**6.1.3. Corollary.** *Let  $X$  be a variety. When  $\mathcal{O}_X(U)$  is defined to be the algebra of regular functions on an open subset  $U$ , the structure sheaf  $\mathcal{O}_X$  on a variety  $X$  extends to a functor*

$$(\text{opens})^\circ \xrightarrow{\mathcal{O}_X} (\text{algebras})$$

$\square$

If it is clear which variety is being studied, we may write  $\mathcal{O}$  for  $\mathcal{O}_X$ .

For any open set  $U$ , the algebra of regular functions on  $U$  is denoted by  $\mathcal{O}_X(U)$ . Its elements are called *sections* of the structure sheaf  $\mathcal{O}_X$  on  $U$ .

When  $V \rightarrow U$  is a morphism in  $(\text{opens})$ , Lemma 6.1.2 tells us that  $\mathcal{O}_X(U)$  is contained in  $\mathcal{O}_X(V)$ . This gives us the homomorphism, an inclusion,

$$\mathcal{O}_X(U) \rightarrow \mathcal{O}_X(V)$$

that makes  $\mathcal{O}_X$  into a functor. Note that arrows are reversed by  $\mathcal{O}_X$ . If  $V \rightarrow U$ , then  $\mathcal{O}_X(U) \rightarrow \mathcal{O}_X(V)$ . A functor that reverses arrows is a *contravariant* functor. The superscript  $\circ$  in (6.1.1) and (6.1.3) is a customary notation to indicate that a functor is contravariant.

**6.1.4. Proposition** *The (extended) structure sheaf has the following **sheaf property**:*

- *Let  $Y$  be an open subset of  $X$ , and let  $U^i = \text{Spec } A_i$  be affine open subsets that cover  $Y$ . Then*

$$\mathcal{O}_X(Y) = \bigcap \mathcal{O}_X(U^i) \quad (= \bigcap A_i)$$

This sheaf property is especially simple because regular functions are elements of the function field. It is more complicated for  $\mathcal{O}$ -modules, which will be defined in the next section.

By definition, if  $f$  is a regular function on  $X$ , there is a covering by affine open sets  $U^i$  such that  $f$  is regular on each of them, i.e., that  $f$  is in  $\bigcup \mathcal{O}(U^i)$ . Therefore the next lemma proves the proposition.

**6.1.5. Lemma.** *Let  $Y$  be an open subset of a variety  $X$ . The intersection  $\bigcap \mathcal{O}_X(U^i)$  is the same for every affine open covering  $\{U^i\}$  of  $Y$ .*

We prove the lemma first in the case of a covering of an affine open set by localizations.

**6.1.6. Sublemma.** *Let  $U = \text{Spec } A$  be an affine variety, and let  $\{U^i\}$  be a covering of  $U$  by localizations, say  $U^i = \text{Spec } A_{s_i}$ . Then  $A = \bigcap A_{s_i}$ , i.e.,  $\mathcal{O}(U) = \bigcap \mathcal{O}(U^i)$ .*

*proof.* A finite subset of the set  $\{U^i\}$  will cover  $U$ , so we may assume that the index set is finite.

It is clear that  $A$  is a subset of  $\bigcap A_{s_i}$ . Let  $\alpha$  be an element of  $\bigcap A_{s_i}$ . So  $\alpha = s_i^{-r} a_i$ , or  $s_i^r \alpha = a_i$  for some  $a_i$  in  $A$  and some integer  $r$ , and we can use the same  $r$  for every  $i$ . Because  $\{U^i\}$  covers  $U$ , the elements  $s_i$



generate the unit ideal in  $A$ , and so do their powers  $s_i^r$ . There are elements  $b_i$  in  $A$  such that  $\sum b_i s_i^r = 1$ . Then  $\alpha = \sum b_i s_i^r \alpha = \sum b_i a_i$  is in  $A$ .  $\square$

*proof of Lemma 6.1.5.* Say that  $Y$  is covered by affine open sets  $\{U^i\}$  and also by affine open sets  $\{V^j\}$ . We cover the intersections  $U^i \cap V^j$  by open sets  $W^{ij\nu}$  that are localizations of  $U^i$  and also localizations of  $V^j$ . Fixing  $i$  and letting  $j$  and  $\nu$  vary, the set  $\{W^{ij\nu}\}_{j,\nu}$  will be a covering of  $U^i$  by localizations, and the sublemma shows that  $\mathcal{O}(U^i) = \bigcap_{j,\nu} \mathcal{O}(W^{ij\nu})$ . Then  $\bigcap_i \mathcal{O}(U^i) = \bigcap_{i,j,\nu} \mathcal{O}(W^{ij\nu})$ . Similarly,  $\bigcap_j \mathcal{O}(V^j) = \bigcap_{i,j,\nu} \mathcal{O}(W^{ij\nu})$ .  $\square$

### 6.1.7. Example.

Let  $A$  denote the polynomial ring  $\mathbb{C}[x, y]$ , and let  $Y$  be the complement of a point  $p$  in affine space  $X = \text{Spec } A$ . We cover  $Y$  by two localizations of  $X$ ,  $X_x = \text{Spec } A[x^{-1}]$  and  $X_y = \text{Spec } A[y^{-1}]$ . A regular function on  $Y$  will be regular on  $X_x$  and on  $X_y$ , so it will be in the intersection of their coordinate algebras. The intersection  $A[x^{-1}] \cap A[y^{-1}]$  is  $A$ . So the sections of the structure sheaf  $\mathcal{O}_X$  on  $Y$  are the same as the sections on  $X$ . They are the elements of  $A$ .  $\square$

## 6.2 $\mathcal{O}$ -Modules

An  $\mathcal{O}_X$ -module on a variety  $X$  is a structure that associates a module to every affine open subset.

**6.2.1. Definition.** An  $\mathcal{O}$ -module  $\mathcal{M}$  on a variety  $X$  is a (contravariant) functor

$$(\text{affines})^\circ \xrightarrow{\mathcal{M}} (\text{modules})$$

such that  $\mathcal{M}(U)$  is an  $\mathcal{O}(U)$ -module for every affine open set  $U$ , and such that, if  $s$  is a nonzero element of  $\mathcal{O}(U)$ , the module  $\mathcal{M}(U_s)$  is the localization of  $\mathcal{M}(U)$ :

$$\mathcal{M}(U_s) = \mathcal{M}(U)_s$$

If  $\mathcal{M}(U)$  is a finite  $\mathcal{O}(U)$ -module for every affine open set  $U$ ,  $\mathcal{M}$  is called a *finite  $\mathcal{O}$ -module*.

A *section* of an  $\mathcal{O}$ -module  $\mathcal{M}$  on an affine open set  $U$  is an element of  $\mathcal{M}(U)$ .

A *homomorphism*  $\mathcal{M} \xrightarrow{\varphi} \mathcal{N}$  of  $\mathcal{O}$ -modules consists of homomorphisms of  $\mathcal{O}(U)$ -modules

$$\mathcal{M}(U) \xrightarrow{\varphi(U)} \mathcal{N}(U)$$

for each affine open subset  $U$  of  $X$  such that, if  $s$  is a nonzero element of  $\mathcal{O}(U)$ , the homomorphism  $\varphi(U_s)$  is the localization of  $\varphi(U)$ .

A sequence of homomorphisms

$$(6.2.2) \quad \mathcal{M} \rightarrow \mathcal{N} \rightarrow \mathcal{P}$$

of  $\mathcal{O}$ -modules on a variety  $X$  is *exact* if the sequence of sections  $\mathcal{M}(U) \rightarrow \mathcal{N}(U) \rightarrow \mathcal{P}(U)$  is exact for every affine open subset  $U$  of  $X$ .  $\square$

At first glance, the definition of  $\mathcal{O}$ -module will seem complicated. However, when a module has a natural definition, the data involved in the definition take care of themselves. This will become clear as we go along.

**Note.** When stating that  $\mathcal{M}(U_s)$  is the localization of  $\mathcal{M}(U)$ , it would be more correct to say that  $\mathcal{M}(U_s)$  and  $\mathcal{M}(U)_s$  are canonically isomorphic. Let's not worry about this.

One example of an  $\mathcal{O}$ -module is the *free module*  $\mathcal{O}^k$ . The sections of the free module on an affine open set  $U$  are the elements of the free  $\mathcal{O}(U)$ -module  $\mathcal{O}(U)^k$ . In particular,  $\mathcal{O}$  can be considered as an  $\mathcal{O}$ -module.

The *kernel*, *image*, and *cokernel* of a homomorphism  $\mathcal{M} \xrightarrow{\varphi} \mathcal{N}$  are among the operations that can be made on  $\mathcal{O}$ -modules. The kernel  $\mathcal{K}$  of  $\varphi$  is the  $\mathcal{O}$ -module defined by  $\mathcal{K}(U) = \ker(\mathcal{M}(U) \xrightarrow{\varphi(U)} \mathcal{N}(U))$  for every affine open set  $U$ , and the image and cokernel are defined analogously. Many operations, including these, are compatible with localization.

### 6.3 The Sheaf Property

In this section, we extend an  $\mathcal{O}$ -module  $\mathcal{M}$  on a variety  $X$  to a functor  $(\text{opens})^\circ \xrightarrow{\widetilde{\mathcal{M}}} (\text{modules})$  on all open subsets of  $X$ , such that  $\widetilde{\mathcal{M}}(Y)$  is an  $\mathcal{O}(Y)$ -module for every open subset  $Y$ , and when  $U$  is an affine open set,  $\widetilde{\mathcal{M}}(U) = \mathcal{M}(U)$ .

The tilde is used for clarity here. We will drop it when we have finished with the discussion, and use the same notation  $\mathcal{M}$  for the functor on *(affines)* and for its extension to *(opens)*.

**6.3.1. Terminology.** If  $(\text{opens})^\circ \xrightarrow{\widetilde{\mathcal{M}}} (\text{modules})$  is a functor and let  $U$  be an open subset, an element of  $\widetilde{\mathcal{M}}(U)$  is a *section* of  $\widetilde{\mathcal{M}}$  on  $U$ . If  $V \xrightarrow{j} U$  is an inclusion of open subsets, the associated homomorphism  $\widetilde{\mathcal{M}}(U) \rightarrow \widetilde{\mathcal{M}}(V)$  is called the *restriction* from  $U$  to  $V$ .

The restriction to  $V$  of a section  $m$  on  $U$  may be denoted by  $j^\circ m$ . However, the operation of restriction occurs very often. Because of this, we often abbreviate, using the same symbol  $m$  for a section and for its restriction. Also, if an open set  $V$  is contained in two open sets  $U$  and  $U'$ , and if  $m, m'$  are sections of  $\widetilde{\mathcal{M}}$  on  $U$  and  $U'$ , respectively, we may say that  $m$  and  $m'$  are *equal on  $V$*  if their restrictions to  $V$  are equal.  $\square$

**6.3.2. Theorem.** An  $\mathcal{O}$ -module  $\mathcal{M}$  extends uniquely to a functor

$$(\text{opens})^\circ \xrightarrow{\widetilde{\mathcal{M}}} (\text{modules})$$

that has the sheaf property described below. Moreover, for every open set  $U$ ,  $\widetilde{\mathcal{M}}(U)$  is an  $\mathcal{O}(U)$ -module, and for every inclusion  $V \rightarrow U$  of nonempty open sets, the map  $\widetilde{\mathcal{M}}(U) \rightarrow \widetilde{\mathcal{M}}(V)$  is compatible with scalar multiplication in this sense:

Let  $m$  be a section of  $\widetilde{\mathcal{M}}$  on  $U$ , let  $\alpha$  be a regular function on  $U$ , and let  $m'$  and  $\alpha'$  denote the restrictions to  $V$ . The restriction of  $\alpha m$  is  $\alpha' m'$ .

In order not to break up the discussion, we have put the proof of this theorem into Section 6.7 at the end of the chapter.

#### (6.3.3) the sheaf property

The sheaf property is the key requirement that determines the extension of an  $\mathcal{O}$ -module  $\mathcal{M}$  to a functor  $\widetilde{\mathcal{M}}$  on *(opens)*.

Let  $Y$  be an open subset of  $X$ , and let  $\{U^i\}$  be a covering of  $Y$  by affine open sets. The intersections  $U^{ij} = U^i \cap U^j$  are also affine open sets, so  $\mathcal{M}(U^i)$  and  $\mathcal{M}(U^{ij})$  are defined. The sheaf property asserts that an element  $m$  of  $\widetilde{\mathcal{M}}(Y)$  corresponds to a set of elements  $m_i$  in  $\mathcal{M}(U^i)$  such that the restrictions of  $m_j$  and  $m_i$  to  $U^{ij}$  are equal.

If the affine open subsets  $U^i$  are indexed by  $i = 1, \dots, n$ , the sheaf property asserts that an element of  $\widetilde{\mathcal{M}}(Y)$  is determined by a vector  $(m_1, \dots, m_n)$  with  $m_i$  in  $\mathcal{M}(U^i)$ , such that the restrictions of  $m_i$  and  $m_j$  to  $U^{ij}$  are equal. This means that  $\widetilde{\mathcal{M}}(Y)$  is the kernel of the map

$$(6.3.4) \quad \prod_i \mathcal{M}(U^i) \xrightarrow{\beta} \prod_{i,j} \mathcal{M}(U^{ij})$$

that sends the vector  $(m_1, \dots, m_n)$  to the  $n \times n$  matrix  $(z_{ij})$ , where  $z_{ij}$  is the difference  $m_j - m_i$  of the restrictions of  $m_j$  and  $m_i$  to  $U^{ij}$ . The analogous description is true when the index set is infinite.

In short, the sheaf property tells us that sections of  $\widetilde{\mathcal{M}}$  are determined locally: A section on an open set  $Y$  is determined by its restrictions to the open subsets  $U^i$  of an affine covering of  $Y$ .

**Note.** The morphisms  $U^{ij} \rightarrow U^i$  needn't be localizations, and if not the restriction maps  $\mathcal{M}(U^i) \rightarrow \mathcal{M}(U^{ij})$  aren't a part of the structure of an  $\mathcal{O}$ -module. We need a definition of the restriction map for an arbitrary inclusion  $V \rightarrow U$  of affine open subsets. This point will be taken care of by the proof of Theorem 6.3.2. (See Step 2 in Section 6.7.) Let's not worry about it here.  $\square$

We drop the tilde now.

The next corollary follows from Theorem 6.3.2.

**6.3.5. Corollary.** *Let  $\{U^i\}$  be an affine open covering of a variety  $X$ .*

**(i)** *An  $\mathcal{O}$ -module  $\mathcal{M}$  is zero if and only if  $\mathcal{M}(U^i) = 0$  for every  $i$ .*

**(ii)** *A homomorphism  $\mathcal{M} \xrightarrow{\varphi} \mathcal{N}$  of  $\mathcal{O}$ -modules is injective, surjective, or bijective if and only if the maps  $\mathcal{M}(U^i) \xrightarrow{\varphi(U^i)} \mathcal{N}(U^i)$  are injective, surjective, or bijective, respectively, for every  $i$ .*

*proof.* **(i)** Let  $V$  be any open subset of  $X$ . We can cover the intersections  $V \cap U^i$  by affine open sets  $V^{i\nu}$  that are localizations of  $U^i$ , and these sets, taken together, cover  $V$ . If  $\mathcal{M}(U^i) = 0$ , then the localizations  $\mathcal{M}(V^{i\nu})$  are zero too. The sheaf property shows that the map  $\mathcal{M}(V) \rightarrow \prod \mathcal{M}(V^{i\nu})$  is injective, and therefore  $\mathcal{M}(V) = 0$ .

**(ii)** This follows from **(i)** because a homomorphism  $\varphi$  is injective or surjective if and only if its kernel or its cokernel is zero.  $\square$

### (6.3.6) families of open sets

It is convenient to have a compact notation for the sheaf property. For this, one can use symbols to represent families of open sets. Say that  $\mathbb{U}$  and  $\mathbb{V}$  represent families of open sets  $\{U^i\}$  and  $\{V^\nu\}$ , respectively. A *morphism* of families  $\mathbb{V} \rightarrow \mathbb{U}$  consists of a morphism from each  $V^\nu$  to one of the subsets  $U^i$ . Such a morphism will be given by a map  $\nu \rightsquigarrow i_\nu$  of index sets, such that  $V^\nu \subset U^{i_\nu}$ .

There may be more than one morphism  $\mathbb{V} \rightarrow \mathbb{U}$ , because a subset  $V^\nu$  may be contained in more than one of the subsets  $U^i$ . To define a morphism, one must make a choice among those subsets. For example, let  $\mathbb{U} = \{U^i\}$  be a family of open sets, and let  $V$  be another open set. There is a morphism  $V \rightarrow \mathbb{U}$  that sends  $V$  to  $U^i$  whenever  $V \subset U^i$ . In the other direction, there is a unique morphism  $\mathbb{U} \rightarrow V$  provided that  $U^i \subset V$  for all  $i$ .

A functor  $(opens)^\circ \xrightarrow{\mathcal{M}} (modules)$  can be extended to families  $\mathbb{U} = \{U^i\}$  by defining

$$(6.3.7) \quad \mathcal{M}(\mathbb{U}) = \prod \mathcal{M}(U^i).$$

Then a morphism of families  $\mathbb{V} \xrightarrow{f} \mathbb{U}$  defines a map  $\mathcal{M}(\mathbb{U}) \xrightarrow{f^\circ} \mathcal{M}(\mathbb{V})$  in a way that is fairly obvious, though notation for it is clumsy. Say that  $f$  is given by a map  $\nu \rightsquigarrow i_\nu$  of index sets, with  $V^\nu \rightarrow U^{i_\nu}$ . A section of  $\mathcal{M}$  on  $\mathbb{U}$ , an element of  $\mathcal{M}(\mathbb{U})$ , can be thought of as a vector  $(u_i)$  with  $u_i \in \mathcal{M}(U^i)$ , and a section of  $\mathcal{M}(\mathbb{V})$  as a vector  $(v_\nu)$  with  $v_\nu \in \mathcal{M}(V^\nu)$ . If  $v_\nu$  is the restriction of  $u_{i_\nu}$  to  $V^\nu$ , the map  $f^\circ$  sends  $(u_i) \rightarrow (v_\nu)$ .

We can write the sheaf property in terms of families of open sets. Let  $\mathbb{U}_0 = \{U^i\}$  be an affine open covering of an open set  $Y$ , and let  $\mathbb{U}_1$  denote the family  $\{U^{ij}\}$  of intersections:  $U^{ij} = U^i \cap U^j$ , which are also affine. Then we have a morphism  $\mathbb{U}_0 \rightarrow Y$ , and the two sets of inclusions

$$U^{ij} \subset U^i \quad \text{and} \quad U^{ij} \subset U^j$$

define two morphisms of families  $\mathbb{U}_1 \xrightarrow{d_0, d_1} \mathbb{U}_0$  of affine open sets,  $U^{ij} \xrightarrow{d_0} U^j$  and  $U^{ij} \xrightarrow{d_1} U^i$ . The two composed morphisms  $\mathbb{U}_1 \xrightarrow{d_i} \mathbb{U}_0 \rightarrow Y$  are equal. They form what we call a *covering diagram*

$$(6.3.8) \quad Y \longleftarrow \mathbb{U}_0 \rightrightarrows \mathbb{U}_1$$

When we apply a functor  $(opens)^\circ \xrightarrow{\mathcal{M}} (modules)$  to this diagram, we obtain a sequence

$$(6.3.9) \quad 0 \rightarrow \mathcal{M}(Y) \xrightarrow{\alpha_{\mathbb{U}}} \mathcal{M}(\mathbb{U}_0) \xrightarrow{\beta_{\mathbb{U}}} \mathcal{M}(\mathbb{U}_1)$$

where  $\alpha_{\mathbb{U}}$  is the restriction map and  $\beta_{\mathbb{U}}$  is the difference  $\mathcal{M}(d_0) - \mathcal{M}(d_1)$  of the maps induced by the two morphisms  $\mathbb{U}_1 \rightrightarrows \mathbb{U}_0$ . The sheaf property for the covering  $\mathbb{U}_0$  of  $Y$  is the assertion that this sequence is exact, which means that  $\alpha_{\mathbb{U}}$  is injective, and that its image is the kernel of  $\beta_{\mathbb{U}}$ .

**6.3.10. Note.** One can suppose that the open sets  $U^i$  that make a covering are distinct. However, the intersections won't be distinct, because  $U^{ij} = U^{ji}$ . Also,  $U^{ii} = U^i$ . These coincidences lead to redundancy in the statement (6.3.9) of the sheaf property. If the indices are  $i = 1, \dots, k$ , we only need to look at intersections  $U^{ij}$  with  $i < j$ . The product  $\mathcal{M}(\mathbb{U}_1) = \prod_{i,j} \mathcal{M}(U^{ij})$  that appears in the sheaf property can be replaced by the product  $\prod_{i < j} \mathcal{M}(U^{ij})$  with increasing pairs of indices. For instance, suppose that an open set  $Y$  is covered by two affine open sets  $U$  and  $V$ . Then, with  $\mathbb{U}_0 = \{U, V\}$ , the sheaf property is the exact sequence

$$0 \rightarrow \mathcal{M}(Y) \xrightarrow{\alpha} \mathcal{M}(U) \times \mathcal{M}(V) \xrightarrow{\beta} \mathcal{M}(U \cap V) \times \mathcal{M}(U \cap V) \times \mathcal{M}(V \cap U) \times \mathcal{M}(V \cap V)$$

is equivalent with the exact sequence

$$(6.3.11) \quad 0 \rightarrow \mathcal{M}(\rightarrow \mathcal{M}(U) \times \mathcal{M}(V) \xrightarrow{-, +} \mathcal{M}(U \cap V))$$

□

### 6.3.12. Example.

We go back to Proposition 6.4.18, which describes the correspondence between an  $\mathcal{O}$ -module  $\mathcal{M}$  on an affine variety  $X = \text{Spec } A$  and an  $A$ -module  $M$ . Namely, if  $U = \text{Spec } B$  is an affine open subset of  $X$ , then  $\mathcal{M}(U) = B \otimes_A M$ . This example shows that, when a subset  $U$  isn't affine, defining  $\mathcal{M}(U) = B \otimes_A M$  may be wrong.

Let  $X$  be the affine plane  $\text{Spec } A$ ,  $A = \mathbb{C}[x, y]$ , let  $U$  be the complement of the origin in  $X$ , and let  $M$  be the  $A$ -module  $A/yA$ . This module can be identified with  $\mathbb{C}[x]$ , which becomes an  $A$ -module when scalar multiplication by  $y$  is defined to be zero. Here  $\mathcal{O}(U) = \mathcal{O}(X) = A$  (6.1.7). If we followed the method used for affine open sets, we would set  $\mathcal{M}(U) = A \otimes_A M = \mathbb{C}[x]$ .

To identify  $\mathcal{M}(U)$  correctly, we cover  $U$  by the two localizations  $U_x = \text{Spec } A[x^{-1}]$  and  $U_y = \text{Spec } A[y^{-1}]$ . Then  $\mathcal{M}(U_x) = M[x^{-1}]$  while  $\mathcal{M}(U_y) = 0$ . The sheaf property of  $\mathcal{M}$  shows that  $\mathcal{M}(U) \approx \mathcal{M}(U_x) = M[x^{-1}] = \mathbb{C}[x, x^{-1}]$ . □

We have been working with nonempty open sets. The next lemma takes care of the empty set.

**6.3.13. Lemma.** *The only section of an  $\mathcal{O}$ -module  $\mathcal{M}$  on the empty set is the zero section:  $\mathcal{M}(\emptyset) = \{0\}$ . In particular,  $\mathcal{O}(\emptyset)$  is the zero ring.*

*proof.* The empty set is covered by the empty covering, the covering indexed by the empty set. Since we want the sheaf property to be true,  $\mathcal{M}(\emptyset)$  will be contained in an empty product. We want both the empty product and  $\mathcal{M}(\emptyset)$  to be modules, so we have no choice but to set them equal to  $\{0\}$ .

If you find this reasoning pedantic, you can take  $\mathcal{M}(\emptyset) = \{0\}$  as an axiom. □

### (6.3.14) Interlude: a useful diagram

We consider a commutative diagram of abelian groups of the form

$$\begin{array}{ccccccc} 0 & \longrightarrow & A & \xrightarrow{f} & B & \xrightarrow{g} & C \\ & & \alpha \downarrow & & \beta \downarrow & & \gamma \downarrow \\ 0 & \longrightarrow & A' & \xrightarrow{f'} & B' & \xrightarrow{g'} & C \end{array}$$

**6.3.15. Lemma.** (i) *Suppose that the rows of the diagram are exact. If  $\beta$  and  $\gamma$  are bijective, so is  $\alpha$ .*

(ii) *Suppose that the bottom row of the diagram is exact. If  $\alpha$  is bijective and  $\beta$  and  $\gamma$  are injective, the top row is exact.*

It is customary to leave the proofs of such statements to the reader. Since this sort of reasoning may be new, we'll give the proof of the most interesting part of (ii), the exactness of the top row at  $B$ , which is the statement that  $\ker g = \text{im } f$ . We show that if  $x$  is an element of  $B$  such that  $gx = 0$ , then  $x = fy$  for some element  $y$  of  $A$ . The hypothesis that  $\gamma$  is injective won't be used here.

Let  $x' = \beta x$ . Then  $g'x' = g'\beta x = \gamma gx = 0$ . Since the bottom row is exact,  $x' = f'y'$  for some element  $y'$  of  $A'$ . Let  $y$  be the element of  $A$  such that  $\alpha y = y'$ . Then  $\beta fy = f'\alpha y = f'y' = x' = \beta x$ . Since  $\beta$  is injective,  $fy = x$ .  $\square$

### (6.3.16) the coherence property

In addition to the sheaf property, an  $\mathcal{O}$ -module on a variety  $X$  has a property called *coherence*.

**6.3.17. Proposition.** (the coherence property) Let  $Y$  be an open subset of a variety  $X$ , let  $s$  be a nonzero regular function on  $Y$ , and let  $\mathcal{M}$  be an  $\mathcal{O}_X$ -module. Then  $\mathcal{M}(Y_s)$  is the localization  $\mathcal{M}(Y)_s$  of  $\mathcal{M}(Y)$ .

Compatibility with localization is a requirement for an  $\mathcal{O}$ -module when  $Y$  is affine. The coherence property is the extension to all open subsets.

*proof of Proposition 6.3.17.* Let  $\mathbb{U}_0 = \{U^i\}$  be a family of affine open sets that covers the open set  $Y$ . The intersections  $U^{ij}$  will be affine open sets too. We inspect the covering diagram  $Y \leftarrow \mathbb{U}_0 \rightrightarrows \mathbb{U}_1$ . If  $s$  is a nonzero regular function on  $Y$ , the localization of this diagram forms a covering diagram  $Y_s \leftarrow \mathbb{U}_{0,s} \rightrightarrows \mathbb{U}_{1,s}$ , in which  $\mathbb{U}_{0,s} = \{U_s^i\}$  is an affine covering of  $Y_s$ . Therefore  $\mathcal{M}(\mathbb{U}_0)_s \approx \mathcal{M}(\mathbb{U}_{0,s})$ . The sheaf property gives us exact sequences

$$0 \rightarrow \mathcal{M}(Y) \rightarrow \mathcal{M}(\mathbb{U}_0) \rightarrow \mathcal{M}(\mathbb{U}_1) \quad \text{and} \quad 0 \rightarrow \mathcal{M}(Y_s) \rightarrow \mathcal{M}(\mathbb{U}_{0,s}) \rightarrow \mathcal{M}(\mathbb{U}_{1,s})$$

and the localization of the first sequence maps to the second one:

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{M}(Y)_s & \longrightarrow & \mathcal{M}(\mathbb{U}_0)_s & \longrightarrow & \mathcal{M}(\mathbb{U}_1)_s \\ & & a \downarrow & & b \downarrow & & c \downarrow \\ 0 & \longrightarrow & \mathcal{M}(Y_s) & \longrightarrow & \mathcal{M}(\mathbb{U}_{0,s}) & \longrightarrow & \mathcal{M}(\mathbb{U}_{1,s}) \end{array}$$

The bottom row is exact, and since localization is an exact operation, the top row of the diagram is exact too. Since  $\mathbb{U}_0$  and  $\mathbb{U}_1$  are families of affine open sets, the vertical arrows  $b$  and  $c$  are bijections. Therefore  $a$  is a bijection. This is the coherence property.  $\square$

## 6.4 Some $\mathcal{O}$ -Modules

### 6.4.1. modules on a point

Let's denote a point, the affine variety  $\text{Spec } \mathbb{C}$ , by  $p$ . The point has only one nonempty open set: the whole space  $p$ , and  $\mathcal{O}_p(p) = \mathbb{C}$ . Let  $\mathcal{M}$  be an  $\mathcal{O}_p$ -module. The space of global sections  $\mathcal{M}(p)$  is an  $\mathcal{O}_p(p)$ -module, a complex vector space. To define  $\mathcal{M}$ , that vector space can be assigned arbitrarily. One may say that a module on the point *is* a complex vector space.  $\square$

### 6.4.2. the residue field module $\kappa_p$ .

Let  $p$  be a point of a variety  $X$ . A residue field module  $\kappa_p$  is defined as follows: If  $U$  is an affine open subset of  $X$  that contains  $p$ , then  $\mathcal{O}(U)$  has a residue field  $k(p)$  at  $p$ , and  $\kappa_p(U) = k(p)$ . If  $U$  doesn't contain  $p$ , then  $\kappa_p(U) = 0$ .

### 6.4.3. ideals.

An ideal  $\mathcal{I}$  of the structure sheaf is an  $\mathcal{O}$ -submodule of  $\mathcal{O}$ .

Let  $p$  be a point of a variety  $X$ . The *maximal ideal* at  $p$ , which we denote by  $\mathfrak{m}_p$ , is an ideal. If an affine open subset  $U$  contains  $p$ , its coordinate algebra  $\mathcal{O}(U)$  will have a maximal ideal consisting of the elements that vanish at  $p$ . That maximal ideal is the module of sections  $\mathfrak{m}_p(U)$ . If  $U$  doesn't contain  $p$ , then  $\mathfrak{m}_p(U) = \mathcal{O}(U)$ .

When  $\mathcal{I}$  is an ideal of  $\mathcal{O}$ , we denote by  $V_X(\mathcal{I})$  the closed set of points  $p$  such that  $\mathcal{I} \subset \mathfrak{m}_p$  – the set of points at which all elements of  $\mathcal{I}$  vanish at  $p$ .

### 6.4.4. examples of homomorphisms

- (i) There is a homomorphism of  $\mathcal{O}$ -modules  $\mathcal{O} \rightarrow \kappa_p$  whose kernel is the maximal ideal  $\mathfrak{m}_p$ .
- (ii) Homomorphisms  $\mathcal{O}^n \rightarrow \mathcal{O}^m$  of free  $\mathcal{O}$ -modules correspond to  $m \times n$ -matrices of global sections of  $\mathcal{O}$ .

(iii) Let  $\mathcal{M}$  be an  $\mathcal{O}$ -module. The  $\mathcal{O}$ -module homomorphisms  $\mathcal{O} \xrightarrow{\varphi} \mathcal{M}$  correspond bijectively to global sections of  $\mathcal{M}$ . This is analogous to the fact that, when  $M$  is a module over a ring  $A$ , homomorphisms  $A \rightarrow M$  correspond to elements of  $M$ . To be explicit: If  $m$  is a global section of  $\mathcal{M}$ , the homomorphism  $\mathcal{O}(U) \xrightarrow{\varphi} \mathcal{M}(U)$  is multiplication by the restriction of  $m$  to  $U$ .

(iv) If  $f$  is a global section of  $\mathcal{O}$ , scalar multiplication by  $f$  defines a homomorphism  $\mathcal{M} \xrightarrow{f} \mathcal{M}$ .

#### 6.4.5. kernel

As we have remarked, many operations that one makes on modules over a ring are compatible with localization, and therefore can be made on  $\mathcal{O}$ -modules. However, when applied to sections over non-affine open sets the operations are almost *never* compatible with localization. One important exception is the kernel of a homomorphism.

**6.4.6. Proposition.** *Let  $X$  be a variety, and let  $0 \rightarrow \mathcal{K} \rightarrow \mathcal{M} \rightarrow \mathcal{N}$  be an exact sequence of  $\mathcal{O}$ -modules. For every open subset  $Y$  of  $X$ , the sequence of sections*

$$(6.4.7) \quad 0 \rightarrow \mathcal{K}(Y) \rightarrow \mathcal{M}(Y) \rightarrow \mathcal{N}(Y)$$

*is exact.*

*proof.* We choose a covering diagram  $Y \leftarrow \mathbb{U}_0 \hookrightarrow \mathbb{U}_1$ , and we inspect the diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{K}(\mathbb{U}_0) & \longrightarrow & \mathcal{M}(\mathbb{U}_0) & \longrightarrow & \mathcal{N}(\mathbb{U}_0) \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathcal{K}(\mathbb{U}_1) & \longrightarrow & \mathcal{M}(\mathbb{U}_1) & \longrightarrow & \mathcal{N}(\mathbb{U}_1) \end{array}$$

where the vertical maps are the maps  $\beta_{\mathbb{U}}$  described in (6.3.9). The rows are exact because  $\mathbb{U}_0$  and  $\mathbb{U}_1$  are families of affines, and the sheaf property asserts that the kernels of the vertical maps form the sequence (6.4.7). It is exact because taking kernels is a left exact operation.  $\square$

The section functor isn't right exact. When

$$0 \rightarrow \mathcal{K} \rightarrow \mathcal{M} \rightarrow \mathcal{N} \rightarrow 0$$

is a short exact sequence of  $\mathcal{O}$ -modules and  $Y$  is a non-affine open set, the sequence (6.4.7) may fail to be exact when a zero is added on the right. There is an example below. Cohomology, which will be discussed in the next chapter, is a substitute for right exactness.

#### 6.4.8. modules on the projective line

The projective line  $\mathbb{P}^1$  is covered by the standard open sets  $\mathbb{U}^0$  and  $\mathbb{U}^1$ , and the intersection  $\mathbb{U}^{01} = \mathbb{U}^0 \cap \mathbb{U}^1$  is a localization of  $\mathbb{U}^0$  and of  $\mathbb{U}^1$ . The coordinate algebras of these affine open sets are  $\mathbb{C}[u] = A_0$  and  $\mathbb{C}[v] = A_1$ , respectively, with  $v = u^{-1}$ , and  $\mathcal{O}(\mathbb{U}^{01}) = \mathbb{C}[u, u^{-1}] = A_{01}$ . The algebra  $A_{01}$  is the Laurent polynomial ring whose elements are (finite) combinations of powers of  $u$ , negative powers included. The sheaf property asserts that a global section of  $\mathcal{O}$  is determined by polynomials  $f(u)$  in  $A_0$  and  $g(v)$  in  $A_1$  such that  $f(u) = g(u^{-1})$  in  $A_{01}$ . The only such polynomials  $f, g$  are the constants. The constants are the only rational functions that are regular everywhere on  $\mathbb{P}^1$ . I think we knew this.

If  $\mathcal{M}$  is an  $\mathcal{O}$ -module,  $\mathcal{M}(\mathbb{U}^0) = M_0$  and  $\mathcal{M}(\mathbb{U}^1) = M_1$  will be modules over the algebras  $A_0$  and  $A_1$ , and the  $A_{01}$ -module  $\mathcal{M}(\mathbb{U}^{01}) = M_{01}$  can be obtained by localizing  $M_0$  and also by localizing  $M_1$ :  $M_0[u^{-1}] \approx M_{01} \approx M_1[v^{-1}]$ . A global section of  $\mathcal{M}$  is determined by a pair of elements  $m_1, m_2$  in  $M_1, M_2$  that become equal in the common localization  $M_{01}$ .

Suppose that  $M_0$  and  $M_1$  are free modules of rank  $r$  over  $A_0$  and  $A_1$ ,. Then  $M_{01}$  will be a free  $A_{01}$ -module of rank  $r$ . A basis  $\mathbf{B}_0$  of the free  $A_0$ -module  $M_0$  will also be a basis of the  $A_{01}$ -module  $M_{01}$ , and a basis  $\mathbf{B}_1$  of  $M_1$  will be a basis of  $M_{01}$ . When regarded as bases of  $M_{01}$ ,  $\mathbf{B}_0$  and  $\mathbf{B}_1$  will be related by an  $r \times r$  invertible  $A_{01}$ -matrix  $P$ , and that matrix determines  $\mathcal{M}$  up to isomorphism. When  $r = 1$ ,  $P$  will be an invertible  $1 \times 1$  matrix in the Laurent polynomial ring  $A_{01}$  – a unit of that ring. The units in  $A_{01}$  are scalar multiples of powers of  $u$ . Since the scalar can be absorbed into one of the bases, an  $\mathcal{O}$ -module of rank 1 is determined, up to isomorphism, by a power of  $u$ . It is one of the twisting modules that will be described in Section 6.6.

The Birkhoff-Grothendieck Theorem, which will be proved in Chapter 8, describes the  $\mathcal{O}$ -modules on the projective line whose sections on  $\mathbb{U}^0$  and on  $\mathbb{U}^1$  are free, as direct sums of free  $\mathcal{O}$ -modules of rank one. This means that by changing the bases  $\mathbf{B}_i$ , one can diagonalize the matrix  $P$ . Such a change of basis is given by an invertible  $A_0$ -matrix  $Q_0$  and an invertible  $A_1$ -matrix  $Q_1$ , respectively. In down-to-Earth terms, the Birkhoff-Grothendieck Theorem asserts that, for any invertible  $A_{01}$ -matrix  $P$ , there exist an invertible  $A_0$ -matrix  $Q_0$  and an invertible  $A_1$ -matrix  $Q_1$ , such that  $Q_0^{-1}PQ_1$  is diagonal. This can be proved by matrix operations.  $\square$

#### 6.4.9. tensor products

Tensor products are compatible with localization. If  $M$  and  $N$  are modules over a domain  $A$  and  $s$  is a nonzero element of  $A$ , the canonical map  $(M \otimes_A N)_s \rightarrow M_s \otimes_{A_s} N_s$  is an isomorphism. Therefore the tensor product  $\mathcal{M} \otimes_{\mathcal{O}} \mathcal{N}$  of  $\mathcal{O}$ -modules  $\mathcal{M}$  and  $\mathcal{N}$  can be defined.

Let  $\mathcal{M}$  and  $\mathcal{N}$  be  $\mathcal{O}$ -modules, let  $\mathcal{M} \otimes_{\mathcal{O}} \mathcal{N}$  be the tensor product module, and let  $V$  be an open subset of  $X$ . For every open set  $V$ , there is a canonical map

$$(6.4.10) \quad \mathcal{M}(V) \otimes_{\mathcal{O}(V)} \mathcal{N}(V) \rightarrow [\mathcal{M} \otimes_{\mathcal{O}} \mathcal{N}](V)$$

By definition of the tensor product module, this map is an equality when  $V$  is affine. For arbitrary  $V$ , we cover by a family  $\mathbb{U}_0$  of affine open sets. The family  $\mathbb{U}_1$  of intersections also consists of affine open sets. We form a diagram

$$\begin{array}{ccccccc} \mathcal{M}(V) \otimes_{\mathcal{O}(V)} \mathcal{N}(V) & \longrightarrow & \mathcal{M}(\mathbb{U}_0) \otimes_{\mathcal{O}(\mathbb{U}_0)} \mathcal{N}(\mathbb{U}_0) & \longrightarrow & \mathcal{M}(\mathbb{U}_1) \otimes_{\mathcal{O}(\mathbb{U}_1)} \mathcal{N}(\mathbb{U}_1) \\ \downarrow a & & \downarrow b & & \downarrow c \\ 0 \longrightarrow & [\mathcal{M} \otimes_{\mathcal{O}} \mathcal{N}](V) & \longrightarrow & [\mathcal{M} \otimes_{\mathcal{O}} \mathcal{N}](\mathbb{U}_0) & \longrightarrow & [\mathcal{M} \otimes_{\mathcal{O}} \mathcal{N}](\mathbb{U}_1) \end{array}$$

The composition of the two arrows in the top row is zero, the bottom row is exact, and the vertical maps  $b$  and  $c$  are equalities. The canonical map  $a$  is induced by the diagram. It is bijective when  $V$  is affine, but when  $V$  isn't affine, it may be neither injective nor surjective.

**6.4.11. Examples. (i)** Let  $p$  and  $q$  be distinct points of the projective line  $X$ , and let  $\kappa_p$  and  $\kappa_q$  be the residue field modules on  $X$ . Then  $\kappa_p(X) = \kappa_q(X) = \mathbb{C}$ , so  $\kappa_p(X) \otimes_{\mathcal{O}(X)} \kappa_q(X) \approx \mathbb{C} \otimes_{\mathbb{C}} \mathbb{C} = \mathbb{C}$ . But  $\kappa_p \otimes_{\mathcal{O}} \kappa_q = 0$ . The canonical map (6.4.10) is the zero map. It isn't injective.

**(ii)** Let  $p$  a point of a variety  $X$ , and let  $\mathfrak{m}_p$  and  $\kappa_p$  be the maximal ideal and residue field modules at  $p$ . There is an exact sequence of  $\mathcal{O}$ -modules

$$(6.4.12) \quad 0 \rightarrow \mathfrak{m}_p \rightarrow \mathcal{O} \xrightarrow{\pi_p} \kappa_p \rightarrow 0$$

The sequence of global sections is exact too.

**(iii)** Let  $p_0$  and  $p_1$  be the points  $(1, 0)$  and  $(0, 1)$  of the projective line  $\mathbb{P}^1$ . We define a homomorphism

$$\mathfrak{m}_{p_0} \times \mathfrak{m}_{p_1} \xrightarrow{\varphi} \mathcal{O}$$

by  $\varphi(a, b) = b - a$ . On the open set  $\mathbb{U}^0$ , the inclusion  $\mathfrak{m}_{p_1} \rightarrow \mathcal{O}$  is bijective and therefore surjective. Similarly,  $\mathfrak{m}_{p_0} \rightarrow \mathcal{O}$  is surjective on  $\mathbb{U}^1$ . Therefore  $\varphi$  is surjective. The only global section of  $\mathfrak{m}_{p_0} \times \mathfrak{m}_{p_1}$  is zero, while  $\mathcal{O}$  has the nonzero global section 1. So the map  $\varphi$  isn't surjective on global sections.  $\square$

#### 6.4.13. the function field module

Let  $F$  be the function field of a variety  $X$ . The module of sections of the function field module  $\mathcal{F}$  on any nonempty open set is the field  $F$ . This is an  $\mathcal{O}$ -module. It is called a *constant  $\mathcal{O}$ -module* because the modules of sections  $\mathcal{F}(U)$  are the same for every nonempty open set  $U$ . It isn't a finite module unless  $X$  is a point.

*Tensoring with the function field module:* Let  $\mathcal{M}$  be an  $\mathcal{O}$ -module on a variety  $X$ , and let  $\mathcal{F}$  be the function field module. Then  $\mathcal{M} \otimes_{\mathcal{O}} \mathcal{F}$  is a constant  $\mathcal{O}$ -module whose sections on any affine open set  $U$  form an  $F$ -vector space (that might be zero).

#### (6.4.14) limits of $\mathcal{O}$ -modules

**6.4.15.** A *directed set*  $M_\bullet$  is a sequence of maps of sets  $M_0 \rightarrow M_1 \rightarrow M_2 \rightarrow \cdots$ . Its *limit*  $\varinjlim M_\bullet$  is the set of equivalence classes on the union  $\bigcup M_k$ , the equivalence relation being that elements  $m$  in  $M_i$  and  $m'$  in  $M_j$  are equivalent if they have the same image in  $M_n$  when  $n$  is sufficiently large. Any element of  $\varinjlim M_\bullet$  will be represented by an element of  $M_i$  for some  $i$ .

**6.4.16. Example.** Let  $R = \mathbb{C}[x]$  and let  $\mathfrak{m}$  be the maximal ideal  $xR$ . Repeated multiplication by  $x$  defines a directed set

$$R \xrightarrow{x} R \xrightarrow{x} R \xrightarrow{x} R \cdots$$

whose limit is isomorphic to the Laurent Polynomial Ring  $R[x^{-1}] = \mathbb{C}[x, x^{-1}]$ . Proving this is an exercise.  $\square$

A *directed set* of  $\mathcal{O}$ -modules on a variety  $X$  is a sequence  $\mathcal{M}_\bullet = \{\mathcal{M}_0 \rightarrow \mathcal{M}_1 \rightarrow \mathcal{M}_2 \rightarrow \cdots\}$  of homomorphisms of  $\mathcal{O}$ -modules. So, for every affine open set  $U$ , the  $\mathcal{O}(U)$ -modules  $\mathcal{M}_n(U)$  form a directed set, as defined in (6.4.15). The *direct limit*  $\varinjlim \mathcal{M}_\bullet$  is defined simply, by taking the limit for each affine open set:  $[\varinjlim \mathcal{M}_\bullet](U) = \varinjlim [\mathcal{M}_\bullet(U)]$ . This limit operation is compatible with localization, so  $\varinjlim \mathcal{M}_\bullet$  is an  $\mathcal{O}$ -module. In fact, the equality  $[\varinjlim \mathcal{M}_\bullet](U) = \varinjlim [\mathcal{M}_\bullet(U)]$  is true for every open set.

**6.4.17. Lemma.** (i) *The limit operation is exact. If  $\mathcal{M}_\bullet \rightarrow \mathcal{N}_\bullet \rightarrow \mathcal{P}_\bullet$  is an exact sequence of directed sets of  $\mathcal{O}$ -modules, the limits form an exact sequence.*

(ii) *Tensor products are compatible with limits: If  $\mathcal{N}_\bullet$  is a directed set of  $\mathcal{O}$ -modules and  $\mathcal{M}$  is another  $\mathcal{O}$ -module, then  $\varinjlim [\mathcal{M} \otimes \mathcal{N}_\bullet] \approx \mathcal{M} \otimes \varinjlim \mathcal{N}_\bullet$ .*  $\square$

**6.4.18. Proposition.** *Let  $X = \text{Spec } A$  be an affine variety. Sending an  $\mathcal{O}$ -module  $\mathcal{M}$  to the  $A$ -module  $\mathcal{M}(X)$  of its global sections defines a bijective correspondence between  $\mathcal{O}$ -modules and  $A$ -modules.*

*proof.* We must invert the functor  $\mathcal{O}(\text{modules}) \rightarrow A\text{-(modules)}$  that sends  $\mathcal{M}$  to  $\mathcal{M}(X)$ . Given an  $A$ -module  $M$ , the corresponding  $\mathcal{O}$ -module  $\mathcal{M}$  is defined as follows: Let  $U = \text{Spec } B$  be an affine open subset of  $X$ . The inclusion  $U \subset X$  corresponds to an algebra homomorphism  $A \rightarrow B$ . We define  $\mathcal{M}(U)$  to be the  $B$ -module  $B \otimes_A M$ . This gives us an  $\mathcal{O}$ -module because, when  $s$  is a nonzero element of  $B$ , then  $B_s \otimes_A M$  is the localization  $(B \otimes_A M)_s$  of  $B \otimes_A M$ .  $\square$

## 6.5 Direct Image

Let  $Y \xrightarrow{f} X$  be a morphism of varieties, and let  $\mathcal{N}$  be an  $\mathcal{O}_Y$ -module. The *direct image*  $f_*\mathcal{N}$  is an  $\mathcal{O}_X$ -module that is defined as follows: Let  $U$  be an affine open subset of  $X$ . The sections of  $f_*\mathcal{N}$  on  $U$  are the sections of  $\mathcal{N}$  on  $f^{-1}U$ .

and  $V = f^{-1}U$ , then

$$[f_*\mathcal{N}](U) = \mathcal{N}(V)$$

In particular, the direct image  $f_*\mathcal{O}_Y$  of the structure sheaf  $\mathcal{O}_Y$  is the functor

$$\mathcal{O}_Y\text{-modules} \xrightarrow{f_*} \mathcal{O}_X\text{-modules}$$

defined by  $[f_*\mathcal{O}_Y](U) = \mathcal{O}_Y(f^{-1}U)$ .

The direct image generalizes restriction of scalars in modules over rings. If  $A \xrightarrow{\varphi} B$  is an algebra homomorphism and  $N$  is a  $B$ -module, one can *restrict scalars* to make  $N$  into an  $A$ -module. Scalar multiplication by an element  $a$  of  $A$  on the restricted module  $N$  is defined to be scalar multiplication by its image  $\varphi(a)$  in  $B$ . For clarity, we may sometimes denote the given  $B$ -module by  $N_B$  and the  $A$ -module obtained by restriction of scalars by  $N_A$ . The additive groups  $N_B$  and  $N_A$  are the same.

When one replaces the algebras  $A$  and  $B$  by their spectra  $X = \text{Spec } A$  and  $Y = \text{Spec } B$ , the algebra homomorphism  $\varphi$  defines a morphism  $Y \xrightarrow{f} X$ . An  $\mathcal{O}_Y$ -module  $\mathcal{N}$  is determined by a  $B$ -module  $N_B$ . Then  $f_*\mathcal{N}$  is the  $\mathcal{O}_X$ -module determined by the  $A$ -module  $N_A$ .

**6.5.1. Lemma.** *Let  $Y \xrightarrow{f} X$  be a morphism of varieties. The direct image  $f_*\mathcal{N}$  of an  $\mathcal{O}_Y$ -module  $\mathcal{N}$  is an  $\mathcal{O}_X$ -module. Moreover, for all open subsets  $U$  of  $X$ , not only for affine open subsets,  $[f_*\mathcal{N}](U) = \mathcal{N}(f^{-1}U)$ .*



*proof.* Let  $U' \rightarrow U$  be an inclusion of affine open subsets of  $X$ , and let  $V = f^{-1}U$  and  $V' = f^{-1}U'$ . These inverse images are open subsets of  $X$ , but they aren't necessarily affine. The inclusion  $V' \rightarrow V$  gives us a homomorphism  $\mathcal{N}(V) \rightarrow \mathcal{N}(V')$ , and therefore a homomorphism  $f_*\mathcal{N}(U) \rightarrow f_*\mathcal{N}(U')$ . So  $f_*\mathcal{N}$  is a functor. Its  $\mathcal{O}_X$ -module structure is explained as follows: Composition with  $f$  defines a homomorphism  $\mathcal{O}_X(U) \rightarrow \mathcal{O}_Y(V)$ , and  $\mathcal{N}(V)$  is an  $\mathcal{O}_Y(V)$ -module. Restriction of scalars makes  $[f_*\mathcal{N}](U) = \mathcal{N}(V)$  into a module over  $\mathcal{O}_X(U)$ .

To show that  $f_*\mathcal{N}$  is an  $\mathcal{O}_X$ -module, we must show that if  $s$  is a nonzero element of  $\mathcal{O}_X(U)$ , then  $[f_*\mathcal{N}](U_s)$  is obtained by localizing  $[f_*\mathcal{N}](U)$ . Let  $s'$  be the image of  $s$  in  $\mathcal{O}_Y(V)$ . Scalar multiplication by  $s$  on  $[f_*\mathcal{N}](U)$  is given by restriction of scalars, so it is the same as scalar multiplication by  $s'$  on  $\mathcal{N}(V)$ . If  $s' \neq 0$ , the localization  $V_{s'}$  is the inverse image of  $U_s$ . So  $[f_*\mathcal{N}](U_s) = \mathcal{N}(V_{s'})$ . The coherence property (6.3.16) tells us that  $\mathcal{N}(V_{s'}) = \mathcal{N}(V)_{s'}$ . Then  $[f_*\mathcal{N}](U_s) = \mathcal{N}(V)_{s'} = [[f_*\mathcal{N}](U)]_s$ .

If  $s' = 0$ , then  $\mathcal{N}(V)_{s'} = 0$ . In this case, because scalar multiplication is defined by restricting scalars,  $s$  annihilates  $[f_*\mathcal{N}](U)$ , and therefore  $[f_*\mathcal{N}](U)_s = 0$  too.  $\square$

**6.5.2. Lemma.** *Direct images are compatible with limits: If  $\mathcal{M}_\bullet$  is a directed set of  $\mathcal{O}$ -modules, then  $\varinjlim (f_*\mathcal{M}_\bullet) \approx f_*(\varinjlim \mathcal{M}_\bullet)$ .*  $\square$

### (6.5.3) extension by zero

When  $Y \xrightarrow{i} X$  is the inclusion of a **closed** subvariety into a variety  $X$ , the direct image  $i_*\mathcal{N}$  of an  $\mathcal{O}_Y$ -module  $\mathcal{N}$  is also called the *extension of  $\mathcal{N}$  by zero*. If  $U$  is an open subset of  $X$  then, because  $i$  is an inclusion map,  $i^{-1}U = U \cap Y$ . Therefore

$$[i_*\mathcal{N}](U) = \mathcal{N}(U \cap Y)$$

The term “extension by zero” refers to the fact that, when an open set  $U$  of  $X$  doesn't meet  $Y$ , the intersection  $U \cap Y$  will be empty, and the module of sections of  $[i_*\mathcal{N}](U)$  will be zero. One may say that  $i_*\mathcal{N}$  is zero outside of the closed set  $Y$ .

### 6.5.4. Examples.

(i) Let  $p \xrightarrow{i} X$  be the inclusion of a point into a variety. We may view the residue field  $k(p)$  as an  $\mathcal{O}$ -module on  $p$ . Its extension by zero  $i_*k(p)$  is the residue field module  $\kappa_p$ .

(ii) Let  $Y \xrightarrow{i} X$  be the inclusion of a closed subvariety, and let  $\mathcal{I}$  be the ideal of  $Y$ . The extension by zero of the structure sheaf on  $Y$  fits into an exact sequence of  $\mathcal{O}_X$ -modules

$$0 \rightarrow \mathcal{I} \rightarrow \mathcal{O}_X \rightarrow i_*\mathcal{O}_Y \rightarrow 0$$

So the extension by zero  $i_*\mathcal{O}_Y$  is isomorphic to the quotient module  $\mathcal{O}_X/\mathcal{I}$ .  $\square$

**6.5.5. Proposition.** *Let  $Y \xrightarrow{i} X$  be the inclusion of a closed subvariety  $Y$  into a variety  $X$ , and let  $\mathcal{I}$  be the ideal of  $Y$ . Let  $\mathbb{M}$  denote the category of  $\mathcal{O}_X$ -modules that are annihilated by  $\mathcal{I}$ . Extension by zero defines an equivalence of categories*

$$(\mathcal{O}_Y\text{-modules}) \xrightarrow{i_*} \mathbb{M}$$

*proof.* Let  $f$  be a section of  $\mathcal{O}_X$  on an affine open set  $U$ , let  $\bar{f}$  be its restriction to  $U \cap Y$ , and let  $\alpha$  be an element of  $[i_*\mathcal{N}](U)$  ( $= \mathcal{N}(U \cap Y)$ ). If  $f$  is in  $\mathcal{I}(U)$ , then  $\bar{f} = 0$  and therefore  $f\alpha = \bar{f}\alpha = 0$ . So the extension by zero of an  $\mathcal{O}_Y$ -module is annihilated by  $\mathcal{I}$ . The direct image  $i_*\mathcal{N}$  is an object of  $\mathbb{M}$ .

We construct a quasi-inverse to the direct image. Starting with an  $\mathcal{O}_X$ -module  $\mathcal{M}$  that is annihilated by  $\mathcal{I}$ , we construct an  $\mathcal{O}_Y$ -module  $\mathcal{N}$  such that  $i_*\mathcal{N}$  is isomorphic to  $\mathcal{M}$ .

Let  $Y'$  be an open subset of  $Y$ . The topology on  $Y$  is induced from the topology on  $X$ , so  $Y' = X_1 \cap Y$  for some open subset  $X_1$  of  $X$ . We try to set  $\mathcal{N}(Y') = \mathcal{M}(X_1)$ . To show that this is well-defined, we show that if  $X_2$  is another open subset of  $X$ , and if  $Y' = X_2 \cap Y$ , then  $\mathcal{M}(X_2)$  is isomorphic to  $\mathcal{M}(X_1)$ . Let  $X_3 = X_1 \cap X_2$ . Then it is also true that  $Y' = X_3 \cap Y$ . Since  $X_3 \subset X_1$ , we have a map  $\mathcal{M}(X_1) \rightarrow \mathcal{M}(X_3)$ ,

and It suffices to show that this map is an isomorphism. The same reasoning will give us an isomorphism  $\mathcal{M}(X_2) \rightarrow \mathcal{M}(X_3)$ .

The complement  $U = X_1 - Y'$  of  $Y'$  in  $X_1$  is an open subset of  $X_1$  and of  $X$ , and  $U \cap Y = \emptyset$ . We cover  $U$  by a set  $\{U^i\}$  of affine open sets. Then  $X_1$  is covered by the open sets  $\{U^i\}$  together with  $X_3$ . The restriction of  $\mathcal{I}$  to each of the sets  $U^i$  is the unit ideal, and since  $\mathcal{I}$  annihilates  $\mathcal{M}$ ,  $\mathcal{M}(U^i) = 0$ . The sheaf property shows that  $\mathcal{M}(X_1)$  is isomorphic to  $\mathcal{M}(X_3)$ . The rest of the proof is boring.  $\square$

### (6.5.6) inclusion of an open set

Let  $Y \xrightarrow{j} X$  be the inclusion of an **open** subvariety  $Y$  into a variety  $X$ .

First, let  $\mathcal{M}$  be an  $\mathcal{O}_X$ -module. Since open subsets of  $Y$  are also open subsets of  $X$ , we can *restrict*  $\mathcal{M}$  from  $X$  to  $Y$ . By definition, the sections of the restricted module on a subset  $U$  of  $Y$  are simply the elements of  $\mathcal{M}(U)$ . For example, the restriction of the structure sheaf  $\mathcal{O}_X$  is the structure sheaf  $\mathcal{O}_Y$  on  $Y$ . We extend the subscript notation to  $\mathcal{O}$ -modules, writing  $\mathcal{M}_Y$  for the restriction of an  $\mathcal{O}_X$ -module  $\mathcal{M}$  to  $Y$  and denoting the given module  $\mathcal{M}$  by  $\mathcal{M}_X$ . Then if  $U$  is an open subset of  $Y$ ,

$$\mathcal{M}_X(U) = \mathcal{M}_Y(U)$$

Now the direct image: Let  $Y \xrightarrow{j} X$  be the inclusion of an open subvariety  $Y$ , and let  $\mathcal{N}$  be an  $\mathcal{O}_Y$ -module. The inverse image of an open subset  $U$  of  $X$  is the intersection  $Y \cap U$ , so

$$[j_*\mathcal{N}](U) = \mathcal{N}(Y \cap U)$$

For example,  $[j_*\mathcal{O}_Y](U)$  is the algebra of rational functions on  $X$  that are regular on  $Y \cap U$ . They needn't be regular on  $U$ .

**6.5.7. Example.** Let  $X_s \xrightarrow{j} X$  be the inclusion of a localization into an affine variety  $X = \text{Spec } A$ . Modules on  $X$  correspond to their global sections, which are  $A$ -modules. Similarly, modules on  $X_s$  correspond to  $A_s$ -modules. We can restrict an  $\mathcal{O}_X$ -module  $\mathcal{M}_X$  to the open set  $X_s$ , obtaining an  $\mathcal{O}_{X_s}$ -module  $\mathcal{M}_{X_s}$ . Then if  $M$  denotes the  $A$ -module of global sections  $\mathcal{M}_X(X)$ , the module of sections of the restriction  $\mathcal{M}_{X_s}$  on  $X_s$  is the localization  $M_s$ , and  $M_s$  is also the module of global sections of  $j_*\mathcal{M}_{X_s}$ :

$$[j_*\mathcal{M}_{X_s}](X) = M_{x_s}(X_s) = M_s$$

The localization  $M_s$  is made into an  $A$ -module by restriction of scalars.  $\square$

**6.5.8. Proposition.** Let  $Y \xrightarrow{j} X$  be the inclusion of an open subvariety  $Y$  into a variety  $X$ .

- (i) The restriction  $\mathcal{O}_X\text{-modules} \rightarrow \mathcal{O}_Y\text{-modules}$  is an exact operation.
- (ii) If  $Y$  is an **affine** open subvariety of  $X$ , the direct image functor  $j_*$  is exact.
- (iii) Let  $\mathcal{M}_X$  be an  $\mathcal{O}_X$ -module. There is a canonical homomorphism  $\mathcal{M}_X \rightarrow j_*[\mathcal{M}_Y]$ .

*proof.* (ii) Let  $U$  be an affine open subset of  $X$ , and let  $\mathcal{M} \rightarrow \mathcal{N} \rightarrow \mathcal{P}$  be an exact sequence of  $\mathcal{O}_Y$ -modules. The sequence  $j_*\mathcal{M}(U) \rightarrow j_*\mathcal{N}(U) \rightarrow j_*\mathcal{P}(U)$  is the same as the sequence  $\mathcal{M}(U \cap Y) \rightarrow \mathcal{N}(U \cap Y) \rightarrow \mathcal{P}(U \cap Y)$ , except that the scalars have changed. Since  $U$  and  $Y$  are affine,  $U \cap Y$  is affine. By definition of exactness, this last sequence is exact.

(iii) Let  $U$  be open in  $X$ . Then  $j_*\mathcal{M}_Y(U) = \mathcal{M}_X(U \cap Y)$ . Since  $U \cap Y \subset U$ ,  $\mathcal{M}_X(U)$  maps to  $\mathcal{M}_X(U \cap Y)$ .  $\square$

**6.5.9. Example.** Let  $X = \mathbb{P}^n$  and let  $j$  denote the inclusion  $\mathbb{U}^0 \subset X$  of the standard affine open subset into  $X$ . The direct image  $j_*\mathcal{O}_{\mathbb{U}^0}$  is the algebra of rational functions that are allowed to have poles on the hyperplane at infinity.

The inverse image of an open subset  $W$  of  $X$  is its intersection with  $\mathbb{U}^0$ :  $j^{-1}W = W \cap \mathbb{U}^0$ . The sections of the direct image  $j_*\mathcal{O}_{\mathbb{U}^0}$  on an open subset  $W$  of  $X$  are the regular functions on  $W \cap \mathbb{U}^0$ :

$$[j_*\mathcal{O}_{\mathbb{U}^0}](W) = \mathcal{O}_{\mathbb{U}^0}(W \cap \mathbb{U}^0) = \mathcal{O}_X(W \cap \mathbb{U}^0)$$

Say that we write a rational function  $\alpha$  as a fraction  $g/h$  of relatively prime polynomials. Then  $\alpha$  is an element of  $\mathcal{O}_X(W)$  if  $h$  doesn't vanish at any point of  $W$ , and  $\alpha$  is a section of  $[j_*\mathcal{O}_{\mathbb{U}^0}](W) = \mathcal{O}_X(W \cap \mathbb{U}^0)$  if  $h$  doesn't vanish on  $W \cap \mathbb{U}^0$ . Arbitrary powers of  $x_0$  can appear in the denominator  $h$ .  $\square$

## 6.6 Twisting

The twisting modules that we define here are among the most important modules on projective space.

Let  $X$  denote the projective space  $\mathbb{P}^n$  with coordinates  $x_0, \dots, x_n$ . As before, a homogeneous fraction of degree  $d$  is a fraction  $g/h$  of homogeneous polynomials with  $\deg g - \deg h = d$ . When  $g$  and  $h$  are relatively prime, the fraction  $g/h$  is *regular* on an open set  $V$  if  $h$  isn't zero at any point of  $V$ .

The definition of the twisting module is this: The sections of  $\mathcal{O}(d)$  on an open subset  $V$  of  $\mathbb{P}^n$  are the homogeneous fractions of degree  $d$  that are regular on  $V$ .

### 6.6.1. Proposition.

(i) Let  $V$  be an affine open subset of  $\mathbb{P}^n$  that is contained in the standard affine open set  $\mathbb{U}^0$ . The sections of  $\mathcal{O}(d)$  on  $V$  form a free module of rank one with basis  $x_0^d$ , over the coordinate algebra  $\mathcal{O}(V)$ .

(ii) The twisting module  $\mathcal{O}(d)$  is an  $\mathcal{O}$ -module.

*proof.* (i) Let  $\alpha$  be a section of  $\mathcal{O}(d)$  on an affine open set  $V$  that is contained in  $\mathbb{U}^0$ . Then  $f = \alpha x_0^{-d}$  has degree zero. It is a rational function. Since  $V \subset \mathbb{U}^0$ ,  $x_0$  doesn't vanish at any point of  $V$ . Since  $\alpha$  is regular on  $V$ ,  $f$  is a regular function on  $V$ , and  $\alpha = f x_0^d$ .

(ii) It is clear that  $\mathcal{O}(d)$  is a contravariant functor. We verify compatibility with localization. Let  $V = \text{Spec } A$  be an affine open subset of  $X$  and let  $s$  be a nonzero element of  $A$ . We must show that  $[\mathcal{O}(d)](V_s)$  is the localization of  $[\mathcal{O}(d)](V)$ , and it is true that  $[\mathcal{O}(d)](V)$  is a subset of  $[\mathcal{O}(d)](V_s)$ . What has to be shown is that if  $\beta$  is a section of  $\mathcal{O}(d)$  on  $V_s$ , then  $s^k \beta$  is a section on  $V$ , if  $k$  is sufficiently large.

We cover  $V$  by the affine open sets  $V^i = V \cap \mathbb{U}^i$ . To show that  $s^k \beta$  is a section on  $V$ , it suffices to show that it is a section on  $V \cap \mathbb{U}^i$  for every  $i$ . This is the sheaf property. We apply (i) to the open subset  $V_s^0$  of  $V^0$ . Since  $V_s^0$  is contained in  $\mathbb{U}^0$ ,  $\beta$  can be written (uniquely) in the form  $f x_0^d$ , where  $f$  is a rational function that is regular on  $V_s^0$ . We know already that  $\mathcal{O}$  has the localization property. Therefore  $s^k f$  is a regular function on  $V^0$  if  $k$  is large, and then  $s^k \alpha = s^k f x_0^d$  is a section of  $\mathcal{O}(d)$  on  $V^0$ . The analogous statement is true for every index  $i$ .  $\square$

Part (i) of the proposition shows that  $\mathcal{O}(d)$  is quite similar to the structure sheaf. However,  $\mathcal{O}(d)$  is only *locally* free. Its sections on the standard open set  $\mathbb{U}^1$  form a free  $\mathcal{O}(\mathbb{U}^1)$ -module with basis  $x_1^d$ . That basis is related to the basis  $x_0^d$  on  $\mathbb{U}^0$  by the factor  $(x_0/x_1)^d$ , a rational function that isn't invertible on  $\mathbb{U}^0$  or on  $\mathbb{U}^1$ .

**6.6.2. Proposition.** When  $d \geq 0$ , the global sections of the twisting module  $\mathcal{O}(d)$  on  $\mathbb{P}^n$  ( $n > 0$ ) are the homogeneous polynomials of degree  $d$ . When  $d < 0$ , the only global section of  $\mathcal{O}(d)$  is zero.

*proof.* A nonzero global section  $u$  of  $\mathcal{O}(d)$  will restrict to a section on the standard affine open set  $\mathbb{U}^0$ . Since  $[\mathcal{O}(d)](\mathbb{U}^0)$  is a free module over  $\mathcal{O}(\mathbb{U}^0)$  with basis  $x_0^d$ , and  $u = g/x_0^m$  for some homogeneous polynomial  $g$  not divisible by  $x_0$  and some  $m$ . Similarly, restriction to  $\mathbb{U}^1$  shows that  $u = h/x_1^n$ . It follows that  $m = n = 0$  and that  $u = g$ . Since  $u$  has degree  $d$ ,  $g$  will be a polynomial of degree  $d$ .  $\square$

### 6.6.3. Examples.

The product  $uv$  of homogeneous fractions of degrees  $r$  and  $s$  is a homogeneous fraction of degree  $r+s$ , and if  $u$  and  $v$  are regular on an open set  $V$ , so is their product  $uv$ . Therefore multiplication defines a homomorphism of  $\mathcal{O}$ -modules

$$(6.6.4) \quad \mathcal{O}(r) \times \mathcal{O}(s) \rightarrow \mathcal{O}(r+s)$$

Multiplication by a homogeneous polynomial  $f$  of degree  $d$  defines an injective homomorphism

$$(6.6.5) \quad \mathcal{O}(k) \xrightarrow{f} \mathcal{O}(k+d).$$

When  $k = -d$ , this becomes a homomorphism  $\mathcal{O}(-d) \xrightarrow{f} \mathcal{O}$ .  $\square$

The twisting modules  $\mathcal{O}(n)$  have a second interpretation. They are isomorphic to the modules that we denote by  $\mathcal{O}(nH)$ , of rational functions on projective space with poles of order at most  $n$  on the hyperplane  $H : \{x_0 = 0\}$  at infinity.

By definition, the nonzero sections of  $\mathcal{O}(nH)$  on an open set  $V$  are the rational functions  $f$  such that  $x_0^n f$  is a section of  $\mathcal{O}(n)$  on  $V$ . Thus multiplication by  $x_0^n$  defines an isomorphism

$$(6.6.6) \quad \mathcal{O}(nH) \xrightarrow{x_0^n} \mathcal{O}(n)$$

If  $f$  is a section of  $\mathcal{O}(nH)$  on an open set  $V$ , and if we write  $f$  as a homogeneous fraction  $g/h$  of degree zero, with  $g, h$  relatively prime, the denominator  $h$  may have  $x_0^k$ , with  $k \leq n$ , as factor. The other factors of  $h$  cannot vanish anywhere on  $V$ . If  $f = g/h$  is a global section of  $\mathcal{O}(nH)$ , then  $h = cx_0^k$ , with  $k \leq n$ , so a global section can be represented as a fraction  $g/x_0^k$ .

Since  $x_0$  doesn't vanish at any point of the standard affine open set  $\mathbb{U}^0$ , the sections of  $\mathcal{O}(nH)$  on an open subset  $V$  of  $\mathbb{U}^0$  are simply the regular functions on  $V$ . The restrictions of  $\mathcal{O}(nH)$  and  $\mathcal{O}$  to  $\mathbb{U}^0$  are equal. Using the subscript notation (6.5.6) for restriction to an open set,

$$(6.6.7) \quad \mathcal{O}(nH)_{\mathbb{U}^0} = \mathcal{O}_{\mathbb{U}^0}$$

Let  $V$  be an open subset of one of the other standard affine open sets, say of  $\mathbb{U}^1$ . The ideal of  $H \cap \mathbb{U}^1$  in  $\mathbb{U}^1$  is principal, generated by  $v_0 = x_0/x_1$ , and  $v_0$  generates the ideal of  $H \cap V$  in  $V$  too. If  $f$  is a rational function, then because  $x_1$  doesn't vanish on  $\mathbb{U}^1$ , the function  $fv_0^n$  will be regular on  $V$  if and only if the homogeneous fraction  $fx_0^n$  is regular there. So  $f$  will be a section of  $\mathcal{O}(nH)$  on  $V$  if and only if  $fv_0^n$  is a regular function. Because  $v_0$  generates the ideal of  $H$  in  $V$ , we say that such a function  $f$  has a pole of order at most  $n$  on  $H$ .

The isomorphic  $\mathcal{O}$ -modules  $\mathcal{O}(n)$  and  $\mathcal{O}(nH)$  are interchangeable. The twisting module  $\mathcal{O}(n)$  is often better because its definition is independent of coordinates. On the other hand,  $\mathcal{O}(nH)$  can be convenient because its restriction to  $\mathbb{U}^0$  is the structure sheaf  $\mathcal{O}_{\mathbb{U}^0}$ .

**6.6.8. Proposition.** *Let  $Y$  be the zero locus of an irreducible homogeneous polynomial  $f$  of degree  $d$ , a hypersurface of degree  $d$  in  $\mathbb{P}^n$ , let  $\mathcal{I}$  be the ideal of  $Y$ , and let  $\mathcal{O}(-d)$  be the twisting module on  $X$ . Multiplication by  $f$  defines an isomorphism  $\mathcal{O}(-d) \xrightarrow{f} \mathcal{I}$ .*

*proof.* If  $\alpha$  is a section of  $\mathcal{O}(-d)$  on an open set  $V$ , then  $f\alpha$  will be a rational function that is regular on  $V$  and vanishes on  $Y$ . Therefore the image of the multiplication map  $\mathcal{O}(-d) \xrightarrow{f} \mathcal{O}$  is contained in  $\mathcal{I}$ . This map is injective because  $\mathbb{C}[x_0, \dots, x_n]$  is a domain. To show that it is an isomorphism, it suffices to show that its restrictions to the standard affine open sets  $\mathbb{U}^i$  are isomorphisms (6.3.5). As usual, we work with  $\mathbb{U}^0$ .

We choose coordinates in  $X$  so that the coordinate variables  $x_i$  don't divide  $f$ . Then  $Y \cap \mathbb{U}^0$  will be a dense open subset of  $Y$ . The sections of  $\mathcal{O}$  on  $\mathbb{U}^0$  are the homogeneous fractions  $g/x_0^k$  of degree zero. Such a fraction is a section of  $\mathcal{I}$  on  $\mathbb{U}^0$  if and only if  $g$  vanishes on  $Y \cap \mathbb{U}^0$ . If so, then since  $Y \cap \mathbb{U}^0$  is dense in  $Y$ , it will vanish on  $Y$ , and therefore it will be divisible by  $f$ :  $g = fq$ . The sections of  $\mathcal{I}$  on  $\mathbb{U}^0$  have the form  $fq/x_0^k$ . They are in the image of  $\mathcal{O}(-d)$ .  $\square$

The proposition has an interesting corollary:

**6.6.9. Corollary.** *The ideals of all hypersurfaces of degree  $d$  are isomorphic, when they are regarded as  $\mathcal{O}$ -modules.*  $\square$

### (6.6.10) twisting a module

**6.6.11. Definition** Let  $\mathcal{M}$  be an  $\mathcal{O}$ -module on projective space  $\mathbb{P}^d$ , and let  $\mathcal{O}(n)$  be the twisting module. The  $n$ th twist of  $\mathcal{M}$  is defined to be the tensor product  $\mathcal{M}(n) = \mathcal{M} \otimes_{\mathcal{O}} \mathcal{O}(n)$ , and similarly,  $\mathcal{M}(nH) = \mathcal{M} \otimes_{\mathcal{O}} \mathcal{O}(nH)$ . Twisting is a functor on  $\mathcal{O}$ -modules.

If  $X$  is a closed subvariety of  $\mathbb{P}^d$  and  $\mathcal{M}$  is an  $\mathcal{O}_X$ -module,  $\mathcal{M}(n)$  and  $\mathcal{M}(nH)$  are obtained by twisting the extension of  $\mathcal{M}$  by zero. (See the equivalence of categories (6.5.5)).

A section of  $\mathcal{M}(n)$  on an open subset  $V$  of  $\mathbb{U}^0$  can be written in the form  $s = m \otimes fx_0^n$ , where  $f$  is a regular function on  $V$  and  $m$  is a section of  $\mathcal{M}$  on  $V$  (6.6.1). The function  $f$  can be moved over to  $m$ , so a section can be written in the form  $s = m \otimes x_0^n$ . This expression is unique.

**6.6.12.** The modules  $\mathcal{O}(n)$  and  $\mathcal{O}(nH)$  form directed sets that are related by a diagram

$$(6.6.13) \quad \begin{array}{ccccccc} \mathcal{O} & \xrightarrow{\subset} & \mathcal{O}(H) & \xrightarrow{\subset} & \mathcal{O}(2H) & \xrightarrow{\subset} & \dots \\ \parallel & & x_0 \downarrow & & x_0^2 \downarrow & & \\ \mathcal{O} & \xrightarrow{x_0} & \mathcal{O}(1) & \xrightarrow{x_0} & \mathcal{O}(2) & \longrightarrow & \dots \end{array}$$

In this diagram, the vertical arrows are bijections. The limit of the upper directed set is the module whose sections are allowed to have arbitrary poles on  $H$ . This is also the module  $j_*\mathcal{O}_{\mathbb{U}}$ , where  $j$  denotes the inclusion of the standard affine open set  $\mathbb{U} = \mathbb{U}^0$  into  $X$  (see (6.5.8) (iii)):

$$(6.6.14) \quad \varinjlim \mathcal{O}(nH) = j_*\mathcal{O}_{\mathbb{U}}$$

The next diagram is obtained by tensoring Diagram 6.6.13 with  $\mathcal{M}$ .

$$(6.6.15) \quad \begin{array}{ccccccc} \mathcal{M} & \longrightarrow & \mathcal{M}(H) & \longrightarrow & \mathcal{M}(2H) & \longrightarrow & \dots \\ \parallel & & x_0 \downarrow & & x_0^2 \downarrow & & \\ \mathcal{M} & \xrightarrow{1 \otimes x_0} & \mathcal{M}(1) & \xrightarrow{1 \otimes x_0} & \mathcal{M}(2) & \longrightarrow & \dots \end{array}$$

Because  $\mathcal{M}$  may have torsion, the horizontal maps in these two directed sets needn't be injective. However, since tensor products are compatible with limits,

$$(6.6.16) \quad \varinjlim \mathcal{M}(nH) = \varinjlim \mathcal{M} \otimes_{\mathcal{O}} \mathcal{O}(nH) = \mathcal{M} \otimes_{\mathcal{O}} j_*\mathcal{O} = j_*\mathcal{M}_{\mathbb{U}}$$

### (6.6.17) generating an $\mathcal{O}$ -module

A set  $m = (m_1, \dots, m_k)$  of global sections of an  $\mathcal{O}$ -module  $\mathcal{M}$  defines a map

$$(6.6.18) \quad \mathcal{O}^k \xrightarrow{m} \mathcal{M}$$

that sends a section  $(\alpha_1, \dots, \alpha_k)$  of  $\mathcal{O}^k$  on an open set to the combination  $\sum \alpha_i m_i$ . The set of global sections  $\{m_1, \dots, m_k\}$  *generates*  $\mathcal{M}$  if this map is surjective. If the sections generate  $\mathcal{M}$ , then they (more precisely, their restrictions to  $U$ ) generate the  $\mathcal{O}(U)$ -module  $\mathcal{M}(U)$  for every affine open set  $U$ . They may fail to generate  $\mathcal{M}(U)$  when  $U$  isn't affine.

**6.6.19. Example.** Let  $X = \mathbb{P}^1$ . For  $n \geq 0$ , the global sections of the twisting module  $\mathcal{O}(n)$  are the polynomials of degree  $n$  in the coordinate variables  $x_0, x_1$  (6.6.2). Consider the map  $\mathcal{O}^2 \xrightarrow{(x_0^n, x_1^n)} \mathcal{O}(n)$ . On  $\mathbb{U}^0$ ,  $\mathcal{O}(n)$  has basis  $x_0^n$ . Therefore this map is surjective on  $\mathbb{U}^0$ . Similarly, it is surjective on  $\mathbb{U}^1$ . So it is a surjective map on all of  $X$  (6.3.5). The global sections  $x_0^n, x_1^n$  generate  $\mathcal{O}(n)$ . However, the global sections of  $\mathcal{O}(n)$  are the homogeneous polynomials of degree  $n$ . When  $n > 1$ , the two sections  $x_0^n, x_1^n$  don't span the space of global sections, and the map  $\mathcal{O}^2 \xrightarrow{(x_0^n, x_1^n)} \mathcal{O}(n)$  isn't surjective.  $\square$

The next theorem explains the importance of the twisting operation.

**6.6.20. Theorem.** *Let  $\mathcal{M}$  be a finite  $\mathcal{O}$ -module on a projective variety  $X$ . For large  $k$ , the twist  $\mathcal{M}(k)$  is generated by global sections.*

*proof.* We may assume that  $X$  is projective space  $\mathbb{P}^n$ .

We are to show that if  $\mathcal{M}$  is a finite  $\mathcal{O}$ -module, the global sections generate  $\mathcal{M}(k)$  when  $k$  is large, and it suffices to show that for each  $i = 0, \dots, n$ , the restrictions of those global sections to  $\mathbb{U}^i$  generate the  $\mathcal{O}(\mathbb{U}^i)$ -module  $[\mathcal{M}(k)](\mathbb{U}^i)$  (6.3.5). We work with the index 0 as usual.

Let  $\mathbb{U} = \mathbb{U}^0$ . We replace  $\mathcal{M}(k)$  by the isomorphic module  $\mathcal{M}(kH)$ . We recall that the restrictions of  $\mathcal{M}(k)$  to  $\mathbb{U}$  are equal to  $\mathcal{M}_{\mathbb{U}}$ , and that  $\varinjlim \mathcal{M}(k) = j_*\mathcal{M}_{\mathbb{U}}$ .

Let  $A_0 = \mathcal{O}(\mathbb{U})$ , and  $M_0 = \mathcal{M}(\mathbb{U})$ .

The global sections of  $j_*\mathcal{M}_{\mathbb{U}}$  are the sections of  $\mathcal{M}$  on  $\mathbb{U}$ , which we are denoting by  $M_0$ . We choose a finite set of generators  $m_1, \dots, m_r$  for the finite  $A_0$ -module  $M_0$ . These generators are global sections of  $j_*\mathcal{M}_{\mathbb{U}}$ , and therefore they are represented by global sections  $m'_1, \dots, m'_r$  of  $\mathcal{M}(k)$  if  $k$  is large. The restrictions of  $m_i$  and  $m'_i$  to  $\mathbb{U}$  are equal, so the restrictions of  $m'_1, \dots, m'_r$  generate  $M_0$  too, and  $M_0$  is generated by global sections of  $\mathcal{M}(k)$ , as was to be shown.  $\square$

## 6.7 Proof of Theorem 6.3.2.

The statement to be proved is that an  $\mathcal{O}$ -module  $\mathcal{M}$  on a variety  $X$  has a unique extension to a functor

$$(\text{opens}) \xrightarrow{\widetilde{\mathcal{M}}} (\text{modules})$$

having the sheaf property, and that a homomorphism  $\mathcal{M} \rightarrow \mathcal{N}$  of  $\mathcal{O}$ -modules has a unique extension to a homomorphism  $\widetilde{\mathcal{M}} \rightarrow \widetilde{\mathcal{N}}$ .

The proof has the following steps:

1. Verification of the sheaf property for a covering of an affine open set by localizations.
2. Extension of the functor  $\mathcal{M}$  to all morphisms between affine open sets.
3. Definition of  $\widetilde{\mathcal{M}}$ .

**Step 1.** (*the sheaf property for a covering of an affine open set by localizations*)

Suppose that an affine open set  $Y = \text{Spec } A$  is covered by a family of localizations  $\mathbb{U}_0 = \{U^{s_i}\}$ . Then if  $\mathcal{M}$  is an  $\mathcal{O}_Y$ -module and if we let  $M = \mathcal{M}(Y)$ ,  $M_i = \mathcal{M}(U^{s_i})$  and  $M_{ij} = \mathcal{M}(U^{s_i s_j})$  the sequence (6.3.9) for the covering diagram  $Y \leftarrow \mathbb{U}_0 \rightrightarrows \mathbb{U}_1$  that we obtain is

$$(6.7.1) \quad 0 \rightarrow M \xrightarrow{\alpha} \prod M_i \xrightarrow{\beta} \prod M_{ij}$$

In this sequence, the map  $\alpha$  sends an element  $m$  of  $M$  to the vector  $(m, \dots, m)$  of its images in  $\prod_i M_i$ , and  $\beta$  sends a vector  $(m_1, \dots, m_k)$  in  $\prod_i M_i$  to the matrix  $(z_{ij})$ , with  $z_{ij} = m_j - m_i$  in  $M_{ij}$ . To be precise,  $M_i$  and  $M_j$  map to  $M_{ij}$ , and  $z_{ij}$  is the difference of their images.

We must show that the sequence (6.7.1) is exact. Since  $U^i$  cover  $Y$ , the elements  $s_1, \dots, s_k$  generate the unit ideal.

*exactness at  $M$ :* Let  $m$  be an element of  $M$  that maps to zero in every  $M_i$ . Then there exists an  $n$  such that  $s_i^n m = 0$ , and we can use the same exponent  $n$  for all  $i$ . The elements  $s_i^n$  generate the unit ideal. Writing  $\sum a_i s_i^n = 1$ , we have  $m = \sum a_i s_i^n m = \sum a_i 0 = 0$ .

*exactness at  $\prod M_i$ :* Let  $m_i$  be elements of  $M_i$  such that  $m_i = m_j$  in  $M_{ij}$  for all  $i, j$ . We must find an element  $w$  in  $M$  that maps to  $m_j$  in  $M_j$  for every  $j$ .

We write  $m_i$  as a fraction:  $m_i = s_i^{-n} x_i$ , or  $x_i = s_i^n m_i$ , with  $x_i$  in  $M$ , using the same integer  $n$  for all  $i$ . The equation  $m_i = m_j$  in  $M_{ij}$  tells us that  $s_j^n x_i = s_i^n x_j$  is true in  $M_{ij}$ , and then  $(s_i s_j)^r s_j^n x_i = (s_i s_j)^r s_i^n x_j$  will be true in  $M$ , if  $r$  is large (see 5.1.19).

We adjust the notation. Let  $\tilde{x}_i = s_i^r x_i$ , and  $\tilde{s}_i = s_i^{r+n}$ . Then in  $M$ ,  $\tilde{x}_i = \tilde{s}_i m_i$  and  $\tilde{s}_j \tilde{x}_i = \tilde{s}_i \tilde{x}_j$ . Since the elements  $s_i$  generate the unit ideal, so do their powers  $\tilde{s}_i$ . There is an equation in  $A$ , of the form  $\sum a_i \tilde{s}_i = 1$ . Let  $w = \sum a_i \tilde{x}_i$ . This is an element of  $M$ , and

$$\tilde{x}_j = \left( \sum_i a_i \tilde{s}_i \right) \tilde{x}_j = \sum_i a_i \tilde{s}_j \tilde{x}_i = \tilde{s}_j w$$

Since  $m_j = \tilde{s}_j^{-1} \tilde{x}_j$ ,  $m_j = w$  is true in  $M_j$ . Since  $j$  is arbitrary,  $w$  is the required element of  $M$ .  $\square$

**Step 2.** (*extending an  $\mathcal{O}$ -module to all morphisms between affine open sets*)

The  $\mathcal{O}$ -module  $\mathcal{M}$  comes with localization maps  $\mathcal{M}(U) \rightarrow \mathcal{M}(U_s)$ . It doesn't come with homomorphisms  $\mathcal{M}(U) \rightarrow \mathcal{M}(V)$  when  $V \rightarrow U$  is an arbitrary inclusion of affine open sets. We define those maps here.

Let  $\mathcal{M}$  be an  $\mathcal{O}$ -module and let  $V \rightarrow U$  be an inclusion of affine open sets. To describe the canonical homomorphism  $\mathcal{M}(U) \rightarrow \mathcal{M}(V)$ , we cover  $V$  by a family  $\mathbb{V}_0 = \{V^i\}$  of open sets that are localizations of  $U$  and therefore also localizations of  $V$ , and we inspect the covering diagram  $V \leftarrow \mathbb{V}_0 \rightrightarrows \mathbb{V}_1$  and the corresponding exact sequence  $0 \rightarrow \mathcal{M}(V) \xrightarrow{\alpha} \mathcal{M}(\mathbb{V}_0) \xrightarrow{\beta} \mathcal{M}(\mathbb{V}_1)$ . The two maps  $\mathbb{V}_1 \rightarrow U$  obtained by composition from the maps

$$U \leftarrow V \leftarrow \mathbb{V}_0 \rightrightarrows \mathbb{V}_1$$

are equal. Since  $V^i$  are localizations of  $U$  and  $V^{ij}$  are localizations of  $V^i$  and of  $V^j$ , the  $\mathcal{O}$ -module  $\mathcal{M}$  comes with maps  $\mathcal{M}(U) \xrightarrow{\psi} \mathcal{M}(\mathbb{V}_0) \rightrightarrows \mathcal{M}(\mathbb{V}_1)$ . The two composed maps  $\mathcal{M}(U) \rightarrow \mathcal{M}(\mathbb{V}_1)$  are equal, so their difference  $\beta\psi$  is zero. Therefore  $\psi$  maps  $\mathcal{M}(U)$  to the kernel of  $\beta$  which, according to Step 1, is  $\mathcal{M}(V)$ . This defines a map  $\mathcal{M}(U) \xrightarrow{\eta} \mathcal{M}(V)$  making a diagram

$$\begin{array}{ccc} \mathcal{M}(U) & \xrightarrow{\eta} & \mathcal{M}(V) \\ \parallel & & \downarrow \lambda \\ \mathcal{M}(U) & \xrightarrow{\psi} & \mathcal{M}(\mathbb{V}_0) \end{array}$$

Both  $\psi$  and  $\lambda$  are compatible with multiplication by a regular function  $f$  on  $U$ , and  $\lambda$  is injective. So  $\eta$  is also compatible with multiplication by  $f$ .

We must check that  $\eta$  is independent of the covering  $\mathbb{V}_0$ . Let  $\mathbb{V}'_0 = \{V'^j\}$  be another covering of  $V$  by localizations of  $U$ . We cover each of the open sets  $V^i \cap V'^j$  by localizations  $W^{ij\nu}$  of  $U$ . Taken together, these open sets form a covering  $\mathbb{W}_0$  of  $V$ . We have a map  $\mathbb{W}_0 \xrightarrow{\epsilon} \mathbb{V}_0$  that gives us a diagram

$$\begin{array}{ccccc} V & \longleftarrow & \mathbb{V}_0 & \rightrightarrows & \mathbb{V}_1 \\ \parallel & & \downarrow & & \downarrow \\ V & \longleftarrow & \mathbb{W}_0 & \rightrightarrows & \mathbb{W}_1 \end{array}$$

and therefore a diagram

$$(6.7.2) \quad \begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{M}(V) & \longrightarrow & \mathcal{M}(\mathbb{V}_0) & \xrightarrow{\beta_V} & \mathcal{M}(\mathbb{V}_1) \\ & & \parallel & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathcal{M}(V) & \longrightarrow & \mathcal{M}(\mathbb{W}_0) & \xrightarrow{\beta_W} & \mathcal{M}(\mathbb{W}_1) \end{array}$$

whose rows are exact sequences. Here  $\mathcal{M}(U)$  maps to the kernels of  $\beta_V$  and  $\beta_W$ , both of which are equal to  $\mathcal{M}(V)$ . Looking at the diagram, one sees that the map  $\mathcal{M}(U) \rightarrow \mathcal{M}(\mathbb{W}_0)$  is the composition of the maps  $\mathcal{M}(U) \rightarrow \mathcal{M}(\mathbb{V}_0) \rightarrow \mathcal{M}(\mathbb{W}_0)$ . Therefore the two maps  $\mathcal{M}(U) \rightarrow \mathcal{M}(V)$  are equal, and they are also equal to the map defined by the covering  $\mathbb{V}'_0$ .

We show that this extended functor has the sheaf property for an affine covering  $\mathbb{U}_0 = \{U^i\}$  of an affine variety  $U$ . We let  $\mathbb{V}_0$  be the covering of  $U$  obtained by covering each  $U^i$  by localizations of  $U$ . This gives us a diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{M}(U) & \longrightarrow & \mathcal{M}(\mathbb{U}_0) & \longrightarrow & \mathcal{M}(\mathbb{U}_1) \\ \parallel & & \beta \downarrow & & \gamma \downarrow & & \\ 0 & \longrightarrow & \mathcal{M}(U) & \longrightarrow & \mathcal{M}(\mathbb{V}_0) & \longrightarrow & \mathcal{M}(\mathbb{V}_1) \end{array}$$

Because  $\mathbb{V}_0$  covers  $\mathbb{U}_0$ ,  $\mathbb{V}_1$  covers  $\mathbb{U}_1$  as well. So the maps  $\beta$  and  $\gamma$  are injective. **Step 1** tells us that the bottom row is exact. Then Lemma 6.3.15 (ii) shows that the top row is exact too.

### Step 3. (definition of $\widetilde{\mathcal{M}}$ )

Let  $Y$  be an open subset of  $X$ . We use the sheaf property to define  $\widetilde{\mathcal{M}}(Y)$ . We choose a (finite) covering  $\mathbb{U}_0 = \{U^i\}$  of  $Y$  by affine open sets, and we define  $\widetilde{\mathcal{M}}(Y)$  to be the kernel  $K_{\mathbb{U}}$  of the map  $\mathcal{M}(\mathbb{U}_0) \xrightarrow{\beta_{\mathbb{U}}} \mathcal{M}(\mathbb{U}_1)$ , where  $\beta_{\mathbb{U}}$  is the map described in (??). When we show that this kernel is independent of the covering  $\mathbb{U}_0$ , it will follow that  $\widetilde{\mathcal{M}}$  is well-defined, and that it has the sheaf property.

Let  $\mathbb{W}_0 = \{W^\nu\}$  be another covering of  $Y$  by affine open sets. We can go from  $\mathbb{U}_0$  to  $\mathbb{V}_0$  and back in a finite number of steps, each of which consists in adding or deleting one affine open set.

We consider a family  $\mathbb{W}_0 = \{U^i, V\}$  obtained by adding one affine open subset  $V$  of  $Y$  to  $\mathbb{U}_0$ , and we let  $\mathbb{W}_1$  be the family of intersections of pairs of elements of  $\mathbb{W}_0$ . Then we have a map  $K_{\mathbb{W}} \rightarrow K_{\mathbb{U}}$ . We show that, for any element  $(u_i)$  in the kernel  $K_{\mathbb{U}}$ , there is a unique element  $v$  in  $\mathcal{M}(V)$  such that  $((u_i), v)$  is in the kernel  $K_{\mathbb{W}}$ . This will show that  $K_{\mathbb{W}} = K_{\mathbb{U}}$ .

To define the element  $v$ , we let  $V^i = U^i \cap V$ . Since  $\mathbb{U}_0 = \{U^i\}$  is a covering of  $Y$ ,  $\mathbb{V}_0 = \{V^i\}$  is a covering of  $V$  by affine open sets. Let  $v_i$  be the restriction of the section  $u_i$  to  $V^i$ . Since  $(u_i)$  is in the kernel of  $\beta_{\mathbb{U}}$ ,  $u_i = u_j$  on  $U^{ij}$ . Then it is also true that  $v_i = v_j$  on the smaller open set  $V^{ij}$ . So  $(v_i)$  is in the kernel of the map  $\mathcal{M}(\mathbb{V}_0) \xrightarrow{\beta_{\mathbb{V}}} \mathcal{M}(\mathbb{V}_1)$ , and since  $\mathbb{V}_0$  is a covering of the affine variety  $V$  by affine open sets, Step 2 tells us that the kernel is  $\mathcal{M}(V)$ . So there is a unique element  $v$  in  $\mathcal{M}(V)$  that restricts to  $v_i$  on  $V^i$ . With this element  $V$ ,  $(u_i, v)$ , is in the kernel of  $\beta_{\mathbb{W}}$ , as is easily verified.

When the subsets in the family  $\mathbb{W}_1$  are listed in the order

$$\mathbb{W}_1 = \{U^i \cap U^j\}, \{V \cap U^j\}, \{U^i \cap V\}, \{V \cap V\}$$

the map  $\beta_{\mathbb{W}}$  sends  $((u_i), v)$  to  $((u_j - u_i), (u_j - v), (v - u_i), 0)$ , restricted appropriately. Here  $u_i = u_j$  on  $U^i \cap U^j$  because  $(u_i)$  is in the kernel of  $\beta_{\mathbb{U}}$ , and  $u_j = v_j = v$  on  $U^j \cap V = V^j$  by definition.

This completes the proof of Theorem 6.3.2. □



## Chapter 7 COHOMOLOGY

- 7.1 Cohomology
- 7.2 Complexes
- 7.3 Characteristic Properties of Cohomology
- 7.4 Existence of Cohomology
- 7.5 Cohomology of the Twisting Modules
- 7.6 Cohomology of Hypersurfaces
- 7.7 Annihilators and Support
- 7.8 Three Theorems about Cohomology
- 7.9 Bézout's Theorem

### 7.1 Cohomology

To simplify the construction, we define cohomology only for  $\mathcal{O}$ -modules. Anyway, the Zariski topology has limited use for cohomology with other coefficients.

Let  $\mathcal{M}$  be an  $\mathcal{O}$ -module on a variety  $X$ . The *zero-dimensional cohomology* of  $\mathcal{M}$  is the space  $\mathcal{M}(X)$  of its global sections. When speaking of cohomology, one denotes this space by  $H^0(X, \mathcal{M})$ .

The functor

$$(\mathcal{O}\text{-modules}) \xrightarrow{H^0} (\text{vector spaces})$$

that carries an  $\mathcal{O}$ -module  $\mathcal{M}$  to  $H^0(X, \mathcal{M})$  is left exact: If

$$(7.1.1) \quad 0 \rightarrow \mathcal{M} \rightarrow \mathcal{N} \rightarrow \mathcal{P} \rightarrow 0$$

is an exact sequence of  $\mathcal{O}$ -modules, the associated sequence of global sections

$$(7.1.2) \quad 0 \rightarrow H^0(X, \mathcal{M}) \rightarrow H^0(X, \mathcal{N}) \rightarrow H^0(X, \mathcal{P})$$

is exact. But unless  $X$  is affine, the map  $H^0(X, \mathcal{N}) \rightarrow H^0(X, \mathcal{P})$  needn't be surjective. The *cohomology* of  $\mathcal{M}$  is a sequence of functors  $(\mathcal{O}\text{-modules}) \xrightarrow{H^q} (\text{vector spaces})$ ,

$$H^0(X, \mathcal{M}), H^1(X, \mathcal{M}), H^2(X, \mathcal{M}), \dots$$

beginning with  $H^0$ , one for each *dimension*, that compensates for the lack of exactness in the following way: Every short exact sequence (7.1.1) of  $\mathcal{O}$ -modules has an associated long exact *cohomology sequence*

$$(7.1.3) \quad \begin{aligned} 0 \rightarrow H^0(X, \mathcal{M}) \rightarrow H^0(X, \mathcal{N}) \rightarrow H^0(X, \mathcal{P}) &\xrightarrow{\delta^0} \\ &\xrightarrow{\delta^0} H^1(X, \mathcal{M}) \rightarrow H^1(X, \mathcal{N}) \rightarrow H^1(X, \mathcal{P}) \xrightarrow{\delta^1} \\ &\dots \dots \dots \\ &\xrightarrow{\delta^{q-1}} H^q(X, \mathcal{M}) \rightarrow H^q(X, \mathcal{N}) \rightarrow H^q(X, \mathcal{P}) \xrightarrow{\delta^q} \dots \end{aligned}$$

And, given a diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & \mathcal{M} & \longrightarrow & \mathcal{N} & \longrightarrow & \mathcal{P} \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & \mathcal{M}' & \longrightarrow & \mathcal{N}' & \longrightarrow & \mathcal{P}' \longrightarrow 0 \end{array}$$

whose rows are short exact sequences of  $\mathcal{O}$ -modules, the diagrams

$$(7.1.4) \quad \begin{array}{ccc} H^q(X, \mathcal{P}) & \xrightarrow{\delta^q} & H^{q+1}(X, \mathcal{M}) \\ \downarrow & & \downarrow \\ H^q(X, \mathcal{P}') & \xrightarrow{\delta^q} & H^{q+1}(X, \mathcal{M}') \end{array}$$

that are obtained from the map of cohomology sequences commute. The other diagrams commute because  $H^q$  are functors. Thus a map of short exact sequences induces a map of cohomology sequences. The maps  $\delta^q$  are the *coboundary maps*.

A sequence  $H^q$ ,  $q = 0, 1, \dots$  of functors from  $\mathcal{O}$ -modules to vector spaces that comes with long cohomology sequences for every short exact sequence (7.1.1) is called a *cohomological functor*.

Unfortunately, there is no really natural construction of the cohomology. Sometimes one needs to look at an explicit construction, but it is usually best to work with the characteristic properties that are described in the Section 7.3. We present a construction in Section 7.4, but it isn't canonical.

The one-dimensional cohomology  $H^1$  has an interesting interpretation that you can read about if you like. We won't use it. The higher cohomology  $H^q$  has no useful direct interpretation.

## 7.2 Complexes

We need complexes because they are used in the definition of cohomology.

A *complex*  $V^\bullet$  of vector spaces is a sequence of homomorphisms of vector spaces

$$(7.2.1) \quad \dots \rightarrow V^{n-1} \xrightarrow{d^{n-1}} V^n \xrightarrow{d^n} V^{n+1} \xrightarrow{d^{n+1}} \dots$$

indexed by the integers, such that the composition  $d^n d^{n-1}$  of adjacent maps is zero – the image of  $d^{n-1}$  is contained in the kernel of  $d^n$ . The  $q$ -dimensional *cohomology* of a complex  $V^\bullet$  is the quotient

$$(7.2.2) \quad \mathbf{C}^q(V^\bullet) = (\ker d^q) / (\operatorname{im} d^{q-1}).$$

An exact sequence is a complex whose cohomology is zero.

If a finite sequence of homomorphisms (7.2.1) is given, say  $V^k \rightarrow V^{k+1} \rightarrow \dots \rightarrow V^\ell$ , it can be made into a complex by defining  $V^n = 0$  for all other integers  $n$ . In our applications  $V^q$  will be zero when  $q < 0$ .

A homomorphism of vector spaces  $V^0 \xrightarrow{d^0} V^1$  can be made into the complex

$$\dots \rightarrow 0 \rightarrow V^0 \xrightarrow{d^0} V^1 \rightarrow 0 \rightarrow \dots$$

For this complex,  $\mathbf{C}^0 = \ker d^0$ ,  $\mathbf{C}^1 = \operatorname{coker} d^0$ , and  $\mathbf{C}^q = 0$  for all other  $q$ .

A *map*  $V^\bullet \xrightarrow{\varphi} V'^\bullet$  of complexes is a collection of homomorphisms  $V^n \xrightarrow{\varphi^n} V'^n$  making a diagram

$$\begin{array}{ccccccc} \longrightarrow & V^{n-1} & \xrightarrow{d^{n-1}} & V^n & \xrightarrow{d^n} & V^{n+1} & \longrightarrow \dots \\ & \varphi^{n-1} \downarrow & & \varphi^n \downarrow & & \varphi^{n+1} \downarrow & \\ \longrightarrow & V'^{n-1} & \xrightarrow{d'^{n-1}} & V'^n & \xrightarrow{d'^n} & V'^{n+1} & \longrightarrow \dots \end{array}$$

A map of complexes induces maps on the cohomology

$$\mathbf{C}^q(V^\bullet) \rightarrow \mathbf{C}^q(V'^\bullet)$$

because  $\ker d^q$  maps to  $\ker d'^q$  and  $\operatorname{im} d^q$  maps to  $\operatorname{im} d'^q$ .

A sequence

$$(7.2.3) \quad \cdots \rightarrow V^\bullet \xrightarrow{\varphi} V'^\bullet \xrightarrow{\psi} V''^\bullet \rightarrow \cdots$$

of maps of complexes is *exact* if the sequences

$$\cdots \rightarrow V^q \xrightarrow{\varphi^q} V'^q \xrightarrow{\psi^q} V''^q \rightarrow \cdots$$

are exact for every  $q$ .

#### 7.2.4. Proposition.

Let  $0 \rightarrow V^\bullet \rightarrow V'^\bullet \rightarrow V''^\bullet \rightarrow 0$  be a short exact sequence of complexes. For every  $q$ , there are maps  $C^q(V''^\bullet) \xrightarrow{\delta^q} C^{q+1}(V^\bullet)$  such that the sequence

$$\rightarrow C^0(V^\bullet) \rightarrow C^0(V'^\bullet) \rightarrow C^0(V''^\bullet) \xrightarrow{\delta^0} C^1(V^\bullet) \rightarrow C^1(V'^\bullet) \rightarrow C^1(V''^\bullet) \xrightarrow{\delta^1} C^2(V^\bullet) \rightarrow \cdots$$

is exact.

The exact sequence displayed above is the *cohomology sequence* associated to the short exact sequence of complexes. This property makes the set of functors  $\{C^q\}$  into a *cohomological functor* on the category of complexes.

**7.2.5. Example.** We make the Snake Lemma into a cohomology sequence. Suppose given a diagram

$$\begin{array}{ccccccc} V & \xrightarrow{u} & V' & \longrightarrow & V'' & \longrightarrow & 0 \\ f \downarrow & & f' \downarrow & & f'' \downarrow & & \\ 0 & \longrightarrow & W & \longrightarrow & W' & \xrightarrow{v} & W'' \end{array}$$

with exact rows. We form the complex  $0 \rightarrow V \xrightarrow{f} W \rightarrow 0$  with  $V$  in degree zero, and we do the analogous thing for the maps  $f'$  and  $f''$ , so that the diagram becomes a short exact sequence of complexes. Its cohomology sequence is the one given by the Snake Lemma.  $\square$

*proof of Proposition 7.2.4.* Let

$$V^\bullet = \{ \cdots \rightarrow V^{q-1} \xrightarrow{d^{q-1}} V^q \xrightarrow{d^q} V^{q+1} \xrightarrow{d^{q+1}} \cdots \}$$

be a complex, let  $B^q$  be the image of  $d^{q-1}$  in  $V^q$ , and let  $Z^q$  be the kernel of  $d^q$ . So  $B^q \subset Z^q \subset V^q$ , and the cohomology of the complex is  $C^q(V^\bullet) = Z^q/B^q$ . Also, let  $D^q$  be the cokernel of  $d^{q-1}$ . So  $D^q = V^q/B^q$ , and there is an exact sequence

$$0 \rightarrow B^q \rightarrow V^q \rightarrow D^q \rightarrow 0$$

Again since  $B^q \subset Z^q$ , the map  $d^q$  can be written as the composition of three maps

$$V^q \xrightarrow{\pi^q} D^q \xrightarrow{f^q} Z^{q+1} \xrightarrow{i^{q+1}} V^{q+1}$$

where  $\pi^q$  is the projection from  $V^q$  to its quotient  $D^q$  and  $i^{q+1}$  is the inclusion of  $Z^{q+1}$  into  $V^{q+1}$ . Studying these maps, one sees that

$$(7.2.6) \quad C^q(V^\bullet) = \ker f^q \quad \text{and} \quad C^{q+1}(V^\bullet) = \operatorname{coker} f^q.$$

Suppose given a short exact sequence of complexes  $0 \rightarrow V^\bullet \rightarrow V'^\bullet \rightarrow V''^\bullet \rightarrow 0$  as in the proposition. In the diagram below, the rows are exact because cokernel is a right exact functor and kernel is a left exact functor.

$$\begin{array}{ccccccc} D^q & \longrightarrow & D'^q & \longrightarrow & D''^q & \longrightarrow & 0 \\ f^q \downarrow & & f'^q \downarrow & & f''^q \downarrow & & \\ 0 & \longrightarrow & Z^{q+1} & \longrightarrow & Z'^{q+1} & \longrightarrow & Z''^{q+1} \end{array}$$

When we apply (7.2.6) and the Snake Lemma to this diagram, we obtain an exact sequence

$$\mathbf{C}^q(V^\bullet) \rightarrow \mathbf{C}^q(V'^\bullet) \rightarrow \mathbf{C}^q(V''^\bullet) \xrightarrow{\delta^q} \mathbf{C}^{q+1}(V^\bullet) \rightarrow \mathbf{C}^{q+1}(V'^\bullet) \rightarrow \mathbf{C}^{q+1}(V''^\bullet)$$

The cohomology sequence is obtained by splicing these sequences together.  $\square$

The coboundary maps  $\delta^q$  in cohomology sequences are related in a natural way. If

$$\begin{array}{ccccccccc} 0 & \longrightarrow & U^\bullet & \longrightarrow & U'^\bullet & \longrightarrow & U''^\bullet & \longrightarrow & 0 \\ & & \downarrow & & \downarrow & & \downarrow & & \\ 0 & \longrightarrow & V^\bullet & \longrightarrow & V'^\bullet & \longrightarrow & V''^\bullet & \longrightarrow & 0 \end{array}$$

is a diagram of maps of complexes whose rows are short exact sequences, the diagrams

$$\begin{array}{ccc} \mathbf{C}^q(U''^\bullet) & \xrightarrow{\delta^q} & \mathbf{C}^{q+1}(U^\bullet) \\ \downarrow & & \downarrow \\ \mathbf{C}^q(V''^\bullet) & \xrightarrow{\delta^q} & \mathbf{C}^{q+1}(V^\bullet) \end{array}$$

commute. Thus a map of short exact sequences induces a map of cohomology sequences.

### 7.3 Characteristic Properties of Cohomology

The cohomology  $H^q(X, \cdot)$  of  $\mathcal{O}$ -modules, the sequence of functors  $H^0(X, \cdot), H^1(X, \cdot), H^2(X, \cdot), \dots$  from ( $\mathcal{O}$ -modules) to (vector spaces), is characterized by the three properties below, the first two of which have already been mentioned.

#### (7.3.1) characteristic properties

- $H^0(X, \mathcal{M})$  is the space  $\mathcal{M}(X)$  of global sections of  $\mathcal{M}$ .
- The sequence  $H^0, H^1, H^2, \dots$  is a cohomological functor on  $\mathcal{O}$ -modules: A short exact sequence of  $\mathcal{O}$ -modules produces a long exact cohomology sequence.
- Let  $Y \xrightarrow{f} X$  be the inclusion of an *affine* open subset  $Y$  into  $X$ , let  $\mathcal{N}$  be an  $\mathcal{O}_Y$ -module, and let  $f_*\mathcal{N}$  be its direct image on  $X$ . The cohomology  $H^q(X, f_*\mathcal{N})$  is zero for all  $q > 0$ .

**7.3.2. Example.** Let  $j$  be the inclusion of the standard affine open set  $\mathbb{U}^0$  into projective space  $X$ . The third property tells us that the cohomology  $H^q(X, j_*\mathcal{O}_{\mathbb{U}^0})$  of the direct image  $j_*\mathcal{O}_{\mathbb{U}^0}$  is zero when  $q > 0$ . The direct image is isomorphic to the limit  $\varinjlim \mathcal{O}_X(nH)$  (6.6.13). We will see below (7.4.28) that cohomology commutes with direct limits. Therefore the limits of  $H^q(X, \mathcal{O}_X(nH))$  and of  $H^q(X, \mathcal{O}_X(n))$  are zero when  $X$  is projective space and  $q > 0$ . This will be useful.

Intuitively, the third property tells us that allowing poles on the complement of an affine open set kills cohomology in positive dimension.  $\square$

**7.3.3. Theorem.** *There exists a cohomology theory with the properties (7.3.1), and it is unique up to unique isomorphism.*

The proof is in the next section.

**7.3.4. Corollary.** *If  $X$  is an affine variety,  $H^q(X, \mathcal{M}) = 0$  for all  $\mathcal{O}$ -modules  $\mathcal{M}$  and all  $q > 0$ .*

This follows when one applies the third characteristic property to the identity map  $X \rightarrow X$ .  $\square$

## 7.4 Existence of Cohomology

The proof of existence of cohomology and its uniqueness are based on the following facts:

- The intersection of two affine open subsets of a variety is an affine open set.
- A sequence  $\cdots \rightarrow \mathcal{M} \rightarrow \mathcal{N} \rightarrow \mathcal{P} \rightarrow \cdots$  of  $\mathcal{O}$ -modules on a variety  $X$  is exact if and only if, for every affine open subset  $U$ , the sequence of sections  $\cdots \rightarrow \mathcal{M}(U) \rightarrow \mathcal{N}(U) \rightarrow \mathcal{P}(U) \rightarrow \cdots$  is exact. (This is the definition of exactness of a sequence of  $\mathcal{O}$ -modules.)

We begin by choosing an arbitrary affine covering  $\mathbb{U} = \{U^\nu\}$  of our variety  $X$  by finitely many affine open sets  $U^\nu$ , and we use this covering to describe the cohomology. When we have shown that the cohomology is unique, we will know that it doesn't depend on our choice of covering.

Let  $\mathbb{U}$  denote our chosen covering of  $X$ , and let  $\mathbb{U} \xrightarrow{j} X$  denote the family of inclusions  $U^\nu \xrightarrow{j^\nu} X$ . If  $\mathcal{M}$  is an  $\mathcal{O}$ -module,  $\mathcal{R}_{\mathcal{M}}$  will denote the  $\mathcal{O}$ -module  $j_*\mathcal{M}_{\mathbb{U}} = \prod j_*^\nu \mathcal{M}_{U^\nu}$ , where  $\mathcal{M}_{U^\nu}$  is the restriction of  $\mathcal{M}$  to the open set  $U^\nu$ . As has been noted (6.5.8), there is a canonical map  $\mathcal{M} \rightarrow j_*^\nu \mathcal{M}_{U^\nu}$ , and therefore a canonical map  $\mathcal{M} \rightarrow \mathcal{R}_{\mathcal{M}}$ .

**7.4.1. Lemma. (i)** *Let  $X'$  be an open subset of  $X$ . The module of sections  $\mathcal{R}_{\mathcal{M}}(X')$  of  $\mathcal{R}_{\mathcal{M}}$  on  $X'$  is the product  $\prod_\nu \mathcal{M}(X' \cap U^\nu)$ . In particular, the space of global sections  $\mathcal{R}_{\mathcal{M}}(X)$  is the product  $\prod_\nu \mathcal{M}(U^\nu)$ .*

**(ii)** *The canonical map  $\mathcal{M} \rightarrow \mathcal{R}_{\mathcal{M}}$  is injective. Thus, if  $\mathcal{S}_{\mathcal{M}}$  denotes the cokernel of that map, there is a short exact sequence of  $\mathcal{O}$ -modules*

$$(7.4.2) \quad 0 \rightarrow \mathcal{M} \rightarrow \mathcal{R}_{\mathcal{M}} \rightarrow \mathcal{S}_{\mathcal{M}} \rightarrow 0$$

**(iii)** *For any cohomology theory with the characteristic properties and for any  $q > 0$ ,  $H^q(X, \mathcal{R}_{\mathcal{M}}) = 0$ .*

*proof.* **(i)** This is seen by going through the definitions:

$$\mathcal{R}(X') = \prod_\nu [j_*^\nu \mathcal{M}_{U^\nu}](X') = \prod_\nu \mathcal{M}_{U^\nu}(X' \cap U^\nu) = \prod_\nu \mathcal{M}(X' \cap U^\nu).$$

**(ii)** Let  $X'$  be an open subset of  $X$ . The map  $\mathcal{M}(X') \rightarrow \mathcal{R}_{\mathcal{M}}(X')$  is the product of the restriction maps  $\mathcal{M}(X') \rightarrow \mathcal{M}(X' \cap U^\nu)$ . Because the open sets  $U^\nu$  cover  $X$ , the intersections  $X' \cap U^\nu$  cover  $X'$ . The sheaf property of  $\mathcal{M}$  tells us that the map  $\mathcal{M}(X') \rightarrow \prod_\nu \mathcal{M}(X' \cap U^\nu)$  is injective.

**(iii)** This follows from the third characteristic property. □

**7.4.3. Lemma. (i)** *A short exact sequence  $0 \rightarrow \mathcal{M} \rightarrow \mathcal{N} \rightarrow \mathcal{P} \rightarrow 0$  of  $\mathcal{O}$ -modules embeds into a diagram*

$$(7.4.4) \quad \begin{array}{ccccc} \mathcal{M} & \longrightarrow & \mathcal{N} & \longrightarrow & \mathcal{P} \\ \downarrow & & \downarrow & & \downarrow \\ \mathcal{R}_{\mathcal{M}} & \longrightarrow & \mathcal{R}_{\mathcal{N}} & \longrightarrow & \mathcal{R}_{\mathcal{P}} \\ \downarrow & & \downarrow & & \downarrow \\ \mathcal{S}_{\mathcal{M}} & \longrightarrow & \mathcal{S}_{\mathcal{N}} & \longrightarrow & \mathcal{S}_{\mathcal{P}} \end{array}$$

*whose rows and columns are short exact sequences. (We have suppressed the surrounding zeros.)*

**(ii)** *The sequence of global sections  $0 \rightarrow \mathcal{R}_{\mathcal{M}}(X) \rightarrow \mathcal{R}_{\mathcal{N}}(X) \rightarrow \mathcal{R}_{\mathcal{P}}(X) \rightarrow 0$  is exact.*

*proof.* **(i)** We are given that the top row of the diagram is a short exact sequence, and we have seen that the columns are short exact sequences. To show that the middle row

$$(7.4.5) \quad 0 \rightarrow \mathcal{R}_{\mathcal{M}} \rightarrow \mathcal{R}_{\mathcal{N}} \rightarrow \mathcal{R}_{\mathcal{P}} \rightarrow 0$$

is exact, we must show that if  $X'$  is an affine open subset, the sections on  $X'$  form a short exact sequence. The sections are explained in Lemma 7.4.1 (i). Since products of exact sequences are exact, we must show that the sequence

$$0 \rightarrow \mathcal{M}(X' \cap U^\nu) \rightarrow \mathcal{N}(X' \cap U^\nu) \rightarrow \mathcal{P}(X' \cap U^\nu) \rightarrow 0$$

is exact. This is true because  $X' \cap U^\nu$  is an intersection of affine opens, and is therefore affine.

Now that we know that the first two rows of the diagram are short exact sequences, the Snake Lemma tells us that the bottom row of the diagram is a short exact sequence.

(ii) The sequence of global sections is the product of the sequences

$$0 \rightarrow \mathcal{M}(U^\nu) \rightarrow \mathcal{N}(U^\nu) \rightarrow \mathcal{P}(U^\nu) \rightarrow 0$$

These sequences are exact because the open sets  $U^\nu$  are affine.  $\square$

#### (7.4.6) uniqueness of cohomology

Suppose that a cohomology with the characteristic properties (7.3.1) is given, and let  $\mathcal{M}$  be an  $\mathcal{O}$ -module. Then  $H^q(X, \mathcal{R}_{\mathcal{M}}) = 0$  if  $q > 0$  (Lemma 7.4.1 (iii)). The cohomology sequence associated to the sequence  $0 \rightarrow \mathcal{M} \rightarrow \mathcal{R}_{\mathcal{M}} \rightarrow \mathcal{S}_{\mathcal{M}} \rightarrow 0$  is

$$0 \rightarrow H^0(X, \mathcal{M}) \rightarrow H^0(X, \mathcal{R}_{\mathcal{M}}) \rightarrow H^0(X, \mathcal{S}_{\mathcal{M}}) \xrightarrow{\delta^0} H^1(X, \mathcal{M}) \rightarrow H^1(X, \mathcal{R}_{\mathcal{M}}) \rightarrow \cdots$$

Since  $H^q(X, \mathcal{R}_{\mathcal{M}}) = 0$  when  $q > 0$ , this sequence breaks up into an exact sequence

$$(7.4.7) \quad 0 \rightarrow H^0(X, \mathcal{M}) \rightarrow H^0(X, \mathcal{R}_{\mathcal{M}}) \rightarrow H^0(X, \mathcal{S}_{\mathcal{M}}) \xrightarrow{\delta^0} H^1(X, \mathcal{M}) \rightarrow 0$$

and isomorphisms

$$(7.4.8) \quad 0 \rightarrow H^q(X, \mathcal{S}_{\mathcal{M}}) \xrightarrow{\delta^q} H^{q+1}(X, \mathcal{M}) \rightarrow 0$$

for every  $q > 0$ . The first three terms of the sequence (7.4.7), and the arrows connecting them, depend on our choice of covering of  $X$ , but the important point is that they don't depend on the cohomology. So that sequence determines  $H^1(X, \mathcal{M})$  up to unique isomorphism as the cokernel of a map that is independent of the cohomology, and this is true for every  $\mathcal{O}$ -module  $\mathcal{M}$ , including for the module  $\mathcal{S}_{\mathcal{M}}$ . Therefore it is also true that  $H^1(X, \mathcal{S}_{\mathcal{M}})$  is determined uniquely. This being so,  $H^2(X, \mathcal{M})$  is determined uniquely for every  $\mathcal{M}$ , by the isomorphism (7.4.8), with  $q = 1$ . The isomorphisms (7.4.8) determine the rest of the cohomology up to unique isomorphism by induction on  $q$ .

#### (7.4.9) construction of cohomology

One can use the sequence (7.4.2) and induction to construct cohomology as well as to prove uniqueness, but it will be clearer to proceed by iterating the construction of  $\mathcal{R}_{\mathcal{M}}$ .

Let  $\mathcal{M}$  be an  $\mathcal{O}$ -module. We rewrite the exact sequence (7.4.2), labeling  $\mathcal{R}_{\mathcal{M}}$  as  $\mathcal{R}_{\mathcal{M}}^0$ , and  $\mathcal{S}_{\mathcal{M}}$  as  $\mathcal{M}^1$ :

$$(7.4.10) \quad 0 \rightarrow \mathcal{M} \rightarrow \mathcal{R}_{\mathcal{M}}^0 \rightarrow \mathcal{M}^1 \rightarrow 0$$

and we repeat the construction with  $\mathcal{M}^1$ . Let  $\mathcal{R}_{\mathcal{M}}^1 = \mathcal{R}_{\mathcal{M}^1}^0 (= j_* \mathcal{M}_{\mathbb{U}}^1)$ , so that there is an exact sequence

$$(7.4.11) \quad 0 \rightarrow \mathcal{M}^1 \rightarrow \mathcal{R}_{\mathcal{M}}^1 \rightarrow \mathcal{M}^2 \rightarrow 0$$

analogous to the sequence (7.4.10), with  $\mathcal{M}^2 = \mathcal{R}_{\mathcal{M}}^1 / \mathcal{M}^1$ . We combine the sequences (7.4.10) and (7.4.11) into an exact sequence

$$(7.4.12) \quad 0 \rightarrow \mathcal{M} \rightarrow \mathcal{R}_{\mathcal{M}}^0 \rightarrow \mathcal{R}_{\mathcal{M}}^1 \rightarrow \mathcal{M}^2 \rightarrow 0$$

Then we let  $\mathcal{R}_{\mathcal{M}}^2 = \mathcal{R}_{\mathcal{M}^2}^0$ . We continue in this way, to construct modules  $\mathcal{R}_{\mathcal{M}}^k$  that form an exact sequence

$$(7.4.13) \quad 0 \rightarrow \mathcal{M} \rightarrow \mathcal{R}_{\mathcal{M}}^0 \rightarrow \mathcal{R}_{\mathcal{M}}^1 \rightarrow \mathcal{R}_{\mathcal{M}}^2 \rightarrow \cdots$$

The next lemma follows by induction from Lemmas 7.4.1 and 7.4.3.

**7.4.14. Lemma.**

(i) Let  $0 \rightarrow \mathcal{M} \rightarrow \mathcal{N} \rightarrow \mathcal{P} \rightarrow 0$  be a short exact sequence of  $\mathcal{O}$ -modules. For every  $n$ , the sequences

$$0 \rightarrow \mathcal{R}_{\mathcal{M}}^n \rightarrow \mathcal{R}_{\mathcal{N}}^n \rightarrow \mathcal{R}_{\mathcal{P}}^n \rightarrow 0$$

are exact, and so are the sequences of global sections

$$0 \rightarrow \mathcal{R}_{\mathcal{M}}^n(X) \rightarrow \mathcal{R}_{\mathcal{N}}^n(X) \rightarrow \mathcal{R}_{\mathcal{P}}^n(X) \rightarrow 0$$

(ii) If  $H^q$ ,  $q = 0, 1, 2, \dots$  is a cohomology theory, then  $H^q(X, \mathcal{R}_{\mathcal{M}}^n) = 0$  for all  $n$  and all  $q > 0$ .  $\square$

An exact sequence such as (7.4.13) is called a *resolution* of  $\mathcal{M}$ , and because  $H^q(X, \mathcal{R}_{\mathcal{M}}^n) = 0$  when  $q > 0$ , it is an *acyclic resolution*.

Continuing with the proof of existence, we consider the complex of  $\mathcal{O}$ -modules  $\mathcal{R}_{\mathcal{M}}^\bullet$  that is obtained by omitting the first term from (7.4.13):

$$(7.4.15) \quad 0 \rightarrow \mathcal{R}_{\mathcal{M}}^0 \rightarrow \mathcal{R}_{\mathcal{M}}^1 \rightarrow \mathcal{R}_{\mathcal{M}}^2 \rightarrow \dots$$

and the complex  $\mathcal{R}_{\mathcal{M}}^\bullet(X)$  of its global sections:

$$(7.4.16) \quad 0 \rightarrow \mathcal{R}_{\mathcal{M}}^0(X) \rightarrow \mathcal{R}_{\mathcal{M}}^1(X) \rightarrow \mathcal{R}_{\mathcal{M}}^2(X) \rightarrow \dots$$

which we could also write as

$$0 \rightarrow H^0(X, \mathcal{R}_{\mathcal{M}}^0) \rightarrow H^0(X, \mathcal{R}_{\mathcal{M}}^1) \rightarrow H^0(X, \mathcal{R}_{\mathcal{M}}^2) \rightarrow \dots$$

The sequence  $\mathcal{R}_{\mathcal{M}}^\bullet$  becomes the resolution (7.4.13) when the module  $\mathcal{M}$  is inserted. So the complex (7.4.15) is exact except at  $\mathcal{R}_{\mathcal{M}}^0$ , but because the global section functor is only left exact, the sequence (7.4.16) of global sections  $\mathcal{R}_{\mathcal{M}}^\bullet(X)$  needn't be exact anywhere. However,  $\mathcal{R}_{\mathcal{M}}^\bullet(X)$  is a complex because  $\mathcal{R}_{\mathcal{M}}^\bullet$  is a complex. The composition of adjacent maps is zero.

Recall that the cohomology of a complex  $0 \rightarrow V^0 \xrightarrow{d^0} V^1 \xrightarrow{d^1} \dots$  of vector spaces is defined to be  $\mathbf{C}^q(V^\bullet) = (\ker d^q) / (\text{im } d^{q-1})$ , and that  $\{\mathbf{C}^q\}$  is a cohomological functor on complexes (7.2.4).

**7.4.17. Definition.** The cohomology of an  $\mathcal{O}$ -module  $\mathcal{M}$  is the cohomology of the complex  $\mathcal{R}_{\mathcal{M}}^\bullet(X)$ :  $H^q(X, \mathcal{M}) = \mathbf{C}^q(\mathcal{R}_{\mathcal{M}}^\bullet(X))$ .

Thus if we denote the maps in the complex (7.4.16) by  $d^q$ ,

$$0 \rightarrow \mathcal{R}_{\mathcal{M}}^0(X) \xrightarrow{d^0} \mathcal{R}_{\mathcal{M}}^1(X) \xrightarrow{d^1} \mathcal{R}_{\mathcal{M}}^2(X) \rightarrow \dots$$

then  $H^q(X, \mathcal{M}) = (\ker d^q) / (\text{im } d^{q-1})$ .

**7.4.18. Lemma.** Let  $X$  be an affine variety. With cohomology defined as above,  $H^q(X, \mathcal{M}) = 0$  for all  $\mathcal{O}$ -modules  $\mathcal{M}$  and all  $q > 0$ .

*proof.* When  $X$  is affine, the sequence of global sections of the exact sequence (7.4.13) is exact.  $\square$

To show that our definition gives the (unique) cohomology, we verify the characteristic properties. Since the sequence (7.4.13) is exact and since the global section functor is left exact,  $\mathcal{M}(X)$  is the kernel of the map  $\mathcal{R}_{\mathcal{M}}^0(X) \rightarrow \mathcal{R}_{\mathcal{M}}^1(X)$ , and this kernel is also equal to  $\mathbf{C}^0(\mathcal{R}_{\mathcal{M}}^\bullet(X))$ . So our cohomology has the first property:  $H^0(X, \mathcal{M}) = \mathcal{M}(X)$ .

To show that we obtain a cohomological functor, we apply Lemma 7.4.14 to conclude that, for a short exact sequence  $0 \rightarrow \mathcal{M} \rightarrow \mathcal{N} \rightarrow \mathcal{P} \rightarrow 0$ , the global sections

$$(7.4.19) \quad 0 \rightarrow \mathcal{R}_{\mathcal{M}}^\bullet(X) \rightarrow \mathcal{R}_{\mathcal{N}}^\bullet(X) \rightarrow \mathcal{R}_{\mathcal{P}}^\bullet(X) \rightarrow 0,$$

form an exact sequence of complexes. Cohomology  $H^q(X, \cdot)$  is a cohomological functor because cohomology of complexes is a cohomological functor.

We make a digression before verifying the third characteristic property.

**(7.4.20) affine morphisms**

Let  $Y \xrightarrow{f} X$  be a morphism of varieties. Let  $U \xrightarrow{j} X$  be the inclusion of an open subvariety into  $X$  and let  $V$  be the inverse image  $f^{-1}U$ , which is an open subvariety of  $Y$ . These varieties and maps form a diagram

$$(7.4.21) \quad \begin{array}{ccc} V & \xrightarrow{i} & Y \\ g \downarrow & & \downarrow f \\ U & \xrightarrow{j} & X \end{array}$$

We use the notation  $\mathcal{M}_U$  for the restriction of  $\mathcal{M}$  to an open subset  $U$  of (6.5.6).

**7.4.22. Lemma.** *With notation as above, let  $\mathcal{N}$  be an  $\mathcal{O}_Y$ -module. The  $\mathcal{O}_U$ -modules  $g_*[\mathcal{N}_V]$  and  $[f_*\mathcal{N}]_U$  are canonically isomorphic.*

*proof.* Let  $U'$  be an open subset of  $U$ , and let  $V' = g^{-1}U'$ . Then

$$[f_*\mathcal{N}]_U(U') = [f_*\mathcal{N}](U') = \mathcal{N}(V') = \mathcal{N}_V(V') = [g_*[\mathcal{N}_V]](U') \quad \square$$

**7.4.23. Definition.** An *affine morphism* is a morphism  $Y \xrightarrow{f} X$  of varieties with the property that the inverse image  $f^{-1}(U)$  of every affine open subset  $U$  of  $X$  is an affine open subset of  $Y$ .  $\square$

The following are examples of affine morphisms:

- the inclusion of an affine open subset  $Y$  into  $X$ ,
- the inclusion of a closed subvariety  $Y$  into  $X$ ,
- a finite morphism, or an integral morphism.

But, if  $Y$  is a closed subset of  $\mathbb{P}^n \times X$ , the projection  $Y \rightarrow X$  will not be an affine morphism unless its fibres are finite, in which case Chevalley's Finiteness Theorem tells us that it is a finite morphism.

**7.4.24. Lemma.** *If  $Y \xrightarrow{f} C$  is an affine morphism and if  $\mathcal{N} \rightarrow \mathcal{N}' \rightarrow \mathcal{N}''$  is an exact sequence of  $\mathcal{O}_Y$ -modules, the sequence of direct images  $f_*\mathcal{N} \rightarrow f_*\mathcal{N}' \rightarrow f_*\mathcal{N}''$  is exact.*  $\square$

Let  $Y \xrightarrow{f} X$  be an affine morphism, let  $j$  be the map from our chosen affine covering  $\mathbb{U} = \{U^\nu\}$  to  $X$ , and let  $\mathbb{V}$  denote the family  $\{V^\nu\} = \{f^{-1}U^\nu\}$  of inverse images. Then  $\mathbb{V}$  is an affine covering of  $Y$ , and there is a morphism  $\mathbb{V} \xrightarrow{g} \mathbb{U}$ . We form a diagram analogous to (7.4.21), in which  $\mathbb{V}$  and  $\mathbb{U}$  replace  $V$  and  $U$ , respectively:

$$\begin{array}{ccc} \mathbb{V} & \xrightarrow{i} & Y \\ g \downarrow & & \downarrow f \\ \mathbb{U} & \xrightarrow{j} & X \end{array}$$

**7.4.25. Proposition.** *Let  $Y \xrightarrow{f} X$  be an affine morphism, and let  $\mathcal{N}$  be an  $\mathcal{O}_Y$ -module. Let  $H^q(X, \cdot)$  be cohomology defined in (7.4.17), and let  $H^q(Y, \cdot)$  be cohomology defined in the analogous way, using the covering  $\mathbb{V}$  of  $Y$ . Then  $H^q(X, f_*\mathcal{N})$  is isomorphic to  $H^q(Y, \mathcal{N})$ .*

*proof.* To compute the cohomology of  $f_*\mathcal{N}$  on  $X$ , we substitute  $\mathcal{M} = f_*\mathcal{N}$  into (7.4.17):

$$H^q(X, f_*\mathcal{N}) = \mathbf{C}^q(\mathcal{R}_{f_*\mathcal{N}}^\bullet(X)).$$

To compute the cohomology of  $\mathcal{N}$  on  $Y$ , we let

$$\mathcal{R}'^0_{\mathcal{N}} = i_*[\mathcal{N}_V]$$



and we continue, to construct a resolution  $\mathcal{R}'^\bullet_{\mathcal{N}} = 0 \rightarrow \mathcal{N} \rightarrow \mathcal{R}'^0_{\mathcal{N}} \rightarrow \mathcal{R}'^1_{\mathcal{N}} \rightarrow \cdots$  and the complex of its global sections  $\mathcal{R}'^\bullet_{\mathcal{N}}(Y)$ . (The prime is there to remind us that  $\mathcal{R}'$  is defined using the covering  $\mathbb{V}$  of  $Y$ .) Then

$$H^q(Y, \mathcal{N}) = \mathbb{C}^q(\mathcal{R}'^\bullet_{\mathcal{N}}(Y)).$$

It suffices to show that the complexes  $\mathcal{R}^\bullet_{f_*\mathcal{N}}(X)$  and  $\mathcal{R}'^\bullet_{\mathcal{N}}(Y)$  are isomorphic, and because  $\mathcal{R}'^q_{\mathcal{N}}(Y) = [f_*\mathcal{R}'^q_{\mathcal{N}}](X)$ , it suffices to show that  $\mathcal{R}^q_{f_*\mathcal{N}} \approx f_*\mathcal{R}'^q_{\mathcal{N}}$ .

##reread this##

We look back at the definition (7.4.11) of the  $\mathcal{O}_X$ -modules  $\mathcal{R}^0$ . We have  $\mathcal{R}'^0_{\mathcal{N}} = i_*\mathcal{N}_{\mathbb{V}}$ . So the sequence for  $\mathcal{N}$  analogous to (7.4.10) can be written as

$$0 \rightarrow \mathcal{N} \rightarrow i_*\mathcal{N}_{\mathbb{V}} \rightarrow \mathcal{N}^1 \rightarrow 0$$

and since  $fi = jg$ , its direct image can be written as

$$(7.4.26) \quad 0 \rightarrow f_*\mathcal{N} \rightarrow j_*g_*[\mathcal{N}_{\mathbb{V}}] \rightarrow f_*[\mathcal{N}^1] \rightarrow 0$$

The sequence for  $f_*\mathcal{N}$  analogous to (7.4.10) is

$$0 \rightarrow f_*\mathcal{N} \rightarrow j_*[f_*\mathcal{N}]_{\mathbb{U}} \rightarrow [f_*\mathcal{N}]^1 \rightarrow 0$$

According to Lemma 7.4.22,  $[f_*\mathcal{N}]_{\mathbb{U}}$  is isomorphic to  $g_*[\mathcal{N}_{\mathbb{V}}]$ . So this sequence can also be written as

$$(7.4.27) \quad 0 \rightarrow f_*\mathcal{N} \rightarrow j_*g_*[\mathcal{N}_{\mathbb{V}}] \rightarrow [f_*\mathcal{N}]^1 \rightarrow 0$$

Combining reffstarN) and (7.4.27), one sees that  $\mathcal{R}^0_{f_*\mathcal{N}} \approx f_*\mathcal{R}'^0_{\mathcal{N}}$  and that  $f_*[\mathcal{N}^1] \approx [f_*\mathcal{N}]^1$ . Then induction applies.  $\square$

We go back to the proof of existence of cohomology to verify the third characteristic property, that when  $Y \xrightarrow{f} X$  is the inclusion of an affine open subset,  $H^q(X, f_*\mathcal{N}) = 0$  for all  $\mathcal{O}_Y$ -modules  $\mathcal{N}$  and all  $q > 0$ . The inclusion of an affine open set is an affine morphism, so  $H^q(Y, \mathcal{N}) = H^q(X, f_*\mathcal{N})$  (7.4.25), and since  $Y$  is affine,  $H^q(Y, \mathcal{N}) = 0$  for all  $q > 0$  (7.4.18).  $\square$

Proposition 7.4.25 is one of the places where a specific construction of cohomology is used. The characteristic properties don't apply directly. The next proposition is another such place.

**7.4.28. Lemma.** *Cohomology is compatible with limits of directed sets of  $\mathcal{O}$ -modules:  $H^q(X, \varinjlim \mathcal{M}_\bullet) \approx \varinjlim H^q(X, \mathcal{M}_\bullet)$  for all  $q$ .*

*proof.* The direct and inverse image functors and the global section functor are all compatible with  $\varinjlim$ , and  $\varinjlim$  is exact (??). So the module  $\mathcal{R}^q_{\varinjlim \mathcal{M}_\bullet}$  that is used to compute the cohomology of  $\varinjlim \mathcal{M}_\bullet$  is isomorphic to  $\varinjlim [\mathcal{R}^q_{\mathcal{M}_\bullet}]$  and  $\mathcal{R}^q_{\varinjlim \mathcal{M}_\bullet}(X)$  is isomorphic to  $\varinjlim [\mathcal{R}^q_{\mathcal{M}_\bullet}](X)$ .  $\square$

#### (7.4.29) uniqueness of the coboundary maps

We have constructed a cohomology  $\{H^q\}$  that has the characteristic properties, and we have shown that the functors  $H^q$  are unique. We haven't shown that the coboundary maps  $\delta^q$  that appear in the cohomology sequences (7.1.3) are unique. To make it clear that there is something to show, we note that the cohomology sequence (7.1.3) remains exact when some of the coboundary maps  $\delta^q$  are multiplied by  $-1$ . Why can't we define a new collection of coboundary maps by changing some signs? The reason we can't do this is that we used the coboundary maps  $\delta^q$  in (7.4.7) and (7.4.8) to identify  $H^q(X, \mathcal{M})$ . Having done that, we aren't allowed to change  $\delta^q$  for the particular short exact sequences (7.4.2). We show that the coboundary maps for those particular sequences determine the coboundary maps for every short exact sequence of  $\mathcal{O}$ -modules

$$(A) \quad 0 \rightarrow \mathcal{M} \rightarrow \mathcal{N} \rightarrow \mathcal{P} \rightarrow 0$$

The sequences (7.4.2) were rewritten as (7.4.10). We will use that form.

To show that the coboundaries for the sequence (A) are determined uniquely, we relate it to a sequence (B) for which the coboundary maps are fixed:

$$(B) \quad 0 \rightarrow \mathcal{M} \longrightarrow \mathcal{R}_{\mathcal{M}}^0 \longrightarrow \mathcal{M}^1 \rightarrow 0$$

We map (A) and (B) to a third short exact sequence

$$(C) \quad 0 \rightarrow \mathcal{M} \xrightarrow{\psi} \mathcal{R}_{\mathcal{N}}^0 \longrightarrow \mathcal{Q} \rightarrow 0$$

where  $\psi$  is the composition of the injective maps  $\mathcal{M} \rightarrow \mathcal{R}_{\mathcal{M}}^0 \rightarrow \mathcal{R}_{\mathcal{N}}^0$  and  $\mathcal{Q}$  is the cokernel of  $\psi$ .

First, we inspect the diagram

$$\begin{array}{ccccc} (A) & \mathcal{M} & \longrightarrow & \mathcal{N} & \longrightarrow & \mathcal{P} \\ & \parallel & & \downarrow & & \downarrow \\ (C) & \mathcal{M} & \xrightarrow{\psi} & \mathcal{R}_{\mathcal{N}}^0 & \longrightarrow & \mathcal{Q} \end{array}$$

and its diagram of coboundary maps

$$\begin{array}{ccc} (A) & H^q(X, \mathcal{P}) & \xrightarrow{\delta_A^q} H^{q+1}(X, \mathcal{M}) \\ & \downarrow & \parallel \\ (C) & H^q(X, \mathcal{Q}) & \xrightarrow{\delta_C^q} H^{q+1}(X, \mathcal{M}) \end{array}$$

This diagram shows that the coboundary map  $\delta_A^q$  for the sequence (A) is determined by the coboundary map  $\delta_C^q$  for (C).

Next, we inspect the diagram

$$(7.4.30) \quad \begin{array}{ccccc} (B) & \mathcal{M} & \longrightarrow & \mathcal{R}_{\mathcal{M}}^0 & \longrightarrow & \mathcal{M}^1 \\ & \parallel & & u \downarrow & & v \downarrow \\ (C) & \mathcal{M} & \xrightarrow{\psi} & \mathcal{R}_{\mathcal{N}}^0 & \longrightarrow & \mathcal{Q} \end{array}$$

and its diagram of coboundary maps

$$\begin{array}{ccc} (B) & H^q(X, \mathcal{M}^1) & \xrightarrow{\delta_B^q} H^{q+1}(X, \mathcal{M}) \\ & v \downarrow & \parallel \\ (C) & H^q(X, \mathcal{Q}) & \xrightarrow{\delta_C^q} H^{q+1}(X, \mathcal{M}) \end{array}$$

When  $q > 0$ ,  $\delta_C^q$  and  $\delta_B^q$  are bijective because the cohomology of  $\mathcal{R}_{\mathcal{M}}^0$  and  $\mathcal{R}_{\mathcal{N}}^0$  is zero in positive dimension. Then  $\delta_C^q$  is uniquely determined by  $\delta_B^q$ , and so is  $\delta_A^q$ .

We have to look more closely to settle the case  $q = 0$ . The maps labeled  $u$  and  $v$  in (7.4.30) are injective, and the Snake Lemma shows that their cokernels are isomorphic. We write both of them as  $\mathcal{R}_{\mathcal{P}}^0$ . When we add the cokernels to the diagram, we obtain a cohomology diagram whose relevant part is

$$\begin{array}{ccccccc} (B) & H^0(X, \mathcal{R}_{\mathcal{M}}^0) & \longrightarrow & H^0(X, \mathcal{M}^1) & \xrightarrow{\delta_B^0} & H^1(X, \mathcal{M}) & \\ & u \downarrow & & \downarrow v & & \parallel & \\ (C) & H^0(X, \mathcal{R}_{\mathcal{N}}^0) & \xrightarrow{\beta} & H^0(X, \mathcal{Q}) & \xrightarrow{\delta_C^0} & H^1(X, \mathcal{M}) & \\ & \downarrow \gamma & & \downarrow & & & \\ & H^0(X, \mathcal{R}_{\mathcal{P}}^0) & \xlongequal{\quad} & H^0(X, \mathcal{R}_{\mathcal{P}}^0) & & & \end{array}$$

The rows and columns in the diagram are exact. We want to show that the map  $\delta_C^0$  is determined uniquely by  $\delta_B^0$ . It is determined by  $\delta_B^0$  on the image of  $v$  and it is zero on the image of  $\beta$ . To show that  $\delta_C^0$  is determined by  $\delta_B^0$ , it suffices to show that the images of  $v$  and  $\beta$  together span  $H^0(X, \mathcal{Q})$ . This follows from the fact that  $\gamma$  is surjective. Thus  $\delta_C^0$  is determined uniquely by  $\delta_B^0$ , and so is  $\delta_A^0$ .  $\square$

## 7.5 Cohomology of the Twisting Modules

We determine the cohomology of the twisting modules  $\mathcal{O}(d)$  on  $\mathbb{P}^n$  here. As we will see,  $H^q(\mathbb{P}^n, \mathcal{O}(d))$  is zero for most values of  $q$ . This will help to determine the cohomology of other modules.

Lemma 7.4.18 about vanishing of cohomology on an affine variety, and Lemma 7.4.25 about the direct image via an affine morphism, were stated using a particular affine covering. Since we know that cohomology is unique, that particular covering is irrelevant. Though it isn't necessary, we restate the lemmas here as a corollary:

**7.5.1. Corollary. (i)** *On an affine variety  $X$ ,  $H^q(X, \mathcal{M}) = 0$  for all  $\mathcal{O}$ -modules  $\mathcal{M}$  and all  $q > 0$ .*

**(ii)** *Let  $Y \xrightarrow{f} X$  be an affine morphism. If  $\mathcal{N}$  is an  $\mathcal{O}_Y$ -module, then  $H^q(X, f_*\mathcal{N})$  and  $H^q(Y, \mathcal{N})$  are isomorphic. If  $Y$  is an affine variety,  $H^q(X, f_*\mathcal{N}) = 0$  for all  $q > 0$ .  $\square$*

One case in which **(ii)** applies is that  $f$  is the inclusion of a closed subvariety  $Y$  into  $X$ .

Let  $\mathcal{M}$  be a finite  $\mathcal{O}$ -module on projective space  $\mathbb{P}^n$ . The twisting modules  $\mathcal{O}(d)$  and the twists  $\mathcal{M}(d) = \mathcal{M} \otimes_{\mathcal{O}} \mathcal{O}(d)$  are isomorphic to the modules  $\mathcal{O}(dH)$  and  $\mathcal{M}(dH) = \mathcal{M} \otimes_{\mathcal{O}} \mathcal{O}(dH)$ , respectively. They form maps of directed sets

$$\begin{array}{ccccccc} \mathcal{O} & \xrightarrow{\subset} & \mathcal{O}(H) & \xrightarrow{\subset} & \mathcal{O}(2H) & \xrightarrow{\subset} & \dots \\ 1 \downarrow & & x_0 \downarrow & & x_0^2 \downarrow & & \\ \mathcal{O} & \xrightarrow{x_0} & \mathcal{O}(1) & \xrightarrow{x_0} & \mathcal{O}(2) & \xrightarrow{x_0} & \dots \end{array}, \quad \begin{array}{ccccccc} \mathcal{M} & \longrightarrow & \mathcal{M}(H) & \longrightarrow & \mathcal{M}(2H) & \longrightarrow & \dots \\ 1 \downarrow & & x_0 \downarrow & & x_0^2 \downarrow & & \\ \mathcal{M} & \xrightarrow{x_0} & \mathcal{M}(1) & \xrightarrow{x_0} & \mathcal{M}(2) & \xrightarrow{x_0} & \dots \end{array}$$

(See (??)). The second diagram is obtained from the first one by tensoring with  $\mathcal{M}$ . Let  $\mathbb{U}$  denote the standard affine open subset  $\mathbb{U}^0$  of  $\mathbb{P}^n$ , and let  $j$  be the inclusion of  $\mathbb{U}$  into  $\mathbb{P}^n$ . Then  $\varinjlim \mathcal{O}(dH) \approx j_*\mathcal{O}_{\mathbb{U}}$  (??) and because  $\varinjlim$  is compatible with tensor products,  $\varinjlim \mathcal{M}(dH) \approx j_*\mathcal{M}_{\mathbb{U}}$ . Since  $j$  is an affine morphism and  $\mathbb{U}^0$  is an affine open set,  $H^q(\mathbb{P}^n, j_*\mathcal{O}_{\mathbb{U}}) = 0$  and  $H^q(\mathbb{P}^n, j_*\mathcal{M}_{\mathbb{U}}) = 0$  for all  $q > 0$ .

The next corollary follows from the facts that  $\mathcal{M}(d)$  is isomorphic to  $\mathcal{M}(dH)$ , and that cohomology is compatible with direct limits (7.4.28).

**7.5.2. Corollary.** *For all projective varieties  $X$  and all  $\mathcal{O}$ -modules  $\mathcal{M}$ ,  $\varinjlim H^q(X, \mathcal{O}(d)) = 0$  and  $\varinjlim H^q(X, \mathcal{M}(d)) = 0$  when  $q > 0$ .  $\square$*

**7.5.3. Notation.** If  $\mathcal{M}$  is an  $\mathcal{O}$ -module, we denote the dimension of  $H^q(X, \mathcal{M})$  by  $\mathbf{h}^q(\mathcal{M})$  or by  $\mathbf{h}^q(X, \mathcal{M})$ . We can write  $\mathbf{h}^q(\mathcal{M}) = \infty$  if the dimension is infinite. However, in Section 7.8, we will see that when  $\mathcal{M}$  is a finite  $\mathcal{O}$ -module on a projective variety  $X$ ,  $H^q(X, \mathcal{M})$  has finite dimension for every  $q$ .

**7.5.4. Theorem.**

**(i)** *For  $d \geq 0$ ,  $\mathbf{h}^0(\mathbb{P}^n, \mathcal{O}(d)) = \binom{d+n}{n}$  and  $\mathbf{h}^q(\mathbb{P}^n, \mathcal{O}(d)) = 0$  if  $q \neq 0$ .*

**(ii)** *For  $r > 0$ ,  $\mathbf{h}^n(\mathbb{P}^n, \mathcal{O}(-r)) = \binom{r-1}{n}$  and  $\mathbf{h}^q(\mathbb{P}^n, \mathcal{O}(-r)) = 0$  if  $q \neq n$ .*

In particular, part **(ii)** implies that  $\mathbf{h}^q(\mathbb{P}^n, \mathcal{O}(-1)) = 0$  for all  $q$ .

*proof.* We have described the global sections of  $\mathcal{O}(d)$  before: If  $d \geq 0$ ,  $H^0(X, \mathcal{O}(d))$  is the space of homogeneous polynomials of degree  $d$  in the coordinate variables, and if  $d < 0$ ,  $H^0(X, \mathcal{O}(d)) = 0$  (see (6.6.2)).

**(i)** *(the case  $d \geq 0$ )*

Let  $X = \mathbb{P}^n$ , and let  $Y \xrightarrow{i} X$  be the inclusion of the hyperplane at infinity into  $X$ . By induction on  $n$ , we may assume that the theorem has been proved for  $Y$ , which is a projective space of dimension  $n-1$ . We consider the exact sequence

$$(7.5.5) \quad 0 \rightarrow \mathcal{O}_X(-1) \xrightarrow{x_0} \mathcal{O}_X \rightarrow i_*\mathcal{O}_Y \rightarrow 0$$

and its twists

$$(7.5.6) \quad 0 \rightarrow \mathcal{O}_X(d-1) \xrightarrow{x_0} \mathcal{O}_X(d) \rightarrow i_*\mathcal{O}_Y(d) \rightarrow 0$$

The twisted sequences are exact because they are obtained by tensoring (7.5.5) with the invertible  $\mathcal{O}$ -modules  $\mathcal{O}(d)$ . Because the inclusion  $i$  of  $Y$  into  $X$  is an affine morphism,  $H^q(X, i_*\mathcal{O}_Y(d)) \approx H^q(Y, \mathcal{O}_Y(d))$ .

The monomials of degree  $d$  in  $n+1$  variables form a basis of the space of global sections of  $\mathcal{O}_X(d)$ . Setting  $x_0 = 0$  and deleting terms that become zero gives us a basis of  $\mathcal{O}_Y(d)$ . Therefore every global section of  $\mathcal{O}_Y(d)$  is the restriction of a global section of  $\mathcal{O}_X(d)$ . The sequence of global sections

$$0 \rightarrow H^0(X, \mathcal{O}_X(d-1)) \xrightarrow{x_0} H^0(X, \mathcal{O}_X(d)) \rightarrow H^0(Y, \mathcal{O}_Y(d))$$

is exact, and it remains exact when a zero is added on the right. This tells us that the map

$$H^1(X, \mathcal{O}_X(d-1)) \rightarrow H^1(X, \mathcal{O}_X(d))$$

is injective. By induction on  $n$ ,  $H^q(Y, \mathcal{O}_Y(d)) = 0$  for  $d \geq 0$  and  $q > 0$ . When combined with the injectivity noted above, the cohomology sequence of (7.5.6) gives us bijections  $H^q(X, \mathcal{O}_X(d-1)) \rightarrow H^q(X, \mathcal{O}_X(d))$  for every  $q > 0$ . Since the limits are zero (7.5.2),  $H^q(X, \mathcal{O}_X(d)) = 0$  for all  $d \geq 0$  and all  $q > 0$ .

(ii) (the case  $d < 0$ )

We use induction on the integers  $r$  and  $n$ . We suppose the theorem proved for  $r$ , and we substitute  $d = -r$  into the sequence (7.5.6):

$$(7.5.7) \quad 0 \rightarrow \mathcal{O}_X(-(r+1)) \xrightarrow{x_0} \mathcal{O}_X(-r) \rightarrow i_*\mathcal{O}_Y(-r) \rightarrow 0$$

The base case  $r = 0$  is the exact sequence (7.5.5). In the cohomology sequence associated to that sequence, the terms  $H^q(X, \mathcal{O}_X)$  and  $H^q(Y, \mathcal{O}_Y)$  are zero when  $q > 0$ , and  $H^0(X, \mathcal{O}_X) = H^0(Y, \mathcal{O}_Y) = \mathbb{C}$ . Therefore

$$(7.5.8) \quad H^q(X, \mathcal{O}_X(-1)) = 0 \text{ for every } q.$$

This proves (ii) for  $r = 1$ .

Our induction hypothesis is that,  $\mathbf{h}^n(\mathbb{P}^n, \mathcal{O}(-r)) = \binom{r-1}{n}$  and  $\mathbf{h}^q = 0$  if  $q \neq n$ . By induction on  $n$ , we may suppose that  $\mathbf{h}^{n-1}(\mathbb{P}^{n-1}, \mathcal{O}(-r)) = \binom{r-1}{n-1}$  and  $\mathbf{h}^q = 0$  if  $q \neq n-1$ . Instead of displaying the cohomology sequence associated to (7.5.7), we assemble the dimensions of cohomology into a table in which the asterisks stand for entries that are to be determined:

$$(7.5.9) \quad \begin{array}{cccc} & \mathcal{O}_X(-(r+1)) & \mathcal{O}_X(-r) & i_*\mathcal{O}_Y(-r) \\ \hline \mathbf{h}^0 & : & * & 0 \\ & \vdots & \vdots & \vdots \\ \mathbf{h}^{n-2} & : & * & 0 \\ \mathbf{h}^{n-1} & : & * & \binom{r-1}{n-1} \\ \mathbf{h}^n & : & \mathbf{h}^n(\mathcal{O}(-(r+1))) & \binom{r-1}{n} \end{array}$$

The second column is determined by induction on  $r$  and the third by induction on  $n$ . The cohomology sequence shows that the entries labeled with an asterisk are zero, and that

$$\mathbf{h}^n(\mathbb{P}^n, \mathcal{O}(-(r+1))) = \binom{r-1}{n-1} + \binom{r-1}{n}$$

The right side of this equation is equal to  $\binom{r}{n}$ . □

## 7.6 Cohomology of Hypersurfaces

We determine the cohomology of a plane projective curve first. Let  $X = \mathbb{P}^2$  and let  $C \xrightarrow{i} X$  denote the inclusion of a plane curve of degree  $k$ . The ideal  $\mathcal{I}$  of functions that vanish on  $C$  is isomorphic to the twisting module  $\mathcal{O}_X(-k)$  (6.6.8), so one has an exact sequence

$$(7.6.1) \quad 0 \rightarrow \mathcal{O}_X(-k) \rightarrow \mathcal{O}_X \rightarrow i_*\mathcal{O}_C \rightarrow 0$$

We form a table showing dimensions of the cohomology. Theorem 7.5.4 determines the first two columns, and the cohomology sequence determines the last column.

$$(7.6.2) \quad \begin{array}{rclcl} & \mathcal{O}_X(-k) & \mathcal{O}_X & i_*\mathcal{O}_C & \\ \mathbf{h}^0 : & 0 & 1 & 1 & \\ \mathbf{h}^1 : & 0 & 0 & \binom{k-1}{2} & \\ \mathbf{h}^2 : & \binom{k-1}{2} & 0 & 0 & \end{array}$$

Since the inclusion of the curve  $C$  into the projective plane  $X$  is an affine morphism,  $\mathbf{h}^q(X, i_*\mathcal{O}_C) = \mathbf{h}^q(C, \mathcal{O}_C)$ . Therefore

$$\mathbf{h}^0(C, \mathcal{O}_C) = 1, \quad \mathbf{h}^1(C, \mathcal{O}_C) = \binom{k-1}{2}, \quad \text{and } \mathbf{h}^q = 0 \text{ when } q > 1.$$

The dimension  $\mathbf{h}^1(C, \mathcal{O}_C)$ , which is  $\binom{k-1}{2}$ , is called the *arithmetic genus* of  $C$ . It is denoted by  $p_a$ . We will see later (8.9.2) that when  $C$  is a smooth curve, its arithmetic genus is equal to its topological genus:  $p_a = g$ , but the arithmetic genus of a plane curve of degree  $k$  is  $\binom{k-1}{2}$  also when  $C$  is singular.

We restate the results as a corollary.

**7.6.3. Corollary.** *Let  $C$  be a plane curve of degree  $k$ . Then  $\mathbf{h}^0(C, \mathcal{O}_C) = 1$ ,  $\mathbf{h}^1(C, \mathcal{O}_C) = \binom{k-1}{2} = p_a$ , and  $\mathbf{h}^q = 0$  if  $q \neq 0, 1$ .  $\square$*

The fact that  $\mathbf{h}^0(C, \mathcal{O}_C) = 1$  tells us that the only rational functions that are regular everywhere on  $C$  are the constants. This reflects a fact that will be proved later: A plane curve is compact and connected in the classical topology. However, it isn't a proof of that fact.

We will need more technique in order to compute cohomology for curves in higher dimensional projective spaces. In the next section we will see that the cohomology on any projective curve is zero except in dimensions 0 and 1. Cohomology of projective curves is the topic of Chapter 8.

One can make a similar computation for the hypersurface  $Y$  in  $X = \mathbb{P}^n$  defined by an irreducible homogeneous polynomial  $f$  of degree  $k$ . The ideal of  $Y$  is isomorphic to  $\mathcal{O}_X(-k)$ , and there is an exact sequence

$$0 \rightarrow \mathcal{O}_X(-k) \xrightarrow{f} \mathcal{O}_X \rightarrow i_*\mathcal{O}_Y \rightarrow 0$$

Since we know the cohomology of  $\mathcal{O}_X(-k)$  and  $\mathcal{O}_X$ , and since  $H^q(X, i_*\mathcal{O}_Y) \approx H^q(Y, \mathcal{O}_Y)$ , we can use this sequence to compute the dimensions of the cohomology of  $\mathcal{O}_Y$ .

**7.6.4. Corollary.** *Let  $Y$  be a hypersurface of dimension  $d$  and degree  $k$  in a projective space of dimension  $d+1$ . Then  $\mathbf{h}^0(Y, \mathcal{O}_Y) = 1$ ,  $\mathbf{h}^d(Y, \mathcal{O}_Y) = \binom{k-1}{d+1}$ , and  $\mathbf{h}^q(Y, \mathcal{O}_Y) = 0$  for all other  $q$ .  $\square$*

If  $S$  is a surface in  $\mathbb{P}^3$  defined by an irreducible polynomial of degree  $k$ , then  $\mathbf{h}^0(S, \mathcal{O}_S) = 1$ ,  $\mathbf{h}^1(S, \mathcal{O}_S) = 0$ ,  $\mathbf{h}^2(S, \mathcal{O}_S) = \binom{k-1}{3}$ , and  $\mathbf{h}^q = 0$  if  $q > 2$ . When a projective surface  $S$  isn't embedded into  $\mathbb{P}^3$ , it is still true that  $\mathbf{h}^q = 0$  when  $q > 2$ , but  $\mathbf{h}^1(S, \mathcal{O}_S)$  may be nonzero. The dimensions  $\mathbf{h}^1(S, \mathcal{O}_S)$  and  $\mathbf{h}^2(S, \mathcal{O}_S)$  are invariants of the surface somewhat analogous to the genus of a curve. In classical terminology,  $\mathbf{h}^2(S, \mathcal{O}_S)$  is the *geometric genus*  $p_g$  and  $\mathbf{h}^1(S, \mathcal{O}_S)$  is the *irregularity*  $q$ . The *arithmetic genus*  $p_a$  is

$$(7.6.5) \quad p_a = \mathbf{h}^2(S, \mathcal{O}_S) - \mathbf{h}^1(S, \mathcal{O}_S) = p_g - q$$

Therefore the irregularity is  $q = p_g - p_a$ . When  $S$  is a surface in  $\mathbb{P}^3$ ,  $q = 0$  and  $p_g = p_a$ .

In modern terminology, it is more natural to replace the arithmetic genus by the Euler characteristic  $\chi(\mathcal{O}_S) = \sum_q (-1)^q \mathbf{h}^q(\mathcal{O}_S)$ . The Euler characteristic of a curve is

$$\chi(\mathcal{O}_C) = \mathbf{h}^0(C, \mathcal{O}_C) - \mathbf{h}^1(C, \mathcal{O}_C) = 1 - p_a$$

and the Euler characteristic of a surface  $S$  is

$$\chi(\mathcal{O}_S) = \mathbf{h}^0(S, \mathcal{O}_S) - \mathbf{h}^1(S, \mathcal{O}_S) + \mathbf{h}^2(S, \mathcal{O}_S) = 1 + p_a$$

But because of tradition, the arithmetic genus is still used quite often.

## 7.7 Annihilators and Support

### (7.7.1) annihilators

Let  $A$  be a ring, and let  $m$  be an element of an  $A$ -module  $M$ . The *annihilator*  $I$  of an element  $m$  of  $M$  is the set of elements  $\alpha$  of  $A$  such that  $\alpha m = 0$ . This is an ideal of  $A$  that we may denote by  $\text{ann}(m)$ .

The annihilator of the  $A$ -module  $M$  is the set of elements of  $A$  such that  $aM = 0$ . This annihilator is also an ideal. We will use the annihilator of a module only in the definition of support, which is below.

**7.7.2. Lemma.** *Let  $I$  be the annihilator of an element  $m$  of  $M$ , and let  $s$  be a nonzero element of  $A$ . The annihilator of the image of  $m$  in the localized module  $M_s$  is the localized ideal  $I_s$*   $\square$

This allows us to extend the concept of annihilator to sections of a finite  $\mathcal{O}$ -module on a variety  $X$ .

### (7.7.3) maximal annihilators

Let  $m$  be an element of a module  $M$  over a noetherian ring  $A$ , and let  $I = \text{ann}(m)$ :  $I = \{a \in A \mid am = 0\}$ . The same ideal  $I$  will annihilate every element of the submodule  $Am$  spanned by  $m$ .

Let  $\mathcal{S}$  be the set whose members are the annihilators of nonzero elements of  $M$ . A *maximal annihilator* is a maximal member of  $\mathcal{S}$ . Because  $A$  is noetherian, the annihilator of any nonzero element will be contained in a maximal annihilator.

**7.7.4. Proposition.** *Let  $M$  be a finite, module over a noetherian ring  $A$ , not the zero module..*

(i) *Let  $m$  be a nonzero element of  $M$  whose annihilator  $P$  is a maximal annihilator, and let  $b$  be an element of  $A$ . Then  $P$  is also the annihilator of  $bm$ , if  $bm \neq 0$ .*

(ii) *The maximal annihilators are prime ideals.*

(iii) *Let  $P_i$  be distinct maximal annihilators, let  $m_i$  be elements of  $M$  whose annihilators are  $P_i$ , and let  $N_i = Am_i$ . The submodules  $N_i$  of  $M$  are independent, i.e., their sum  $\sum N_i$  is the direct sum  $\bigoplus N_i$ .*

(iv) *The set of maximal annihilators is finite and nonempty.*

*proof.* Let's denote by  $\text{ann}(m)$  the annihilator of an element  $m$  of  $M$ .

(i) An element  $a$  that annihilates  $m$  also annihilates  $bm$ . So  $\text{ann}(m) \subset \text{ann}(bm)$ . If  $\text{ann}(m)$  is maximal and  $bm \neq 0$ , then  $\text{ann}(m) = \text{ann}(bm)$ .

(ii) Let  $P = \text{ann}(m)$  be a maximal annihilator, and let  $a, b$  be elements of  $A$  such that  $ab \in P$ . If  $b \notin P$ , then  $bm \neq 0$  but  $abm = 0$ . So  $a \in \text{ann}(bm) = P$ .

(iii) We must show that if  $n_i$  are elements of  $N_i$  such that  $n_1 + \cdots + n_k = 0$ , then  $n_i = 0$  or all  $i$ . We use induction on  $k$ . Let  $a$  be a nonzero element of the annihilator  $P_k$  of  $N_k$ . Then  $an_1 + \cdots + an_{k-1} + 0 = 0$ . By induction  $an_i = 0$  for all  $i$ . Therefore  $a$  is in  $P_i$  for all  $i$ . Unless  $k = 1$ , this contradicts the assumption that  $P_i$  are distinct maximal annihilators.

(iv) The submodules  $N_i$  are nonzero and independent. Since  $\bigoplus N_i$  is a submodule of the finite module  $A$  and since  $A$  is noetherian,  $\bigoplus N_i$  is a finite module. So there can be only finitely many indices  $i$ .  $\square$

**7.7.5. Corollary.** *Let  $M$  be a finite module over a noetherian domain  $A$ , and let  $s$  be an element of  $A$  that isn't contained in any of the maximal annihilators of  $M$ . The multiplication map  $M \xrightarrow{s} M$  is injective, and therefore the map from  $M$  to its localization  $M_s$  is injective.*  $\square$

**7.7.6. Support** Let  $M$  be a finite module over a finite-type domain  $A$  and  $X = \text{Spec } A$ . The *support* of  $M$  is the locus  $C = V_X(I)$  of zeros of its annihilator  $I$  in  $X$ . The support is a closed subset of  $X$ .

and the support of the localization  $M_s$  is the intersection  $C_s = C \cap X_s$ .

Let  $A$  be a finite-type domain and let  $X = \text{Spec } A$ . The *support* of a finite  $A$ -module  $M$  is the locus of zeros  $V_X(I)$  of its annihilator  $I$  in  $X$ . It is the set of points  $p$  of  $X$  such that  $I \subset \mathfrak{m}_p$ .

The annihilator of a localized module  $M_s$  is the localization  $I_s$  of the annihilator  $I$  of  $M$ , and the support of the localization  $M_s$  is the intersection  $C_s = C \cap X_s$ .

The *support* of a finite  $\mathcal{O}_X$ -module is the closed subset  $V_X(\mathcal{I})$  of points such that  $\mathcal{I} \subset \mathfrak{m}_p$ .

For example, the support of the residue field module  $\kappa_p$  is the point  $p$ . The support of the maximal ideal  $\mathfrak{m}_p$  at  $p$  is the whole variety  $X$ .

### (7.7.7) $\mathcal{O}$ -modules with support of dimension zero

**7.7.8. Proposition.** *Let  $\mathcal{M}$  be a finite  $\mathcal{O}$ -module on a variety  $X$ .*

(i) *Suppose that the support of  $\mathcal{M}$  is a single point  $p$ , let  $M = \mathcal{M}(X)$ , and let  $U$  be an affine open subset of  $X$ . If  $U$  contains  $p$ , then  $\mathcal{M}(U) = M$ , and if  $U$  doesn't contain  $p$ , then  $\mathcal{M}(U) = 0$ .*

(ii) *(Chinese Remainder Theorem) If the support of  $\mathcal{M}$  is a finite set  $\{p_1, \dots, p_k\}$ , then  $\mathcal{M}$  is the direct sum  $\mathcal{M}_1 \oplus \dots \oplus \mathcal{M}_k$  of  $\mathcal{O}$ -modules supported at the points  $p_i$ .*

*proof.* (i) Let  $\mathcal{I}$  be the annihilator of  $\mathcal{M}$ . The locus  $V_X(\mathcal{I})$  is  $p$ . If  $p$  isn't contained in  $U$ , then when we restrict  $\mathcal{M}$  to  $U$ , we obtain an  $\mathcal{O}_U$ -module whose support is empty. Therefore the restriction to  $U$  is the zero module.

Next, suppose that  $p$  is contained in  $U$ , and let  $V$  denote the complement of  $p$  in  $X$ . We cover  $X$  by a set  $\{U^i\}$  of affine open sets with  $U = U^1$ , and such that  $U^i \subset V$  if  $i > 1$ . By what has been shown,  $\mathcal{M}(U^i) = 0$  if  $i > 1$  and  $\mathcal{M}(U^{ij}) = 0$  if  $j \neq i$ . The sheaf axiom for this covering shows that  $\mathcal{M}(X) \approx \mathcal{M}(U)$ .

(ii) This follows from the ordinary Chinese Remainder Theorem.  $\square$

## 7.8 Three Theorems about Cohomology

We will use the concept of the *support* of an  $\mathcal{O}$ -module  $\mathcal{M}$ , the zero set of its annihilator.

**7.8.1. Theorem.** *Let  $X$  be a projective variety, and let  $\mathcal{M}$  be a finite  $\mathcal{O}_X$ -module.*

(i) *If the support of  $\mathcal{M}$  has dimension  $k$ , then  $H^q(X, \mathcal{M}) = 0$  for all  $q > k$ . In particular, if the dimension of  $X$  is  $n$ , then  $H^q(X, \mathcal{M}) = 0$  for all  $q > n$ .*

(ii) *Let  $\mathcal{M}(d)$  be the twist of the finite  $\mathcal{O}_X$ -module  $\mathcal{M}$ . For sufficiently large  $d$ ,  $H^q(X, \mathcal{M}(d)) = 0$  for all  $q > 0$ .*

(iii) *For every  $q$ , the cohomology  $H^q(X, \mathcal{M})$  is a finite-dimensional vector space.*

**7.8.2. Notes.** (a) The structure of the proofs is interesting. The first part allows us to use *descending* induction to prove the second and third parts, beginning with the fact that  $\mathbf{h}^k(\mathcal{M}) = 0$  when  $k$  is larger  $\dim X$ . The descending induction step is to prove that if a statement  $S_k$  is true when  $k = r + 1$ , then it is true when  $k = r$ .

The third part of the theorem tells us that, when  $\mathcal{M}$  is a finite  $\mathcal{O}$ -module, the space  $H^0(X, \mathcal{M})$  of global sections is finite-dimensional. This is one of the most important consequences of the theorem, and it isn't easy to prove directly.

(b) Let  $X$  be a projective variety. The highest dimension in which cohomology of an  $\mathcal{O}_X$ -module can be nonzero is called the *cohomological dimension* of  $X$ . Theorem 7.8.1 shows that its cohomological dimension is at most its algebraic dimension. In fact, it is equal to the algebraic dimension. On the other hand,  $X$  has dimension  $2n$  in the classical topology, and the constant coefficient cohomology  $H_{class}^{2n}(X, \mathbb{Z})$  in the classical topology will be nonzero. In the classical topology, the cohomological dimension of a projective variety  $X$  is its topological dimension  $2n$ . In the Zariski topology,  $H^q(X, \mathbb{Z})$  is zero for every  $q > 0$ .  $\square$

In the theorem, we are given that  $X$  is a closed subvariety of a projective space  $\mathbb{P}^n$ . We can replace an  $\mathcal{O}_X$ -module by its extension by zero (7.5.1). This doesn't change the cohomology. So we may assume that  $X$  is a projective space.

The proofs are based on the cohomology of the twisting modules (7.5.4), the vanishing of the limit  $\varinjlim H^q(X, \mathcal{M}(d))$  for  $q > 0$  (7.5.2), and on two exact sequences. As we know,  $\mathcal{M}(r)$  is generated by global sections if  $r$  is sufficiently large (6.6.20). Choosing generators gives us a surjective map  $\mathcal{O}^m \rightarrow \mathcal{M}(r)$ . Let  $\mathcal{N}$  be the kernel of this map. When we twist the sequence  $0 \rightarrow \mathcal{N} \rightarrow \mathcal{O}^m \rightarrow \mathcal{M}(r) \rightarrow 0$ , we obtain short exact sequences

$$(7.8.3) \quad 0 \rightarrow \mathcal{N}(d) \rightarrow \mathcal{O}(d)^m \rightarrow \mathcal{M}(d+r) \rightarrow 0$$

for every  $d \geq 0$ . These sequences are useful because we know that  $H^q(X, \mathcal{O}(d)) = 0$  when  $q > 0$ .

Next, Lemma ?? tells us that, with coordinates in general position, there will be an exact sequence  $0 \rightarrow \mathcal{M}(-1) \xrightarrow{x_0} \mathcal{M} \rightarrow \overline{\mathcal{M}} \rightarrow 0$ , where  $\overline{\mathcal{M}}$  is the quotient  $\mathcal{M}/x_0\mathcal{M}(-1)$ . Twisting this sequence gives us exact sequences

$$(7.8.4) \quad 0 \rightarrow \mathcal{M}(d-1) \xrightarrow{x_0} \mathcal{M}(d) \rightarrow \overline{\mathcal{M}}(d) \rightarrow 0$$

Since the zero locus of  $x_0$  is the hyperplane  $H$  at infinity, the support  $S$  of  $\overline{\mathcal{M}}$  will be contained in  $S \cap H$ . If  $S$  has dimension  $k$  and  $x_0$  is generic, the support of  $\overline{\mathcal{M}}$  will have dimension less than  $k$ . This will allow us to use induction on  $k$  and  $d$ .

*proof of Theorem 7.8.1 (i) (vanishing in large dimension)*

We inspect the sequence (7.8.4). Let  $k$  be the dimension of the support of  $\mathcal{M}$ . If  $k = 0$ , then  $\overline{\mathcal{M}} = 0$  and  $H^q(X, \overline{\mathcal{M}}(d)) = 0$  for all  $q$ . With coordinates in general position, the support of  $\overline{\mathcal{M}}$  will have dimension at most  $k-1$ , if  $k > 0$ . So by induction on  $k$ , we may assume that  $H^q(X, \overline{\mathcal{M}}(d)) = 0$  for all  $q > k-1$  and all  $d$ .

The cohomology sequence associated to the sequence (7.8.4) is

$$(7.8.5) \quad \cdots \rightarrow H^{q-1}(X, \overline{\mathcal{M}}(d)) \xrightarrow{\delta^{q-1}} H^q(X, \mathcal{M}(d-1)) \xrightarrow{x_0} H^q(X, \mathcal{M}(d)) \rightarrow H^q(X, \overline{\mathcal{M}}(d)) \xrightarrow{\delta^q} \cdots$$

When  $q > k$ , the terms on the left and right of this display are zero, and therefore the map

$$H^q(X, \mathcal{M}(d-1)) \xrightarrow{x_0} H^q(X, \mathcal{M}(d))$$

is an isomorphism. According to (7.5.2),  $\varinjlim H^q(X, \mathcal{M}(d)) = 0$ . It follows that  $H^q(X, \mathcal{M}(d)) = 0$  for all  $d$ , and in particular,  $H^q(X, \mathcal{M}) = 0$  when  $q > k$ .

*proof of Theorem 7.8.1 (ii) (vanishing for a large twist)*

We must show this:

(\*) Let  $\mathcal{M}$  be a finite  $\mathcal{O}$ -module. For every  $q > 0$  and for sufficiently large  $d$ ,  $H^q(X, \mathcal{M}(d)) = 0$ .

By part (i), we know that (\*) is true for every  $q > n = \dim X$ , because all cohomology in dimension  $q$  is zero when  $q > n$ . This leaves a finite set of integers  $q = 1, \dots, n$  to consider, and it suffices to consider them one at a time. If (\*) is true for each individual  $q$  there will be a single  $d$  such that it is true for  $q = 1, \dots, n$ , and therefore for all positive integers  $q$ , as the theorem asserts.

We use descending induction on  $q$ , the base case being  $q = n + 1$ , for which (\*) is true. We suppose that (\*) is true for every finite  $\mathcal{O}$ -module  $\mathcal{M}$  when  $q = p + 1$ , and that  $p > 0$ , and we show that (\*) is true for every finite  $\mathcal{O}$ -module  $\mathcal{M}$  when  $q = p$ .

We substitute  $q = p$  into the cohomology sequence associated to the sequence (7.8.3). The relevant part of that sequence is

$$H^p(X, \mathcal{O}(d))^m \rightarrow H^p(X, \mathcal{M}(d+r)) \xrightarrow{\delta^p} H^{p+1}(X, \mathcal{N}(d))$$

Since  $p$  is positive,  $H^p(X, \mathcal{O}(d)) = 0$  for all  $d \geq 0$ , and therefore the map  $\delta^p$  is injective. Our induction hypothesis, applied to the  $\mathcal{O}$ -module  $\mathcal{N}$ , shows that  $H^{p+1}(X, \mathcal{N}(d)) = 0$  for large  $d$ , and then

$$H^p(X, \mathcal{M}(d+r)) = 0$$

The particular integer  $d+r$  isn't useful. Our conclusion is that, for every finite  $\mathcal{O}$ -module  $\mathcal{M}$ ,  $H^p(X, \mathcal{M}(k)) = 0$  when  $k$  is large enough.  $\square$

*proof of Theorem 7.8.1 (iii) (finiteness of cohomology)*

This proof also uses descending induction on  $q$ . As was mentioned above, it isn't easy to prove directly that the space  $H^0(X, \mathcal{M})$  of global sections is finite-dimensional.

We go back to the sequence (7.8.4) and its cohomology sequence (7.8.5). Induction on the dimension of the support of  $\mathcal{M}$  allows us to assume that  $H^r(X, \overline{\mathcal{M}}(d))$  is finite-dimensional for all  $r$ . So, in the part of the cohomology sequence that is depicted in (7.8.5), the terms on the left and right are finite-dimensional.



Therefore  $H^q(X, \mathcal{M}(d-1))$  and  $H^q(X, \mathcal{M}(d))$  are either both finite-dimensional, or else they are both infinite-dimensional, and this is true for every  $d$ .

Suppose that  $q > 0$ . Then  $H^q(X, \mathcal{M}(d)) = 0$  when  $d$  is large enough. Since the zero space is finite-dimensional, we can use the sequence together with descending induction, to conclude that  $H^q(X, \mathcal{M}(d))$  is finite-dimensional for every finite module  $\mathcal{M}$  and every  $d$ . In particular,  $H^q(X, \mathcal{M})$  is finite-dimensional.

This leaves the case that  $q = 0$ . To prove that  $H^0(X, \mathcal{M})$  is finite-dimensional, we set  $d = -r$  in the sequence (7.8.3):

$$0 \rightarrow \mathcal{N}(-r) \rightarrow \mathcal{O}(-r)^m \rightarrow \mathcal{M} \rightarrow 0$$

The corresponding cohomology sequence is

$$0 \rightarrow H^0(X, \mathcal{N}(-r)) \rightarrow H^0(X, \mathcal{O}(-r)^m) \rightarrow H^0(X, \mathcal{M}) \xrightarrow{\delta^0} H^1(X, \mathcal{N}(-r)) \rightarrow \cdots$$

Here  $H^0(X, \mathcal{O}(-r)^m) = 0$ , and we've shown that  $H^1(X, \mathcal{N}(-r))$  is finite-dimensional. It follows that  $H^0(X, \mathcal{M})$  is finite-dimensional, and this completes the proof.  $\square$

Notice that the finiteness of  $H^0$  comes out only at the end. The higher cohomology is essential for the proof.

### (7.8.6) Euler characteristic

Theorem 7.8.1 allows us to define the Euler characteristic of a finite module on projective variety.

**7.8.7. Definition.** Let  $X$  be a projective variety. The *Euler characteristic* of a finite  $\mathcal{O}$ -module  $\mathcal{M}$  is the alternating sum of the dimensions of the cohomology:

$$(7.8.8) \quad \chi(\mathcal{M}) = \sum (-1)^q h^q(X, \mathcal{M}).$$

This makes sense because  $h^q(X, \mathcal{M})$  finite for every  $q$ , and is zero when  $q$  is large.

**7.8.9. Proposition. (i)** If  $0 \rightarrow \mathcal{M} \rightarrow \mathcal{N} \rightarrow \mathcal{P} \rightarrow 0$  is a short exact sequence of finite  $\mathcal{O}$ -modules on a projective variety  $X$ , then  $\chi(\mathcal{M}) - \chi(\mathcal{N}) + \chi(\mathcal{P}) = 0$ .

**(ii)** If  $0 \rightarrow \mathcal{M}_0 \rightarrow \mathcal{M}_1 \rightarrow \cdots \rightarrow \mathcal{M}_n \rightarrow 0$  is an exact sequence of finite  $\mathcal{O}$ -modules on  $X$ , the alternating sum  $\sum (-1)^i \chi(\mathcal{M}_i)$  is zero.

**7.8.10. Lemma.** Let  $0 \rightarrow V^0 \rightarrow V^1 \rightarrow \cdots \rightarrow V^n \rightarrow 0$  be an exact sequence of finite dimensional vector spaces. The alternating sum  $\sum (-1)^q \dim V^q$  is zero.  $\square$

*proof of Proposition 7.8.9. (i)* Since the cohomology sequence associated to the given sequence is exact, the lemma tells us that the alternating sum of its dimensions is zero. That alternating sum is also equal to  $\chi(\mathcal{M}) - \chi(\mathcal{N}) + \chi(\mathcal{P})$ .

**(ii)** Let's denote the given sequence by  $\mathbb{S}_0$  and the alternating sum  $\sum_i \chi(\mathcal{M}_i)$  by  $\chi(\mathbb{S}_0)$ .

Let  $\mathcal{N} = \mathcal{M}_1/\mathcal{M}_0$ . The sequence  $\mathbb{S}_0$  decomposes into the two exact sequences

$$\mathbb{S}_1: 0 \rightarrow \mathcal{M}_0 \rightarrow \mathcal{M}_1 \rightarrow \mathcal{N} \rightarrow 0 \quad \text{and} \quad \mathbb{S}_2: 0 \rightarrow \mathcal{N} \rightarrow \mathcal{M}_2 \rightarrow \cdots \rightarrow \mathcal{M}_k \rightarrow 0 \rightarrow$$

Then  $\chi(\mathbb{S}_0) = \chi(\mathbb{S}_1) - \chi(\mathbb{S}_2)$ , so the assertion follows from **(i)** by induction on  $n$ .  $\square$

## 7.9 Bézout's Theorem

As an application of cohomology, we use it to prove Bézout's Theorem.

Recall that, if  $f(x) = p_1(x)^{e_1} \cdots p_k(x)^{e_k}$  is a factorization of a homogeneous polynomial in  $x = x_0, x_1, x_2$  into irreducible polynomials, the *divisor* of  $f$  is defined to be the integer combination  $e_1 C_1 + \cdots + e_k C_k$ , where  $C_i$  is the curve of zeros of  $p_i$ .

We restate the theorem to be proved.

**7.9.1. Bézout's Theorem.** Let  $Y$  and  $Z$  be the divisors in the projective plane  $X$  defined by relatively prime homogeneous polynomials  $f$  and  $g$  of degrees  $m$  and  $n$ , respectively. The number of intersection points  $Y \cap Z$ , counted with an appropriate multiplicity, is equal to  $mn$ . Moreover, the multiplicity is 1 at a point at which  $Y$  and  $Z$  intersect transversally.

The definition of the multiplicity will emerge during the proof.

**7.9.2. Example.** Suppose that  $f$  and  $g$  are products of linear polynomials, so that  $Y$  is the union of  $m$  lines and  $Z$  is the union of  $n$  lines, and suppose that those lines are distinct. Since two distinct lines intersect transversally in a single point, there are  $mn$  intersection points of multiplicity 1.  $\square$

*proof of Bézout's Theorem.* We will suppress the notation for the extension by zero from a closed subset.

Multiplication by  $f$  defines a short exact sequence

$$0 \rightarrow \mathcal{O}_X(-m) \xrightarrow{f} \mathcal{O}_X \rightarrow \mathcal{O}_Y \rightarrow 0$$

where  $\mathcal{O}_Y$  stands for  $i_*\mathcal{O}_Y$ ,  $i$  being the inclusion  $Y \rightarrow X$ . This sequence describes  $\mathcal{O}_X(-m)$  as the ideal  $\mathcal{I}$  of  $Y$ , and there is a similar sequence describing the module  $\mathcal{O}_X(-n)$  as the ideal  $\mathcal{J}$  of  $Z$ . The zero locus of the ideal  $\mathcal{I} + \mathcal{J}$  is the intersection  $Y \cap Z$ .

We denote the quotient  $\mathcal{O}_X/(\mathcal{I} + \mathcal{J})$  by  $\overline{\mathcal{O}}$ . Since  $f$  and  $g$  have no common factor,  $Y \cap Z$  is a finite set of points  $\{p_1, \dots, p_k\}$ , and  $\overline{\mathcal{O}}$  is isomorphic to a direct sum  $\bigoplus \overline{\mathcal{O}}_i$ , where  $\overline{\mathcal{O}}_i$  is a finite-dimensional algebra whose support is  $p_i$  (7.7.7). We define the *intersection multiplicity* of  $Y$  and  $Z$  at  $p_i$  to be the dimension of  $\overline{\mathcal{O}}_i$ , which is also equal to  $h^0(X, \overline{\mathcal{O}}_i)$ , and we denote the multiplicity by  $\mu_i$ . The dimension of  $H^0(X, \overline{\mathcal{O}})$  is the sum  $\mu_1 + \dots + \mu_k$ , and  $H^q(X, \overline{\mathcal{O}}) = 0$  for all  $q > 0$  (Theorem 7.8.1 (i)). So the Euler characteristic  $\chi(\overline{\mathcal{O}})$  is equal to  $h^0(X, \overline{\mathcal{O}})$ . We'll show that  $\chi(\overline{\mathcal{O}}) = mn$ , and therefore that  $\mu_1 + \dots + \mu_k = mn$ . This will prove Bézout's Theorem.

We form an exact sequence of  $\mathcal{O}$ -modules, in which  $\mathcal{O} = \mathcal{O}_X$ :

$$(7.9.3) \quad 0 \rightarrow \mathcal{O}(-m-n) \xrightarrow{(g,f)^t} \mathcal{O}(-m) \times \mathcal{O}(-n) \xrightarrow{(f,-g)} \mathcal{O} \xrightarrow{\pi} \overline{\mathcal{O}} \rightarrow 0$$

In order to interpret the maps in this sequence as matrix multiplication with homomorphisms acting on the left, sections of  $\mathcal{O}(-m) \times \mathcal{O}(-n)$  should be represented as column vectors  $(u, v)^t$ ,  $u$  and  $v$  being sections of  $\mathcal{O}(-m)$  and  $\mathcal{O}(-n)$ , respectively.

**7.9.4. Lemma.** The sequence (7.9.3) is exact.

*proof.* To prove exactness, it suffices to show that the sequence of sections on each of the standard affine open sets is exact. We look at  $\mathbb{U}^0$ , as usual. Let's suppose  $s$  coordinates are chosen so that none of the points making up  $Y \cap Z$  lie on the coordinate axes. Let  $A$  be the algebra of regular functions on  $\mathbb{U}^0$ , the polynomial algebra  $\mathbb{C}[u_1, u_2]$ , with  $u_i = x_i/x_0$ . We identify  $\mathcal{O}(k)$  with  $\mathcal{O}(kH)$ ,  $H$  being the hyperplane at infinity. The restriction of the module  $\mathcal{O}(kH)$  to  $\mathbb{U}^0$  is isomorphic to  $\mathcal{O}_{\mathbb{U}^0}$ . Its sections on  $\mathbb{U}^0$  are the elements of  $A$ . Let  $\overline{A}$  be the algebra of sections of  $\overline{\mathcal{O}}$  on  $\mathbb{U}^0$ . Since  $f$  and  $g$  are relatively prime, so are their dehomogenizations  $F = f(1, u_1, u_2)$  and  $G = g(1, u_1, u_2)$ . The sequence of sections of (7.9.3) on  $\mathbb{U}^0$  is

$$0 \rightarrow A \xrightarrow{(G,F)^t} A \times A \xrightarrow{(F,-G)} A \rightarrow \overline{A} \rightarrow 0$$

and the only place at which exactness of this sequence isn't obvious is at  $A \times A$ . Suppose that  $(u, v)^t$  is in the kernel of the map  $(F, -G)$ , i.e., that  $Fu = Gv$ . Since  $F$  and  $G$  are relatively prime,  $F$  divides  $v$ ,  $G$  divides  $u$ , and  $v/F = u/G$ . Let  $w = v/F = u/G$ . Then  $(u, v)^t = (G, F)^t w$ .  $\square$

Since cohomology is compatible with products,  $\chi(\mathcal{M} \times \mathcal{N}) = \chi(\mathcal{M}) + \chi(\mathcal{N})$ . Proposition 7.8.9(ii), applied to the exact sequence (7.9.3), tells us that the alternating sum

$$(7.9.5) \quad \chi(\mathcal{O}(-m-n)) - (\chi(\mathcal{O}(-m)) + \chi(\mathcal{O}(-n))) + \chi(\mathcal{O}) - \chi(\overline{\mathcal{O}})$$

is zero. Solving for  $\chi(\overline{\mathcal{O}})$  and applying Theorem 7.5.4,

$$\chi(\overline{\mathcal{O}}) = \binom{n+m-1}{2} - \binom{m-1}{2} - \binom{n-1}{2} + 1$$

This equation shows that the term  $\chi(\overline{\mathcal{O}})$  depends only on the integers  $m$  and  $n$ . Since we know that the answer is  $mn$  when  $Y$  and  $Z$  are unions of distinct lines, it is  $mn$  in every case. This completes the proof.

If you are suspicious of this reasoning, you can evaluate the right side of the equation.  $\square$

We still need to explain the assertion that the multiplicity at a transversal intersection  $p$  is equal to 1. This will be true if and only if  $\mathcal{I} + \mathcal{J}$  generates the maximal ideal at  $p$  locally, and it is obvious when  $Y$  and  $Z$  are lines. In that case we may choose affine coordinates so that  $p$  is the origin in  $\mathbb{A}^2 = \text{Spec } A$ ,  $A = \mathbb{C}[y, z]$  and the curves are the coordinate axes  $\{z = 0\}$  and  $\{y = 0\}$ . The variables  $u, v$  generate the maximal ideal at the origin, so the quotient algebra  $A/(y, z)$  has dimension 1.

Suppose that  $Y$  and  $Z$  intersect transversally at  $p$ , but that they aren't lines. We choose affine coordinates so that  $p$  is the origin and that the tangent directions are the coordinate axes. The affine equations of  $Y$  and  $Z$  will have the form  $y' = 0$  and  $z' = 0$ , where  $y' = y + g(y, z)$  and  $z' = z + h(y, z)$ ,  $g$  and  $h$  being polynomials all of whose terms have degree at least 2. Because  $Y$  and  $Z$  may intersect at points other than  $p$ , the elements  $y'$  and  $z'$  may not generate the maximal ideal at  $p$ . However, it suffices to show that they generate the maximal ideal locally.

Let  $\tilde{A}$  be the local ring of the polynomial ring  $\mathbb{C}[y, z]$  at the origin, and let  $\tilde{\mathfrak{m}}$  and  $\tilde{k}$  be the maximal ideal and residue field of  $\tilde{A}$ , respectively. To show that  $y', z'$  generate  $\tilde{\mathfrak{m}} = (y, z)\tilde{A}$ , the Local Nakayama Lemma 5.1.22 tells us that it suffices to show that their images generate  $\tilde{\mathfrak{m}}/\tilde{\mathfrak{m}}^2$ . The images of  $g$  and  $h$  in  $\tilde{\mathfrak{m}}^2$  are zero, so  $y'$  and  $z'$  are congruent to  $y$  and  $z$  modulo  $\tilde{\mathfrak{m}}^2$ . They do generate  $\tilde{\mathfrak{m}}/\tilde{\mathfrak{m}}^2$ , so they generate  $\tilde{\mathfrak{m}}$ .  $\square$



## Chapter 8 THE RIEMANN-ROCH THEOREM FOR CURVES

- 8.1 Branched Coverings
- 8.2 Modules on a Smooth Curve
- 8.3 Divisors
- 8.4 The Riemann-Roch Theorem I
- 8.5 The Birkhoff-Grothendieck Theorem
- 8.6 Differentials
- 8.7 Trace
- 8.8 The Riemann-Roch Theorem II
- 8.9 Using Riemann-Roch

The topic of this chapter is a classical problem of algebraic geometry, to determine the rational functions on a smooth projective curve with given poles. This can be difficult, and one is usually happy if one can determine the dimension of the space of such functions. The most important tool for this is the Riemann-Roch Theorem.

### 8.1 Branched Coverings

Smooth affine curves were discussed in Chapter ???. An affine curve is smooth if its local rings are valuation rings, or if its coordinate ring is a normal domain. An arbitrary curve is smooth if it has an open covering by smooth affine curves.

An integral morphism  $Y \xrightarrow{\pi} X$  of smooth curves will be called a *branched covering*. It follows from Chevalley's Finiteness Theorem that every nonconstant morphism of smooth projective curves is a branched covering.

If  $Y \rightarrow X$  is a branched covering, the function field  $K$  of  $Y$  will be a finite extension of the function field  $F$  of  $X$ . The *degree* of the covering is the degree  $[K : F]$  of the field extension. It will be denoted by  $[Y : X]$ .

If a branched covering  $Y \rightarrow X$  is given, and if  $X' = \text{Spec } A$  is an affine open subset of  $X$ , its inverse image  $Y'$  will be a smooth affine curve,  $Y' = \text{Spec } B$ , and if the degree  $[Y : X]$  of the covering is  $n$ ,  $B$  will be a locally free  $A$ -module of rank  $[Y : X]$ .

To describe the fibre of a branched covering  $Y \xrightarrow{\pi} X$  over a point  $p$  of  $X$ , we may localize. So we may assume that  $X$  and  $Y$  are affine, say  $X = \text{Spec } A$  and  $Y = \text{Spec } B$ , and that the maximal ideal  $\mathfrak{m}_p$  of  $A$  at a point  $p$  is a principal ideal, generated by an element  $x$  of  $A$ . If a point  $q$  of  $Y$  lies over  $p$ , the *ramification index* at  $q$  is defined to be  $e = v_q(x)$ , where  $v_q$  is the valuation of the function field  $K$  corresponding to  $q$ . Then, if  $y$  is a local generator for the maximal ideal  $\mathfrak{m}_q$  of  $B$  at  $q$ , we will have

$$x = uy^e$$

where  $u$  is a local unit.

The next lemma follows from Lemma 8.2.2 and the Chinese Remainder Theorem.

**8.1.1. Lemma.** *Let  $Y \xrightarrow{\pi} X$  be a branched covering, with  $X = \text{Spec } A$  and  $Y = \text{Spec } B$ . Let  $q_1, \dots, q_k$  be the points of  $Y$  that lie over a point  $p$  of  $X$ , let  $x$  be a generator for the maximal ideal  $\mathfrak{m}_p$  at  $p$ , and let  $\mathfrak{m}_i$  and  $e_i$  be the maximal ideal and ramification index at  $q_i$ , respectively.*

- (i) The extended ideal  $\mathfrak{m}_p B = xB$  is the product ideal  $\mathfrak{m}_1^{e_1} \cdots \mathfrak{m}_k^{e_k}$  of  $B$ .
- (ii) Let  $\overline{B}_i = B/\mathfrak{m}_i^{e_i}$ . The quotient  $\overline{B} = B/xB$  is isomorphic to the product  $\overline{B}_1 \times \cdots \times \overline{B}_k$ .
- (iii) The degree  $[Y:X]$  of the covering is the sum  $e_1 + \cdots + e_k$  of the ramification indices at the points  $q_i$  in the fibre over  $p$ .  $\square$

Points  $q$  of  $Y$  whose ramification indices are greater than one are called *branch points*. We also call a point  $p$  of  $X$  a *branch point* of the covering if there is a branch point  $q$  that lies over  $p$ .

**8.1.2. Lemma.** *A branched covering  $Y \rightarrow X$  has finitely many branch points. If a point  $p$  is not a branch point, the fibre over  $p$  consists of  $n = [Y:X]$  points with ramification indices equal to 1.*

*proof.* We can delete finite sets of points, so we may suppose that  $X$  and  $Y$  are affine,  $X = \text{Spec } A$  and  $Y = \text{Spec } B$ . Then  $B$  is a finite  $A$ -module of rank  $n$ . Let  $F$  and  $K$  be the fraction fields of  $A$  and  $B$ , respectively, and let  $\beta$  be an element of  $B$  that generates the field extension  $K/F$ . Then  $A[\beta] \subset B$ , and since these two rings have the same fraction field, there will be a nonzero element  $s \in A$  such that  $A_s[\beta] = B$ . We may suppose that  $B = A[\beta]$ . Let  $g$  be the monic irreducible polynomial for  $\beta$  over  $A$ . The discriminant of  $g$  is nonzero (1.7.19), so for all but finitely many points  $p$  of  $X$ , there will be  $n$  points of  $Y$  over  $p$  with ramification indices equal to 1.  $\square$

**8.1.3. Corollary.** *A branched covering  $Y \xrightarrow{\pi} X$  of degree one is an isomorphism.*

*proof.* When  $[Y:X] = 1$ , the function fields of  $Y$  and  $X$  are equal. Then, because  $Y \rightarrow X$  is an integral morphism and  $X$  is normal,  $Y = X$ .  $\square$

figure: a branched covering

#### (8.1.4) local analytic structure

The local analytic structure of a branched covering  $Y \xrightarrow{\pi} X$  in the classical topology is very simple. We explain it there because it is useful and helpful for intuition.

Let  $q$  be a point of  $Y$ , let  $p$  be its image in  $X$ , and let  $x$  be a local generator for the maximal ideal of  $X$  at  $p$ . Also, let  $e = v_q(x)$  be the ramification index at  $q$ .

**8.1.5. Proposition.** *In the classical topology,  $Y$  is locally isomorphic to the  $e$ -th root covering  $y^e = x$ .*

*proof.* Let  $z$  be a local generator for the maximal ideal  $\mathfrak{m}_q$  of  $\mathcal{O}_Y$ . If the ramification index is  $e$ , then  $x$  has the form  $uz^e$ , where  $u$  is a local unit at  $q$ . In a neighborhood of  $q$  in the classical topology,  $u$  will have an analytic  $e$ -th root  $w$ . Then  $y = wz$  also generates  $\mathfrak{m}_q$  locally, and  $x = y^e$ . It follows from the implicit function theorem that  $x$  and  $y$  are local analytic coordinate functions on  $X$  and  $Y$  (see (??)).  $\square$

**8.1.6. Corollary.** *Let  $Y \xrightarrow{\pi} X$  be a branched covering, let  $\{q_1, \dots, q_k\}$  be the fibre over a point  $p$  of  $X$ , and let  $e_i$  be the ramification index at  $q_i$ . As a point  $p'$  of  $X$  approaches  $p$ ,  $e_i$  points of the fibre over  $p'$  approach  $q_i$ .  $\square$*

## 8.2 Modules on a Smooth Curve

A *torsion element* of a module  $M$  over a domain  $A$  is an element that is annihilated by some nonzero element  $a$  of  $A$ :  $am = 0$ . The set of torsion elements of  $M$  is its *torsion submodule*, and a module whose torsion submodule is zero is *torsion-free*. These definitions are extended to  $\mathcal{O}$ -modules by applying them to affine open sets.

**8.2.1. Lemma.** *Let  $Y$  be a smooth curve.*

- (i) *A finite  $\mathcal{O}$ -module  $\mathcal{M}$  is locally free if and only if it is torsion-free.*
- (ii) *If an  $\mathcal{O}$ -module  $\mathcal{M}$  is not torsion-free, it has a nonzero global section.*

*proof.* (i) We may assume that  $Y$  is affine,  $Y = \text{Spec } B$ , and that  $\mathcal{M}$  is the  $\mathcal{O}$ -module associated to a  $B$ -module  $M$ . Let  $\widetilde{B}$  and  $\widetilde{M}$  be the localizations of  $B$  and  $M$  at a point  $q$ , respectively. Then  $\widetilde{M}$  is a finite, torsion-free

module over the local ring  $\widetilde{B}$ . It suffices to show that, for every point  $q$  of  $Y$ ,  $\widetilde{M}$  is a free  $\widetilde{B}$ -module (5.1.15). The local ring  $\widetilde{B}$  is a valuation ring. A valuation ring is a principal ideal domain because the nonzero ideals of  $\widetilde{B}$  are powers of the maximal ideal  $\widetilde{m}$ , which is a principal ideal. Every finite, torsion-free module over a principal ideal domain is free.

(ii) If the torsion submodule of  $\mathcal{M}$  isn't zero, there will be an affine open set  $U$ , and there will be nonzero elements  $m$  in  $\mathcal{M}(U)$  and  $a$  in  $\mathcal{O}(U)$ , such that  $am = 0$ . Let  $C$  be the finite set of zeros of  $a$  in  $U$ , and let  $V = Y - C$  be the complement of  $C$  in  $Y$ . Then  $a$  is invertible on the intersection  $W = U \cap V$ , and since  $am = 0$ , the restriction of  $m$  to  $W$  is zero.

The open sets  $U$  and  $V$  cover  $Y$ , and the sheaf property for this covering can be written as an exact sequence

$$0 \rightarrow \mathcal{M}(Y) \rightarrow \mathcal{M}(U) \times \mathcal{M}(V) \xrightarrow{+,-} \mathcal{M}(W)$$

(Lemma 6.3.10). In this sequence, the section  $(m, 0)$  of  $\mathcal{M}(U) \times \mathcal{M}(V)$  is mapped to zero in  $\mathcal{M}(W)$ . Therefore it is the image of a nonzero global section of  $\mathcal{M}$ .  $\square$

**8.2.2. Lemma.** *Let  $Y$  be a smooth curve. Every nonzero ideal  $\mathcal{I}$  of  $\mathcal{O}_Y$  is a product of powers of maximal ideals of  $\mathcal{O}_Y$ :  $\mathcal{I} = \mathfrak{m}_1^{e_1} \cdots \mathfrak{m}_k^{e_k}$ .*

*proof.* This follows for any smooth curve from the case that  $Y$  is affine, which is Proposition 5.3.7.  $\square$

**8.2.3. Notation.** When considering a branched covering  $Y \xrightarrow{\pi} X$  of smooth curves, we will often pass between an  $\mathcal{O}_Y$ -module  $\mathcal{M}$  and its direct image  $\pi_*\mathcal{M}$ , and it will be convenient to work primarily on  $X$ . Recall that if  $Y'$  is the inverse image of an open subset  $X'$  of  $X$ , then

$$[\pi_*\mathcal{M}](X') = \mathcal{M}(Y')$$

One can think of the direct image  $\pi_*\mathcal{M}$  as working with  $\mathcal{M}$ , but looking only at open subsets  $Y'$  of  $Y$  that are inverse images of open subsets  $X'$  of  $X$ . If we look only at such open subsets, the only significant difference between  $\mathcal{M}$  and its direct image will be that the  $\mathcal{O}_Y(Y')$ -module  $\mathcal{M}(Y')$  is made into an  $\mathcal{O}_X(X')$ -module by restriction of scalars. To simplify notation, we will often drop the symbol  $\pi_*$ , and write  $\mathcal{M}$  instead of  $\pi_*\mathcal{M}$ . If  $X'$  is an open subset of  $X$ ,  $\mathcal{M}(X')$  will stand for  $\mathcal{M}(\pi^{-1}X')$ . When thinking of an  $\mathcal{O}_Y$ -module  $\mathcal{M}$  as the direct image, we may refer to it as an  $\mathcal{O}_X$ -module. In accordance with this convention, we may also write  $\mathcal{O}_Y$  for  $\pi_*\mathcal{O}_Y$ , but we must be careful to include the subscript  $Y$ .

If you find this abbreviation confusing, you can put the symbol  $\pi_*$  into the text where appropriate.  $\square$

**8.2.4. Lemma.** *Let  $Y \rightarrow X$  be a branched covering of smooth curves.*

(i) *A finite  $\mathcal{O}_Y$ -module  $\mathcal{N}$  is a torsion  $\mathcal{O}_Y$ -module if and only if it is a torsion  $\mathcal{O}_X$ -module.*

(ii) *A finite  $\mathcal{O}_Y$ -module  $\mathcal{N}$  is a locally free  $\mathcal{O}_Y$ -module if and only if it is a locally free  $\mathcal{O}_X$ -module. If  $\mathcal{N}$  is a locally free  $\mathcal{O}_Y$ -module of rank  $r$ , then it is a locally free  $\mathcal{O}_X$ -module of rank  $nr$ , where  $n$  is the degree  $[Y : X]$  of the covering.  $\square$*

### (8.2.5) the module $\text{Hom}$

Let  $M$  and  $N$  be modules over a ring  $A$ . We are going to need the  $A$ -module that is usually denoted by  $\text{Hom}_A(M, N)$ , of homomorphisms  $M \rightarrow N$ . The set of such homomorphisms becomes an  $A$ -module with some fairly obvious laws of composition: If  $\varphi$  and  $\psi$  are homomorphisms and  $a$  is an element of  $A$ , then  $\varphi + \psi$  and  $a\varphi$  are defined by

$$(8.2.6) \quad [\varphi + \psi](m) = \varphi(m) + \psi(m) \quad \text{and} \quad [a\varphi](m) = a\varphi(m)$$

If  $\varphi$  is a module homomorphism, we also have  $\varphi(m_1 + m_2) = \varphi(m_1) + \varphi(m_2)$ , and  $a\varphi(m) = \varphi(am)$ .

**8.2.7. Lemma.** *An  $A$ -module  $N$  is canonically isomorphic to  $\text{Hom}_A(A, N)$ . The homomorphism  $A \xrightarrow{\varphi} N$  that corresponds to an element  $n$  of  $N$  is multiplication by  $n$ :  $\varphi(a) = an$ . Conversely, the element of  $N$  that corresponds to a homomorphism  $A \xrightarrow{\varphi} N$  is  $n = \varphi(1)$ .*

*Similarly, an  $\mathcal{O}$ -module  $\mathcal{M}$  on a smooth curve  $Y$  is isomorphic to  $\underline{\text{Hom}}_{\mathcal{O}}(\mathcal{O}, \mathcal{M})$ .  $\square$*

Thus  $\text{Hom}_A(A^k, N)$  is isomorphic to  $N^k$ , and  $\text{Hom}_A(A^\ell, A^k)$  is isomorphic to the module  $A^{\ell \times k}$  of  $k \times \ell$   $A$ -matrices.

**8.2.8. Lemma.** *Let  $A$  be a noetherian ring.*

(i) *For every finite  $A$ -module  $M$ , there is an exact sequence  $A^\ell \rightarrow A^k \rightarrow M \rightarrow 0$ .*

(ii) *If  $M$  and  $N$  are finite  $A$ -modules, then  ${}_A(M, N)$  is a finite  $A$ -module.*  $\square$

An exact sequence of the form  $A^\ell \rightarrow A^k \rightarrow M \rightarrow 0$  is called a *presentation* of module  $M$ .

The module  $\text{Hom}$  is compatible with localization:

**8.2.9. Lemma.** *Let  $M$  and  $N$  be modules over a noetherian domain  $A$ , and suppose that  $M$  is a finite module. Let  $S$  be a multiplicative system in  $A$ . The localization  $S^{-1} \text{Hom}_A(M, N)$  is canonically isomorphic to  $\text{Hom}_{S^{-1}A}(S^{-1}M, S^{-1}N)$ .*

*proof.* We choose a presentation  $A^\ell \rightarrow A^k \rightarrow M \rightarrow 0$ . Its localization  $(S^{-1}A)^\ell \rightarrow (S^{-1}A)^k \rightarrow S^{-1}M \rightarrow 0$  is a presentation of  $S^{-1}M$ . Because  $\text{Hom}_A(\cdot, \cdot)$  is a left exact, contravariant functor of the first variable, the sequence

$$0 \rightarrow \text{Hom}_A(M, N) \rightarrow \text{Hom}_A(A^k, N) \rightarrow \text{Hom}_A(A^\ell, N)$$

is exact, as is its localization. This it suffices to prove the lemma in the case that  $M = A$ . It is true in that case.  $\square$

This lemma shows that when  $\mathcal{M}$  and  $\mathcal{N}$  are finite  $\mathcal{O}$ -modules on a variety  $X$ , there is an  $\mathcal{O}$ -module of homomorphisms  $\mathcal{M} \rightarrow \mathcal{N}$ , which will be denoted by  $\underline{\text{Hom}}_{\mathcal{O}}(\mathcal{M}, \mathcal{N})$ . If  $U = \text{Spec } A$  is an affine open set,  $M = \mathcal{M}(U)$  and  $N = \mathcal{N}(U)$ , the module of sections of  $\underline{\text{Hom}}_{\mathcal{O}}(\mathcal{M}, \mathcal{N})$  on  $U$  is  $\text{Hom}_A(M, N)$ . We use a new symbol  $\underline{\text{Hom}}$  here because the vector space of homomorphisms  $\mathcal{M} \rightarrow \mathcal{N}$  defined on all of  $X$ , which is the space of global sections of  $\underline{\text{Hom}}_{\mathcal{O}}(\mathcal{M}, \mathcal{N})$ , is customarily denoted by  $\text{Hom}_{\mathcal{O}}(\mathcal{M}, \mathcal{N})$ .

**8.2.10. Notation.** The notation  $\text{Hom}_A(M, N)$  is cumbersome. It seems permissible to drop the symbol  $\text{Hom}$ , and to write  ${}_A(M, N)$  for  $\text{Hom}_A(M, N)$ . Similarly, if  $\mathcal{M}$  and  $\mathcal{N}$  are  $\mathcal{O}$ -modules on a variety  $X$ , we will write  ${}_{\mathcal{O}}(\mathcal{M}, \mathcal{N})$  or  ${}_X(\mathcal{M}, \mathcal{N})$  for  $\underline{\text{Hom}}_{\mathcal{O}}(\mathcal{M}, \mathcal{N})$ .

**8.2.11. Lemma.** *Let  $A \subset B$  be rings, let  $M$  be an  $A$ -module, and let  $N$  be a  $B$ -module. Then  ${}_A(M, N)$  becomes a  $B$ -module.*

When we write  ${}_A(M, N)$ , we are interpreting the  $B$ -module  $N$  as an  $A$ -module by restriction of scalars.

*proof.* Let  $M \xrightarrow{\varphi} N$  be a homomorphism of  $A$ -modules, and let  $b$  be an element of  $B$ . Then multiplication by  $b$  is defined by the rule  $[b\varphi](m) = \varphi(bm)$ . There are several things to check. We list here as a reminder:

The map  $[b\varphi]$  is a homomorphism of  $A$ -modules  $M \rightarrow N$ :

$$[b\varphi](m_1 + m_2) = [b\varphi](m_1) + [b\varphi](m_2) \text{ and } [b\varphi](am) = a[b\varphi](m)$$

The  $A$ -module  ${}_A(M, N)$  has the structure of a  $B$ -module:

$$[b(\varphi_1 + \varphi_2)] = [b\varphi_1] + [b\varphi_2], [(b_1 + b_2)\varphi] = [b_1\varphi] + [b_2\varphi], [1\varphi] = \varphi, \text{ and } [b_1[b_2\varphi]] = [b_1b_2\varphi] \quad \square$$

**8.2.12. Lemma.**

(i) *The functor  $\text{Hom}_A$  is a left exact and **contravariant** in the first variable. An exact sequence  $M_1 \rightarrow M_2 \rightarrow M_3 \rightarrow 0$  of  $A$ -modules induces, for any  $A$ -module  $N$ , an exact sequence*

$$0 \rightarrow {}_A(M_3, N) \rightarrow {}_A(M_2, N) \rightarrow {}_A(M_1, N)$$

(ii) *The functor  $\text{Hom}_A$  is a left exact and **covariant** in the second variable. An exact sequence  $0 \rightarrow N_1 \rightarrow N_2 \rightarrow N_3 \rightarrow 0$  of  $A$ -modules induces, for any  $A$ -module  $M$ , an exact sequence*

$$0 \rightarrow {}_A(M, N_1) \rightarrow {}_A(M, N_2) \rightarrow {}_A(M, N_3)$$

*The analogous statements are true for  $\underline{\text{Hom}}_{\mathcal{O}}$ .*  $\square$

**(8.2.13) the dual module**



The dual module  $\mathcal{M}^*$  of a locally free  $\mathcal{O}$ -module  $\mathcal{M}$  is the  $\mathcal{O}$ -module  ${}_{\mathcal{O}}(\mathcal{M}, \mathcal{O})$ . A section of  $\mathcal{M}^*$  on an open set  $U$  is a homomorphism  $\mathcal{M}(U) \rightarrow \mathcal{O}(U)$ . The dual is contravariant. A homomorphism  $\mathcal{M} \rightarrow \mathcal{N}$  of locally free  $\mathcal{O}$ -modules induces a homomorphism  $\mathcal{M}^* \leftarrow \mathcal{N}^*$ .

If  $\mathcal{M}$  is a free module with basis  $v_1, \dots, v_k$ , then  $\mathcal{M}^*$  will also be free, with the dual basis  $v_i^*$  defined by  $v_i^*(v_i) = 1$  and  $v_i^*(v_j) = 0$  if  $i \neq j$ . Therefore, when  $\mathcal{M}$  is locally free,  $\mathcal{M}^*$  is also locally free. The dual  $\mathcal{O}^*$  of the structure sheaf  $\mathcal{O}$  is  $\mathcal{O}$  itself. If  $\mathcal{M}$  and  $\mathcal{N}$  are locally free  $\mathcal{O}$ -modules, the dual  $(\mathcal{M} \otimes_{\mathcal{O}} \mathcal{N})^*$  is isomorphic to the tensor product  $\mathcal{M}^* \otimes_{\mathcal{O}} \mathcal{N}^*$ .

There is a canonical  $\mathcal{O}$ -bilinear map  $\mathcal{M}^* \times \mathcal{M} \rightarrow \mathcal{O}$ . If  $\alpha$  and  $m$  are sections of  $\mathcal{M}^*$  and  $\mathcal{M}$ , respectively, the bilinear map evaluates  $\alpha$  at  $m$ :  $\langle \alpha, m \rangle = \alpha(m)$ .

**8.2.14. Corollary.** A locally free  $\mathcal{O}$ -module  $\mathcal{M}$  is canonically isomorphic to its bidual:  $(\mathcal{M}^*)^* \approx \mathcal{M}$ . □

**8.2.15. Proposition.** Let  $0 \rightarrow \mathcal{M} \rightarrow \mathcal{N} \rightarrow \mathcal{P} \rightarrow 0$  be an exact sequence of  $\mathcal{O}$ -modules on a variety  $X$ .

(i) If  $\mathcal{P}$  is a free  $\mathcal{O}$ -module, and if the map of global sections  $H^0(\mathcal{N}) \rightarrow H^0(\mathcal{P})$  is surjective, the sequence splits:  $\mathcal{N}$  is isomorphic to the direct sum  $\mathcal{M} \oplus \mathcal{P}$ .

(ii) If  $\mathcal{P}$  is locally free, the dual modules form an exact sequence  $0 \rightarrow \mathcal{P}^* \rightarrow \mathcal{N}^* \rightarrow \mathcal{M}^* \rightarrow 0$ .

*proof.* (i) Let  $\{p_i\}$  be a basis of global sections of  $\mathcal{P}$ , let  $p'_i$  be global sections of  $\mathcal{N}$  that map to  $p_i$ , and let  $\mathcal{P}'$  be the free  $\mathcal{O}$ -submodule of  $\mathcal{N}$  spanned by  $\{p'_i\}$ . So  $\mathcal{P}'$  is isomorphic to  $\mathcal{P}$ , and  $\mathcal{N} \approx \mathcal{M} \oplus \mathcal{P}'$ .

(ii) The sequence  $0 \rightarrow \mathcal{P}^* \rightarrow \mathcal{M}^* \rightarrow \mathcal{N}^*$  is exact whether or not the modules are locally free (8.2.12(ii)). The zero on the right comes from the fact that, when  $\mathcal{P}$  is locally free, it is free on some affine covering. Thus the given sequence splits locally. □

invertmod

### (8.2.16) invertible modules

An invertible  $\mathcal{O}$ -module is a locally free module of rank one – a module that is isomorphic to the free module  $\mathcal{O}$  in a neighborhood of any point.

The tensor product  $\mathcal{L} \otimes_{\mathcal{O}} \mathcal{M}$  of invertible modules is invertible. The dual  $\mathcal{L}^*$  of an invertible module  $\mathcal{L}$  is invertible. Part (i) of the next lemma explains the adjective 'invertible'.

**8.2.17. Lemma.** Let  $\mathcal{L}$  be an invertible  $\mathcal{O}$ -module.

(i) Let  $\mathcal{L}^*$  be the dual module. The canonical map  $\mathcal{L}^* \otimes_{\mathcal{O}} \mathcal{L} \rightarrow \mathcal{O}$  defined by  $\gamma \otimes \alpha \mapsto \gamma(\alpha)$  is an isomorphism.

(ii) The map  $\mathcal{O} \rightarrow {}_{\mathcal{O}}(\mathcal{L}, \mathcal{L})$  that sends a regular function  $\alpha$  to multiplication by  $\alpha$  is an isomorphism.

(iii) Every nonzero homomorphism  $\mathcal{L} \xrightarrow{\varphi} \mathcal{M}$  to a locally free module  $\mathcal{M}$  is injective.

*proof.* (i),(ii) It is enough to verify these assertions in the case that  $\mathcal{L}$  is free, isomorphic to  $\mathcal{O}$ , in which case they are clear.

(iii) The problem is local, so we may assume that the variety is affine, say  $Y = \text{Spec } A$ , and that  $\mathcal{L}$  and  $\mathcal{M}$  are free. Then  $\varphi$  becomes a nonzero homomorphism  $A^1 \rightarrow A^k$ . Such a homomorphism is injective because  $A$  is a domain. □

## 8.3 Divisors

A divisor on a smooth curve  $Y$  is a finite integer combination of points:

$$D = r_1 q_1 + \cdots + r_k q_k$$

with  $r_i \in \mathbb{Z}$ . The terms  $r_i q_i$  whose integer coefficients  $r_i$  are zero can be omitted or not, as desired.

The support of  $D$  is the set of points  $q_i$  of  $Y$  such that  $r_i \neq 0$ . The degree of  $D$  is the sum  $r_1 + \cdots + r_k$  of the coefficients.

Let  $Y'$  be an open subset of  $Y$ . The restriction of a divisor  $D = r_1 p_1 + \cdots + r_k p_k$  to  $Y'$  is the divisor on  $Y'$  obtained from  $D$  by deleting points that aren't in  $Y'$ . Thus, if  $D = q$ , the restriction of  $D$  to  $Y'$  is  $q$  when  $q \in Y'$ , and is zero when  $q \notin Y'$ .

A divisor  $D = \sum r_i q_i$  is *effective* if all of its coefficients  $r_i$  are non-negative, and  $D$  is *effective on an open subset*  $Y'$  if its restriction to  $Y'$  is effective – if  $r_i \geq 0$  for every  $i$  such that  $q_i$  is a point of  $Y'$ .

### (8.3.1) the divisor of a function

divfn

Let  $f$  be a rational function on a smooth curve  $Y$ . The *divisor of  $f$*  is

$$\operatorname{div}(f) = \sum_{q \in Y} v_q(f) q$$

where  $v_q$  denotes the valuation of  $K$  that corresponds to the point  $q$  of  $Y$ .

This divisor is written here as a sum over all points  $q$ , but it becomes a finite sum when we disregard terms with coefficient zero, because  $f$  has finitely many zeros and poles. The coefficients will be zero at all other points.

The map

$$K^\times \rightarrow (\operatorname{divisors})^+$$

that sends a rational function to its divisor is a homomorphism from the multiplicative group  $K^\times$  of nonzero elements of  $K$  to the additive group of divisors:

$$\operatorname{div}(fg) = \operatorname{div}(f) + \operatorname{div}(g)$$

As before, a rational function  $f$  has a *zero* of order  $r$  at  $q$  if  $v_q(f) = r$  with  $r > 0$ , and it has a *pole* of order  $r$  at  $q$  if  $v_q(f) = -r$ . Thus the divisor of  $f$  is the difference of two effective divisors:

$$\operatorname{div}(f) = \operatorname{zeros}(f) - \operatorname{poles}(f)$$

A rational function  $f$  is regular on  $Y$  if and only if its divisor is effective – if and only if  $\operatorname{poles}(f) = 0$ .

The divisor of a rational function is called a *principal divisor*, and two divisors  $D$  and  $E$  are *linearly equivalent* if their difference  $D - E$  is a principal divisor. For instance, the divisors  $\operatorname{zeros}(f)$  and  $\operatorname{poles}(f)$  of a rational function  $f$  are linearly equivalent.

**8.3.2. Lemma.** *Let  $f$  be a rational function on a smooth curve  $Y$ . For all complex numbers  $c$ , the divisors of zeros of  $f - c$ , the level sets of  $f$ , are linearly equivalent.*

*proof.* The functions  $f - c$  have the same poles as  $f$ . □

### (8.3.3) the module $\mathcal{O}(D)$

To analyze the space of functions with given poles on a smooth curve  $Y$ , we associate an  $\mathcal{O}$ -module  $\mathcal{O}(D)$  to a divisor  $D$ . The nonzero sections of  $\mathcal{O}(D)$  on an open subset  $V$  of  $Y$  are the rational functions  $f$  such that the divisor  $\operatorname{div}(f) + D$  is effective on  $V$  – such that its restriction to  $V$  is effective.

$$(8.3.4) \quad [\mathcal{O}(D)](V) = \{f \mid \operatorname{div}(f) + D \text{ is effective on } V\} \cup \{0\}$$

Points that aren't in the open set  $V$  impose no conditions on the sections on  $V$ .

When  $D$  is an effective divisor, a rational function  $f$  is a global section of  $\mathcal{O}(D)$  if  $\operatorname{poles}(f) \leq D$ .

Say that  $D = \sum r_i q_i$ . If  $q_i$  is a point of an open set  $V$  and if  $r_i > 0$ , a section of  $\mathcal{O}(D)$  on  $V$  may have a pole of order at most  $r_i$  at  $q_i$ , and if  $r_i < 0$  a section must have a zero of order at least  $-r_i$  at  $q_i$ . For example, the module  $\mathcal{O}(-q)$  is the maximal ideal  $\mathfrak{m}_q$ . The sections of  $\mathcal{O}(-q)$  on an open set  $V$  that contains  $q$  are the regular functions on  $V$  that are zero at  $q$ . Similarly, the sections of  $\mathcal{O}(q)$  on an open set  $V$  that contains  $q$  are the rational functions that have a pole of order at most 1 at  $q$  and are regular at every other point of  $V$ . The sections of  $\mathcal{O}(-q)$  and of  $\mathcal{O}(q)$  on an open set  $V$  that doesn't contain  $p$  are the regular functions on  $V$ . For any  $D$ , sections of  $\mathcal{O}(D)$  on  $V$  can have arbitrary zeros or poles at points that aren't in  $V$ .

The fact that a section of  $\mathcal{O}(D)$  is allowed to have a pole at  $q_i$  if  $r_i > 0$  contrasts with the divisor of a function. If  $\operatorname{div}(f) = \sum r_i q_i$ , then  $r_i > 0$  means that  $f$  has a zero at  $q_i$ . If  $\operatorname{div}(f) = D$ , then  $f$  will be a global section of  $\mathcal{O}(-D)$ .

**8.3.5. Lemma.** (i) If  $D$  and  $E$  are divisors and if  $E - D$  is effective, then  $\mathcal{O}(D) \subset \mathcal{O}(E)$ .

(ii) The function field module  $\mathcal{K}$  of a smooth curve  $Y$  is the union of the modules  $\mathcal{O}(D)$ .  $\square$

The next lemma follows from Lemma 8.2.2.

**8.3.6. Lemma.** Let  $Y$  be a smooth curve. The product ideal  $\mathcal{I} = \mathfrak{m}_1^{r_1} \cdots \mathfrak{m}_k^{r_k}$  of  $\mathcal{O}_Y$  is isomorphic to the  $\mathcal{O}$ -module  $\mathcal{O}_Y(-D)$ , where  $D$  is the effective divisor  $\sum r_i p_i$ . Thus nonzero ideals of  $\mathcal{O}_Y$  correspond bijectively to divisors  $-D$ , where  $D$  is effective.  $\square$

**8.3.7. Proposition.** Let  $D$  and  $E$  be divisors on a smooth curve  $Y$ .

(i) The  $\mathcal{O}$ -module  $\mathcal{O}(D)$  is invertible.

(ii) The map  $\mathcal{O}(D) \otimes_{\mathcal{O}} \mathcal{O}(E) \rightarrow \mathcal{O}(D+E)$  that sends  $f \otimes g$  to the product  $fg$  is an isomorphism.

(iii) The dual module  $\mathcal{O}(D)^*$  is  $\mathcal{O}(-D)$ .

(iv) Every invertible  $\mathcal{O}$ -module  $\mathcal{L}$  is isomorphic to a module of the form  $\mathcal{O}(D)$ .

The only difference between an invertible module  $\mathcal{L}$  and a module of the form  $\mathcal{O}(D)$  is that  $\mathcal{O}(D) \otimes_{\mathcal{O}} \mathcal{K}$  is equal to  $\mathcal{K}$ , whereas  $\mathcal{L}_{\mathcal{K}}$  can be a one-dimensional  $K$ -vector space without chosen basis.

It is important to note that, though every invertible module  $\mathcal{M}$  is isomorphic to one of the form  $\mathcal{O}(D)$ , the divisor  $D$  isn't uniquely determined by  $\mathcal{M}$ . (See (8.3.11) below.)

**8.3.8. Definition.** Let  $\mathcal{L}$  be an invertible  $\mathcal{O}$ -module on a smooth projective curve  $Y$ . If  $\mathcal{L}$  is isomorphic to  $\mathcal{O}(D)$ , we call the degree of  $D$  the *degree* of  $\mathcal{L}$ . With this definition,  $\chi(\mathcal{L}) = \deg \mathcal{L} + 1 - p_a$ .

*proof of Proposition 8.3.7.* (i) We may assume that  $Y$  is affine and that the support of  $D$  contains at most one point:  $D = rp$ . We may also assume that the maximal ideal at  $p$  is a principal ideal, generated by an element  $x$ . In that case,  $\mathcal{O}(D)$  will be the free module with basis  $x^r$ .

(ii),(iii) Proceeding as in the proof of (i), we may assume that  $D = rp$  and  $E = sp$ . Then  $\mathcal{O}(D)$ ,  $\mathcal{O}(-D)$ ,  $\mathcal{O}(E)$ , and  $\mathcal{O}(D+E)$  have bases  $x^r$ ,  $x^{-r}$ ,  $x^s$  and  $x^{r+s}$ , respectively.

(iv) Let  $K$  be the function field of  $Y$ , and let  $\mathcal{K}$  be the function field module. When  $\mathcal{L}$  is an invertible  $\mathcal{O}$ -module,  $\mathcal{L}_{\mathcal{K}} = \mathcal{L} \otimes_{\mathcal{O}} \mathcal{K}$  will be a one-dimensional  $K$ -vector space (see (??)). Since the function field module  $\mathcal{K}$  of  $Y$  is the union  $\mathcal{K} = \bigcup \mathcal{O}(D)$ , we also have  $\mathcal{L}_{\mathcal{K}} = \bigcup \mathcal{L}(D)$ , where  $\mathcal{L}(D)$  denotes the tensor product  $\mathcal{L} \otimes_{\mathcal{O}} \mathcal{O}(D)$ . A nonzero global section  $\alpha$  of  $\mathcal{L}_{\mathcal{K}}$  will be a global section of  $\mathcal{L}(D)$  for some  $D$ . It will define a map  $\mathcal{O} \xrightarrow{\alpha} \mathcal{L}(D)$ . Passing to duals,  $\mathcal{L}(D)^* = \mathcal{L}^* \otimes_{\mathcal{O}} \mathcal{O}(D)^* \approx \mathcal{L}^*(-D)$ . The dual of the map  $\alpha$  is a nonzero and therefore injective map  $\mathcal{L}^*(-D) \rightarrow \mathcal{O}$  whose image is an ideal of  $\mathcal{O}$ . So  $\mathcal{L}^*(-D)$  is isomorphic to  $\mathcal{O}(-E)$  for some effective divisor  $E$ , and therefore  $\mathcal{L}^*$  is isomorphic to  $\mathcal{O}(D-E)$ . Dualizing once more,  $\mathcal{L}$  is isomorphic to  $\mathcal{O}(E-D)$ .  $\square$

If  $\mathcal{L}$  is an invertible module, we denote by  $\mathcal{L}(D)$  the invertible module  $\mathcal{L} \otimes_{\mathcal{O}} \mathcal{O}(D)$ .

**8.3.9. Proposition.** Let  $\mathcal{L} \subset \mathcal{M}$  be an inclusion of invertible  $\mathcal{O}$ -modules. Then  $\mathcal{M} = \mathcal{L}(E)$  for some effective divisor  $E$ .

*proof.*  $\mathcal{L}$  is isomorphic to  $\mathcal{O}(D)$  for some  $D$ . Then  $\mathcal{L}(-D) \subset \mathcal{O}$  so  $\mathcal{L}(-D)$  is an ideal, isomorphic to  $\mathcal{O}(-E)$  for some effective divisor  $E$ . Then  $\mathcal{L}(E) \approx \mathcal{O}(D) \approx \mathcal{M}$ .  $\square$

If  $D$  and  $E$  are divisors,  $\mathcal{O}(D)$  is a submodule of  $\mathcal{O}(E)$  only when  $E - D$  is effective. But as the next proposition explains, there may be homomorphisms from  $\mathcal{O}(D)$  to  $\mathcal{O}(E)$  that aren't inclusions.

**8.3.10. Proposition.** Let  $D$  and  $E$  be divisors on a smooth curve  $Y$ . Multiplication by a rational function  $f$  such that  $\operatorname{div}(f) + E - D \geq 0$  defines a homomorphism of  $\mathcal{O}$ -modules  $\mathcal{O}(D) \rightarrow \mathcal{O}(E)$ , and every homomorphism  $\mathcal{O}(D) \rightarrow \mathcal{O}(E)$  is multiplication by such a function.

*proof.* For any  $\mathcal{O}$ -module  $\mathcal{M}$ , a homomorphism  $\mathcal{O} \rightarrow \mathcal{M}$  is multiplication by a global section of  $\mathcal{M}$  (6.4.4). Then a homomorphism  $\mathcal{O} \rightarrow \mathcal{O}(E-D)$  will be multiplication by a rational function  $f$  such that  $\operatorname{div}(f) + E - D \geq 0$ . If  $f$  is such a function, one obtains a homomorphism  $\mathcal{O}(D) \rightarrow \mathcal{O}(E)$  by tensoring with  $\mathcal{O}(D)$ .  $\square$

**8.3.11. Corollary.**

(i) The modules  $\mathcal{O}(D)$  and  $\mathcal{O}(E)$  are isomorphic if and only if the divisors  $D$  and  $E$  are linearly equivalent.

(ii) Let  $f$  be a rational function on  $Y$ , and let  $D = \operatorname{div}(f)$ . Multiplication by  $f$  defines an isomorphism  $\mathcal{O}(D) \rightarrow \mathcal{O}$ .  $\square$

## 8.4 The Riemann-Roch Theorem I

Let  $Y$  be a smooth projective curve. In Chapter 7, we learned that when  $\mathcal{M}$  is a finite  $\mathcal{O}_Y$ -module, the cohomology  $H^q(Y, \mathcal{M})$  is a finite-dimensional vector space for all  $q$ , and is zero if  $q \neq 0, 1$ . As before, we denote the dimension of the space  $H^q(Y, \mathcal{M})$  by  $h^q(\mathcal{M})$  or, if there is ambiguity about the variety, by  $h^q(Y, \mathcal{M})$ .

The Euler characteristic (7.6.5) of a finite  $\mathcal{O}$ -module  $\mathcal{M}$  is

$$(8.4.1) \quad \chi(\mathcal{M}) = h^0(\mathcal{M}) - h^1(\mathcal{M})$$

In particular,

$$\chi(\mathcal{O}_Y) = h^0(\mathcal{O}_Y) - h^1(\mathcal{O}_Y)$$

The dimension  $h^1(\mathcal{O}_Y)$  is the *arithmetic genus* of  $Y$ . It is denoted by  $p_a$ . We will see below, in (8.4.8)(iv), that  $h^0(\mathcal{O}_Y) = 1$ . So

$$(8.4.2) \quad \chi(\mathcal{O}_Y) = 1 - p_a$$

**8.4.3. Riemann-Roch Theorem (version 1).** *Let  $D = \sum r_i p_i$  be a divisor on a smooth projective curve  $Y$ . Then*

$$\chi(\mathcal{O}(D)) = \chi(\mathcal{O}) + \deg D \quad (= \deg D + 1 - p_a)$$

*proof.* We analyze the effect on cohomology when a divisor is changed by adding or subtracting a point by inspecting the inclusion  $\mathcal{O}(D-p) \subset \mathcal{O}(D)$ . Let  $\epsilon$  be the cokernel of the inclusion map, so that there is a short exact sequence in which  $\epsilon$  is a one-dimensional vector space supported at  $p$ , with  $h^0(\epsilon) = 1$ , and  $h^1(\epsilon) = 0$ .

$$(8.4.4) \quad 0 \rightarrow \mathcal{O}(D-p) \rightarrow \mathcal{O}(D) \rightarrow \epsilon \rightarrow 0$$

Since  $\mathfrak{m}_p$  is isomorphic to  $\mathcal{O}(-p)$ , this sequence can be obtained by tensoring the sequence

$$(8.4.5) \quad 0 \rightarrow \mathfrak{m}_p \rightarrow \mathcal{O} \rightarrow \kappa_p \rightarrow 0$$

with the invertible module  $\mathcal{O}(D)$ .

Let's denote the one-dimensional vector space  $H^0(Y, \epsilon)$  by  $[1]$ . The cohomology sequence associated to (8.4.4) is

$$(8.4.6) \quad 0 \rightarrow H^0(Y, \mathcal{O}(D-p)) \rightarrow H^0(Y, \mathcal{O}(D)) \xrightarrow{\gamma} [1] \xrightarrow{\delta} H^1(Y, \mathcal{O}(D-p)) \rightarrow H^1(Y, \mathcal{O}(D)) \rightarrow 0$$

In this exact sequence, one of the two maps,  $\gamma$  or  $\delta$ , must be zero. Either

(1)  $\gamma$  is zero and  $\delta$  is injective. In this case

$$h^0(\mathcal{O}(D-p)) = h^0(\mathcal{O}(D)) \quad \text{and} \quad h^1(\mathcal{O}(D-p)) = h^1(\mathcal{O}(D)) + 1, \quad \text{or}$$

(2)  $\delta$  is zero and  $\gamma$  is surjective. In this case

$$h^0(\mathcal{O}(D-p)) = h^0(\mathcal{O}(D)) - 1 \quad \text{and} \quad h^1(\mathcal{O}(D-p)) = h^1(\mathcal{O}(D))$$

In either case,

$$(8.4.7) \quad \chi(\mathcal{O}(D)) = \chi(\mathcal{O}(D-p)) + 1$$

The Riemann-Roch theorem follows from this, because we can get from  $\mathcal{O}$  to  $\mathcal{O}(D)$  by a finite number of operations, each of which changes the divisor by adding or subtracting a point.  $\square$

Because  $h^0 \geq h^0 - h^1 = \chi$ , this version of the Riemann-Roch Theorem gives reasonably good control of  $H^0$ . It is less useful for controlling  $H^1$ . To do that, one wants the full Riemann-Roch Theorem. That theorem requires some preparation, so we have put it into Section 8.8. However, version 1 has important consequences:

**8.4.8. Corollary.** *Let  $Y$  be a smooth projective curve.*

- (i) *The divisor of a rational function has degree zero: The number of zeros is equal to the number of poles.*
- (ii) *Linearly equivalent divisors have equal degrees.*
- (iii) *A nonconstant rational function takes every value, including infinity, the same number of times.*
- (iv) *A rational function that is regular at every point is a constant:  $H^0(Y, \mathcal{O}) = \mathbb{C}$ .*
- (v) *Let  $D$  be a divisor. If  $\deg D \geq p_a$ , then  $h^0(\mathcal{O}(D)) > 0$ .*
- (vi) *If  $h^0(\mathcal{O}(D)) > 0$ , then  $\deg D \geq 0$ .*

*proof.* (i) Let  $D = \text{div}(f)$ . Multiplication by the rational function  $f$  defines an isomorphism  $\mathcal{O}(D) \rightarrow \mathcal{O}$ , so  $\chi(\mathcal{O}(D)) = \chi(\mathcal{O})$ . On the other hand, by Riemann-Roch,  $\chi(\mathcal{O}(D)) = \chi(\mathcal{O}) + \deg D$ . Therefore  $\deg D = 0$ .

(ii) If two divisors  $D$  and  $E$  are linearly equivalent, say  $D - E = \text{div}(f)$ , then  $D - E$  has degree zero, and  $\deg D = \deg E$ .

(iii) The zeros of the functions  $f - c$  are linearly equivalent to the poles of  $f$  (8.3.2).

(iv) According to (iii), a nonconstant function must have a pole.

(v)  $h^0 \geq h^0 - h^1 = \chi = \deg D + 1 - p_a$ .

(vi) Suppose that  $\mathcal{O}(D)$  has a nonzero global section  $f$ , a rational function such that  $\text{div}(f) + D = E$  is effective. Then  $\deg E \geq 0$ . Since the degree of  $\text{div}(f)$  is zero,  $\deg D \geq 0$ .  $\square$

**8.4.9. Theorem.** *With its classical topology, a smooth projective curve  $Y$  is a connected, compact, orientable two-dimensional manifold.*

*proof.* All points except connectedness have been discussed before (Theorem 1.7.24). A nonempty topological space is connected if it isn't the union of two disjoint, nonempty, closed subsets. We argue by contradiction. Suppose that, in the classical topology,  $Y$  is the union of disjoint, nonempty closed subsets  $Y_1$  and  $Y_2$ . Both  $Y_1$  and  $Y_2$  will be compact manifolds. Let  $q$  be a point of  $Y_1$ . Part (v) of Corollary 8.4.8 shows that  $h^0(\mathcal{O}(nq)) > 1$  when  $n$  is large. A nonconstant global section  $f$  of  $\mathcal{O}(nq)$  will be a regular function on the complement  $Y - q$  of  $q$ . Then  $f$  is analytic, and it has no pole on the compact manifold  $Y_2$ . It will map  $Y_2$  to a compact subset of the complex plane. A nonconstant analytic function maps open sets to open sets. So if  $f$  weren't constant on  $Y_2$ , its image would be open. A compact subset of  $\mathbb{C}$  can't be open, so  $f$  must be constant on  $Y_2$ . When we subtract that constant from  $f$ , we obtain a nonconstant rational function  $g$  that is zero on  $Y_2$ . But since  $Y$  has dimension 1, the zero locus of a rational function is finite. This is a contradiction.  $\square$

## 8.5 The Birkhoff-Grothendieck Theorem

This theorem describes finite, torsion-free modules on the projective line.

**8.5.1. Birkhoff-Grothendieck Theorem.** *A finite, torsion-free  $\mathcal{O}$ -module  $\mathcal{M}$  on the projective line  $\mathbb{P}^1$  is isomorphic to a direct sum of twisting modules:  $\mathcal{M} \approx \bigoplus \mathcal{O}(n_i)$ .*

We recall the cohomology of the twisting modules on  $\mathbb{P}^1$ : If  $n \geq 0$ , then  $h^0(\mathcal{O}(n)) = n+1$  and  $h^1(\mathcal{O}(n)) = 0$ , and if  $r > 0$ , then  $h^0(\mathcal{O}(-r)) = 0$  and  $h^1(\mathcal{O}(-r)) = r-1$  (Theorem 7.5.4).

**8.5.2. Lemma.** *Let  $X$  denote the projective line, and let  $\mathcal{M}$  be a finite, torsion-free  $\mathcal{O}$ -module on  $X$ .*

- (i) *The integers  $r$  for which there exists a nonzero map  $\mathcal{O}(r) \rightarrow \mathcal{M}$  are bounded above.*
- (ii) *For large  $r$ ,  $h^0(X, \mathcal{M}(-r)) = 0$ .*

*proof.* (i) Since  $\mathcal{M}$  is torsion-free, any nonzero map  $\mathcal{O} \rightarrow \mathcal{M}$ , which is multiplication by a global section of  $\mathcal{M}$ , will be injective. Since  $\mathcal{O}(r)$  is locally isomorphic to  $\mathcal{O}$ , a nonzero map  $\mathcal{O}(r) \rightarrow \mathcal{M}$  will be injective too, and the associated map  $H^0(X, \mathcal{O}(r)) \rightarrow H^0(X, \mathcal{M})$  will be injective. Then  $h^0(X, \mathcal{O}(r)) \leq h^0(X, \mathcal{M})$ . Since  $h^0(X, \mathcal{O}(r)) = r+1$  and  $h^0(X, \mathcal{M})$  is finite,  $r$  is bounded.

(ii) A global section of  $\mathcal{M}(-r)$  defines a map  $\mathcal{O} \rightarrow \mathcal{M}(-r)$ . Its twist by  $r$  will be a map  $\mathcal{O}(r) \rightarrow \mathcal{M}$ .  $\square$

By the way, the conclusions of the lemma are true for any projective variety  $X$ .

*proof of the Birkhoff-Grothendieck Theorem.* This is Grothendieck's proof. A version of Birkhoff's proof, which uses matrices, is suggested as an exercise.

Lemma 8.2.1 tells us that  $\mathcal{M}$  is locally free. We use induction on the rank of  $\mathcal{M}$ . We suppose that the theorem has been proved for locally free  $\mathcal{O}$ -modules of rank less than  $r$ , that  $\mathcal{M}$  has rank  $r$ , and that  $r > 0$ . The plan is to show that  $\mathcal{M}$  has a twisting module as a direct summand, so that  $\mathcal{M} = \mathcal{W} \oplus \mathcal{O}(n)$  for some  $\mathcal{W}$ . Then we can apply induction on the rank to  $\mathcal{W}$ .

Since twisting is compatible with direct sums, we may replace  $\mathcal{M}$  by a twist  $\mathcal{M}(n)$ . Instead of showing that  $\mathcal{M}$  has a twisting module  $\mathcal{O}(n)$  as a direct summand, we show that, after we replace  $\mathcal{M}$  by a suitable twist, the structure sheaf  $\mathcal{O}$  will be a direct summand.

As we know (6.6.20), the twist  $\mathcal{M}(n)$  will have a nonzero global section when  $n$  is sufficiently large, and by Lemma 8.5.2 (ii), it will have no nonzero global section when  $n$  is sufficiently negative. Therefore, when we replace  $\mathcal{M}$  by a suitable twist, we will have  $H^0(X, \mathcal{M}) \neq 0$  but  $H^0(X, \mathcal{M}(-1)) = 0$ . We assume that this is true for  $\mathcal{M}$ .

We choose a nonzero global section  $s$  of  $\mathcal{M}$  and consider the injective multiplication map  $\mathcal{O} \xrightarrow{s} \mathcal{M}$ . Let  $\mathcal{W}$  be the cokernel of this map, so that we have a short exact sequence

$$(8.5.3) \quad 0 \rightarrow \mathcal{O} \xrightarrow{s} \mathcal{M} \rightarrow \mathcal{W} \rightarrow 0$$

**8.5.4. Lemma.** *Let  $\mathcal{W}$  be the  $\mathcal{O}$ -module that appears in the sequence (8.5.3).*

- (i)  $H^0(X, \mathcal{W}(-1)) = 0$ .
- (ii)  $\mathcal{W}$  is torsion-free, and therefore locally free.
- (iii)  $\mathcal{W}$  is a direct sum  $\bigoplus_{i=1}^{r-1} \mathcal{O}(n_i)$  of twisting modules on  $\mathbb{P}^1$ , with  $n_i \leq 0$ .

*proof.* (i) This follows from the cohomology sequence associated to the twisted sequence

$$0 \rightarrow \mathcal{O}(-1) \rightarrow \mathcal{M}(-1) \rightarrow \mathcal{W}(-1) \rightarrow 0$$

because  $H^0(X, \mathcal{M}(-1)) = 0$  and  $H^1(X, \mathcal{O}(-1)) = 0$ .

(ii) If  $\mathcal{W}$  had a nonzero torsion submodule, so would  $\mathcal{W}(-1)$ , and then  $\mathcal{W}(-1)$  would have a nonzero global section (8.2.1).

(iii) The fact that  $\mathcal{W}$  is a direct sum of twisting modules follows by induction on the rank:  $\mathcal{W} \approx \bigoplus \mathcal{O}(n_i)$ . Since  $H^0(X, \mathcal{W}(-1)) = 0$ , we must have  $H^0(X, \mathcal{O}(n_i - 1)) = 0$  too. Therefore  $n_i - 1 < 0$ , and  $n_i \leq 0$ .  $\square$

We go back to the proof of Theorem 8.5.1. Lemma 8.2.15 tells us that the dual of the sequence (8.5.3) is an exact sequence

$$0 \rightarrow \mathcal{W}^* \rightarrow \mathcal{M}^* \rightarrow \mathcal{O}^* \rightarrow 0$$

and  $\mathcal{W}^* \approx \bigoplus \mathcal{O}(-n_i)$  with  $-n_i \geq 0$ . Therefore  $\mathbf{h}^1(\mathcal{W}^*) = 0$ . The map  $H^0(\mathcal{M}) \rightarrow H^0(\mathcal{O}^*)$  is surjective. Lemma 8.2.15 tells us that  $\mathcal{M}^*$  is isomorphic to  $\mathcal{W}^* \oplus \mathcal{O}^*$ . Then  $\mathcal{M}$  is isomorphic to  $\mathcal{W} \oplus \mathcal{O}$ .  $\square$

## 8.6 Differentials

Why differentials enter into the Riemann-Roch Theorem is a mystery, but they do, so we introduce them here.

Let  $A$  be an algebra and let  $M$  be an  $A$ -module. A *derivation*  $A \xrightarrow{\delta} M$  is a  $\mathbb{C}$ -linear map that satisfies the product rule for differentiation – a map with these properties:

$$(8.6.1) \quad \delta(ab) = a\delta b + b\delta a, \quad \delta(a+b) = \delta a + \delta b, \quad \text{and} \quad \delta c = 0$$

for all  $a, b$  in  $A$  and all  $c$  in  $\mathbb{C}$ . The fact that  $\delta$  is  $\mathbb{C}$ -linear, i.e., that  $\delta(cb) = c\delta b$ , follows.

For example, differentiation  $\frac{d}{dt}$  is a derivation  $\mathbb{C}[t] \rightarrow \mathbb{C}[t]$ .

The module of differentials  $\Omega_A$  of an algebra  $A$  is an  $A$ -module generated by elements denoted by  $da$ , one for each element  $a$  of  $A$ . Its elements are (finite) combinations  $\sum b_i da_i$ , with  $a_i$  and  $b_i$  in  $A$ . The defining relations among the generators  $da$  are the ones that make the map  $A \xrightarrow{d} \Omega_A$  that sends  $a$  to  $da$  a derivation: For all  $a, b$  in  $A$  and all  $c$  in  $\mathbb{C}$ ,

$$(8.6.2) \quad d(ab) = a db + b da, \quad d(a+b) = da + db, \quad \text{and} \quad dc = 0$$

The elements of  $\Omega_A$  are called *differentials*.

### 8.6.3. Lemma.

(i) Let  $\Omega_A \xrightarrow{\varphi} M$  be a homomorphism of  $\mathcal{O}$ -modules. When we compose  $\varphi$  with the derivation  $A \xrightarrow{d} \Omega_A$ , we obtain a derivation  $A \xrightarrow{\varphi \circ d} M$ . Composition with  $d$  defines a bijection between homomorphisms  $\Omega_A \rightarrow M$  and derivations  $A \xrightarrow{\delta} M$ .

(ii)  $\Omega$  is a functor: An algebra homomorphism  $A \xrightarrow{u} B$  induces a homomorphism  $\Omega_A \xrightarrow{v} \Omega_B$  that is compatible with the ring homomorphism  $u$ , and that makes a diagram

$$\begin{array}{ccc} B & \xrightarrow{d} & \Omega_B \\ u \uparrow & & \uparrow v \\ A & \xrightarrow{d} & \Omega_A \end{array}$$

*proof.* (i) When we compose the derivation  $d$  with a homomorphism  $\varphi$ , we do get a derivation  $A \xrightarrow{\delta} M$ . In the other direction, given a derivation  $A \xrightarrow{\delta} M$ , we define a map  $\Omega_A \xrightarrow{\varphi} M$  by  $\varphi(r da) = r\delta(a)$ . It follows from the defining relations for  $\Omega_A$  that  $\varphi$  is a homomorphism of  $A$ -modules.

(ii) When  $\Omega_B$  is made into an  $A$ -module by restriction of scalars, the composed map  $A \xrightarrow{u} B \xrightarrow{d} \Omega_B$  will be a derivation to which (i) applies.  $\square$

**8.6.4. Lemma.** Let  $R$  be the polynomial ring  $\mathbb{C}[x_1, \dots, x_n]$ . The  $R$ -module of differentials  $\Omega_R$  is free, with basis  $dx_1, \dots, dx_n$ .

*proof.* The formula  $df = \sum \frac{df}{dx_i} dx_i$  follows from the defining relations. It shows that the elements  $dx_1, \dots, dx_n$  generate the  $R$ -module  $\Omega_R$ .

Let  $V$  be a free  $R$ -module with basis  $v_1, \dots, v_n$ . The map  $R \xrightarrow{\delta} V$  defined by  $\delta(f) = \sum \frac{\partial f}{\partial x_i} v_i$  is a derivation. It induces a surjective module homomorphism  $\Omega_R \xrightarrow{\varphi} V$  that sends  $dx_i$  to  $v_i$ . Since  $dx_1, \dots, dx_n$  generate  $\Omega_R$  and since  $v_1, \dots, v_n$  is a basis,  $\varphi$  is an isomorphism.  $\square$

**8.6.5. Proposition.** Let  $I$  be an ideal of an algebra  $R$ , let  $A$  be the quotient algebra  $R/I$ , and let  $dI$  denote the set of differentials  $df$  with  $f$  in  $I$ . The subset  $N = dI + I\Omega_R$  is a submodule of  $\Omega_R$ , and  $\Omega_A$  is isomorphic to the quotient module  $\Omega_R/N$ .

The proposition can be interpreted this way: Suppose that the ideal  $I$  is generated by elements  $f_1, \dots, f_r$  of  $R$ . Then  $\Omega_A$  is the quotient of  $\Omega_R$  obtained from  $\Omega_R$  by introducing these two rules:

- $df_i = 0$ , and
- multiplication by  $f_i$  is zero.

For example, let  $A$  be the quotient  $\mathbb{C}[x]/(x^n)$  of a polynomial ring in one variable and let  $\bar{x}$  be the residue of  $x$  in  $A$ . Then  $\Omega_A$  is generated by an element  $d\bar{x}$ , with the relation  $n\bar{x}^{n-1}d\bar{x} = 0$ .

*proof of Proposition 8.6.5.* First,  $I\Omega_R$  is a submodule of  $\Omega_R$ , and  $dI$  is an additive subgroup of  $\Omega_R$ . To show that  $N$  is a submodule, we must show that scalar multiplication by an element of  $R$  carries  $dI$  to  $N$ , i.e., that if  $g$  is in  $R$  and  $f$  is in  $I$ , then  $g df$  is in  $N$ . By the product rule,  $g df = d(fg) - f dg$ . Since  $I$  is an ideal,  $fg$  is in  $I$ . Then  $d(fg)$  is in  $dI$  and  $f dg$  is in  $I\Omega_R$ . So  $g df$  is in  $N$ .

The two rules shown above hold in  $\Omega_A$  because the generators  $f_i$  of  $I$  are zero in  $A$ . Therefore  $N$  is in the kernel of the surjective map  $\Omega_R \xrightarrow{v} \Omega_A$  defined by the homomorphism  $R \rightarrow A$ . The quotient module

$\overline{\Omega} = \Omega_R/N$ , is an  $A$ -module, and  $v$  defines a surjective map of  $A$ -modules  $\overline{\Omega} \xrightarrow{\overline{v}} \Omega_A$ . We show that  $\overline{v}$  is bijective. Let  $x$  be an element of  $R$ , let  $a$  be its image in  $A$ , and let  $\overline{dx}$  be the image of  $dx$  in  $\overline{\Omega}$ . The composed map  $R \xrightarrow{d} \Omega_R \rightarrow \overline{\Omega}$  is a derivation that sends  $x$  to  $\overline{dx}$ , and  $I$  is in its kernel. It defines a derivation  $R/I = A \xrightarrow{\delta} \overline{\Omega}$  that sends  $a$  to  $\overline{dx}$ . This derivation corresponds to a homomorphism of  $A$ -modules  $\Omega_A \rightarrow \overline{\Omega}$  that sends  $da$  to  $\overline{dx}$ , and that inverts  $\overline{v}$  (8.6.3).  $\square$

**8.6.6. Corollary.** *If  $A$  is a finite-type algebra, then  $\Omega_A$  is a finite  $A$ -module.*

This follows from Proposition 8.6.5 because the module of differentials on the polynomial ring  $\mathbb{C}[x_1, \dots, x_n]$  is a finite module.  $\square$

**8.6.7. Lemma.** *Let  $S$  be a multiplicative system in a domain  $A$ , and let  $S^{-1}\Omega_A$  be the module of fractions of  $\Omega_A$ . The modules  $S^{-1}\Omega_A$  and  $\Omega_{S^{-1}A}$  are canonically isomorphic. In particular, if  $K$  is the field of fractions of  $A$ , then  $K \otimes_A \Omega_A \approx \Omega_K$ .*

We have moved the symbol  $S^{-1}$  to the left for clarity.

*proof of Lemma 8.6.7.* The composition  $A \rightarrow S^{-1}A \xrightarrow{d} \Omega_{S^{-1}A}$  is a derivation that defines an  $A$ -module homomorphism  $\Omega_A \rightarrow \Omega_{S^{-1}A}$ . This map extends to an  $S^{-1}A$ -homomorphism  $S^{-1}\Omega_A \xrightarrow{\varphi} \Omega_{S^{-1}A}$  because scalar multiplication by the elements of  $S$  is invertible in  $\Omega_{S^{-1}A}$ . The relation  $ds^{-k} = -ks^{k-1}ds$  follows from the definition of a differential, and it shows that  $\varphi$  is surjective. We use the quotient rule

$$\delta(s^{-k}a) = -ks^{-k-1}a ds + s^{-k}da$$

to define a derivation  $S^{-1}A \xrightarrow{\delta} S^{-1}\Omega_A$ . That derivation will correspond to a homomorphism  $\Omega_{S^{-1}A} \rightarrow S^{-1}\Omega_A$  that inverts  $\varphi$ . However, we must show that  $\delta$  is well-defined, that  $\delta(s_1^{-k}a_1) = \delta(s_2^{-\ell}a_2)$  if  $s_1^{-\ell}a_1 = s_2^{-k}a_2$ , and that  $\delta$  is a derivation. You will be able to do this.  $\square$

Lemma 8.6.7 shows that a finite  $\mathcal{O}$ -module  $\Omega_Y$  of differentials on a variety  $Y$  is defined, such that, when  $U = \text{Spec } A$  is an affine open subset of  $Y$ ,  $\Omega_Y(U) = \Omega_A$ .

**8.6.8. Proposition.** *The module  $\Omega_Y$  of differentials on a smooth curve  $Y$  is invertible. If  $y$  is a local generator for the maximal ideal at a point  $q$ , then in a suitable neighborhood of  $q$ ,  $\Omega_Y$  will be a free  $\mathcal{O}$ -module with basis  $dy$ .*

*proof.* We may assume that  $Y$  is affine, say  $Y = \text{Spec } B$ . Let  $q$  be a point of  $Y$ , and let  $y$  be an element of  $B$  with  $v_q(y) = 1$ . To show that  $dy$  generates  $\Omega_B$  locally, we may localize, so we may suppose that  $y$  generates the maximal ideal  $\mathfrak{m}$  at  $q$ . We must show that after we localize  $B$  once more, every differential  $df$  with  $f$  in  $B$  will be a multiple of  $dy$ . Let  $c$  be the value of the function  $f$  at  $q$ : Then  $f = c + yg$  for some  $g$  in  $B$ , and because  $dc = 0$ ,  $df = g dy + y dg$ . Here  $g dy$  is in  $B dy$  and  $y dg$  is in  $\mathfrak{m}\Omega_B$ . So

$$\Omega_B = B dy + \mathfrak{m}\Omega_B$$

Let  $M$  denote the quotient module  $\Omega_B/(B dy)$ . Then  $M = \mathfrak{m}M$ . The Nakayama Lemma applies. It tells us that there is an element  $z$  in  $\mathfrak{m}$  such that  $s = 1 - z$  annihilates  $M$ . When we replace  $B$  by its localization  $B_s$ , we will have  $M = 0$  and  $\Omega_B = B dy$ , as required.

###ugh##

We must still verify that  $dy$  isn't a torsion element. If it were, say  $b dy = 0$ , then because  $dy$  is a local generator,  $\Omega_B$  would be the zero module except at the finite set of zeros of  $b$ . Since we can take for  $q$  an arbitrary point of  $Y$ , it suffices to show that the local generator  $dy$  for  $\Omega_B$  isn't zero. Let  $R = \mathbb{C}[y]$  and  $A = \mathbb{C}[y]/(y^2)$ . The module  $\Omega_R$  is free, with basis  $dy$ , and as noted above, if  $\overline{y}$  is the residue of  $y$  in  $A$ , the  $A$ -module  $\Omega_A$  is generated by  $d\overline{y}$ , with the relation  $2\overline{y} d\overline{y} = 0$ . It isn't the zero module. Proposition 5.3.7 tells us that, at our point  $q$ , the algebra  $B/\mathfrak{m}_q^2$  is isomorphic to  $A$ , and Proposition 8.6.5 tells us that  $\Omega_A$  is a quotient of  $\Omega_B$ . Since  $\Omega_A$  isn't zero, neither is  $\Omega_B$ .  $\square$

## 8.7 Trace

### (8.7.1) trace of a function



Let  $Y \xrightarrow{\pi} X$  be a branched covering of smooth curves, and let  $F$  and  $K$  be the function fields of  $X$  and  $Y$ , respectively.

The trace map  $K \xrightarrow{\text{tr}} F$  for a field extension of finite degree has been defined before (4.3.10). If  $\alpha$  is an element of  $K$ , multiplication by  $\alpha$  on the  $F$ -vector space  $K$  is an  $F$ -linear operator, and  $\text{tr}(k)$  is the trace of that operator. The trace is  $F$ -linear: If  $f_i$  are in  $F$  and  $\alpha_i$  are in  $K$ , then  $\text{tr}(\sum f_i \alpha_i) = \sum f_i \text{tr}(\alpha_i)$ . Moreover, the trace carries regular functions to regular functions: If  $X' = \text{Spec } A'$  is an affine open subset of  $X$  whose inverse image is  $Y' = \text{Spec } B'$ , then because  $A'$  is a normal algebra, the trace of an element of  $B'$  will be in  $A'$  (4.3.6). Using our abbreviated notation  $\mathcal{O}_Y$  for  $\pi_* \mathcal{O}_{Y'}$ , the trace defines a homomorphism of  $\mathcal{O}_X$ -modules

$$(8.7.2) \quad \mathcal{O}_Y \xrightarrow{\text{tr}} \mathcal{O}_X$$

Analytically, the trace can be described as a sum over the sheets of the covering. Let  $n = [Y : X]$ . Over a point  $p$  of  $X$  that isn't a branch point, there will be  $n$  points  $q_1, \dots, q_n$  of  $Y$ . If  $U$  is a small neighborhood of  $p$  in  $X$  in the classical topology, its inverse image  $V$  will consist of disjoint neighborhoods  $V_i$  of  $q_i$ , each of which maps bijectively to  $U$ . On  $V_i$ , the ring  $\mathcal{B}$  of analytic functions will be isomorphic to the ring  $\mathcal{A}$  of analytic functions on  $U$ . So  $\mathcal{B}$  is the direct sum  $\mathcal{A}_1 \oplus \dots \oplus \mathcal{A}_n$  of  $n$  copies of  $\mathcal{A}$ . If a rational function  $g$  on  $Y$  is regular on  $V$ , its restriction to  $V$  can be written as  $g = g_1 \oplus \dots \oplus g_n$ , with  $g_i$  in  $\mathcal{A}_i$ . The matrix of left multiplication by  $g$  on  $\mathcal{A}_1 \oplus \dots \oplus \mathcal{A}_n$  is the diagonal matrix with entries  $g_i$ , so

$$(8.7.3) \quad \text{tr}(g) = g_1 + \dots + g_n$$

**8.7.4. Lemma.** *Let  $Y \xrightarrow{\pi} X$  be a branched covering of smooth curves, let  $p$  be a point of  $X$ , let  $q_1, \dots, q_k$  be the fibre over  $p$ , and let  $e_i$  be the ramification index at  $q_i$ . If a rational function  $g$  on  $Y$  is regular at the points  $q_1, \dots, q_k$ , its trace is regular at  $p$ , and its value at  $p$  is  $[\text{tr}(g)](p) = e_1 g(q_1) + \dots + e_k g(q_k)$ .*

*proof.* The regularity was discussed above. If  $p$  isn't a branch point, we will have  $k = n$  and  $e_i = 1$  for all  $i$ . In this case, the lemma follows by evaluating (8.7.3). It follows by continuity for any point  $p$ . As a point  $p'$  approaches  $p$ ,  $e_i$  points  $q'$  of  $Y$  approach  $q_i$  (8.1.6). For each such point, the limit of  $g(q')$  will be  $g(q_i)$ .  $\square$

#### (8.7.5) trace of a differential

The structure sheaf is naturally contravariant. A branched covering  $Y \xrightarrow{\pi} X$  gives us an  $\mathcal{O}_X$ -module homomorphism  $\mathcal{O}_X \rightarrow \mathcal{O}_Y$ . The trace map for functions is a homomorphism of  $\mathcal{O}_X$ -modules in the opposite direction:  $\mathcal{O}_Y \xrightarrow{\text{tr}} \mathcal{O}_X$ .

Differentials are also naturally contravariant. A morphism  $Y \rightarrow X$  induces an  $\mathcal{O}_X$ -module homomorphism  $\Omega_X \rightarrow \Omega_Y$  that sends a differential  $dx$  on  $X$  to a differential on  $Y$  that we denote by  $dx$  too (8.6.3) (ii). As is true for functions, there is a trace map for differentials in the opposite direction. It is defined below, in (8.7.7), and will be denoted by  $\tau : \Omega_Y \xrightarrow{\tau} \Omega_X$ .

First, a lemma about the contravariant map  $\Omega_X \rightarrow \Omega_Y$ :

**8.7.6. Lemma. (i)** *Let  $p$  be the image in  $X$  of a point  $q$  of  $Y$ , let  $x$  and  $y$  be local generators for the maximal ideals of  $X$  and  $Y$  at  $p$  and  $q$ , respectively, and let  $e$  be the ramification index of the covering at  $q$ . Then  $dx = vy^{e-1}dy$ , where  $v$  is a local unit at  $q$ .*

**(ii)** *The canonical homomorphism  $\Omega_X \rightarrow \Omega_Y$  is injective.*

*proof. (i)* As we have noted before,  $x$  has the form  $uy^e$ , where  $u$  is a local unit. Since  $dy$  generates  $\Omega_Y$  locally, there is a rational function  $z$  that is regular at  $q$  such that  $du = zdy$ . Let  $v = yz + eu$ . Since  $eu$  is a local unit and  $yz$  is zero at  $q$ ,  $v$  is a local unit, and

$$dx = d(uy^e) = y^e z dy + ey^{e-1} u dy = vy^{e-1} dy$$

**(ii)** See (8.2.17).  $\square$

To define the trace for differentials, we begin with differentials of the functions fields. Let  $F$  and  $K$  be the function fields of  $X$  and  $Y$ , respectively. Because the  $\mathcal{O}_Y$ -module  $\Omega_Y$  is invertible, the module  $\Omega_K$  of

$K$ -differentials, which is the localization  $\Omega_Y \otimes_{\mathcal{O}} K$ , is a free  $K$ -module of rank one. Any nonzero differential will form a  $K$ -basis. We choose a nonzero  $F$ -differential  $\alpha$ . Its image in  $\Omega_K$ , which we also denote by  $\alpha$ , will be a  $K$ -basis for  $\Omega_K$ . We can, for example, take  $\alpha = dx$ , where  $x$  is a local coordinate function on  $X$ .

An element  $\beta$  of  $\Omega_K$  can be written uniquely in the form

$$\beta = g\alpha$$

where  $g$  is an element of  $K$ . The trace  $\Omega_K \xrightarrow{\tau} \Omega_F$  is defined by

$$(8.7.7) \quad \tau(\beta) = \text{tr}(g)\alpha$$

where  $\text{tr}(g)$  is the trace of the function  $g$ . Since the trace for functions is  $F$ -linear,  $\tau$  is also an  $F$ -linear map.

We need to check that  $\tau$  is independent of the choice of  $\alpha$ . If  $\alpha'$  is another nonzero  $F$ -differential, then  $f\alpha' = \alpha$  for some nonzero element  $f$  of  $F$ , and  $g\alpha = gf\alpha'$ . Since  $\text{tr}$  is  $F$ -linear,

$$\text{tr}(gf)\alpha' = \text{tr}(g)f\alpha' = \text{tr}(g)\alpha$$

Using  $\alpha'$  in place of  $\alpha$  gives the same value for the trace.

A differential of the function field  $K$  will be called a *rational differential*. A rational differential  $\beta$  is *regular* at a point  $q$  of  $Y$  if there is an affine open neighborhood  $Y' = \text{Spec } B$  of  $q$  such that  $\beta$  is an element of  $\Omega_B$ . If  $y$  is a local generator for the maximal ideal  $\mathfrak{m}_q$  and  $\beta = g dy$ , then  $\beta$  is regular at  $q$  if the rational function  $g$  is regular at  $q$ .

Let  $p$  be a point of  $X$ . Working locally at  $p$ , we may suppose that  $X$  and  $Y$  are affine,  $X = \text{Spec } A$  and  $Y = \text{Spec } B$ , that the maximal ideal at  $p$  is a principal ideal, generated by an element  $x$  of  $A$ , and that the differential  $dx$  generates  $\Omega_A$ . Let  $q_1, \dots, q_k$  be the points of  $Y$  that lie over  $p$ , and let  $e_i$  be the ramification index at  $q_i$ .

**8.7.8. Corollary. (i)** *When viewed as a differential on  $Y$ ,  $dx$  has zeros of orders  $e_i - 1$  at  $q_i$ .*

**(ii)** *If a differential  $\beta$  on  $Y$  is regular at the points  $q_i, \dots, q_k$ , it will have the form  $\beta = g dx$ , where  $g$  is a rational function with poles of orders at most  $e_i - 1$  at  $q_i$ .*

This follows from Lemma 8.7.6 (i). □

**8.7.9. Main Lemma.** *Let  $Y \xrightarrow{\pi} X$  be a branched covering. Let  $p$  be a point of  $X$ , let  $q_1, \dots, q_k$  be the points of  $Y$  that lie over  $p$ , and let  $\beta$  be a rational differential on  $Y$ .*

**(i)** *If  $\beta$  is regular at the points  $q_1, \dots, q_k$ , then its trace  $\tau(\beta)$  is regular at  $p$ .*

**(ii)** *If  $\beta$  has a simple pole at  $q_i$  and is regular at  $q_j$  when  $j \neq i$ , then  $\tau(\beta)$  is not regular at  $p$ .*

*proof.* **(i)** Corollary 8.7.8 tells us that  $\beta = g dx$ , where  $g$  has poles of orders at most  $e_i - 1$  at the points  $q_i$ . Since  $x$  has a zero of order  $e_i$  at  $q_i$ , the function  $xg$  is regular at  $q_i$ , and its value there is zero. Then  $\text{tr}(xg)$  is regular at  $p$ , and its value at  $p$  is zero (8.7.4). So  $x^{-1} \text{tr}(xg)$  is a regular function at  $p$ . Since  $\text{tr}$  is  $F$ -linear and  $x$  is in  $F$ ,  $x^{-1} \text{tr}(xg) = \text{tr}(g)$ . Therefore  $\text{tr}(g)$  and  $\tau(\beta) = \text{tr}(g)dx$  are regular at  $p$ .

**(ii)** With  $\beta = g dx$ , the function  $xg$  will be regular at  $p$ . Its value at  $q_j$  will be zero when  $j \neq i$ , and not zero when  $j = i$ . Then  $\text{tr}(xg)$  will be regular at  $p$ , but not zero there (8.7.4). Therefore  $\tau(\beta) = x^{-1} \text{tr}(xg)dx$  won't be regular at  $p$ . □

**8.7.10. Corollary.** *The trace map defines a homomorphism of  $\mathcal{O}_X$ -modules  $\Omega_Y \xrightarrow{\tau} \Omega_X$ .* □

**8.7.11. Example.** Let  $Y$  be the locus  $y^e = x$  in  $\mathbb{A}_{x,y}^2$ . Multiplication by  $\zeta = e^{2\pi i/e}$  permutes the sheets of  $Y$  over  $X$ . The trace of a power  $y^k$  is

$$(8.7.12) \quad \text{tr}(y^k) = \sum_j \zeta^{kj} y^k$$

The sum  $\sum \zeta^{kj}$  is zero unless  $k \equiv 0$  modulo  $e$ . Then  $\tau(y^r dy) = \tau(y^{r+1-e})dx = 0$  if  $r \not\equiv -1$  modulo  $e$ , but  $\tau(y^{-1} dy) = \text{tr}(e^{-1} x^{-1})dx = x^{-1} dx$  isn't regular at  $x = 0$ . □

Let  $Y \rightarrow X$  be a branched covering, and suppose that  $Y = \operatorname{Spec} B$  and  $X = \operatorname{Spec} A$  are affine. Both  $B$  and  $\Omega_B$  are torsion-free, and therefore locally free  $A$ -modules. Let's assume that they are free  $A$ -modules, that the maximal ideal of  $A$  at  $p$  is generated by an element  $x$ , and that  $\Omega_A$  is a free module of rank one with basis  $dx$ . Then  ${}_A(B, \Omega_A)$  will be a free  $A$ -module too.

As is true for any  $B$ -module,  $\Omega_B$  is isomorphic to  ${}_B(B, \Omega_B)$ . The map  $B \rightarrow \Omega_B$  that corresponds to an element  $\beta$  of  $\Omega_B$  is multiplication by  $\beta$ . It sends an element  $z$  of  $B$  to  $z\beta$ .

## rethink wording##

If  $\beta$  is a  $B$ -linear map  $B \rightarrow \Omega_B$ , then because  $\tau$  is  $A$ -linear, the composed map  $B \xrightarrow{\beta} \Omega_B \xrightarrow{\tau} \Omega_A$  will be  $A$ -linear – a homomorphism of  $A$ -modules. Thus composition with the trace  $\tau$  defines a map

$$(8.7.13) \quad \Omega_B \approx {}_B(B, \Omega_B) \xrightarrow{\tau} {}_A(B, \Omega_A)$$

**8.7.14. Theorem.** *The map (8.7.13) is an isomorphism of  $B$ -modules.*

*proof.* This theorem follows from the Main Lemma 8.7.9, when one looks closely.

Let's denote  ${}_A(B, \Omega_A)$  by  $\mathcal{H}$ . This is an  $A$ -module, but it becomes a  $B$ -module because  $B$  is a  $B$ -module (8.2.11). Scalar multiplication by an element  $b$  of  $B$  is defined as follows: Let  $B \xrightarrow{u} \Omega_A$  be an  $A$ -linear map. Then  $bu$  is the map  $[bu](z) = u(zb)$  for  $z$  in  $B$ .

Next, because  $B$  and  $\Omega_A$  are locally free  $A$ -modules,  $\mathcal{H}$  is a locally free  $A$ -module and a locally free  $B$ -module. Since  $\Omega_A$  has  $A$ -rank 1, the  $A$ -rank of  $\mathcal{H}$  is the same as the  $A$ -rank of  $B$ . Therefore the  $B$ -rank of  $\mathcal{H}$  is 1 (8.2.4)(ii). So  $\mathcal{H}$  is an invertible  $B$ -module.

The trace map  $\Omega_B \xrightarrow{\tau} \mathcal{H}$  isn't the zero map because  $\tau dx \neq 0$ . Since domain and range are invertible  $B$ -modules,  $\tau$  is an injective homomorphism. Its image, which is isomorphic to  $\Omega_B$ , is an invertible submodule of the  $B$ -module  $\mathcal{H}$ . Therefore  $\mathcal{H}$  is isomorphic to the invertible module  $\Omega_B(D)$  for some effective divisor  $D$  (8.3.7). To complete the proof of the theorem, we show that the divisor  $D$  is zero.

Suppose that  $D > 0$  and let  $q$  be a point in the support of  $D$ . We may suppose that  $q$  lies over our chosen point  $p$ . Then  $\Omega_B(q) \subset \Omega_B(D) \approx \mathcal{H}$ . We choose a rational differential  $\beta$  in  $\Omega_K$  that has a simple pole at  $q$ , and is regular at the other points of  $Y$  in the fibre over  $p$ . The Chinese Remainder Theorem allows us to do this. According to Lemma 8.7.9 (ii), the trace  $\tau(\beta)$  isn't regular at  $p$ . It isn't in  $\mathcal{H}$ .  $\square$

**Note.** This is a subtle theorem, and I don't like the proof. It is understandable, but it doesn't give much insight as to why the theorem is true. To get more insight, we would need a better understanding of differentials. My father Emil Artin said "One doesn't really understand differentials, but one can learn to work with them."

**8.7.15. Theorem.** *Let  $Y \rightarrow X$  be a branched covering of affine varieties  $X = \operatorname{Spec} A$  and  $Y = \operatorname{Spec} B$ , and let  $M$  be a finite  $B$ -module. Composition with the trace  $\Omega_B \xrightarrow{\tau} \Omega_A$  defines a bijection*

$$(8.7.16) \quad {}_B(M, \Omega_B) \xrightarrow{\tau \circ} {}_A(M, \Omega_A)$$

*proof.* We choose a resolution

$$B^m \rightarrow B^n \rightarrow M \rightarrow 0$$

of  $M$  and form a diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & {}_B(M, \Omega_B) & \longrightarrow & {}_B(B, \Omega_B)^n & \longrightarrow & {}_B(B, \Omega_B)^m \\ & & a \downarrow & & b \downarrow & & c \downarrow \\ 0 & \longrightarrow & {}_A(M, \Omega_A) & \longrightarrow & {}_A(A, \Omega_A)^n & \longrightarrow & {}_A(A, \Omega_A)^m \end{array}$$

in which the maps  $a, b, c$  are the compositions with the trace  $\tau$ , as was described above. Because the functor  $\operatorname{Hom}$  is left exact and contravariant in the first variable, the rows of this diagram are exact. Theorem 8.7.14 Shows that  $b$  and  $c$  are bijective. Therefore  $a$  is bijective too.  $\square$

Extension of this theorem to branched coverings  $Y \rightarrow X$  in which  $Y$  and  $X$  aren't affine presents no problem.

**8.7.17. Corollary.** *Let  $Y \xrightarrow{\pi} X$  be a branched covering of smooth curves, and let  $\mathcal{M}$  be a finite  $\mathcal{O}_Y$ -module. The map  ${}_Y(\mathcal{M}, \Omega_Y) \xrightarrow{\tau \circ} {}_X(\mathcal{M}, \Omega_X)$  obtained by composition with the trace  $\Omega_Y \xrightarrow{\tau} \Omega_X$  is an isomorphism of  $\mathcal{O}_X$ -modules.*  $\square$

When written without dropping the symbol  $\underline{\text{Hom}}$  or suppressing the notation for the direct image, this isomorphism becomes an isomorphism

$$\pi_* (\underline{\text{Hom}}_{\mathcal{O}_Y}(\mathcal{M}, \Omega_Y)) \xrightarrow{\tau_*} \underline{\text{Hom}}_{\mathcal{O}_X}(\pi_* \mathcal{M}, \Omega_X)$$

## 8.8 The Riemann-Roch Theorem II

### (8.8.1) the Serre dual

Let  $Y$  be a smooth projective curve, and let  $\mathcal{M}$  be a locally free  $\mathcal{O}_Y$ -module. The *Serre dual* of  $\mathcal{M}$ , which we will denote by  $\mathcal{M}^\#$ , is the module

$$(8.8.2) \quad \mathcal{M}^\# = {}_Y(\mathcal{M}, \Omega_Y) = \underline{\text{Hom}}_{\mathcal{O}_Y}(\mathcal{M}, \Omega_Y)$$

For example,  $\mathcal{O}_Y^\# = \Omega_Y$  and  $\Omega_Y^\# = \mathcal{O}_Y$ .

Since the invertible module  $\Omega_Y$  is locally isomorphic to  $\mathcal{O}_Y$ , the Serre dual  $\mathcal{M}^\#$  will be locally isomorphic to the ordinary dual  $\mathcal{M}^*$ . It will be a locally free module with the same rank as  $\mathcal{M}$ , and the bidual  $(\mathcal{M}^\#)^\#$  will be isomorphic to  $\mathcal{M}$ .

**8.8.3. Riemann-Roch Theorem, version 2.** *Let  $\mathcal{M}$  be a locally free  $\mathcal{O}_Y$ -module on a smooth projective curve  $Y$ , and let  $\mathcal{M}^\#$  be its Serre dual. Then  $\mathbf{h}^0(\mathcal{M}) = \mathbf{h}^1(\mathcal{M}^\#)$  and  $\mathbf{h}^1(\mathcal{M}) = \mathbf{h}^0(\mathcal{M}^\#)$ .*

Because  $\mathcal{M}$  and  $(\mathcal{M}^\#)^\#$  are isomorphic, the two assertions of the theorem are equivalent.

For example,  $\mathbf{h}^1(\Omega_Y) = \mathbf{h}^0(\mathcal{O}_Y) = 1$  and  $\mathbf{h}^0(\Omega_Y) = \mathbf{h}^1(\mathcal{O}_Y) = p_a$ .

If  $\mathcal{M}$  is a locally free  $\mathcal{O}_Y$ -module on a smooth projective curve  $Y$ , then

$$(8.8.4) \quad \chi(\mathcal{M}) = \mathbf{h}^0(\mathcal{M}) - \mathbf{h}^0(\mathcal{M}^\#)$$

A more precise statement of the Riemann-Roch Theorem is that  $H^1(Y, \mathcal{M})$  and  $H^0(Y, \mathcal{M}^\#)$  are dual vector spaces in a canonical way. We omit the proof of this. The fact that their dimensions are equal is enough for many applications. The canonical isomorphism becomes important only when one wants to apply the theorem to a cohomology sequence. And of course, any complex vector spaces  $V$  and  $W$  whose dimensions are equal can be made into dual spaces by the choice of a nondegenerate bilinear form  $V \times W \rightarrow \mathbb{C}$ .

Our plan is to prove Theorem 8.8.3 directly for the projective line. The structure of locally free modules on  $\mathbb{P}^1$  is very simple, so this will be easy. Following Grothendieck, we derive it for an arbitrary smooth projective curve  $Y$  by projection to  $\mathbb{P}^1$ .

Let  $Y$  be a smooth projective curve, let  $X = \mathbb{P}^1$ , and let  $Y \xrightarrow{\pi} X$  be a branched covering. Let  $\mathcal{M}$  be a locally free  $\mathcal{O}_Y$ -module, and let the Serre dual of  $\mathcal{M}$ , as defined in (8.8.2), be  $\mathcal{M}_1^\#$ :

$$\mathcal{M}_1^\# = {}_Y(\mathcal{M}, \Omega_Y)$$

The direct image of  $\mathcal{M}$  is a locally free  $\mathcal{O}_X$ -module that we are denoting by  $\mathcal{M}$  too, and we can form the Serre dual on  $X$ . Let

$$\mathcal{M}_2^D = {}_X(\mathcal{M}, \Omega_X)$$

**8.8.5. Corollary.** *The direct image  $\pi_* \mathcal{M}_1^\#$ , which we also denote by  $\mathcal{M}_1^\#$ , is isomorphic to  $\mathcal{M}_2^D$ .*

*proof.* This is Theorem 8.7.17. □

The corollary allows us to drop the subscripts from  $\mathcal{M}^\#$ . Because a branched covering is an affine morphism, the cohomology of  $\mathcal{M}$  and of its Serre dual  $\mathcal{M}^\#$  can be computed, either on  $Y$  or on  $X$ . (See (7.4.25).)

If  $Y \xrightarrow{\pi} X$  is a branched covering of projective curves and  $\mathcal{M}$  is a locally free  $\mathcal{O}_Y$ -module, then  $H^q(Y, \mathcal{M}) \approx H^q(X, \mathcal{M})$  and  $H^q(Y, \mathcal{M}^\#) \approx H^q(X, \mathcal{M}^\#)$ .

Thus it is enough to prove Riemann-Roch for the projective line.

### (8.8.6) Riemann-Roch for the projective line

The Riemann-Roch Theorem for the projective line  $X = \mathbb{P}^1$  is a simple consequence of the Birkhoff-Grothendieck Theorem, which tells us that every locally free  $\mathcal{O}_X$ -module  $\mathcal{M}$  on  $X$  is a direct sum of twisting modules  $\mathcal{O}_X(k)$ . To prove Riemann-Roch for the projective line  $X = \mathbb{P}^1$ , it suffices to the theorem for the twisting modules.

**8.8.7. Lemma.** *The module of differentials  $\Omega_X$  on  $X$  is isomorphic to the twisting module  $\mathcal{O}_X(-2)$ .*

*proof.* Since  $\Omega_X$  is invertible, the Birkhoff-Grothendieck Theorem tells us that it is a twisting module  $\mathcal{O}_X(k)$ . We only need to identify the integer  $k$ . On the standard open subset  $\mathbb{U}^0 = \text{Spec } \mathbb{C}[x]$ , the module of differentials is free, with basis  $dx$ , and  $z = x^{-1}$  is the coordinate on  $\mathbb{U}^1 = \text{Spec } \mathbb{C}[z]$ . Then  $dx = d(z^{-1}) = -z^{-2}dz$  describes the differential  $dx$  on  $\mathbb{U}^1$ . Since the point  $p$  at infinity is  $\{z = 0\}$ ,  $dx$  has a pole of order 2 there. It is a global section of  $\Omega_X(2p)$ , and as a section of that module, it isn't zero anywhere. So multiplication by  $dx$  defines an isomorphism  $\mathcal{O} \rightarrow \Omega_X(2p)$  that sends 1 to  $dx$ . Tensoring with  $\mathcal{O}(-2p)$ , we find that  $\Omega_X$  is isomorphic to  $\mathcal{O}(-2p)$ .  $\square$

**8.8.8. Lemma.** *Let  $\mathcal{M}$  and  $\mathcal{N}$  be locally free  $\mathcal{O}$ -modules on the projective line  $X$ . Then  ${}_X(\mathcal{M}(r), \mathcal{N})$  is canonically isomorphic to  ${}_X(\mathcal{M}, \mathcal{N}(-r))$ .*

*proof.* When we tensor a homomorphism  $\mathcal{M}(r) \xrightarrow{\varphi} \mathcal{N}$  with  $\mathcal{O}(-r)$ , we obtain a homomorphism  $\mathcal{M} \rightarrow \mathcal{N}(-r)$ , and tensoring with  $\mathcal{O}(r)$  is the inverse operation.  $\square$

The Serre dual  $\mathcal{O}(n)^\#$  of  $\mathcal{O}(n)$  is therefore

$$\mathcal{O}(n)^\# = {}_X(\mathcal{O}(n), \Omega_X) \approx \mathcal{O}(-2-n)$$

To prove Riemann-Roch for  $X = \mathbb{P}^1$ , we must show that

$$\mathbf{h}^0(X, \mathcal{O}(n)) = \mathbf{h}^1(X, \mathcal{O}(-2-n)) \quad \text{and} \quad \mathbf{h}^1(X, \mathcal{O}(n)) = \mathbf{h}^0(X, \mathcal{O}(-2-n))$$

This follows from the computation of cohomology of the twisting modules (Theorem 7.5.4).  $\square$

## 8.9 Using Riemann-Roch

### (8.9.1) genus

Three closely related numbers associated to a smooth projective curve  $Y$  are: its *topological genus*  $g$ , its *arithmetic genus*  $p_a = \mathbf{h}^1(\mathcal{O}_Y)$ , and the *degree*  $\delta$  of the module of differentials  $\Omega_Y$ .

**8.9.2. Theorem.** *Let  $Y$  be a smooth projective curve. The topological genus  $g$  and the arithmetic genus  $p_a$  of  $Y$  are equal, and the degree  $\delta$  of the module  $\Omega_Y$  is  $2p_a - 2$ , which is equal to  $2g - 2$ .*

*proof.* Let  $Y \xrightarrow{\pi} X$  be a branched covering of  $X = \mathbb{P}^1$ . The topological Euler characteristic  $e(Y)$ , which is  $2 - 2g$ , can be computed in terms of the branching data for the covering (see (1.7.27)). Let  $q_i$  be the ramification points in  $Y$ , and let  $e_i$  be the ramification index at  $q_i$ . Then  $e_i$  sheets of the covering come together at  $q_i$ . If the degree of  $Y$  over  $X$  is  $n$ , then since  $e(X) = 2$ ,

$$(8.9.3) \quad 2 - 2g = e(Y) = ne(X) - \sum (e_i - 1) = 2n - \sum (e_i - 1)$$

We compute the degree  $\delta$  of  $\Omega_Y$  in two ways. First, the Riemann-Roch Theorem tells us that  $\mathbf{h}^0(\Omega_Y) = \mathbf{h}^1(\mathcal{O}_Y) = p_a$  and  $\mathbf{h}^1(\Omega_Y) = \mathbf{h}^0(\mathcal{O}_Y) = 1$ . So  $\chi(\Omega_Y) = -\chi(\mathcal{O}_Y) = p_a - 1$ . The Riemann-Roch Theorem also tells us that  $\chi(\Omega_Y) = \delta + 1 - p_a$  (8.3.8). Therefore

$$(8.9.4) \quad \delta = 2p_a - 2$$

Next, we compute  $\delta$  by computing the divisor of the differential  $dx$  on  $Y$ ,  $x$  being a coordinate in  $X$ . Let  $q_i$  be one of the ramification points in  $Y$ , and let  $e_i$  be the ramification index at  $q_i$ . Then  $dx$  has a zero of order  $e_i - 1$  at  $q_i$ . On  $X$ ,  $dx$  has a pole of order 2 at  $\infty$ . Let's suppose that the point at infinity isn't a branch point. Then there will be  $n$  points of  $Y$  at which  $dx$  has a pole of order 2,  $n$  being the degree of  $Y$  over  $X$ . The degree of  $\Omega_Y$  is therefore

$$(8.9.5) \quad \delta = \text{zeros} - \text{poles} = \sum (e_i - 1) - 2n$$

Combining (8.9.5) with (8.9.3), one sees that  $\delta = 2g - 2$ . Since we also have  $\delta = 2p_a - 2$ ,  $g = p_a$ .  $\square$

**8.9.6. Corollary.** *Let  $D$  be a divisor on a smooth projective curve  $Y$  of genus  $g$ . If  $\deg D > 2g - 2$  then  $h^1(\mathcal{O}(D)) = 0$ . If  $\deg D \leq g - 2$ , then  $h^1(\mathcal{O}(D)) > 0$ .*

*proof.* This follows from Corollary 8.4.8 (v) and (vi).  $\square$

### (8.9.7) curves of genus zero

Let  $Y$  be a smooth projective curve  $Y$  of genus zero, and let  $p$  be a point of  $Y$ . The exact sequence

$$0 \rightarrow \mathcal{O}_Y \rightarrow \mathcal{O}_Y(p) \rightarrow \epsilon \rightarrow 0$$

where  $\epsilon$  is a one-dimensional module supported at  $p$  (8.4.6), gives us an exact cohomology sequence

$$0 \rightarrow H^0(Y, \mathcal{O}_Y) \rightarrow H^0(Y, \mathcal{O}_Y(p)) \rightarrow H^0(Y, \epsilon) \rightarrow 0$$

The zero on the right is due to the fact that, because  $p_a = 0$ ,  $H^1(Y, \mathcal{O}_Y) = 0$ . We also have  $h^0(\mathcal{O}_Y) = h^0(\epsilon) = 1$ , so  $h^0(\mathcal{O}_Y(p)) = 2$ . We choose a basis  $(1, x)$  for  $H^0(Y, \mathcal{O}_Y(p))$ , 1 being the constant function and  $x$  being a nonconstant function with a single pole of order 1 at  $p$ . This basis defines a point of  $\mathbb{P}^1$  with values in the function field  $K$  of  $Y$ , and therefore a morphism  $Y \xrightarrow{\varphi} \mathbb{P}^1$ . Because  $x$  has just one pole of order 1, it takes every value exactly once. Therefore  $\varphi$  is bijective. It is a map of degree 1, and therefore an isomorphism (8.1.3).

**8.9.8. Corollary.** *Every smooth projective curve of genus zero is isomorphic to the projective line  $\mathbb{P}^1$ .*  $\square$

A *rational curve* is a curve (smooth or not) whose function field is isomorphic to the field  $\mathbb{C}(t)$  of rational functions in one variable. A smooth projective curve of genus zero is a rational curve.

### (8.9.9) curves of genus one

A smooth projective curve of genus 1 is called an *elliptic curve*. The Riemann-Roch Theorem tells us that on an elliptic curve  $Y$ ,

$$\chi(\mathcal{O}(D)) = \deg D$$

Since  $h^0(\Omega_Y) = h^1(\mathcal{O}_Y) = 1$ ,  $\Omega_Y$  has a nonzero global section  $\omega$ . Since  $\Omega_Y$  has degree zero (8.9.2),  $\omega$  doesn't vanish anywhere. Multiplication by  $\omega$  defines an isomorphism  $\mathcal{O} \rightarrow \Omega_Y$ . So  $\Omega_Y$  is a free module of rank one. It follows that the Serre dual  $\mathcal{M}^\#$  of an  $\mathcal{O}$ -module  $\mathcal{M}$  is isomorphic to the ordinary dual  $\mathcal{M}^*$ .

The next lemma follows from Riemann-Roch.

**8.9.10. Lemma.** *Let  $p$  be a point of an elliptic curve  $Y$ . For any  $r > 0$ ,  $h^0(\mathcal{O}(rp)) = r$ , and  $h^1(\mathcal{O}(rp)) = 0$ .*  $\square$

Since  $H^0(Y, \mathcal{O}_Y) \subset H^0(Y, \mathcal{O}_Y(p))$ , and since both spaces have dimension one, they are equal. So (1) is a basis for  $H^0(Y, \mathcal{O}_Y(p))$ . We choose a basis  $(1, x)$  for the two-dimensional space  $H^1(Y, \mathcal{O}_Y(2p))$ . Then  $x$  isn't a section of  $\mathcal{O}(p)$ . It has a pole of order precisely 2 at  $p$ . Next, we choose a basis  $(1, x, y)$  for  $H^1(Y, \mathcal{O}_Y(3p))$ . So  $x$  and  $y$  are functions with poles of orders 2 and 3, respectively, at  $p$ , and no other poles. The point  $(1, x, y)$  of  $\mathbb{P}^2$  with values in  $K$  determines a morphism  $Y \xrightarrow{\varphi} \mathbb{P}^2$ . Let  $u, v, w$  be coordinates in  $\mathbb{P}^2$ . The map  $\varphi$  sends

a point  $q$  distinct from  $p$  to  $(u, v, w) = (1, x(q), y(q))$ . Since  $Y$  has dimension one,  $\varphi$  is a finite morphism. Its image will be a closed subvariety of  $\mathbb{P}^2$  of dimension one. Since  $(1, x, y)$  are independent, the image isn't contained in a line.

To determine the image of the point  $p$ , we multiply  $(1, x, y)$  by  $\lambda = y^{-1}$  to normalize the second coordinate to 1, obtaining the equivalent vector  $(y^{-1}, xy^{-1}, 1)$ . The rational function  $y^{-1}$  has a zero of order 3 at  $p$ , and  $xy^{-1}$  has a simple zero there. Evaluating at  $p$ , we see that the image of  $p$  is the point  $(0, 0, 1)$ .

Let  $Y'$  be the image of  $Y$ , which is a curve in  $\mathbb{P}^2$ . The map  $Y \rightarrow \mathbb{P}^2$  restricts to a finite morphism  $Y \rightarrow Y'$ . Let  $\ell$  be a generic line  $\{au + bv + cw = 0\}$  in  $\mathbb{P}^2$ . The rational function  $a + bx + cy$  on  $Y$  has a pole of order 3 at  $p$  and no other pole. It takes every value, including zero, three times, and the set of points  $q$  of  $Y$  at which  $a + bx + cy$  is zero is the inverse image of the intersection  $Y' \cap \ell$ . The only possibilities for the degree of  $Y'$  are 1 and 3. Since  $1, x, y$  are independent,  $Y'$  isn't a line. So the image  $Y'$  is a cubic curve (Corollary 1.3.9).

To determine the image, we look for a cubic relation among the functions  $1, x, y$  on  $Y$ . The seven monomials  $1, x, y, x^2, xy, x^3, y^2$  have poles at  $p$  of orders  $0, 2, 3, 4, 5, 6, 6$ , respectively, and no other poles. They are sections of  $\mathcal{O}_Y(6p)$ . Riemann-Roch tells us that  $h^0(\mathcal{O}_Y(6p)) = 6$ . So those seven functions are dependent. The linear dependency relation gives us a cubic equation among  $x$  and  $y$ , which we write in the form

$$cy^2 + (a_1x + a_3)y + (a_0x^3 + a_2x^2 + a_4x + a_6) = 0$$

There can be no linear relation among functions whose orders of pole at  $p$  are distinct. So when we delete either  $x^3$  or  $y^2$  from the list of monomials, we obtain an independent set of six functions that form a basis for the six-dimensional space  $H^0(Y, \mathcal{O}(6p))$ . In the cubic relation, the coefficients  $c$  and  $a_0$  aren't zero. We can scale  $y$  and  $x$  to normalize  $c$  and  $a_0$  to 1. We eliminate the linear term in  $y$  from this relation by substituting  $y - \frac{1}{2}(a_1x + a_3)$  for  $y$ . Next, we eliminate the quadratic term in  $x$  by substituting  $x - \frac{1}{3}a_2$  for  $x$ . Bringing the terms in  $x$  to the other side of the equation, we are left with a cubic relation

$$y^2 = x^3 + a_4x + a_6$$

The coefficients  $a_4$  and  $a_6$  have changed, of course.

The cubic curve  $Y'$  defined by the homogenized equation  $y^2z = x^3 + a_4xz^2 + a_6z^3$  is the image of  $Y$ . This curve  $Y'$  meets a generic line  $ax + by + cz = 0$  in three points and, as we saw above, its inverse image in  $Y$  consists of three points too. Therefore the morphism  $Y \xrightarrow{\varphi} Y'$  is generically injective, and  $Y$  is the normalization of  $Y'$ . Corollary 7.6.3 computes the cohomology of  $Y'$ :  $h^0(\mathcal{O}_{Y'}) = h^1(\mathcal{O}_{Y'}) = 1$ . This tells us that  $h^q(\mathcal{O}_{Y'}) = h^q(\mathcal{O}_Y)$  for all  $q$ . Let's denote the direct image of  $\mathcal{O}_Y$  by the same symbol  $\mathcal{O}_Y$ . The quotient  $\mathcal{F} = \mathcal{O}_Y/\mathcal{O}_{Y'}$  is a torsion module with no global sections, so it is zero (8.2.1) (ii).

**8.9.11. Corollary.** *Every elliptic curve is isomorphic to a cubic curve in  $\mathbb{P}^2$ .* □

### (8.9.12) the group law on an elliptic curve

The points of an elliptic curve form an abelian group, once one chooses a point to be the identity element.

We choose a point of  $Y$ , and label it  $o$ . We'll write the law of composition in the group as  $p \oplus q$ , using the symbol  $\oplus$  to distinguish this sum, which is a point of  $Y$ , from the divisor  $p + q$ .

Let  $p$  and  $q$  be points of  $Y$ . To define  $p \oplus q$ , we compute the cohomology of  $\mathcal{O}_Y(p + q - o)$ . It follows from Riemann-Roch that  $h^0(\mathcal{O}_Y(p + q - o)) = 1$  and that  $h^1(\mathcal{O}_Y(p + q - o)) = 0$ . There is a nonzero function  $f$ , unique up to scalar factor, with simple poles at  $p$  and  $q$  and a zero at  $o$ . This function has exactly one other zero. That zero is defined to be the sum  $p \oplus q$  in the group. In terms of linearly equivalent divisors,  $s = p \oplus q$  is the unique point such that  $s$  is linearly equivalent to  $p + q - o$ , or such that  $p + q$  is linearly equivalent to  $o + s$ .

**8.9.13. Proposition.** *The law of composition  $\oplus$  defined above makes an elliptic curve into an abelian group.*

The proof is an exercise. □

### (8.9.14) interlude: maps to projective space

Let  $Y$  be a smooth projective curve. We have seen that any set  $(f_0, \dots, f_n)$  of rational functions on  $Y$  defines a morphism  $Y \xrightarrow{\varphi} \mathbb{P}^n$  (5.3.3). As a reminder, let  $q$  be a point of  $Y$  and let  $g_j = f_j/f_i$ , where  $i$  is an index such that  $f_i$  has the minimum value  $v_q(f_i)$ . Then  $g_j$  are regular at  $q$  for all  $j$ , and the morphism  $\varphi$  sends the point  $q$  to  $(g_0(q), \dots, g_n(q))$ . For example, the inverse image  $Y^0 = \varphi^{-1}(\mathbb{U}^0)$  of the standard open set  $\mathbb{U}^0$  is the set of points at which the functions  $g_j = f_j/f_0$  are regular. If  $q$  is such a point, then  $\varphi(q) = (1, g_1(q), \dots, g_n(q))$ .

**8.9.15. Lemma.** *Let  $Y \xrightarrow{\varphi} \mathbb{P}^n$  be the morphism of a smooth projective curve  $Y$  to projective space that is defined by a set  $(f_0, \dots, f_n)$  of rational functions on  $Y$ .*

(i) *If the space spanned by  $\{f_0, \dots, f_n\}$  has dimension at least two, then  $\varphi$  is not a constant function.*

(ii) *If  $\{f_0, \dots, f_n\}$  are linearly independent, the image isn't contained in any hyperplane.*  $\square$

The *degree*  $d$  of a nonconstant morphism  $\varphi$  from a projective curve  $Y$ , smooth or not, to projective space  $\mathbb{P}^n$ , is the number of points of the inverse image  $\varphi^{-1}(H)$  of a generic hyperplane  $H$  in  $\mathbb{P}^n$ . We check that this number is well-defined. Say that  $H$  is the locus  $h(x) = 0$ , where  $h = \sum a_i x_i$ , and that another generic hyperplane  $G$  is the locus  $g(x) = 0$ , where  $g = \sum b_i x_i$ . Let  $f(x) = h/g$ . The divisor of the rational function  $\tilde{f} = f \circ \varphi$  on  $Y$  is  $\varphi^{-1}H - \varphi^{-1}H'$ .

### (8.9.16) base points

If  $D$  is a divisor on the smooth projective curve  $Y$ , a basis  $(f_0, \dots, f_k)$  of global sections of  $\mathcal{O}(D)$  defines a morphism  $Y \rightarrow \mathbb{P}^{k-1}$ . This is the most common way to construct such a morphism, though one could use any set of rational functions.

If a global section of  $\mathcal{O}(D)$  vanishes at a point  $p$  of  $Y$ , it is a section of  $\mathcal{O}(D - p)$ . A point  $p$  is a *base point* of  $\mathcal{O}(D)$  if every global section of  $\mathcal{O}(D)$  vanishes at  $p$ . A base point can be described in terms of the usual exact sequence

$$0 \rightarrow \mathcal{O}(D-p) \rightarrow \mathcal{O}(D) \rightarrow \epsilon \rightarrow 0$$

The point  $p$  is a base point if  $\mathbf{h}^0(\mathcal{O}(D-p)) = \mathbf{h}^0(\mathcal{O}(D))$ , or if  $\mathbf{h}^1(\mathcal{O}(D-p)) = \mathbf{h}^1(\mathcal{O}(D)) - 1$ .

Let  $Y \xrightarrow{\pi} \mathbb{P}^n$  is a morphism. The *degree*  $d$  of  $\pi$  is the number of points in the inverse image of a generic hyperplane.

**8.9.17. Lemma.** *Let  $D$  be a divisor on a smooth projective curve  $Y$ , and suppose that  $H^0(\mathcal{O}(D)) \neq 0$ . Let  $Y \xrightarrow{\varphi} \mathbb{P}^n$  be the morphism defined by a basis of global sections.*

(i) *The image of  $\varphi$  isn't contained in any hyperplane.*

(ii) *If  $\mathcal{O}(D)$  has no base points, the degree  $r$  of the morphism  $\varphi$  is equal to degree of  $D$ . If there are base points, the degree is lower.*  $\square$

### (8.9.18) canonical divisors

Because the module  $\Omega_Y$  of differentials on a smooth curve  $Y$  is invertible, it is isomorphic to  $\mathcal{O}(K)$  for some divisor  $K$ . Such a divisor  $K$  is called a *canonical divisor*. It is often convenient to represent  $\Omega_Y$  as a module  $\mathcal{O}(K)$ , though the canonical divisor  $K$  isn't unique. It is determined only up to linear equivalence (see (8.3.11)).

When written in terms of a canonical divisor  $K$ , the Serre dual of an invertible module  $\mathcal{O}(D)$  will be  $\mathcal{O}(D)^\# = \mathcal{O}(\mathcal{O}(D), \mathcal{O}(K)) \approx \mathcal{O}(K-D)$ . With this notation, the Riemann-Roch Theorem for  $\mathcal{O}(D)$  becomes

$$(8.9.19) \quad \mathbf{h}^0(\mathcal{O}(D)) = \mathbf{h}^1(\mathcal{O}(K-D)) \quad \text{and} \quad \mathbf{h}^1(\mathcal{O}(D)) = \mathbf{h}^0(\mathcal{O}(K-D)) \quad \square$$



**8.9.20. Proposition.** Let  $K$  be a canonical divisor on a smooth projective curve  $Y$  of genus  $g > 0$ .

(i)  $\mathcal{O}(K)$  has no base point.

(ii) Every point  $p$  of  $Y$  is a base point of  $\mathcal{O}(K+p)$ .

*proof.* (i) Let  $p$  be a point of  $Y$ . We apply Riemann-Roch to the exact sequence

$$0 \rightarrow \mathcal{O}(K-p) \rightarrow \mathcal{O}(K) \rightarrow \epsilon_1 \rightarrow 0$$

where  $\epsilon_1$  denotes a one-dimensional module supported on a point  $p$ . The Serre duals  $\mathcal{O}$  and  $\mathcal{O}(p)$  of  $\mathcal{O}(K)$  and  $\mathcal{O}(K-p)$ , respectively, form an exact sequence

$$0 \rightarrow \mathcal{O} \rightarrow \mathcal{O}(p) \rightarrow \epsilon_2 \rightarrow 0$$

When  $Y$  has positive genus, there is no rational function on  $Y$  with just one simple pole. So  $h^0(\mathcal{O}(p)) = h^0(\mathcal{O}) = 1$ . Riemann-Roch tells us that  $h^1(\mathcal{O}(K-p)) = h^1(\mathcal{O}(K)) = 1$ . The cohomology sequence

$$0 \rightarrow H^0(\mathcal{O}(K-p)) \rightarrow H^0(\mathcal{O}(K)) \rightarrow [1] \rightarrow H^1(\mathcal{O}(K-p)) \rightarrow H^1(\mathcal{O}(K)) \rightarrow 0$$

shows that  $h^0(\mathcal{O}(K-p)) = h^0(\mathcal{O}(K)) - 1$ . So  $p$  is not a base point.

(ii) Here, the relevant sequence is

$$0 \rightarrow \mathcal{O}(K) \rightarrow \mathcal{O}(K+p) \rightarrow \epsilon_3 \rightarrow 0$$

The Serre dual of  $\mathcal{O}(K+p)$  is  $\mathcal{O}(-p)$ , which has no global section. Therefore  $h^1(\mathcal{O}(K+p)) = 0$ , while  $h^1(\mathcal{O}(K)) = h^0(\mathcal{O}) = 1$ . The cohomology sequence

$$0 \rightarrow h^0(\mathcal{O}(K)) \rightarrow h^0(\mathcal{O}(K+p)) \rightarrow [1] \rightarrow h^1(\mathcal{O}(K)) \rightarrow h^1(\mathcal{O}(K+p)) \rightarrow 0$$

shows that  $H^0(\mathcal{O}(K)) = H^0(\mathcal{O}(K+p))$ . So  $p$  is a base point of  $\mathcal{O}(K+p)$ .  $\square$

### (8.9.21) hyperelliptic curves

A *hyperelliptic curve*  $Y$  is a smooth projective curve of genus  $g > 1$  that can be represented as a branched double covering of the projective line. So  $Y$  is hyperelliptic if there is a morphism  $Y \xrightarrow{\pi} X$  of degree two, with  $X = \mathbb{P}^1$ .

The topological Euler characteristic of a hyperelliptic curve  $Y$  can be computed in terms of the covering  $Y \rightarrow X$ , which will be branched at a finite set  $p_1, \dots, p_n$  of  $n$  points. Since  $\pi$  has degree two, the multiplicity of a branch point will be  $e = 2$ . The Euler characteristic is therefore  $e(Y) = 2e(X) - n = 4 - n$ . Since  $e(Y) = 2 - 2g$ , the number of branch points is  $n = 2g + 2$ . So when  $g = 3$ ,  $n = 8$ .

It would take some experimentation to guess that the next remarkable theorem might be true, and some time to find a proof.

**8.9.22. Theorem.** Let  $K$  be a canonical divisor on a hyperelliptic curve  $Y$ , and let  $Y \xrightarrow{\pi} X = \mathbb{P}^1$  be the associated branched covering of degree 2. The morphism  $Y \xrightarrow{\kappa} \mathbb{P}^{g-1}$  defined by the global sections of  $\Omega_Y = \mathcal{O}(K)$  factors through  $X$ . There is a morphism  $X \xrightarrow{u} \mathbb{P}^{g-1}$  such that  $\pi = \kappa \circ u$ :  $Y \xrightarrow{\pi} X \xrightarrow{u} \mathbb{P}^{g-1}$ .

**8.9.23. Corollary.** A curve of genus  $g \geq 2$  can be presented as a branched covering of  $\mathbb{P}^1$  of degree 2 in at most one way.  $\square$

*proof of Theorem 8.9.22.*

Let  $x$  be an affine coordinate in  $X$ , so that the standard affine open subset  $\mathbb{U}^0$  of  $X$  is  $\text{Spec } \mathbb{C}[x]$ . We may suppose that the point  $p_\infty$  at infinity isn't a branch point of the covering. Let  $Y^0 = \pi^{-1}\mathbb{U}^0$ . Then  $Y^0$  will have an equation of the form

$$y^2 = f(x)$$

where  $f$  is a polynomial with  $n = 2g + 2$  simple roots. There will be two points of  $Y$  above the point  $p_\infty$ . They are interchanged by the automorphism  $y \rightarrow -y$ . Let's call those points  $q_1$  and  $q_2$ .

We start with the differential  $dx$ , which we view as a differential on  $Y$ . Then  $2y dy = f'(x)dx$ . Since  $f$  has simple roots,  $f'$  doesn't vanish at any of them. Therefore  $dx$  has simple zeros on  $Y$  above the roots of  $f$ . We also have a regular function on  $Y^0$  with simple roots at those points, namely the function  $y$ . Therefore the differential  $\omega = \frac{dx}{y}$  is regular and nowhere zero on  $Y^0$ . Because the degree of a differential on  $Y$  is  $2g - 2$  (??),  $\omega$  has a total of  $2g - 2$  zeros at infinity. By symmetry,  $\omega$  has zeros of order  $g - 1$  at each of two points  $q_1$  and  $q_2$ . So  $K = (g-1)q_1 + (g-1)q_2$  is a canonical divisor on  $Y$ , i.e.,  $\Omega_Y \approx \mathcal{O}_Y(K)$ .

Since  $K$  has zeros of order  $g - 1$  at infinity, the rational functions  $1, x, x^2, \dots, x^{g-1}$ , viewed as functions on  $Y$ , are among the global sections of  $\mathcal{O}_Y(K)$ . They are independent, and there are  $g$  of them. Since  $h^0(\mathcal{O}_Y(K)) = g$ , they form a basis of  $H^0(\mathcal{O}_Y(K))$ . The map  $Y \rightarrow \mathbb{P}^{g-1}$  defined by the global sections of  $\mathcal{O}_Y(K)$  evaluates these powers of  $x$ , so it factors through the double covering  $Y \xrightarrow{\pi} X$ .  $\square$

### (8.9.24) canonical embedding

Let  $Y$  be a smooth projective curve of genus  $g \geq 2$ , and let  $K$  be a canonical divisor on  $Y$ . Since  $\mathcal{O}(K)$  has no base point (??), its global sections define a morphism  $Y \xrightarrow{\kappa} \mathbb{P}^{g-1}$ , the *canonical map* whose degree is equal to the degree  $2g - 2$  of the canonical divisor.

**8.9.25.** *Let  $Y$  be a smooth projective curve of genus  $g$  at least two. If  $Y$  is not hyperelliptic, the canonical map embeds  $Y$  as a closed subvariety of projective space  $\mathbb{P}^{g-1}$ .*

*proof.* We show first that, if  $\kappa$  isn't an injective map, then  $Y$  is hyperelliptic. Let  $p$  and  $q$  be distinct points such that  $\kappa(p) = \kappa(q)$ . We may assume that the canonical divisor  $K$  is effective, and that  $p$  and  $q$  are not in its support. We inspect the global sections of  $\mathcal{O}(K - p - q)$ . Since  $\kappa(p) = \kappa(q)$ , any global section of  $\mathcal{O}(K)$  that vanishes at  $p$  vanishes at  $q$  too. Therefore  $\mathcal{O}(K - p)$  and  $\mathcal{O}(K - p - q)$  have the same global sections, and  $q$  is a base point of  $\mathcal{O}(K - p)$ . We've computed the cohomology of  $\mathcal{O}(K - p)$ :  $h^0(\mathcal{O}(K - p)) = g - 1$  and  $h^1(\mathcal{O}(K - p)) = 1$ . Then  $h^0(\mathcal{O}(K - p - q)) = g - 1$  and  $h^1(\mathcal{O}(K - p - q)) = 2$ . The Serre dual of  $\mathcal{O}(K - p - q)$  is  $\mathcal{O}(p + q)$ , so by Riemann-Roch,  $h^0(\mathcal{O}(p + q)) = 2$ . If  $D$  is a divisor of degree one on a curve of positive genus, then  $h^0(\mathcal{O}(D)) \leq 1$  (Proposition ??). Therefore  $\mathcal{O}(p + q)$  has no base point. Its global sections define a morphism  $Y \rightarrow \mathbb{P}^1$  of degree 2. So  $Y$  is hyperelliptic. Conversely, if  $Y$  is hyperelliptic, Theorem 8.9.22 shows that  $\kappa$  has degree 2.

If  $Y$  isn't hyperelliptic, the canonical map is injective, so  $Y$  is mapped bijectively to its image  $Y'$  in  $\mathbb{P}^{g-1}$ . This almost proves the theorem, but: Can  $Y'$  have a cusp? We must show that the bijective map  $Y \xrightarrow{\kappa} Y'$  is an isomorphism.

We go over the computation made above for a pair of points  $p, q$ , this time taking  $q = p$ . The computation is the same. It shows that, since  $Y$  isn't hyperelliptic,  $p$  isn't a base point of  $\mathcal{O}_Y(K - p)$ . Therefore  $h^0(\mathcal{O}_Y(K - 2p)) = h^0(\mathcal{O}_Y(K - p)) - 1$ . This tells us that there is a global section  $f$  of  $\mathcal{O}_Y(K)$  that has a zero of order exactly 1 at  $p$ . When properly interpreted, this fact shows that  $\kappa$  doesn't collapse any tangent vectors to  $Y$ , and therefore that  $\kappa$  is an isomorphism. Since we haven't discussed tangent vectors, we prove this directly.

Since  $\kappa$  is a bijective, finite morphism, it is an integral morphism. The function fields of  $Y$  and its image  $Y'$  are equal, and  $Y$  is the normalization of  $Y'$ . Moreover,  $\kappa$  is an isomorphism except on a finite set.

We work locally at a point  $p$  of  $Y'$ . When we restrict the global section  $f$  of  $\mathcal{O}_Y(K)$  found above to the image  $Y'$ , we obtain an element of the maximal ideal  $\mathfrak{m}'$  of  $\mathcal{O}_{Y'}$  at  $p$ , that we denote by  $x$ . On  $Y$ , this element has a zero of order one at  $p$ , and therefore it is a local generator for the maximal ideal  $\mathfrak{m}_p$  of  $\mathcal{O}_Y$ . We may assume that  $x$  generates  $\mathfrak{m}_p$ , and that the quotient  $\mathcal{F} = \mathcal{O}_Y / \mathcal{O}_{Y'}$  is a finite-dimensional vector space supported at  $p$ .

We multiply the short exact sequence  $0 \rightarrow \mathcal{O}_{Y'} \xrightarrow{i} \mathcal{O}_Y \xrightarrow{\pi} \mathcal{F} \rightarrow 0$  by  $x$ . The cokernels of the multiplication maps form an exact sequence

$$\mathcal{O}_{Y'} / x\mathcal{O}_{Y'} \xrightarrow{\bar{i}} \mathcal{O}_Y / x\mathcal{O}_Y \xrightarrow{\bar{\pi}} \mathcal{F} / x\mathcal{F} \rightarrow 0$$

Since  $x$  generates  $\mathcal{M}$ ,  $\mathcal{O}_Y / x\mathcal{O}_Y$  is the residue field  $\kappa(p)$ , which has dimension one. The map  $\bar{i}$  isn't zero because it sends the residue of 1 in  $\mathcal{O}_{Y'} / x\mathcal{O}_{Y'}$  to the residue of 1 in  $\kappa(p)$ . Therefore  $\bar{i}$  is surjective. This shows that  $\mathcal{F} / x\mathcal{F} = 0$ . But since  $\mathcal{F}$  is a finite  $\mathcal{O}_{Y'}$ -module and  $x$  is in the maximal ideal  $\mathfrak{m}'$ , the quotient  $\mathcal{F} / x\mathcal{F}$  can't be zero unless  $\mathcal{F}$  is zero (5.1.22). Therefore  $\mathcal{O}_{Y'} = \mathcal{O}_Y$ .  $\square$

**8.9.26. Examples.** Let  $Y$  be a smooth projective curve of genus  $g$ .

(i) When  $g = 2$ , the canonical morphism  $\kappa$  is a map of degree  $2g - 2 = 2$  from  $Y$  to  $\mathbb{P}^1$ . Every smooth projective curve of genus 2 is hyperelliptic.

(ii) When  $g = 3$ ,  $\kappa$  is a morphism of degree 4 from  $Y$  to  $\mathbb{P}^2$ . If  $Y$  isn't hyperelliptic, its image will be a plane curve of degree 4, isomorphic to  $Y$ . The genus of a smooth projective curve of degree 4 is  $\binom{3}{2} = 3$  (1.7.29), which checks.

The number of moduli of curves of degree 3 (the number of essential parameters) is obtained this way: There are 15 monomials of degree 4 in three variables. The group  $GL_3$  of dimension 9 operates by coordinate changes. So the number of moduli is  $15 - 9 = 6$ . When a hyperelliptic curve of genus 3 is represented as a branched double covering of  $\mathbb{P}^1$ , there will be 8 branch points. The group  $GL_2$  of dimension 4 operates on the branch points, but scalars don't move them. So the number of moduli of hyperelliptic curves of genus three is  $8 - 3 = 5$ . Since  $5 < 6$ , this agrees with the fact that not all curves of genus three are hyperelliptic.

(iii) When  $g = 4$ ,  $\kappa$  is a morphism of degree 6 from  $Y$  to  $\mathbb{P}^3$ , and it becomes harder to count moduli. It is a fact that the number of moduli of curves of any genus  $g$  is  $3g - 3$ . The number of moduli of hyperelliptic curves of genus  $g$  is easy to compute. It is  $2g - 1$ . □

## Index

- $S$ -fractions, 103
- $\mathcal{O}$ -module, 121
- Bézout, Etienne (1730-1783), 153
- Birkhoff, George David (1884-1944), 165
- Chevalley, Claude (1909-1984), 89
- Grothendieck, Alexander (1928-2014), 165, 172
- Hilbert, David (1862-1943), 39
- Serre, Jean-Pierre (1926- ), 172
- Artin, Emil (1898-1962), 171
- Bézout, Etienne (1730-1783), 32
- Betti, Enrico (1823-1892), 26
- Borel, Emile (1871-1956), 58
- Dürer, Albrecht (1471-1528), 8
- Desargues, Girard (1591-1661), 8
- Grassmann, Hermann (1809-1877), 73
- Hausdorff, Felix (1868-1942), 58
- Heine, Eduard (1821-1881), 58
- Hensel, Kurt 1861-1941, 29
- Hesse, Otto (1811-1877), 15
- Hilbert, David (1862-1943), 44
- Jacobi, Carl Gustav Jacob 1804-1851, 29
- Möbius, August Ferdinand (1790-1868), 8
- Nakayama, Tadashi (1912-1964), 79
- Noether, Emmy (1882-1935), 38, 41, 81
- Plücker, Julius (1801-1868), 35
- Schelter, William (1947-2001), 91
- Segre, Corrado (1863-1924), 59
- Veronese, Giuseppe (1854-1917), 59
- Zariski, Oscar (1899-1986), 39
- affine cone, 60
- affine double plane, 95
- affine hypersurface, 43
- affine morphism, 144
- affine open subset, 72
- affine plane, 5
- affine plane curve, 5
- affine space, 5
- affine variety, 42, 71
- algebraic dimension, 6
- algebra, 37
- algebra generators, 37
- algebraically dependent, independent, 17
- annihilator, 150
- arithmetic genus, 149
- ascending, descending chain conditions, 41
- base point, 176
- basis for a topology, 49
- bidual, 20
- bihomogeneous polynomial, 64
- bilinear relations, 102
- Birkhoff-Grothendieck Theorem, 165
- blowup, 69
- blowup map, 28
- branch locus, 95, 97
- branch point, 25, 158
- branch points, 26
- branched covering, 157
- branched covering, 25
- canonical divisor, 176
- canonical map, 178
- center of projection, 25, 69
- characteristic properties of cohomology, 140
- Chinese Remainder Theorem, 38, 151
- classical topology, 12
- closed set, open set, 41
- coboundary map, 138
- cohomological dimension, 151
- cohomological functor, 138, 139
- cohomology of a complex, 138
- cohomology of an  $\mathcal{O}$ -module, 137
- cohomology sequence, 137, 139
- cokernel, 121
- comaximal ideals, 38
- commutative diagram, 38
- commuting matrices, 45
- compact space, 57
- complex, 138
- conic, 6, 9
- connected space, 42
- constructible function, 117
- constructible set, 111
- contracted ideal, 49, 103
- contravariant functor, 120
- coordinate algebra, 46
- coordinate algebra, 43
- Correspondence Theorem, 38
- covering diagram, 123
- curve, 108
- cusp, 28
- decomposable element, 74
- degree of a morphism to  $\mathbb{P}^n$ , 176
- degree of a morphism to projective space, 176

degree of an affine plane curve, 5  
 degree of an invertible module, 163  
 derivation, 166  
 diagonal, 68  
 differential, 167  
 direct image, 128  
 direct limit, 128  
 directed set, 128  
 discriminant, 23  
 divisor, 153, 161  
 divisor of a function, 162  
 divisor of a polynomial, 11  
 domain, 17, 37  
 double point, 28  
 dual  $\mathcal{O}$ -module, 161  
 dual curve, 18  
 dual plane, 18  
  
 effective divisor, 162  
 eigenvector, 79  
 elliptic curve, 174  
 Euler characteristic, 26  
 Euler characteristic, 164  
 Euler characteristic of a finite  $\mathcal{O}$ -module, 153  
 exact sequence, 101, 121  
 extended ideal, 49  
 extension by zero, 129  
 extension of domains, 80  
 extension of scalars, 103  
 exterior algebra, 73  
  
 Fermat curve, 14  
 fibre dimension, 117  
 fibre of a map, 6  
 fibred product, 115  
 finite  $\mathcal{O}$ -module, 121  
 finite module, 38  
 finite morphism, 80, 89  
 finite-type algebra, 37  
 finitely generated ideal, 38  
 flex point, 14  
 function field, 65, 119  
 function field of an affine variety, 50  
  
 general position, generic, 30  
 generators of an  $\mathcal{O}$ -module, 133  
 genus, 26, 164, 173  
 geometric genus, 149  
 good point, 66  
 graded algebra, 73  
 graph of a morphism, 68, 115  
 Grassmanian, 73  
  
 Hausdorff space, 58  
 Heine-Borel Theorem, 58  
 Hessian determinant, 16  
 Hessian divisor, 33  
  
 Hessian matrix, 15  
 Hilbert Nullstellensatz, 44  
 Hilbert Basis Theorem, 39  
 Hilbert Nullstellensatz, 44  
 homogeneous fraction, 70  
 homogeneous ideal, 60  
 homogeneous parts, 10  
 homogeneous polynomial, 9, 91  
 homogenize, dehomogenize, 14  
 homomorphism of  $\mathcal{O}$ -modules, 121  
 hyperelliptic curve, 177  
 hypersurface, 59  
  
 ideal generators, 40  
 ideal in the structure sheaf, 125  
 image, 121  
 increasing, strictly increasing chain, 39  
 induced topology, 41  
 integral closure, 83  
 integral extension, 80  
 integral morphism, 80, 89  
 intersection multiplicity, 11, 154  
 invariant element, 53  
 invertible  $\mathcal{O}$ -module, 161  
 irreducible space, 42  
 irreducible polynomial, 6  
 irregularity, 149  
 irrelevant ideal, 60  
 isolated point, 13  
 isomorphism, 67  
 isomorphism of affine varieties, 50  
  
 kernel, 121  
  
 line, 6, 7, 41  
 line at infinity, 7  
 linear subspace, 58  
 linearly equivalent divisors, 162  
 local domain of dimension one, 105  
 local property, 108  
 local ring, 105  
 local ring at a point, 107  
 localization of an algebra, 48  
 locally closed set, 111  
 locally free module, 108  
 locally principal ideal, 108  
 locus of zeros, 5  
  
 maximal annihilator, 150  
 maximal ideal, 37, 125  
 maximal member, 39  
 module homomorphisms, 104  
 module of differentials, 167  
 morphism, 67  
 morphism of affine varieties, 50  
 morphism of families, 123  
 multiplicative system, 103

multiplicity, 11  
multiplicity of zero, 27  
  
Nagata, Masayoshi (1927-2008), 44  
nilpotent ideal, 48  
nilradical, 48  
node, 28  
noetherian ring, 38  
noetherian space, 41  
normal domain, normal variety, 82  
normalization, 82  
  
open covering, 58  
order of zero, pole, 106  
ordinary bitangent, 34  
ordinary curve, 35  
orientability, 25  
  
Pierre de Fermat (1607-1665), 14  
Plücker Formulas, 35  
plane projective curve, 11  
point at infinity, 7  
point with values in a field, 65  
presentation of a module, 160  
prime ideal, 37  
principal divisor, 162  
product equations, 29  
product ideal, 37  
product topology, 62  
projection, 25, 69  
projective double plane, 97  
projective line, 7  
projective plane, 7  
projective space, 7  
projective variety, 57  
proper variety, 116  
  
quadric, 58  
quasicompact, 58  
quasiprojective variety, 62  
  
radical of an ideal, 40  
Rainich, George Yuri (1886-1968), 44  
ramification index, 157  
rank, 108  
rational curve, 96, 174  
rational function on a variety, 65  
rational function on an affine variety, 50  
real projective plane, 8  
reducible curve, 11  
regular differential, 170  
regular function, 50, 65, 120  
regular function on an affine variety, 46  
regular homogeneous fraction, 131  
residue field, 43  
residue field module, 125  
residue field of a local ring, 105  
  
resolution, acyclic resolution, 143  
restriction of a divisor to an open set, 161  
restriction of a morphism, 89  
restriction of an  $\mathcal{O}$ -module, 130  
restriction of scalars, 104  
resultant, 22  
resultant matrix, 22  
  
scaling, 9  
section, 120, 121  
sections equal on an open set, 122  
Segre embedding, 59  
semicontinuous function, 117  
Serre dual, 172  
short exact sequence, 101  
simple localization, 103  
smooth curve, singular curve, 14  
smooth point, singular point, 14, 108  
special line, 27  
special linear group, 43  
spectrum of an algebra, 46  
square-free polynomial, 84  
standard affine open set, 8  
standard cusp, 28  
Strong Nullstellensatz, 44  
structure sheaf, 119  
support of a divisor, 161  
support of a module, 150  
  
tacnode, 28  
tangent line, 14  
tensor algebra, 76  
tensor product module, 102  
torsion, 158  
torsion-free, 158  
trace of a differential, 170  
trace of a field extension, 84  
transcendence basis, 17  
transcendental element, 17  
transversal intersection, 32  
trefoil knot, 28  
triangulation, 26  
twist of an  $\mathcal{O}$ -module, 132  
twisted cubic, 60  
twisting module, 132  
  
unit ideal, 12  
  
valuation, 105  
valuation ring, 106  
value of a function, 46  
  
weighted degree, 23  
weighted projective space, 97  
  
Zariski closed set, 57  
Zariski closed, Zariski open, 40

Zariski topology, 40, 57  
zero of a polynomial, 11, 40