

2 Elliptic curves as abelian groups

In Lecture 1 we defined an elliptic curve as a smooth projective curve of genus 1 with a distinguished rational point. An equivalent definition is that an elliptic curve is an abelian variety of dimension one. An *abelian variety* is a smooth projective variety that is also a group, where the group operation is defined by rational functions (ratios of polynomials). Remarkably, these constraints force the group to be commutative, which is why they are called abelian varieties.

A variety is (roughly speaking) the zero locus of a set of polynomials, subject to an irreducibility condition. The precise definition won't concern us here, it is enough for us to know that a variety of dimension one is a curve, so an abelian variety of dimension one is a smooth projective curve with a group structure specified by rational functions. We will prove in this lecture that elliptic curves are abelian varieties. In fact the converse holds, every abelian variety of dimension one is an elliptic curve, but we won't prove this.

As mentioned in the first lecture, it is possible to associate an abelian variety to any smooth projective curve; this abelian variety is called the *Jacobian* of the curve. The dimension of the Jacobian is equal to the genus g of the curve, which means that in general the Jacobian is a much more complicated object than the curve itself (which always has dimension one). Writing explicit equations for the Jacobian as a projective variety is quite complicated, in general, but for elliptic curves, the curve and its Jacobian both have dimension one, and in fact the Jacobian is isomorphic to the curve itself.

2.1 The group law for Weierstrass curves

Recall from Lecture 1 that the group law for an elliptic curve defined by a Weierstrass equation $y^2 = x^3 + Ax + B$ is determined by the following rule:

Three points on a line sum to zero, which is the point at infinity.

It is then easy to determine the inverse of a point: negating the y -coordinate of a projective point $P = (x : y : z)$ yields the point $Q = (x : -y : z)$. The line between P and Q is vertical, and like all vertical lines it passes through the point O at infinity. We then have $P + Q + O = O$, which means that $P + Q = O$, so $Q = -P$ (this also works when $P = O$).

We can also check that O acts as the identity: the line between O and any point P intersects the curve at $-P$ (this is a double intersection at a tangent when $P = -P$). We then have $0 + P + (-P) = O$, and therefore $O + P = P$.


Associativity is not obvious, and while it can be rigorously proven algebraically, this is a tedious task that does not yield much insight. So we will give two proofs. The first will only apply to the generic case but it is short and provides some explanation as to *why* the group operation is associative. The second will be algebraic and fully rigorous, but we will let Sage do all the dirty work for us.

2.1.1 A geometric proof of associativity in the generic case

This is an adaptation of the proof in [2, p. 28]. Let P, Q, R be three points on an elliptic curve E over a field k that we may assume is algebraically closed. We shall also assume that

P, Q, R , and the zero point O are all in *general position* (this means that in the diagram below there are no relationships among the points other than those that necessarily exist by construction).

The line ℓ_0 through P and Q meets the curve E at a third point, $-(P + Q)$, and the line m_2 through O and $-(P + Q)$ meets E at $P + Q$. Similarly, the line m_0 through P and R meets E at $-(P + R)$, and the line ℓ_2 through O and $-(P + R)$ meets E at $P + R$. Let S be the third point where the line ℓ_1 through $Q + P$ and R meets E , and let T be the third point where the line m_1 through Q and $P + R$ meets E . See the diagram below.



We have $S = -((Q + P) + R)$ and $T = -(Q + (P + R))$. It suffices to show $S = T$. Suppose not. Let $g(x, y, z)$ be the cubic polynomial formed by the product of the lines ℓ_0, ℓ_1, ℓ_2 in homogeneous coordinates, and similarly let $h(x, y, z) = m_0 m_1 m_2$. We may assume $g(T) \neq 0$ and $h(S) \neq 0$, since the points are in general position and $S \neq T$. Thus g and h are linearly independent elements of the k -vector space V of homogeneous cubic polynomials in $k[x, y, z]$. The space V has dimension 10, thus the subspace of homogeneous cubic polynomials that vanish at the eight points $O, P, Q, R, \pm(Q + P)$, and $\pm(P + R)$ has dimension 2 and is spanned by g and h . The polynomial $f(x, y, z) = x^3 + Axz^2 + Bz^3 - zy^2$ that defines E is a nonzero element of this subspace, so we may write $f = ag + bh$ as a linear combination of g and h . Now $f(S) = f(T) = 0$, since S and T are both points on E , but $g(S) = h(T) = 0$ and $g(T), h(S) \neq 0$, which implies that both a and b are zero. This is a contradiction because f is not the zero polynomial.

2.1.2 The group law in algebraic terms

Let $P = (x_1, y_1, z_1)$ and $Q = (x_2, y_2, z_2)$ be two points on E . We will compute the sum $P + Q = R = (x_3, y_3, z_3)$ by expressing the coordinates of R as rational functions of the coordinates of P and Q . If either P or Q is the point at infinity, then R is simply the other point, so we assume that P and Q are affine points with $z_1 = z_2 = 1$. There are two cases:

Case 1. $x_1 \neq x_2$. The line \overline{PQ} has slope $m = (y_2 - y_1)/(x_2 - x_1)$, which yields the equation $y - y_1 = m(x - x_1)$. The point $-R = (x_3, -y_3, 1)$ is on this line, thus $-y_3 = m(x_3 - x_1) + y_1$. Substituting for y_3 in the Weierstrass equation for E yields

$$(m(x_3 - x_1) + y_1)^2 = x_3^3 + Ax_3 + B.$$

Simplifying, we obtain $0 = x_3^3 - m^2 x_3^2 + \dots$, where the ellipsis hides lower order terms. The values x_1 and x_2 satisfy the same cubic equation, thus its roots are x_1, x_2 , and x_3 ,

and the sum of these roots is equal to the negation of the quadratic coefficient $-m^2$. Thus $x_3 = m^2 - x_1 - x_2$. To sum up, we have

$$\begin{aligned} m &= (y_2 - y_1)/(x_2 - x_1), \\ x_3 &= m^2 - x_1 - x_2, \\ y_3 &= m(x_1 - x_3) - y_1. \end{aligned}$$

Thus to compute $P + Q = R$, we need to perform three multiplications (one of which is a squaring) and one inversion in the field k . We'll denote this cost $3\mathbf{M}+\mathbf{I}$.

Case 2. $x_1 = x_2$. If $y_1 \neq y_2$, then they must be opposite points and $R = 0$. Otherwise $P = Q$, and we compute the slope of the tangent line by implicitly differentiating the Weierstrass equation for E . This yields

$$2y dy = 3x^2 dx + A dx,$$

so

$$m = \frac{dy}{dx} = \frac{3x_1^2 + A}{2y_1}.$$

The formulas for x_3 and y_3 are then the same as before. Note that we require an extra multiplication (a squaring) here, so computing $R = 2P$ has a cost of $4\mathbf{M}+\mathbf{I}$.

With these equations in hand, we can now prove associativity as a formal identity, treating $x_1, y_1, z_1, x_2, y_2, z_2, x_3, y_3, z_3, A, B$ as indeterminants subject to the three relations implied by the fact that P, Q , and R all lie on the curve E . See the Sage worksheet

[Lecture 2 Proof of associativity](#)

for details, which includes checking all the special cases.

The equations above can be converted to projective coordinates by replacing x_1, y_1, x_2 , and y_2 with $x_1/z_1, y_1/z_1, x_2/z_2$, and y_2/z_2 respectively, and then writing the resulting expressions for x_3/z_3 and y_3/z_3 with a common denominator. This has the advantage of avoiding inversions, which are more costly than multiplications (in a finite field of cryptographic size inversions may be 50 or even 100 times more expensive than multiplications). This increases the number of multiplications to $12\mathbf{M}$ in case 1 (general addition) and $14\mathbf{M}$ in case 2 (doubling).

2.2 Edwards curves

There are many alternative representations of elliptic curves that have been proposed. We give just one example here, Edwards curves [1, 3], which have two significant advantages over Weierstrass equations. Let d be a non-square element of a field k whose characteristic is not 2. Then the equation

$$x^2 + y^2 = 1 + dx^2y^2 \tag{1}$$

defines an elliptic curve with distinguished point $(0, 1)$.

Remark 2.1. The plane projective curve defined by equation (1) has two singular points at infinity, violating our requirement that an elliptic curve be smooth. However, this plane curve can be *desingularized* by embedding it in $\mathbb{P}^3(k)$. The points at infinity are then no longer rational, and do not play a role in the group operation on $E(k)$, whose elements can all be uniquely represented as solutions (x, y) to equation (1) above.

The group operation is given by

$$(x_3, y_3) = \left(\frac{x_1 y_2 + x_2 y_1}{1 + dx_1 x_2 y_1 y_2}, \frac{y_1 y_2 - x_1 x_2}{1 - dx_1 x_2 y_1 y_2} \right), \quad (2)$$

which implies that the inverse of (x_1, y_1) is $(-x_1, y_1)$. In contrast to the formulas for curves in Weierstrass form, the formula in (2) is well defined for every pair of points (x_1, y_1) and (x_2, y_2) in $E(k)$. To see this, note that if either of the denominators in (2) is zero then we must have

$$(1 + dx_1 x_2 y_1 y_2)(1 - dx_1 x_2 y_1 y_2) = 1 - d^2 x_1^2 x_2^2 y_1^2 y_2^2 = 0,$$

so $d^2 x_1^2 x_2^2 y_1^2 y_2^2 = 1$ and none of x_1, x_2, y_1, y_2 is zero. Applying this and the curve equation (twice) yields

$$x_1^2 + y_1^2 = 1 + dx_1^2 y_1^2 = 1 + \frac{1}{dx_2^2 y_2^2} = \frac{x_2^2 + y_2^2}{dx_2^2 y_2^2}.$$

By adding or subtracting $2x_1 y_1 = \pm 2/(dx_2 y_2)$ to both sides we can obtain

$$(x_1 \pm y_1)^2 = \frac{(x_2 \pm y_2)^2}{dx_2^2 y_2^2},$$

with either choice of sign on the LHS (the sign on the RHS may vary, but in any case the numerator of the RHS is a square). Since x_1 and y_1 are nonzero, one of $x_1 + y_1$ and $x_1 - y_1$ is nonzero, and this implies that d is a square in k , a contradiction.

As written, the group law involves five multiplications and two inversions (ignoring the multiplication by d , which we can choose to be small), which is greater than the cost of the group operation in Weierstrass form. However, in projective coordinates we have

$$\frac{x_3}{z_3} = \frac{z_1 z_2 (x_1 y_2 + x_2 y_1)}{z_1^2 z_2^2 + dx_1 x_2 y_1 y_2}, \quad \frac{y_3}{z_3} = \frac{z_1 z_2 (y_1 y_2 - x_1 x_2)}{z_1^2 z_2^2 - dx_1 x_2 y_1 y_2}.$$

There are a bunch of common subexpressions here, and in order to compute z_3 , we need a common denominator. Let $r = z_1 z_2$, let $s = x_1 y_2 + x_2 y_1$, let $t = dx_1 y_2 x_2 y_1$, and let $u = y_1 y_2 - x_1 x_2$. We then have

$$x_3 = rs(r^2 - t), \quad y_3 = ru(r^2 + t), \quad z_3 = (r^2 + t)(r^2 - t).$$

This yields a cost of 12M. If we compute s as $s = (x_1 + y_1)(x_2 + y_2) - x_1 x_2 - y_1 y_2$, the cost is reduced to 11M.

A simple Sage implementation of these formulas can be found here:

[Lecture 2 Group law on Edwards curves](#)

Because the formula in (2) is well-defined at every point in $E(k)$, we do not need separate formulas for addition and doubling. Moreover, additions involving a point and its negation, or the identity element, all work exactly the same as in the general case. Such formulas are called *complete*, and they have two distinct advantages. First, they can be implemented very efficiently because there is no branching. Second, they protect against what is known as a *side-channel* attack. If an adversary can distinguish whether you are doubling or adding points, e.g. by monitoring the CPU and noticing the difference in the time required by each operation, they can break a cryptosystem that performs scalar multiplication by an integer that is meant to be secret.

Having said that, if you know you are going to be doubling and are not concerned about a side-channel attack, there are several optimizations that can be made (these include replacing $1 + dx^2y^2$ with $x^2 + y^2$). This reduces the cost of doubling a point on an Edwards curves to **7M**, which is a huge improvement over the **14M** cost of doubling a point in Weierstrass coordinates.

The [explicit formulas database](#) contains optimized formulas for Edwards curves and various generalizations, as well as many other forms of elliptic curves. Operation counts and verification scripts are provided with each set of formulas.

We should note that, unlike Weierstrass equations, not every elliptic curve can be put into Edwards form. In particular, an Edwards curve always has a rational point of order 4, the point $(1, 0)$, but this is not true of many elliptic curves.

References

- [1] Daniel J. Bernstein and Tanja Lange, *Faster addition and doubling on elliptic curves*, Advances in Cryptology - ASIACRYPT 2007, Lecture Notes in Computer Science **4833**, Springer-Verlag, New York (2007), 29–50.
- [2] J. W. S. Cassels, *Lectures on elliptic curves*, London Mathematical Society Student Texts **24**, Cambridge University Press, 1991.
- [3] Harold M. Edwards, *A normal form for elliptic curves*, Bulletin of the American Mathematical Society **44** (2007), 393–422.