

18.783 Elliptic Curves

Lecture 3

Andrew Sutherland

February 24, 2021

Representing finite fields

For $\mathbb{F}_p \simeq \mathbb{Z}/p\mathbb{Z}$ we use integers in $[0, p - 1]$ denoting elements of $\mathbb{Z}/p\mathbb{Z}$.

For $\mathbb{F}_q \simeq \mathbb{F}_p^d \simeq \mathbb{F}_p[x]/(x^d)$ we use vectors in \mathbb{F}_p^d denoting elements of $\mathbb{F}_p[x]/(x^d)$, which can view as elements of $\mathbb{F}_p[x]/(f)$ for some irreducible $f \in \mathbb{F}_p[x]$ of degree d . It does not matter which f we pick, but some choices are better than others.

This reduces all computation in finite fields to integer and polynomial arithmetic.

We should note that there are other choices. If $\mathbb{F}_q^\times = \langle r \rangle$ (so r is a **primitive root**), we could use 0 to denote 0 and $e \in [1, q - 1]$ to denote r^e .

Integer arithmetic

Complexity of ring operations on n -bit integers:

addition/subtraction	$O(n)$
multiplication (FFT)	$O(n \log n)$ 🙌🤖

To multiply polynomials in $\mathbb{F}_p[x]$ we use **Kronecker substitution**.

Let $\hat{f} \in \mathbb{Z}[x]$ denote the **lift** of $f \in \mathbb{F}_p[x]$ to $\mathbb{Z}[x]$. We compute $h = fg \in \mathbb{F}_p[x]$ via

$$\hat{h}(2^m) = \hat{f}(2^m)\hat{g}(2^m)$$

with $m \geq 2 \lg p + \lg(d + 1)$, where $d := \deg f$. The k th coefficient of h can be obtained by extracting the k th block of m bits from $\hat{h}(2^m)$ and reducing it modulo p .

All ring operations in $\mathbb{F}_p[x]$ can thus be reduced to ring operations in \mathbb{Z} , provided we know how to reduce integers modulo p .

Euclidean division

For positive integers a, b we want to compute the unique $q, r \geq 0$ for which

$$a = bq + r \quad (0 \leq r < b),$$

that is, $q = \lfloor a/b \rfloor$ and $r = a \bmod b$. Recall Newton's method to find a root of $f(x)$:

$$x_{i+1} := x_i - \frac{f(x_i)}{f'(x_i)}.$$

To compute $c \approx 1/b$, we apply this to $f(x) = 1/x - b$, using the Newton iteration

$$x_{i+1} = x_i - \frac{f(x_i)}{f'(x_i)} = x_i - \frac{\frac{1}{x_i} - b}{-\frac{1}{x_i^2}} = 2x_i - bx_i^2.$$

We can then compute $q = \lfloor ca \rfloor$ and $r = a - bq$.

Euclidean division

As an example, let us approximate $1/b = 1/123456789$ working in base 10 (in an actual implementation would use base 2, or base 2^w , where w is the word size).

$$x_0 = 1 \times 10^{-8}$$

$$\begin{aligned}x_1 &= 2(1 \times 10^{-8}) - (1.2 \times 10^8)(1 \times 10^{-8})^2 \\ &= 0.80 \times 10^{-8}\end{aligned}$$

$$\begin{aligned}x_2 &= 2(0.80 \times 10^{-8}) - (1.234 \times 10^8)(0.80 \times 10^{-8})^2 \\ &= 0.8102 \times 10^{-8}\end{aligned}$$

$$\begin{aligned}x_3 &= 2(0.8102 \times 10^{-8}) - (1.2345678 \times 10^8)(0.8102 \times 10^{-8})^2 \\ &= 0.81000002 \times 10^{-8}.\end{aligned}$$

We double the precision we are using at each step, and each x_i is correct up to an error in its last decimal place. The value x_3 suffices to correctly compute $\lfloor a/b \rfloor$ for $a \leq 10^{15}$.

Euclidean division

There is an analogous algorithm for Euclidean division in $\mathbb{F}_p[x]$.

Given $a, b \in \mathbb{F}_p[x]$ with b monic we can compute the unique $q, r \in \mathbb{F}_p[x]$ for which

$$a = bq + r \quad (\deg r < \deg b).$$

See the lecture notes for details. In both cases if the divisor b is fixed we can save time by precomputing $c \approx 1/b$ (as on Problem Set 1).

Theorem

Let $q = p^d$ be a prime power and assume $\log d = O(\log p)$ or $p = O(1)$.

The time to multiply two elements in \mathbb{F}_q is $O(M(n)) = O(n \log n)$, where $n = \log q$.

Under a widely believed conjecture we know that multiplication in \mathbb{F}_q takes time $O(n \log n)$ (but not necessarily $O(M(n))$), without any assumptions about p and d .

Inverting elements of a finite field

Given integers $a > b > 0$ the (extended) Euclidean algorithm computes $s, t \in \mathbb{Z}$ with

$$\gcd(a, b) = as + bt \quad (|s| \leq b/\gcd(a, b), |t| \leq a/\gcd(a, b))$$

If $a = p$ is prime, then $ps + bt = 1$ and $t \equiv b^{-1} \pmod{p}$ with $t \in [0, p - 1]$.

The Euclidean algorithm works in any Euclidean ring, including $\mathbb{F}_p[x]$.

But note that $\mathbb{F}_p[x]$ has a larger unit group than \mathbb{Z} and $\gcd(a, b)$ is defined only up to units. More formally, $\gcd(a, b) = (a, b) = (c)$ is a principal ideal. In \mathbb{Z} there is a unique positive choice of c , while in $\mathbb{F}_p[x]$ there is a unique monic choice of c .

The [fast Euclidean algorithm](#) (see lecture notes) yields the following theorem.

Theorem

Let $q = p^d$ be a prime power and assume $\log d = O(\log p)$ or $p = O(1)$.

The time to invert an element of \mathbb{F}_q^\times is $O(M(n) \log n) = O(n \log^2 n)$, where $n = \log q$.

Exponentiation (also known as scalar multiplication)

Given a group element g and a positive integer a we want to compute $g^a = gg \cdots g$ (or if we write the group operation additively, $ag = g + g + \cdots + g$).

We can achieve this using a “square-and-multiply” (or “double-and-add”) algorithm:

1. Let $a = \sum_{i=0}^n 2^i a_i$ and initialize h to g .
2. For i from $n - 1$ down to 0:
 - a. Replace h with h^2
 - b. If $a_i = 1$ then replace h with hg .

At the end of the i th loop we have $h = g^b$ with $b = \sum_{j=0}^{n-i} 2^j a_{i+j}$.

This allows us to compute g^a using at most $2n = O(n)$ group operations. The leading constant 2 can be improved; you will have a chance to explore this on Problem Set 2.

For \mathbb{F}_q^\times each group operation takes time $O(M(n))$, and for $a \leq q - 1$ the time to compute g^a is $O(nM(n)) = O(n^2 \log n)$. Note: we can always reduce a modulo $q - 1$.