

# Explicit Irreducible Decomposition of the Spin Representation of the Temperley-Lieb Algebra

Rachel Chen

Under the direction of

Kenta Suzuki  
Department of Mathematics  
Massachusetts Institute of Technology

Research Science Institute  
July 29th, 2025

## Abstract

The spin representation  $(\mathbb{C}^2)^{\otimes n}$  can be decomposed uniquely into irreducible modules called the Specht Modules, denoted  $W_k^n$ . Lusztig's dual canonical basis of the spin representation can be viewed as diagrams of the Temperley-Lieb algebra, and the Specht Modules also have a diagrammatic basis. We explicitly describe the decomposition of the spin representation into Specht Modules,  $(\mathbb{C}^2)_{k-1}^{\otimes n-2} \cong W_{n-2}^{n-2} \oplus \cdots \oplus W_{n-2k}^{n-2}$ , by computing the images of the diagrammatic basis elements of the Specht Modules. We do this by using induction to compute the image of  $W_n^n$  and reducing the problem of computing the images of other Specht modules to this case. Our results may lead to the notion of a canonical basis for Specht Modules in the future.

## Summary

In both mathematics and physics, a common method of understanding complicated systems is to break them down into smaller pieces. Accordingly, it is known that a multiple-electron system can be viewed as multiple single-particle systems, but mathematicians and physicists do not yet know which single-particle systems a multiple-electron system explicitly breaks into. This paper investigates the question of computing such a decomposition explicitly. To do so, we investigate both the multiple-electron and single-particle systems mathematically and visually. We view multiple-electron systems using a mathematical object called the Temperley-Lieb algebra, an object defined in terms of diagrams with applications in statistical mechanics, representation theory, and topology. Similarly, we view the single-particle systems through certain diagrams of mathematical objects called Specht Modules. We compute how the Temperley-Lieb algebra explicitly relates to the Specht Modules through how the diagrams of both objects explicitly relate to each other. This, in turn, allows us to compute how a certain system with multiple electrons decomposes into systems with only one particle, simplifying computations in quantum mechanics.

# 1 Introduction

The spin representation  $(\mathbb{C}^2)^{\otimes n}$  is an important object in various areas of mathematics and physics. Since it was first introduced by Cartan in 1913 [1] and applied to physics by Dirac in 1928 [2], mathematicians and physicists have studied it extensively. The spin representation parametrizes configurations of  $n$  electrons, each of which has spin-up or spin-down. We focus on  $(\mathbb{C}^2)_k^{\otimes n}$ , which parametrizes configurations with  $k$  spin-down electrons and  $n - k$  spin-up electrons.

If one considers the spin representation  $(\mathbb{C}^2)^{\otimes n}$  as a representation of a quantum group, the algebra of endomorphisms that commute with the action of a quantum group is precisely  $TL_n$ , the *Temperley-Lieb algebra* on  $n$  vertices. First introduced by Temperley and Lieb in 1971 [3], the Temperley-Lieb algebra has applications in statistical mechanics, representation theory, and topology. As a vector space, the Temperley-Lieb algebra is generated by diagrams consisting of two parallel lines with  $n$  vertices on each line, where each vertex is connected to exactly one other vertex and no lines intersect (see Figure 1).




Figure 1: Examples of diagrams in the Temperley-Lieb algebra.

The spin representation with  $k$  spin-down electrons,  $(\mathbb{C}^2)_k^{\otimes n}$ , decomposes uniquely into irreducible modules named Specht modules, which are a generalization of the irreducible representations of  $S_n$  introduced by Specht in 1935 [4]. One method often used to understand algebraic structures, such as modules, is to find a diagrammatic interpretation that allows mathematicians to visualize the object combinatorially. Martin [5] discovered a diagrammatic basis for the Specht modules: the Specht module  $W_k^n$  for  $k \equiv n \pmod{2}$  can be expressed as a vector space spanned by certain diagrams from  $n$  vertices to  $k$  vertices.

Similarly, Khovanov [6] first introduced the possibility that the spin representation may have a diagrammatic interpretation. The author [7] generalized his work, allowing one to view the spin representation, specifically Lusztig's canonical basis [8], in terms of the diagrams of the Temperley-Lieb algebra.

This paper investigates the natural question of how to explicitly compute the decomposition of the spin representation into Specht modules in terms of the diagrams. Specifically,

there exists a known isomorphism  $\varphi_k^n: W_n^n \oplus W_{n-2}^n \oplus \cdots \oplus W_{n-2k}^n \rightarrow (\mathbb{C}^2)_k^{\otimes n}$ , which we describe explicitly. We compute the images of each of the basis elements under  $\varphi_k^n$  in  $W_{n-2i}^n$  for  $0 \leq i \leq k$ , where  $k = 0$  or  $k = 1$ , in Section 3, laying the groundwork for future induction. For  $k = 1$ , the solution to the more complicated case is that the image of the diagram  $\text{id}_n \in W_n^n$  is equal to  $\sum_{i=0}^{n-1} [n-i]_q D_i$ , where the diagrams  $D_i$  are shown in Figure 2.




Figure 2: The diagrams  $D_i$  for  $n = 4$ , the diagrams in the image of  $\text{id}_n$ .

In Section 4, we describe a method to reduce the computation of images of each of the basis elements of  $W_{n-2i}^n$  for  $1 \leq i \leq k$  to the computation of the image of a basis element in  $W_{n-2i}^{n-2i}$ . In Section 5, we explicitly describe the decomposition  $(\mathbb{C}^2)_k^{\otimes n-1} \oplus (\mathbb{C}^2)_{k-1}^{\otimes n-1} \cong (\mathbb{C}^2)_k^{\otimes n} |_{\text{TL}_{n-1}}$  and lay the groundwork for Section 6, in which we describe inductive formulas to explicitly compute the decomposition for general  $k$ . Specifically, the combination of Theorem 4.9 and Theorem 6.9 allows us to explicitly compute  $\varphi_k^n: W_n^n \oplus W_{n-2}^n \oplus \cdots \oplus W_{n-2k}^n \rightarrow (\mathbb{C}^2)_k^{\otimes n}$  for all  $k$  and  $n$ .

As Chen [7] gives a diagrammatic description of Lusztig’s dual canonical basis, a possible application of this research is to use this explicit decomposition to define the notion of a canonical basis for the Specht modules by having the Specht module inherit the canonical basis. Furthermore, the explicit decomposition of the spin representation allows us to view systems with many electrons as multiple one-particle systems, simplifying computations in quantum mechanics.

## 2 Background

### 2.1 The Temperley-Lieb Category

In this section, we introduce the Temperley-Lieb category. We also introduce notation that will be used throughout our paper.

**Definition 2.1.** A  $(m, n)$ -*diagram* comprises two parallel lines with  $m$  vertices on the bottom

line and  $n$  vertices on the top line such that the vertices are connected by edges satisfying the following properties:

- the edges are between the parallel lines,
- the edges do not intersect each other, and
- each vertex is the endpoint of exactly one edge.

Given a  $(\ell, m)$ -diagram and a  $(m, n)$ -diagram, we can obtain a  $(\ell, n)$ -diagram by concatenation. In concatenation, a contractible loop may appear; we remove it and multiply the resulting diagram by formal variable  $\beta = -q - q^{-1}$ , where  $q \in \mathbb{C}$  is not a root of unity. An example of concatenation is shown in Figure 3.




Figure 3: The composition of a  $(3, 5)$ -diagram and a  $(5, 5)$ -diagram.

**Definition 2.2** (Temperley-Lieb Category). The *Temperley-Lieb category*, denoted TL, has objects  $\{[n], n \in \mathbb{Z}_{\geq 0}\}$ , where  $[n]$  is a set of  $n$  vertices. The morphisms  $\text{Hom}([m], [n])$  is the vector space with a basis of diagrams consisting of  $(m, n)$ -diagrams.

We call the edges of a diagram *links*. We say a link is

- *straight* if it connects the  $i$ th vertex on the bottom line to the  $i$ th vertex on the top line,
- *quasi-simple* if it connects two vertices on the same line, and
- *simple* if it connects two adjacent vertices on the same line.

Define  $\epsilon_i^n \in \text{Hom}([n], [n-2])$  to be the diagram connecting the  $i$ th and  $(i+1)$ th vertices on the bottom line and containing no other quasi-simple links. Similarly, define  $\delta_i^n \in \text{Hom}([n-2], [n])$  to be the diagram connecting the  $i$ th and  $(i+1)$ th vertices on the top line with no other quasi-simple links. An example of these diagrams are shown in Figure 4.



Figure 4: Diagrams for  $\epsilon_2^4$  and  $\delta_2^4$ , respectively, in  $\text{Hom}([4], [2])$ .

Every diagram in the Temperley-Lieb category can be decomposed into certain generators, as described by the following theorem.

**Theorem 2.3** ([9]). *The algebra  $\bigoplus_{m,n \in \mathbb{Z}_{\geq 0}} \text{Hom}([m], [n])$  is generated as an algebra by all  $\epsilon_i^n$  and  $\delta_i^n$ , where  $1 \leq i \leq n-1$ , with the following relations:*

- (i)  $\epsilon_i^n \cdot \delta_i^n = -q - q^{-1}$  where  $q \in \mathbb{C}$  is not a root of unity, and
- (ii)  $(\text{id} \otimes \epsilon_i^n) \cdot (\delta_i^n \otimes \text{id}) = \text{id} = (\epsilon_i^n \otimes \text{id}) \cdot (\text{id} \otimes \delta_i^n)$ .

## 2.2 The Temperley-Lieb Algebra

In this section, we introduce the Temperley-Lieb Algebra, the object that is central to our paper. Refer to an  $(n, n)$ -diagram as an  $n$ -diagram.

**Definition 2.4** (Temperley-Lieb Algebra). Let the *Temperley-Lieb algebra*, denoted  $\text{TL}_n$ , be  $\text{Hom}([n], [n])$ , the vector space spanned by diagrams with  $n$  vertices on both the top and bottom lines. Let  $\text{tl}_n$  denote the set of diagrams of  $\text{TL}_n$ .

Let  $e_i$  refer to the diagram with a simple link connecting the  $i$ th and  $(i+1)$ th vertices, counting from the left, on each line and with straight links everywhere else (for example, see Figure 5).

As a corollary of Theorem 2.3, we can state the following Proposition, describing the Temperley-Lieb algebra in terms of generators. Recall that  $\beta = -q - q^{-1}$ .




Figure 5: The diagram  $e_2$  in  $\text{TL}_4$ .

**Proposition 2.5** ([10]). *The algebra  $\text{TL}_n$  is generated by  $\{e_1, \dots, e_{n-1}\}$  defined by the following relations:*

$$e_i^2 = \beta \cdot e_i, \quad e_i e_{i\pm 1} e_i = e_i, \quad \text{and } e_i e_j = e_j e_i \text{ for } |i - j| \geq 2.$$

### 2.3 The Spin Representation of the Temperley-Lieb Algebra

In this section, we discuss the spin representation  $(\mathbb{C}^2)^{\otimes n}$ , an important object in many areas of mathematics and physics, and how the Temperley-Lieb algebra acts on it.

Let  $v_+$  denote  $(1, 0) \in \mathbb{C}^2$  and  $v_-$  denote  $(0, 1) \in \mathbb{C}^2$ . The 4-dimensional vector space  $\mathbb{C}^2 \otimes \mathbb{C}^2$  has basis  $\{v_+ \otimes v_+, v_+ \otimes v_-, v_- \otimes v_+, v_- \otimes v_-\}$ .

Let  $\epsilon: \mathbb{C}^2 \otimes \mathbb{C}^2 \rightarrow \mathbb{C}$  be

$$v_+ \otimes v_+ \mapsto 0, \quad v_+ \otimes v_- \mapsto -q, \quad v_- \otimes v_+ \mapsto 1, \quad v_- \otimes v_- \mapsto 0$$

and let  $\delta: \mathbb{C} \rightarrow \mathbb{C}^2 \otimes \mathbb{C}^2$  be

$$1 \mapsto v_+ \otimes v_- - q^{-1} v_- \otimes v_+.$$

Let  $\epsilon_i^n: (\mathbb{C}^2)^{\otimes n} \rightarrow (\mathbb{C}^2)^{\otimes n-2}$  be  $(\text{id})^{\otimes i-1} \otimes \epsilon \otimes (\text{id})^{\otimes n-i-1}$  and  $\delta_i^n: (\mathbb{C}^2)^{\otimes n-2} \rightarrow (\mathbb{C}^2)^{\otimes n}$  be  $\text{id}^{\otimes i-1} \otimes \delta \otimes \text{id}^{\otimes n-i-1}$ .

Since  $\epsilon_i^n$  and  $\delta_i^n$  generate the algebra  $\bigoplus_{m,n \geq 0} \text{Hom}([m], [n])$  and all the relations in Proposition 2.5 are satisfied, we have an action of  $\bigoplus_{m,n \geq 0} \text{Hom}([m], [n])$  on  $\bigoplus_{n \geq 0} (\mathbb{C}^2)^{\otimes n}$ .

We can consider the case of  $m = n$  to obtain a representation of  $\text{TL}_n$ .

**Definition 2.6.** *The spin representation is the action of  $\text{TL}_n = \text{Hom}([n], [n])$  on  $(\mathbb{C}^2)^{\otimes n}$ .*

The spin representation is the central object of our paper.

**Example 2.7.** We describe an example of how  $\text{TL}_4$  acts on  $(\mathbb{C}^2)^{\otimes 4}$ . Given the first diagram in Figure 6, we write it in terms of the generators of TL by decomposing it as shown in the second and third diagrams. Explicitly, the first diagram decomposes into  $\delta_1^4 \delta_3^2 \epsilon_1^2 \epsilon_2^4$ .




Figure 6: An example of how a  $\text{TL}_4$  diagram is sliced.

We label each tensor product with its original position for convenience purposes. The action of the following diagram (Figure 6) is as follows:

$$\mathbb{C}_1^2 \otimes \mathbb{C}_2^2 \otimes \mathbb{C}_3^2 \otimes \mathbb{C}_4^2 \xrightarrow{\text{id} \otimes \epsilon \otimes \text{id}} \mathbb{C}_1^2 \otimes \mathbb{C}_4^2 \xrightarrow{\text{id} \otimes \text{id}} \mathbb{C}_2^2 \otimes \mathbb{C}_3^2 \xrightarrow{\epsilon} \mathbb{C} \xrightarrow{\delta \otimes \delta} \mathbb{C}_1^2 \otimes \mathbb{C}_2^2 \otimes \mathbb{C}_3^2 \otimes \mathbb{C}_4^2 .$$

## 2.4 A Diagrammatic Interpretation of the Spin Representation

Khovanov [6] introduced the possibility of a diagrammatic interpretation of the spin representation. Chen [7] generalized his work, representing Lusztig's dual canonical basis [8] as diagrams in the Temperley-Lieb algebra.

**Definition 2.8.** Let  $(\mathbb{C}^2)_k^{\otimes n}$  denote the subspace of  $(\mathbb{C}^2)^{\otimes n}$  generated by vectors of the form  $v_{a_n} \otimes \cdots \otimes v_{a_1}$  such that  $k$  of the  $a_i$  are  $-$  and  $n - k$  of the  $a_i$  are  $+$ .

*Remark.* As the spin representation parametrizes the positions of  $n$  electrons, each of which has either spin-down or spin-up, the module  $(\mathbb{C}^2)_k^{\otimes n}$  parametrizes the positions of  $n$  electrons where  $k$  of them have spin-down and  $n - k$  have spin-up.

First, we define the notion of a diagrammatic basis for the induced representation that will be used to view the spin representation in terms of the diagrams of  $\text{tl}_n$ .

**Lemma 2.9** ([7]). *For all  $1 \leq k \leq n$ ,  $\text{Ind}_{\text{TL}_k \otimes \text{TL}_{n-k}}^{\text{TL}_n} \mathbb{C}_{\text{triv}}$  is precisely generated by the diagrams in which the only quasi-simple links on the bottom line connect  $k - i$  to  $k + 1 + i$  for some  $i$ , which is the diagrammatic basis of the left hand side.*

**Definition 2.10.** Let  $C_k^n$  denote the diagrammatic basis of  $\text{Ind}_{\text{TL}_k \otimes \text{TL}_{n-k}}^{\text{TL}_n} \mathbb{C}_{\text{triv}}$ .

The following proposition describes how the spin representation can be viewed through diagrams of  $\text{tl}_n$ .

**Proposition 2.11** ([7]). *There is an isomorphism of  $\text{TL}_n$ -modules:*

$$\text{Ind}_{\text{TL}_k \otimes \text{TL}_{n-k}}^{\text{TL}_n} \mathbb{C}_{\text{triv}} \cong (\mathbb{C}^2)_k^{\otimes n},$$

and the isomorphism identifies the diagrammatic basis of the left hand side to the dual canonical basis of the right hand side.

Therefore,  $(\mathbb{C}^2)_k^{\otimes n}$  is generated by the diagrams in  $C_k^n$ .

We can extend Proposition 2.11 to the entire spin representation to obtain the following, giving a diagrammatic basis of the entire Temperley-Lieb algebra.

**Corollary 2.12** ([7]). *There is an isomorphism of  $\text{TL}_n$ -modules*

$$\bigoplus_{0 \leq k \leq n} \text{Ind}_{\text{TL}_k \otimes \text{TL}_{n-k}}^{\text{TL}_n} \mathbb{C}_{\text{triv}} \cong (\mathbb{C}^2)^{\otimes n}$$

which identifies the diagrammatic basis of the left hand side to the dual canonical basis of the right hand side.

With these results, we can work with  $(\mathbb{C}^2)^{\otimes n}$  in terms of the diagrams in  $\text{tl}_n$ .

## 2.5 Specht Modules

We now study the irreducible representations of  $(\mathbb{C}^2)_k^{\otimes n}$ , called the *Specht modules*, which are another important object in our work.

**Definition 2.13.** A diagram is *monic* when it does not have any simple links on the bottom line. For  $0 \leq \ell \leq n$  such that  $\ell \equiv n \pmod{2}$  let  $w_\ell^n$  denote the set of monic diagrams with  $n$  vertices on the top line and  $\ell$  vertices on the bottom line.

**Definition 2.14** (Specht modules). The *Specht module*  $W_\ell^n$  is the  $\text{TL}_n$ -module which is the quotient of  $\text{Hom}([\ell], [n])$  by the image of left multiplication by  $\epsilon_j^{\ell-2}$ s.

The following lemma gives an explicit description of the action of  $\text{TL}_n$  on the Specht Module.

**Lemma 2.15.** *The Specht module  $W_\ell^n$  has a basis consisting of the diagrams in  $w_\ell^n$ . A diagram  $D \in \text{tl}_n$  sends  $D' \in w_\ell^n$  to the concatenation  $DD'$  if the  $(n, \ell)$ -diagram  $DD'$  remains monic, and to 0 otherwise.*

**Example 2.16.** The composition of the two diagrams in Figure 7 is 0, as the monicity of the diagram in  $w_2^6$  is not preserved.

The following Proposition can be obtained through results in [11].



Figure 7: A diagram in the Specht Module  $W_2^6$  (left) and a diagram in the Temperley-Lieb algebra  $TL_6$  (right).

**Proposition 2.17** ([11]). *The  $TL_n$ -representation  $(\mathbb{C}^2)_k^{\otimes n}$  for  $k \leq n/2$  can be uniquely (up to scalar multiplication) decomposed into a direct sum of Specht modules  $W_n^n \oplus W_{n-2}^n \oplus \cdots \oplus W_{n-2k}^n$ .*

*Proof.* There is a decomposition of  $(\mathbb{C}^2)^{\otimes n}$  as a  $TL_n \otimes \mathcal{U}_q(\mathfrak{sl}_2)$ -module as<sup>1</sup>

$$(\mathbb{C}^2)^{\otimes n} \cong \bigoplus_{0 \leq k \leq n/2} W_{n-2k}^n \boxtimes (\mathbb{C}^2)_{n/2-k}^{\otimes n}. \quad (1)$$

Since  $Kv_{\pm} = q^{\pm 1}v_{\pm}$ , the  $q^{\ell}$ -eigenspace of  $K$  in (1) gives that

$$(\mathbb{C}^2)_{n-2\ell}^{\otimes n} = \bigoplus_{0 \leq k \leq |\ell|} W_{n-2k}^n. \quad \square$$

As both  $(\mathbb{C}^2)_k^{\otimes n}$  and  $W_i^n$  for all  $0 \leq i \leq k$  where  $k \equiv n \pmod{2}$  have a diagrammatic interpretation, a natural question to ask is how the above isomorphism interacts with the diagrammatic perspective. We investigate the images of the diagrams in  $w_i^n$ , leading to an explicit decomposition of the spin representation.

### 3 Computation of $\varphi_k^n$ for $k = 0$ and $k = 1$

In the previous section, we have defined an isomorphism  $\varphi_k^n: W_n^n \oplus W_{n-2}^n \oplus \cdots \oplus W_{n-2k}^n \rightarrow (\mathbb{C}^2)_k^{\otimes n}$ . In this section, we describe  $\varphi_k^n$  explicitly in terms of diagrams for  $k = 0$  and  $k = 1$ , laying the groundwork for an inductive computation for general  $k$  in Section 6. Recall that  $w_k^n$  denotes the diagrammatic basis of  $W_k^n$ ,  $tl_n$  denotes the diagrammatic basis of  $TL_n$ , and  $C_n^k$  denotes the diagrammatic basis of  $(\mathbb{C}^2)_k^{\otimes n}$ . This notation will be used throughout the paper.

<sup>1</sup>where  $\boxtimes$  denotes the box tensor product.

**Lemma 3.1.** *The image of the unique diagram in  $w_n^n$  under  $\varphi_0^n$  is equal to the identity diagram in  $\text{tl}_n$ .*

*Proof.* In both  $W_n$  and  $\text{TL}_n$ , the action of the generators  $e_i$  (see Figure 5) for all  $i$  send the diagram described in the lemma to 0.  $\square$

Recall that  $\epsilon_i^n$  and  $\delta_i^n$  (see Figure 4) are the generators of the Temperley-Lieb category.

**Proposition 3.2.** *We can describe the image of  $\varphi_1^n$  as follows:*

(i) *the image of a diagram  $D \in w_{n-2}^n$  is equal to  $D\epsilon_1^{n-2}$ ; and*

(ii) *the image of the diagram  $\text{id}_n \in w_n^n$  is equal to  $\sum_{i=0}^{n-1} [n-i]_q D_i$ , where  $D_i = \delta_i^n \cdot \epsilon_1^n$  for  $1 \leq i \leq n-1$  and  $D_0$  is the identity diagram in  $\text{tl}_n$ .*

*Proof.* First, statement (i) is true as the actions of  $e_i$  for all  $i$  will be consistent across both  $W_{n-2}^n$  and  $(\mathbb{C}^2)_1^{\otimes n}$ .

Now, we prove statement (ii). The action of  $e_i$  on  $D_0$  will give  $D_1$  for  $i = 1$  and 0 for  $i \neq 1$ . For each  $D_i$  for  $1 \leq i \leq n-1$ , we have that

$$e_j D_i = \begin{cases} \beta D_i & \text{if } j = i \\ D_i & \text{if } j = i \pm 1 \\ 0 & \text{otherwise.} \end{cases}$$

Let  $c_i$  denote the coefficient of  $D_i$  in  $\varphi_1^n(\text{id}_n)$ . It is necessary that  $e_j \varphi_1^n(\text{id}_n) = e_j \sum_i c_i D_i = 0$  for all  $j$ . From the action of the  $e_i$  on the diagrams described above, we obtain that  $c_{n-1} = 1$ ,  $c_{n-2} = -\beta = q + q^{-1}$ , and  $c_{n-i} = (q + q^{-1})c_{n-i+1} - c_{n-i+2}$ . Induction allows us to conclude that that  $c_{n-i} = \frac{q^i - q^{-i}}{q - q^{-1}} = [i]_q$ . Therefore, we obtain that  $\varphi_1^n(\text{id}_n) = \sum_{i=0}^{n-1} [n-i]_q D_i$ .  $\square$

This proposition will be used in Section 6 as a base case for our induction procedure.

## 4 Reduction Procedure For General $k$

In this section, we describe how to calculate the images under  $\varphi_k^n$  of the diagrammatic basis of  $W_{n-2}^n, \dots, W_{n-2k}^n$  by reducing the problem to computing the images of the diagrammatic basis of  $W_{n-2}^{n-2}, \dots, W_{n-2k}^{n-2}$  under  $\varphi_{k-1}^{n-2}$ . This procedure can be applied repeatedly, allowing us to reduce the problem of computing  $\varphi_k^n(W_{n-2i}^n)$  to computing  $\varphi_{k-i}^{n-2i}(W_{n-2i}^{n-2i})$ . We

construct this reduction by proving this reduction for the specific case of diagrams with links in the top left corner in Proposition 4.7, and then extending it to all diagrams in Theorem 4.9.

The commutative diagram in Figure 8 summarizes the reduction procedure we describe in this section.

$$\begin{array}{ccc}
W_{n-2i}^{n-2i} & \xrightarrow{\varphi_{k-i}^{n-2i}} & (\mathbb{C}^2)_{k-i}^{\otimes n-2i} \\
D \mapsto \Pi_j \delta_j \cdot D \downarrow & & \downarrow D \mapsto \Pi_j \delta_k \cdot D \cdot \Pi_{j=0}^i \epsilon_k^{n-2i+2j} \\
W_{n-2i}^n & \xrightarrow{\varphi_k^n} & (\mathbb{C}^2)_k^{\otimes n}
\end{array}$$

Figure 8: How the computation of  $\varphi_k^n(W_{n-2i}^n)$  reduces to the computation of  $\varphi_{k-i}^{n-2i}(W_{n-2i}^{n-2i})$ .

Recall that  $(\mathbb{C}^2)_{k-1}^{\otimes n-2} \cong W_{n-2}^{n-2} \oplus \cdots \oplus W_{n-2k}^{n-2}$  and  $(\mathbb{C}^2)_k^{\otimes n} \cong W_n^n \oplus W_{n-2}^n \oplus \cdots \oplus W_{n-2k}^n$ . The action of  $D \in \text{tl}_{n-2}$  on  $\mathbb{C}_k^{\otimes n}$  is defined using the embedding  $\text{TL}_{n-2} \hookrightarrow \text{TL}_n$  which sends  $e_i$  to  $e_{i+2}$ . An example of the embedding is in Figure 9.




Figure 9: A basis element of  $\text{TL}_3$  embedded into  $\text{TL}_5$ .

**Definition 4.1.** Define the map  $\Phi: (\mathbb{C}^2)_{k-1}^{\otimes n-2} \rightarrow ((\mathbb{C}^2)_k^{\otimes n})^{e_1=\beta}$  by  $\Phi: D \mapsto \delta_1^n \cdot D \cdot \epsilon_k^n$  for diagrams  $D$  in  $C_{k-1}^{n-2}$ .

The map  $\Phi$  is a specific case of the rightmost map in Figure 8.

**Lemma 4.2.** *The map  $\Phi$  is an isomorphism of  $\text{TL}_{n-2}$ -representations.*

*Proof.* We can verify that  $\Phi$  is well-defined and commutes with the action of  $\text{TL}_{n-2}$ , so it is a homomorphism.

Let  $S$  be the set of diagrams  $D$  in  $C_k^n$  that can be expressed in the form  $D = \delta_1^n \cdot D'$ , where  $D'$  is any diagram in  $\text{Hom}([n], [n-2])$  in  $\text{TL}$  (in other words,  $D$  contains a link in the top left corner). Let  $R$  be the set of diagrams in the same set  $C_k^n$  that cannot be expressed in this way (in other words, the diagrams in  $R$  do not have a link in the top left corner).

A vector  $v$  in  $((\mathbb{C}^2)_k^{\otimes n})^{e_1=\beta}$  can be expressed as  $\sum_{D \in S} a_D D + \sum_{A \in R} b_A A$ , where the  $a_D$  and  $b_A$  are coefficients in  $\mathbb{C}$ . As every individual component  $a_D D$  has eigenvector  $e_1$  with

eigenvalue  $\beta$ ,  $\sum_{A \in R} b_A A$  must also be  $e_1$ -invariant with eigenvalue  $\beta$ . However,  $e_1$  sends any diagram  $A \in R$  to a diagram in  $S$ , which implies that  $\sum_{A \in R} b_A A = 0$ , so  $v$  is spanned by diagrams in  $S$ .

We now show that  $\Phi$  is surjective. Every diagram in  $S$  has  $e_1$  as an eigenvector with eigenvalue  $\beta$ , so we conclude that  $((\mathbb{C}^2)_k^{\otimes n})^{e_1=\beta}$  is generated by diagrams of the form  $\delta_1^n \cdot D$ . Furthermore, as such a diagram in  $\text{Ind}_{\text{TL}_k \otimes \text{TL}_{n-k}}^{\text{TL}_n} \mathbb{C}_{\text{triv}}$  must also have a link on the bottom line connecting the  $k$ th and  $(k+1)$ th vertices, it can also be expressed in the form  $\delta_1^n \cdot D \cdot \epsilon_k^n$ .

Injectivity follows from construction, so  $\Phi$  is an isomorphism.  $\square$

*Remark.* The map  $\Phi: D \mapsto \delta_1^n \cdot D \cdot \epsilon_k^n$  in Lemma 4.2 is adding a link to the top left and a link on the bottom line between the  $k$ th and  $(k+1)$ th vertices of  $D$ .

**Example 4.3.** The diagram on the left maps under  $\Phi$  to the diagram in the right in Figure 10.




Figure 10: An example of a map from a diagram in  $(\mathbb{C}^2)_2^{\otimes 3}$  to  $(\mathbb{C}^2)_2^{\otimes 5}$  under  $\Phi$ .

**Definition 4.4.** For each  $1 \leq i \leq n/2$ , define  $\Phi_{n-2i}: W_{n-2i}^{n-2} \rightarrow (W_{n-2i}^n)^{e_1=\beta}$  by

$$\Phi_{n-2i}: D \mapsto \delta_1^n \cdot D.$$

This is a specific case of the leftmost map in Figure 8.

**Lemma 4.5.** *The map  $\Phi_{n-2i}$  is an isomorphism of  $\text{TL}_n$ -modules for all  $i$ .*

*Proof.* Using a method similar to the proof of Lemma 4.2, we can verify that  $\Phi_{n-2i}$  is both well-defined and intertwining with the  $\text{TL}_{n-2}$  action for all  $i$ , so it is a homomorphism.

Let  $S$  be the set of diagrams  $D$  in  $w_{n-2i}^n$  that can be expressed as  $\delta_1^n \cdot D$  where  $D \in w_{n-2i}^{n-2}$  (so  $S$  is the set of diagrams that have a link in the top left corner). Let  $R$  be the set of diagrams in  $w_{n-2i}^n$  that cannot be expressed this way. A vector  $v$  in  $(W_{n-2i}^n)^{e_1=\beta}$  can be expressed as  $\sum_{D \in S} a_D D + \sum_{A \in R} b_A A$ . As every individual component  $a_D D$  has  $e_1$  as an eigenvector with eigenvalue  $\beta$ ,  $\sum_{A \in R} b_A A$  must also have  $\beta$  as the  $e_1$ -eigenvalue. The action

of  $e_1$  on any diagram in  $w_{n-2i}^n$  must map to either zero or a diagram in  $S$ , so it cannot send  $A \in R$  to  $\beta A$ . This implies that  $\sum_{A \in R} b_A A = 0$ , so  $v$  is spanned by diagrams in  $S$ . Furthermore, every diagram in  $S$  has a  $e_1$ -eigenvalue  $\beta$ , so we conclude that  $(W_{n-2i}^n)^{e_1=\beta}$  is generated by diagrams that can be expressed in the form  $\delta_1^n \cdot D$ .

The above implies that  $\Phi_{n-2i}$  is surjective, and injectivity follows from construction, so  $\Phi_{n-2i}$  is an isomorphism.  $\square$

*Remark.* The map  $\Phi_{n-2i}: D \mapsto \delta_1^n \cdot D$  in Lemma 4.5 is adding a link to the top left corner of  $D$ .

In addition to Lemma 4.2 and Lemma 4.5, we can use the following general result, which follows from Schur's Lemma.

**Lemma 4.6** (Schur's Lemma). *Given irreducible representations  $A_1, \dots, A_r$  and  $A'_1, \dots, A'_r$  of  $\text{TL}_n$  such that  $A_i \not\cong A_j$  and  $A'_i \not\cong A'_j$  when  $i \neq j$  and  $\varphi_i: A_i \cong A'_i$  is an isomorphism for all  $i$ , if  $\varphi: A_1 \oplus \dots \oplus A_r \cong A'_1 \oplus \dots \oplus A'_r$  is an isomorphism, then  $\varphi|_{A_i}$  lands in  $A'_i$  and is a multiple of  $\varphi_i$ .*

We can now prove the following proposition, which reduces the problem of computing the images of diagrams with links on the top left corner in  $w_{n-2i}^n$  to computing the images of diagrams in  $w_{n-2i}^{n-2}$ . This proposition is generalized in Theorem 4.9.

**Proposition 4.7.** *The image of the isomorphism  $\Phi: (\mathbb{C}^2)_{k-1}^{\otimes n-2} \rightarrow ((\mathbb{C}^2)_k^{\otimes n})^{e_1=\beta}$  of each component  $W_{n-2i}^{n-2}$  of  $(\mathbb{C}^2)_{k-1}^{\otimes n-2}$  falls in  $W_{n-2i}^n$ . Furthermore, the restriction of  $\Phi$  for each component  $W_{n-2i}^{n-2}$  is equal to the map  $\Phi_{n-2i}$ .*

*Proof.* Lemma 4.2 gives an isomorphism  $\Phi: (\mathbb{C}^2)_{k-1}^{\otimes n-2} \rightarrow ((\mathbb{C}^2)_k^{\otimes n})^{e_1=\beta}$  and Lemma 4.5 give isomorphisms between irreducible representations  $\Phi_{n-2}: W_{n-2}^{n-2} \rightarrow (W_{n-2}^n)^{e_1=\beta}, \dots, \Phi_{n-2k}: W_{n-2k}^{n-2} \rightarrow (W_{n-2k}^n)^{e_1=\beta}$ . Lemma 4.6 implies that the restriction  $\Phi|_{W_{n-2i}^{n-2}}$  is precisely  $\Phi_{n-2i}$ .  $\square$

From this inductive procedure, one can also compute the images of every diagram in  $w_{n-2}, \dots, w_{n-2k}^n$  under  $\varphi_k^n$  given the images of the diagrams in  $w_{n-2}, \dots, w_{n-2k}^{n-2}$  under  $\varphi_{k-1}^{n-1}$ .

**Lemma 4.8.** *Given any diagram  $\delta_i^n \cdot D$  in either a Specht module or the Temperley-Lieb algebra, the action of  $e_{i+1}$  sends it to  $\delta_{i+1}^n \cdot D$ .*

*Proof.* This lemma follows by the fact that  $e_{i+1}\delta_i = \delta_{i+1}$ .  $\square$

The following theorem is the main result of this section, allowing us to reduce the problem of computing  $\varphi_k^n(W_{n-2i}^n)$  to computing  $\varphi_{k-i}^{n-2i}(W_{n-2i}^{n-2i})$ .

**Theorem 4.9.** *Let  $D = \delta_j^n D'$ , where  $D' \in w_{n-2i}^{n-2}$ , be a diagram in  $w_{n-2i}^n$  for some  $1 \leq i \leq k$ . Let  $\varphi_{k-1}^{n-2}(D') = \sum_{A \in \text{tl}_{n-2}} c_A A$ . Then*

$$\varphi_k^{n-2}(D) = \sum_{A \in \text{tl}_{n-2}} c_A \delta_j^n \cdot A \cdot \epsilon_k^n.$$

*Proof.* Lemma 4.8 implies that given  $\varphi_k^n(\delta_i^j \cdot D)$  for any  $\delta_i^j \cdot D \in (W_{n-2i}^n)^{e_j=\beta}$ , one can obtain an element in  $(W_{n-2i}^n)^{e_{j+1}=\beta}$  by  $\delta_j^n D \mapsto \delta_{j+1}^n D$ . Note that this operation gives an isomorphism of vector spaces  $(W_{n-2i}^n)^{e_j=\beta} \cong (W_{n-2i}^n)^{e_{j+1}=\beta}$ , so every element of  $(W_{n-2i}^n)^{e_{j+1}=\beta}$  can be obtained this way.

As Proposition 4.7 gives the image of  $(W_{n-2i}^n)^{e_1=\beta}$  under  $\varphi_k^n$ , we can compute the image of  $(W_{n-2i}^n)^{e_j=\beta}$  under  $\varphi_k^n$  for all  $1 \leq j \leq n-1$ , yielding the explicit irreducible decomposition of  $(\mathbb{C}^2)_k^{\otimes n}$ .  $\square$

With this theorem, we obtain that for  $i = 1$ , Figure 8 commutes. We can repeatedly apply this theorem to obtain the reduction described in the diagram for general  $i$ .

*Remark.* In this section, we reduced the problem of computing the images of the diagrams in  $w_{n-2i}^n$  with links in the top left corner, and we reduced the problem of computing the images of the other diagrams of  $w_{n-2i}^n$  through applying actions of  $e_j$  to the former case. Alternatively, we could have also directly computed these using reduction, as  $(\mathbb{C}^2)_{k-1}^{n-2}$  is isomorphic to  $((\mathbb{C}^2)_k^n)^{e_j=\beta}$  as  $\text{TL}_{i-1} \otimes \text{TL}_{n-i-1}$ -modules.

## 5 The Decomposition $(\mathbb{C}^2)_k^{\otimes n-1} \oplus (\mathbb{C}^2)_{k-1}^{\otimes n-1} \cong (\mathbb{C}^2)_k^{\otimes n} |_{\text{TL}_{n-1}}$

The spin representation can be decomposed into a direct sum of “smaller” spin representations. In this section, we describe the isomorphism  $(\mathbb{C}^2)_k^{\otimes n-1} \oplus (\mathbb{C}^2)_{k-1}^{\otimes n-1} \cong (\mathbb{C}^2)_k^{\otimes n} |_{\text{TL}_{n-1}}$  explicitly, using it to give an inductive procedure to compute  $\varphi_k^n(W_n^n)$  in the next section.

We use the following result from [12] in order to prove Proposition 5.2, which allows us to decompose the spin representation into spin representations of a smaller number of electrons, thus allowing us to describe an inductive procedure to calculate the image of the generator of  $\varphi_k^n(W_n^n)$  in Section 6.

**Lemma 5.1** ([12]). *There is an isomorphism of  $\text{TL}_{n-1}$ -modules  $W_k^n |_{\text{TL}_{n-1}} \cong W_{k-1}^{n-1} \oplus W_{k+1}^{n-1}$ .*

**Proposition 5.2.** *There is an isomorphism of  $\mathrm{TL}_{n-1}$ -modules  $(\mathbb{C}^2)_k^{\otimes n}|_{\mathrm{TL}_{n-1}} \cong (\mathbb{C}^2)_k^{\otimes n-1} \oplus (\mathbb{C}^2)_{k-1}^{\otimes n-1}$ .*

*Proof.* Proposition 2.17 states that  $(\mathbb{C}^2)_k^{\otimes n} \cong W_n^n \oplus \cdots \oplus W_{n-2k}^n$ . Restricting the action on both sides to  $\mathrm{TL}_{n-1}$ , we obtain from Lemma 5.1 that  $(\mathbb{C}^2)_k^{\otimes n}$  is isomorphic to

$$\begin{aligned} W_{n-1}^{n-1} \oplus W_{n+1}^{n-1} \cdots \oplus W_{n-2k-1}^{n-1} \oplus W_{n-2k+1}^{n-1} &= W_{n-1}^{n-1} \oplus W_{n-4}^{n-1} \oplus W_{n-2}^{n-1} \cdots \oplus W_{n-2k-1}^{n-1} \oplus W_{n-2k+1}^{n-1} \\ &= (\mathbb{C}^2)_k^{\otimes n-1} \oplus (\mathbb{C}^2)_{k-1}^{\otimes n-1}. \end{aligned} \quad \square$$

We describe one such isomorphism explicitly.

**Definition 5.3.** We define the map  $\Theta_k^n \oplus \Psi_k^n : (\mathbb{C}^2)_k^{\otimes n-1} \oplus (\mathbb{C}^2)_{k-1}^{\otimes n-1} \rightarrow (\mathbb{C}^2)_k^{\otimes n}|_{\mathrm{TL}_{n-1}}$  on each component.

- Let  $\Theta_k^n$  be defined by mapping a diagram  $D \in C_k^{n-1}$  to the diagram  $D$  with a link added to the right.
- Define  $\Psi_k^n$  as mapping the identity diagram  $\mathrm{id}_{n-1}$  to  $\delta_k^n \epsilon_k^n + [2]_q \delta_{k+1}^n \epsilon_k^n + \cdots + [n-k]_q \delta_{n-1}^n \epsilon_k^n$ . Extend it to be compatible with the  $\mathrm{TL}_{n-1}$ -action.

In this section, we show that  $\Theta_k^n \oplus \Psi_k^n$  is an isomorphism. First, we show that it is a homomorphism.

**Proposition 5.4.** *The maps  $\Theta_k^n$  and  $\Psi_k^n$  are well-defined homomorphisms.*

*Proof.* Firstly,  $\Theta_k^n$  is a homomorphism of  $\mathrm{TL}_{n-1}$ -modules. Indeed, adding a link at the end does not affect any actions or relations.

We prove that the actions of  $e_2, \dots, e_{n-1}$  send  $\Phi_k^n(\mathrm{id}_{n-1})$  to 0. First, we consider  $e_k$ . The action annihilates all of  $\delta_{k+2}^n \epsilon_k^n$  to  $\delta_{n-1}^n \epsilon_k^n$  as it creates a link on the bottom line between the  $(k+2)$ th and  $(k+3)$ th vertices. What remains is  $e_k(\delta_k^n \epsilon_k^n + [2]_q \delta_{k+1}^n \epsilon_k^n) = \beta e_k + [2]_q e_k = 0$ . Now we consider  $e_i$  for  $i > k$ . The action annihilates everything except for  $[i-k]_q \delta_{i-1}^n \epsilon_k^n + [i-k+1]_q \delta_i^n \epsilon_k^n$ . Furthermore,  $e_i([i-k]_q \delta_{i-1}^n \epsilon_k^n + [i-k+1]_q \delta_i^n \epsilon_k^n + [i-k+2]_q \delta_{i+1}^n \epsilon_k^n) = [i-k]_q \delta_i^n \epsilon_k^n + [i-k+1]_q \beta \delta_i^n \epsilon_k^n + [i-k+2]_q \delta_i^n \epsilon_k^n = 0$ .

As  $(\mathbb{C}^2)_k^{\otimes n-1} \cong \mathrm{Ind}_{\mathrm{TL}_k \otimes \mathrm{TL}_{n-k}}^{\mathrm{TL}_n} \mathbb{C}_{\mathrm{triv}}$ , the universal property of induced representations states that there are no further relations.  $\square$

We compute the image of certain diagrams under  $\Psi_k^n$ , which will be used in Section 6.

**Lemma 5.5.** *The map  $\Psi_k^n$  sends  $\delta_i^{n-1} \epsilon_{k-1}^{n-1}$  to  $\delta_i^n \epsilon_{k-1}^n (\delta_k^n + [2]_q \delta_{k+1}^n + \cdots + [n-k]_q \delta_{n-1}^n) \epsilon_k^n$ .*

*Proof.* To obtain the image of  $\delta_{k-1}^{n-1}\epsilon_{k-1}^{n-1}$  under  $\Psi_k^n$ , we apply  $e_{k-1}$  to  $\text{id}_{n-1}$  and compute its image. The image will be  $\delta_{k-1}^n\epsilon_{k-1}^n(\delta_k^n\epsilon_k^n + [2]_q\delta_{k+1}^n\epsilon_k^n + \cdots + [n-k]_q\delta_{n-1}^n\epsilon_k^n)$  by Definition 5.3.  $\square$

Finally, we prove the following result, which allows us to use induction to compute  $\varphi_k^n(W_n^n)$  in Section 6.

**Theorem 5.6.** *The map  $\Theta_k^n \oplus \Psi_k^n$  defines an isomorphism of  $\text{TL}_{n-1}$ -modules.*

*Proof.* Each of the diagrams in  $C_k^{n-1}$  maps to a unique diagram in  $C_k^n$  with a straight link on the  $n$ th vertex. The last term the images of each of the diagrams in  $C_{k-1}^{n-1}$  gives a diagram  $\delta_i^n\epsilon_{k-1}^n\delta_{n-1}^n\epsilon_k^n$  that does not appear in the image of any other diagram under  $\Theta_k^n \oplus \Psi_k^n$ . Therefore, each of the images are linearly independent.  $\square$

## 6 Computation of $\varphi_k^n(W_n^n)$ Using the Decomposition

Recall that  $\Theta_k^n \oplus \Psi_k^n : (\mathbb{C}^2)_k^{\otimes n-1} \oplus (\mathbb{C}^2)_{k-1}^{\otimes n-1} \rightarrow (\mathbb{C}^2)_k^{\otimes n}|_{\text{TL}_{n-1}}$  is an isomorphism by Theorem 5.6.

We state the following proposition, which we use to inductively compute  $\varphi_k^n(W_n^n)$ .

**Proposition 6.1.** *Given vectors  $x_k^{n-1} \in \varphi_k^{n-1}(W_{n-1}^{n-1})$  in  $(\mathbb{C}^2)_k^{\otimes n-1}$  and  $x_{k-1}^{n-1} \in \varphi_{k-1}^{n-1}(W_{n-1}^{n-1})$  in  $(\mathbb{C}^2)_{k-1}^{\otimes n-1}$  that are annihilated by the actions of  $e_1, \dots, e_{n-2}$ , the one-dimensional space  $\varphi_k^n(W_n^n)$  is generated by the unique (up to scalar) vector in  $(\mathbb{C}^2)_k^{\otimes n}$  that is annihilated by the actions of all  $e_i$ s. This vector is the unique (up to scalar) linear combination of  $\Theta_k^n(x_k^{n-1})$  and  $\Psi_k^n(x_{k-1}^{n-1})$  that is also annihilated by  $e_{n-1}$ .*

*Proof.* The image of the identity diagram  $\text{id}_n$  in  $W_n^n$  under the homomorphism  $\varphi$ , is the unique vector (up to scalar) that is annihilated by the actions of all  $e_i$ s. This vector is in the space  $V$  of vectors that are annihilated by the actions of  $e_i$  for  $1 \leq i \leq n-2$ . Due to the  $\text{TL}_{n-1}$ -module isomorphism  $\Theta_k^n \oplus \Psi_k^n$ , the subspace  $V$  is 2-dimensional, generated by  $\Theta_k^n(x_k^{n-1})$  and  $\Psi_k^n(x_{k-1}^{n-1})$ . Therefore, the vector in  $(\mathbb{C}^2)_k^{\otimes n}$  that is annihilated by the actions of all  $e_i$ s is a linear combination of  $\Theta_k^n(x_k^{n-1})$  and  $\Psi_k^n(x_{k-1}^{n-1})$ .  $\square$

We construct a sequence  $x_i^n$ , in which each term is a linear combination of the images of the two before, and each term is annihilated by the action of every generator.

**Definition 6.2.** Let  $x_0^n = x_n^n = \text{id}_n$  and define  $x_k^n$  inductively by  $[n-k]_q x_k^n = [n]_q \Theta_k^n(x_k^{n-1}) + \Psi_k^n(x_{k-1}^{n-1})$ .

We show that  $x_k^n$  is the unique vector that is annihilated by all of  $e_1, \dots, e_{n-1}$  in  $(\mathbb{C}^2)_k^{\otimes n}$ . Note that for  $k = 0$ ,  $\text{id}_n$  is annihilated by the action of every generator, so this statement holds. In this section, we inductively prove that  $x_k^n$  defined this way is annihilated by the action of all the generators, for general  $k$ .

## 6.1 The $k = 1$ Case

We first show that  $x_1^n$  is annihilated by the action of all  $e_{i\text{'s}}$ . To do so, we introduce the following lemma.

**Lemma 6.3.** *We have the relation  $[n]_q[n - i - 1]_q + [i]_q = [n - 1]_q[n - i]_q$ .*

*Proof.* We have the above is equal to

$$\begin{aligned}
[n]_q[n - i - 1]_q + [i]_q &= \frac{q^n - q^{-n}}{q - q^{-1}} \cdot \frac{q^{n-i-1} - q^{i+1-n}}{q - q^{-1}} + \frac{q^i - q^{-i}}{q - q^{-1}} \\
&= \frac{q^{2n-i-1} - q^{i+1} - q^{-i-1} + q^{i+1-2n} + q^{i+1} - q^{-i+1} - q^{-1+i} + q^{-i-1}}{(q - q^{-1})^2} \\
&= \frac{q^{2n-i-1} + q^{i+1-2n} - q^{-i+1} - q^{-1+i}}{(q - q^{-1})^2} \\
&= \frac{(q^{n-1} - q^{1-n})(q^{n-i} - q^{i-n})}{(q - q^{-1})^2} \\
&= [n - 1]_q[n - i]_q. \quad \square
\end{aligned}$$

With this, we can show that the definition of  $x_1^n$  is consistent for the  $k = 1$  case.

**Proposition 6.4.** *The inductive computation of  $x_1^n$  by  $[n - 1]_q x_1^n = [n]_q \Theta_1^n(x_1^{n-1}) + \Psi_1^n(x_0^{n-1})$  yields that  $x_1^n$  is annihilated by the action of all the generators.*

*Proof.* We give a formula for  $x_1^n$  by induction. In the base case, we have that  $\Theta_1^1(x_1^1) = \text{id}_1$ .

By the inductive hypothesis, we have

$$[n]_q \Theta_1^n(x_1^{n-1}) = [n]_q \left( \sum_{i=1}^{n-2} [n - i - 1]_q \delta_i^n \epsilon_1^n + [n - 1]_q \text{id}_n \right)$$

and

$$\Psi_1^n(x_0^{n-1}) = \sum_{i=1}^{n-1} [i]_q \delta_i^n \epsilon_1^n.$$

The sum of these two expressions is

$$\begin{aligned} & [n]_q [n-1]_q \text{id}_n + [n-1]_q \delta_i^n \epsilon_1^n + \sum_{i=1}^{n-2} ([n]_q [n-i-1]_q + [i]_q) \delta_i^n \epsilon_1^n. \\ & = [n-1]_q \left( \sum_{i=0}^{n-1} [n-i]_q \delta_i^n \epsilon_1^n + [n]_q \text{id}_n \right), \end{aligned}$$

due to Lemma 6.3, so  $x_1^n = \sum_{i=0}^{n-1} [n-i]_q \delta_i^n \epsilon_1^n + [n]_q \text{id}_n$  which Proposition 3.2 states is annihilated by all the generators.  $\square$

## 6.2 Inductive Computation for General $k$

In this section, we prove that  $x_k^n$ , as defined inductively by  $[n-k]_q x_k^n = [n]_q \Theta_k^n(x_k^{n-1}) + \Psi_k^n(x_{k-1}^{n-1})$ , is annihilated by the actions of all the generators. The vectors  $\Theta_k^n(x_k^{n-1})$  and  $\Psi_k^n(x_{k-1}^{n-1})$  are both annihilated by the actions of  $e_1, \dots, e_{n-2}$ , so by Proposition 6.1, a linear combination of them must be annihilated by  $e_{n-1}$ . The action of  $e_{n-1}$  is defined by the action of  $\epsilon_{n-1}^n$ , so we consider this action for the sake of simplicity. Computing the coefficients of  $\epsilon_k^n$  allows us to obtain the linear combination, so it is sufficient to compute  $\text{coeff}_{\epsilon_k^n}(\epsilon_{n-1}^n \Theta_k^n(x_k^{n-1}))$  and  $\text{coeff}_{\epsilon_k^n}(\epsilon_{n-1}^n \Psi_k^n(x_{k-1}^{n-1}))$  and show that the coefficient of  $\epsilon_k^n$  of  $[n]_q \epsilon_{n-1}^n \Theta_k^n(x_k^{n-1}) + \epsilon_{n-1}^n \Psi_k^n(x_{k-1}^{n-1})$  is equal to 0.

We reduce the problem of computing  $\text{coeff}_{\epsilon_k^n}(\epsilon_{n-1}^n \Theta_k^n(x_k^{n-1}))$  and  $\text{coeff}_{\epsilon_k^n}(\epsilon_{n-1}^n \Psi_k^n(x_{k-1}^{n-1}))$  to computing the coefficient of  $\text{id}_{n-i}$  in  $x_j^{n-i}$  in Proposition 6.5, Proposition 6.6, and Lemma 6.7. We then compute the coefficient of  $\text{id}_{n-i}$  in  $x_j^{n-i}$  in Lemma 6.8, leading to Theorem 6.9, an inductive formula for computing  $\varphi_k^n(W_n^n)$ .

First, we reduce the problem of computing  $\text{coeff}_{\epsilon_k^n}(\epsilon_{n-1}^n \Theta_k^n(x_k^{n-1}))$ .

**Proposition 6.5.** *We have that  $\text{coeff}_{\epsilon_k^n}(\epsilon_{n-1}^n \Theta_k^n(x_k^{n-1})) = \text{coeff}_{\text{id}_{n-2}}(x_{k-1}^{n-2})$ .*

*Proof.* By the inductive hypothesis,  $[n-k-1]_q x_k^{n-1} = [n-1]_q \Theta_k^{n-1}(x_k^{n-2}) + \Psi_k^{n-1}(x_{k-1}^{n-2})$ . We must compute the coefficient of  $\epsilon_k^n$  of  $\frac{1}{[n-k-1]_q} (\epsilon_{n-1}^n \Theta_k^n \Theta_k^{n-1}(x_k^{n-2}) + \epsilon_{n-1}^n \Theta_k^n \Psi_k^{n-1}(x_{k-1}^{n-2}))$ . Since  $\Theta_i^j$  is a map adding a straight link to the right,  $\Theta_k^n \Theta_k^{n-1}(x_k^{n-2})$  is a diagram with a straight link on both the  $(n-1)$ th and  $n$ th vertices, so the action of  $\epsilon_{n-1}$  sends it to 0. Therefore, we only need to compute the coefficient of  $\epsilon_k^n$  in  $\frac{1}{[n-k-1]_q} \epsilon_{n-1}^n \Theta_k^n \Psi_k^{n-1}(x_{k-1}^{n-2})$ .

Note that  $\epsilon_{n-1}(D)$  is zero if  $D$  has two straight links to its right. As the action of  $\Theta_k^n$  adds a straight link to the right of a diagram,  $\Psi_k^{n-1}(x_{k-1}^{n-2})$  must have a simple link between the  $(n-2)$ th and  $(n-1)$ th vertices on the top row.

We show that  $\Psi_k^{n-1}(D)$  for  $D \in C_{k-1}^{n-2}$  has  $\epsilon_k^n$  as a term if and only if  $D$  is the identity diagram on  $n-2$  vertices. Note that by Theorem 5.6, we obtain that

$$\Psi_k^{n-1}(D) = D\delta_k^{n-1}\epsilon_k^{n-1} + [2]_q D\delta_{k+1}^{n-1}\epsilon_k^{n-1} + \cdots + [n-k-1]_q D\delta_{n-2}^{n-1}\epsilon_k^{n-1}.$$

If  $D$  is a nontrivial diagram in  $C_{k-1}^{n-2}$ , we only need to consider  $D\delta_{n-2}^{n-1}\epsilon_k^{n-1}$  as it is the only possibility where a simple link connecting the  $(n-2)$ th and  $(n-1)$ th vertices on the top line appears. Furthermore,  $D$  must also have a simple link between the  $(k-1)$ th and  $k$ th vertices. In the edge case where  $k \geq n-2$ , there will no longer be a link connecting the  $(n-1)$ th and  $(n-2)$ th vertices in  $D\delta_{n-2}^{n-1}\epsilon_k^{n-1}$ , so it will be sent to 0. If  $k < n-2$ , the link connecting the  $(k-1)$ th and  $k$ th vertices in  $D$  will create a second quasi-simple link on the bottom line in  $D\delta_{n-2}^{n-1}\epsilon_k^{n-1}$ . Therefore, in this case,  $\text{coeff}_{\epsilon_k}(\Psi_k^{n-1}(D)) = 0$ .

If  $D = \text{id}_{n-2}$ , recall that by Definition 5.3,  $\Psi_k^{n-1}(D) = \delta_k^{n-1}\epsilon_k^{n-1} + [2]_q \delta_{k+1}^{n-1}\epsilon_k^{n-1} + \cdots + [n-k-1]_q \delta_{n-2}^{n-1}\epsilon_k^{n-1}$ . In this case, as  $\epsilon_{n-1}^n \delta_i^n \epsilon_k^n = \epsilon_k^n$  when  $k \leq i \leq n-2$  implies that  $i = n-2$ , we obtain that  $\text{coeff}_{\epsilon_k^n}(\epsilon_{n-1}^n \Theta_k^n \Psi_k^{n-1}(\text{id}_{n-2})) = [n-k-1]_q$ . Therefore,

$$\text{coeff}_{\epsilon_k}(\epsilon_{n-1}^n \Theta_k^n(x_k^{n-1})) = \frac{1}{[n-k-1]_q} \text{coeff}_{\epsilon_k^n}(\epsilon_{n-1}^n \Theta_k^n \Psi_k^{n-1}(x_{k-1}^{n-2})) = \text{coeff}_{\text{id}_{n-2}}(x_{k-1}^{n-2}). \quad \square$$

Similarly to the above proposition, in the following proposition, we reduce the problem of computing  $\text{coeff}_{\epsilon_k^n}(\epsilon_{n-1}^n \Psi_k^n(x_{k-1}^{n-1}))$ .

**Proposition 6.6.** *We have the recurrence relation*

$$\text{coeff}_{\epsilon_k^n}(\epsilon_{n-1}^n \Psi_k^n(x_{k-1}^{n-1})) = -[n-k+1] \text{coeff}_{\text{id}_{n-1}}(x_{k-1}^{n-1}) + \text{coeff}_{\delta_{n-2}^{n-1}\epsilon_{k-1}^{n-1}}(x_{k-1}^{n-1}).$$

*Proof.* We show that if  $\Psi_k^n(D)$  has a term that is  $\epsilon_k^n$ ,  $D \in C_{k-1}^{n-1}$  must be either  $\text{id}_{n-1}$  or  $\delta_{n-2}^{n-1}\epsilon_{k-1}^{n-1}$ . Recall that by Theorem 5.6, we obtain that  $\Psi_k^n(D) = D\delta_k^n \epsilon_k^n + [2]_q D\delta_{k+1}^n \epsilon_k^n + \cdots + [n-k]_q D\delta_{n-1}^n \epsilon_k^{n-1}$ .

In the case where  $D = \text{id}_{n-1}$ , only  $\epsilon_{n-1}^n \delta_{n-2}^n \epsilon_k^n$  and  $\epsilon_{n-1}^n \delta_{n-1}^n \epsilon_k^n$  create a coefficient of  $\epsilon_k^n$ . The coefficient of  $\epsilon_k^n$  in  $\Psi_k^n(\text{id}_{n-1})$  is  $[n-k-1] - [2][n-k] = -[n-k+1]$ . In the case where  $D$  is not the identity, for the coefficient of  $\epsilon_k^n$  to be nonzero,  $D$  must be equal to  $\delta_{n-2}^{n-1}\epsilon_{k-1}^{n-1}$  as otherwise there will be a simple link on the top line. If this is the case, the only term where a coefficient of  $\epsilon_k^n$  appears is  $\delta_{n-2}^{n-1}\epsilon_{k-1}^{n-1}\delta_k^n \epsilon_k^n$ . In this case, the coefficient of  $\epsilon_k^n$  in  $\Psi_k^n(\delta_{n-2}^{n-1}\epsilon_{k-1}^{n-1})$  is 1. Combining this information, we obtain this proposition.  $\square$

We further reduce the original problem, proving that computing the coefficient of the

identity diagram is sufficient to compute  $\text{coeff}_{\delta_{n-2}^{n-1}\epsilon_{k-1}^{n-1}}(x_{k-1}^{n-1})$ .

**Lemma 6.7.** *We have the relation  $\text{coeff}_{\delta_{n-2}^{n-1}\epsilon_{k-1}^{n-1}}(x_{k-1}^{n-1}) = \text{coeff}_{\text{id}_{n-2}}(x_{k-2}^{n-2})$ .*

*Proof.* As  $[n-1-k]_q x_{k-1}^{n-1} = [n-1]_q \Theta_{k-1}^{n-1}(x_{k-1}^{n-2}) + \Psi_{k-1}^{n-1}(x_{k-2}^{n-2})$  and each component of  $\Theta_{k-1}^{n-1}(x_{k-1}^{n-2})$  has a rightmost straight link as  $\Theta_{k-1}^{n-1}$  is a map that adds a straight link to the right, we obtain that  $\text{coeff}_{\delta_{n-2}^{n-1}\epsilon_{k-1}^{n-1}}(x_{k-1}^{n-1}) = \text{coeff}_{\delta_{n-2}^{n-1}\epsilon_{k-1}^{n-1}}(\Psi_{k-1}^{n-1}(x_{k-2}^{n-2}))$ . Similarly to the proof of Proposition 6.5,  $\Psi_{k-1}^{n-1}(D)$  has a term of  $\delta_{n-2}^{n-1}\epsilon_{k-1}^{n-1}$  if and only if  $D = \text{id}_{n-2}$ . Therefore,

$$\text{coeff}_{\delta_{n-2}^{n-1}\epsilon_{k-1}^{n-1}}(x_{k-1}^{n-1}) = [n-k]_q \text{coeff}_{\text{id}_{n-2}}(x_{k-2}^{n-2}). \quad \square$$

Define  $[a]_q! = [a]_q [a-1]_q \dots [1]_q$  for integer  $a$ . Furthermore, let  $\begin{bmatrix} n \\ k \end{bmatrix}_q = \frac{[n]_q!}{[k]_q! [n-k]_q!}$ .

We can compute the coefficient of the identity diagram inductively by using the recurrence found in Lemma 6.7.

**Lemma 6.8.** *We obtain the formula  $\text{coeff}_{\text{id}_{n-1}}(x_{k-1}^{n-1}) = \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}_q$ .*

*Proof.* Since  $x_{k-1}^{n-1} = \frac{[n-1]_q}{[n-k]_q} \Theta_{k-1}^{n-1}(x_{k-1}^{n-2}) + \frac{1}{[n-k]!} \Psi_{k-1}^{n-1}(x_{k-2}^{n-2})$  by the inductive hypothesis and every term in  $\Psi_{k-1}^{n-1}(x_{k-2}^{n-2})$  is nontrivial, we conclude that

$$\text{coeff}_{\text{id}_{n-1}}(x_{k-1}^{n-1}) = \frac{[n-1]_q}{[n-k]_q} \text{coeff}_{\text{id}_{n-1}}(\Theta_{k-1}^{n-1}(x_{k-1}^{n-2})).$$

Repeating this process, we obtain that

$$\text{coeff}_{\text{id}_{n-1}}(x_{k-1}^{n-1}) = \frac{[n-1]_q}{[n-k]_q} \dots \frac{[k]_q}{[1]_q} = \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}_q. \quad \square$$

Combining all the results in this section, we can prove the following theorem, one of the main results of our paper.

**Theorem 6.9.** *The vector  $x_k^n$  in  $(\mathbb{C}^2)_k^{\otimes n}$ , defined by the inductive formula  $[n-k]_q x_k^n = [n]_q \Theta_k^n(x_k^{n-1}) + \Psi_k^n(x_{k-1}^{n-1})$ , is annihilated by the actions of all  $e_i$ s.*

*Proof.* We must show that the action of  $\epsilon_{n-1}^n$  on  $[n]_q \Theta_k^n(x_k^{n-1}) + \Psi_k^n(x_{k-1}^{n-1})$  is equal to 0. By Proposition 6.1, this is equivalent to proving that the coefficient of  $\epsilon_k^n$  of  $[n]_q \epsilon_{n-1}^n \Theta_k^n(x_k^{n-1}) + \epsilon_{n-1}^n \Psi_k^n(x_{k-1}^{n-1})$  is equal to 0.

By Proposition 6.5, Proposition 6.6, Lemma 6.7, and Lemma 6.8, we have that

$$\begin{aligned}
\frac{\text{coeff}_{\epsilon_k^n}(\epsilon_{n-1}^n \Psi_k^n(x_{k-1}^{n-1}))}{\text{coeff}_{\epsilon_k^n}(\epsilon_{n-1}^n \Theta_k^n(x_k^{n-1}))} &= \frac{-[n-k+1]_q \text{coeff}_{\text{id}_{n-1}}(x_{k-1}^{n-1}) + \text{coeff}_{\delta_{n-2}^{n-1} \epsilon_{k-1}^{n-1}}(x_{k-1}^{n-1})}{\text{coeff}_{\text{id}_{n-2}}(x_{k-1}^{n-2})} \\
&= \frac{-[n-k+1]_q \begin{bmatrix} n-1 \\ k-1 \end{bmatrix}_q + \begin{bmatrix} n-2 \\ k-2 \end{bmatrix}_q}{\begin{bmatrix} n-2 \\ k-1 \end{bmatrix}_q} \\
&= \frac{-[n-k+1]_q [n-1]_q + [k-1]_q}{[n-k]_q} \\
&= -[n]_q.
\end{aligned}$$

This implies that  $[n]_q \Theta_k^n(x_k^{n-1}) + \Psi_k^n(x_{k-1}^{n-1})$  is a vector that is annihilated by all generators in  $(\mathbb{C}^2)_k^{\otimes n}$ , proving the theorem.  $\square$

Theorem 6.9 gives an inductive procedure for the computation of  $\varphi_k^n(W_n^n)$ , while the results in Section 4, specifically Theorem 4.9 allow us to reduce the computation of  $\varphi_k^n(W_{n-2i}^n)$  to the computation of  $\varphi_k^{n-2i}(W_{n-2i}^{n-2i})$ . These two cases cover all of  $W_n^n \oplus \cdots \oplus W_{n-2k}^n$ , allowing us to explicitly compute the irreducible decomposition of the spin representation into Specht modules.

**Example 6.10.** We explicitly compute  $x_2^4$ . We have that  $[2]_q x_2^4 = [4] \Theta_2^4(x_2^3) + \Psi_2^4(x_1^3)$ . By Proposition 3.2,  $\Theta_2^4(x_2^3) = [1]_q \delta_1^4 \epsilon_2^4 + [2]_q \delta_2^4 \epsilon_2^4 + [3]_q \text{id}_4$  and  $x_1^3 = [1]_q \delta_2^3 \epsilon_1^3 + [2]_q \delta_1^3 \epsilon_1^3 + [3]_q \text{id}_3$ . Therefore,

$$\Psi_2^4(x_1^3) = [1]_q (\delta_2^4 \epsilon_2^4 + [2]_q \delta_2^4 \delta_2^2 \epsilon_2^2 \epsilon_2^4) + [2]_q (\delta_1^4 \epsilon_2^4 + \delta_1^4 \delta_3^4 \epsilon_2^2 \epsilon_2^4) + [3]_q (\delta_2^4 \epsilon_2^4 + [2]_q \delta_3^4 \epsilon_2^4).$$

From these, we obtain that

$$\begin{aligned}
[2]_q x_2^4 &= [4]_q ([1]_q \delta_1^4 \epsilon_2^4 + [2]_q \delta_2^4 \epsilon_2^4 + [3]_q \text{id}_4) \\
&\quad + [1]_q (\delta_2^4 \epsilon_2^4 + [2]_q \delta_2^4 \delta_2^2 \epsilon_2^2 \epsilon_2^4) + [2]_q (\delta_1^4 \epsilon_2^4 + \delta_1^4 \delta_3^4 \epsilon_2^2 \epsilon_2^4) + [3]_q (\delta_2^4 \epsilon_2^4 + [2]_q \delta_3^4 \epsilon_2^4) \\
&= [4]_q [3]_q \text{id}_4 + ([4]_q + [2]_q) \delta_1^4 \epsilon_2^4 + ([4]_q [2]_q + 1 + [3]_q) \delta_2^4 \epsilon_2^4 \\
&\quad + [3]_q [2]_q \delta_3^4 \epsilon_2^4 + [2]_q \delta_2^4 \delta_2^2 \epsilon_2^2 \epsilon_2^4 + [2]_q \delta_1^4 \delta_3^4 \epsilon_2^2 \epsilon_2^4.
\end{aligned}$$

Therefore,

$$x_2^4 = ([5]_q + 1) \text{id}_4 + [3]_q \delta_1^4 \epsilon_2^4 + ([4]_q + [2]_q) \delta_2^4 \epsilon_2^4 + [3]_q \delta_3^4 \epsilon_2^4 + \delta_2^4 \delta_2^2 \epsilon_2^2 \epsilon_2^4 + \delta_1^4 \delta_3^4 \epsilon_2^2 \epsilon_2^4.$$

## 7 Acknowledgements

I am grateful for my mentor Kenta Suzuki for answering my questions and guiding me through my research process. I would also like to thank Vasily Krylov for his help in coming up with the research problem. I am grateful to Tanya Khovanova for giving me both feedback on my paper and insights on how to be a better mathematician. I'm also thankful for Jonathan Bloom and Roman Bezrukavnikov for running the MIT Math program at RSI. Furthermore, I appreciate my tutor AnaMaria Perez for giving me feedback and encouraging me. I would like to give my thanks to Scott Kominers, Austin Luo, Richard Eager, and Akilan Sankaran for their feedback, as well as the first-week TAs Valencia Zhang, Varun Thvar, and Finan Gammell. I really appreciate RSI and all the RSI staff for creating this wonderful research opportunity, and would also like to thank CEE, MIT, and Citadel Securities for sponsoring me.

## References

- [1] É. Cartan. Les groupes projectifs qui ne laissent invariante aucune multiplicité plane. *Bulletin de la Société Mathématique de France*, 41:53–96, 1913.
- [2] P. A. M. Dirac. The quantum theory of the electron. part ii. *Proceedings of the royal society of London. Series A, Containing papers of a mathematical and physical character*, 118(779):351–361, 1928.
- [3] H. N. Temperley and E. H. Lieb. Relations between the ‘percolation’ and ‘colouring’ problem and other graph-theoretical problems associated with regular planar lattices: some exact results for the ‘percolation’ problem. *Condensed Matter Physics and Exactly Soluble Models: Selecta of Elliott H. Lieb*, pages 475–504, 2004.
- [4] W. Specht. Die irreduziblen darstellungen der symmetrischen gruppe. *Mathematische Zeitschrift*, 39(1):696–711, 1935.
- [5] P. Martin. *Potts models and related problems in statistical mechanics*, volume 5 of *Series on Advances in Statistical Mechanics*. World Scientific Publishing Co., Inc., Teaneck, NJ, 1991.
- [6] M. Khovanov. Graphical calculus, canonical bases and kazhdan-lusztig theory. *PhD Thesis*, 1997.
- [7] R. Chen. The dual danonical basis in the spin representation via the temperley-lieb algebra. 2025.
- [8] G. Lusztig. Canonical bases arising from quantized enveloping algebras. *Journal of the American Mathematical Society*, 3(2):447–498, 1990.
- [9] J. Chen. The temperley-lieb categories and skein modules. *arXiv preprint arXiv:1502.06845*, 2014.
- [10] J. de Groot. An introduction to the representation theory of temperley-lieb algebras. *Korteweg-de Vries Instituut voor Wiskunde Universiteit van Amsterdam*, 2015.
- [11] M. Jimbo. A q-difference analogue of  $u(\mathfrak{g})$  and the yang-baxter equation. *Letters in Mathematical Physics*, 10(1):63–69, 1985.
- [12] D. Ridout and Y. Saint-Aubin. Standard modules, induction and the structure of the Temperley-Lieb algebra. *Adv. Theor. Math. Phys.*, 18(5):957–1041, 2014.