

**18.100A Fall 2012: Assignment 23** due Fri. Nov. 30

*The rules are the same as for previous assignments.*

**Reading Mon.: 25.1-.3** Constructing closed and open sets; characterization of compact sets.

1. (2) Work 25.1/4ab, using and citing theorems in Chapter 25.

2. (2) Work 25.2/2; assume the set  $S$  is non-empty.

This is an improvement on the result in 24.7/1 (on Assignment 22). Once again, you are asked to prove there is a point  $\mathbf{a} \in S$  which is closest to  $\mathbf{0}$ , in the sense that no other point is closer.

(You have to adapt the Extremal Value Theorem (24.7B) to the non-compact set  $S$ ; it's analogous to what you had to do in  $\mathbf{R}^1$  for Problem 1, Exam 2. You will need the theorems of 25.1 and 25.2, rather than the definitions.)

3. (2: .5, 1.5)

a) Let  $P$  be the graph in  $\mathbf{R}^2$  of the parabola  $y = 2x^2 - 1$ . Discuss, with proof, whether it is open, closed, compact, or none of these. Use theorems in Chapter 25; don't go back to definitions if you don't have to.

b) Consider the sequence  $\mathbf{x}_n = (\cos n, \cos 2n)$ ,  $n = 0, 1, 2, \dots$  in  $\mathbf{R}^2$ .

Using the theorems in Chapter 25, and part (a), prove it has a subsequence which converges to a point  $\mathbf{a}$  on the parabola in part (a).

(This is like Problem 5 in Ass't. 21, except you now have more powerful theorems in Chapter 25 that you can use directly or adapt to this problem.)

4. (2) We can think of a function  $w = f(\mathbf{x})$  defined for all  $\mathbf{x} \in \mathbf{R}^2$  as giving a map  $f : \mathbf{R}^2 \rightarrow \mathbf{R}^1$ . If  $S \subset \mathbf{R}^1$ , we define

$$f^{-1}(S) = \{\mathbf{x} \in \mathbf{R}^2 : f(\mathbf{x}) \in S\} .$$

Assume  $f(\mathbf{x})$  is continuous; prove that if  $S$  is closed in  $\mathbf{R}^1$ , then  $f^{-1}(S)$  is closed in  $\mathbf{R}^2$ .

(This is a general property of continuous maps; conversely, if the statement is true for all closed subsets  $S$  of  $\mathbf{R}^1$ , then it follows that the function  $f(\mathbf{x})$  is continuous.)

Focus on what you have to prove about  $f^{-1}(S)$ ; use (1c) in Def'n. 25.1A as the definition of cluster point. Some care is needed.

*see over for reading and problems for Wed. class →*

**Reading Wed.:** 26.1-.2 to top third of p. 379

Functions defined by integrals with a parameter: continuity, differentiation under the integral sign.

5. (1.5: 1, .5) For the function  $\phi(x) = \int_0^1 \frac{e^{x+t}}{1+xt} dt$

(a) indicating the reasoning, find the largest  $x$ -interval on which the Continuity Theorem 26.1 predicts the integral will be continuous; your answer should take account of the  $t$ -values being restricted to a certain interval.

(b) Using the information provided by part (a), find  $\lim_{x \rightarrow 0} \int_0^1 \frac{e^{x+t}}{1+xt} dt$ . Cite the theorem(s) I hope you are using.

6. (2.5: 1, .5, 1) Let  $\phi(x) = \int_0^\pi \sin(xt) dt$ .

(a) The Derivative Theorem 26.2 says  $\phi'(x)$  exists for all  $x$  and gives a formula for it.

Using the formula, calculate  $\phi'(x)$  explicitly for all  $x$ , including  $x = 0$ , using standard integration techniques; your final answer should not have any integral signs in it, and should include all values of  $x$ .

(b) Verify your calculation of  $\phi'(x)$  for  $x \neq 0$  by first calculating  $\phi(x)$ ,  $x \neq 0$  explicitly, and then differentiating it by the usual rules.

(c) The Continuity Theorem predicts  $\phi'(x)$  will be continuous for all  $x$ . The explicit formula for it shows it is continuous if  $x \neq 0$ ; show it is continuous also at 0.

Do this, preferably by using the standard quadratic approximations to  $\sin u$  and  $\cos u$  at  $u = 0$  given by their Taylor series; or else by L'Hospital's rule.

7. (3) Work P26-1, showing that the Bessel function  $J_0(x)$  of order 0, given in the form of a definite integral involving a parameter  $x$ , is the solution to Bessel's ODE, with the given initial conditions.

You will need to differentiate  $J_0(x)$  twice; verify hypotheses each time, and verify the initial conditions are satisfied.

After substituting into Bessel's ODE and combining the integrals you get a very big definite integral, which is supposed to have the value 0, if the function  $J_0(x)$  really does solve the ODE.

Stop at this point and check your work carefully – the definite integral must be calculated correctly. Then try showing its value is 0 by using one of the standard techniques for transforming or evaluating integrals in Chapter 20. More than one technique will work.

If nothing strikes you after a few minutes, try sleeping on it and looking at the integral again in the morning.