

§8. $n \times n$ systems of linear ODEs

§8.1. Basics and examples

In this chapter we study $n \times n$ systems of ODEs.

We write them in vector form:

$$\boxed{\vec{y}' = A \vec{y}}$$

$$\text{where } \vec{y}(t) = \begin{pmatrix} y_1(t) \\ y_2(t) \\ \vdots \\ y_n(t) \end{pmatrix}$$

is the unknown vector and
 A is a given $n \times n$ matrix.

In §6 we studied the special case $n=2$

§8.1.1. Example: linear SEIR model

The SEIR model is a nonlinear system of 4 ODEs which can be used to model the spread of an infectious disease.
(As every model, it has its limitations.)

For instance the version here assumes that nobody dies!)

The time variable is t (in days)

and the unknown functions are

$S(t), E(t), I(t), R(t)$ (dimensionless)

$$\left. \begin{array}{l} S(t) = \text{Susceptible} \\ E(t) = \text{Exposed} \\ I(t) = \text{Infected} \\ R(t) = \text{Recovered} \end{array} \right\} \begin{array}{l} \text{Population} \\ \text{fraction at time } t \end{array}$$

The equations are:

$$\left. \begin{array}{l} S' = -\beta S \cdot I \\ E' = \beta S \cdot I - \alpha E \\ I' = \alpha E - \gamma I \\ R' = \gamma I \end{array} \right\} \begin{array}{l} (1) \\ (2) \\ (3) \\ (4) \end{array}$$

- The term $\beta S \cdot I$ in (1), (2)
accounts for susceptible people becoming exposed
via contact with an infected person
- The term αE in (2), (3)
accounts for exposed people becoming infected
(used because of incubation period)
- The term γI in (3), (4)
accounts for infected people recovering

$$S \xrightarrow{\beta} E \xrightarrow{\alpha} I \xrightarrow{\gamma} R$$

The constant parameters are

α (in $\frac{1}{\text{day}}$): inverse of the incubation period

β (in $\frac{1}{\text{day}}$): average contact rate in the population
(Social distancing \rightarrow smaller β)

γ (in $\frac{1}{\text{day}}$): inverse of the mean infectious period

For COVID-19 by one estimate:

$$\alpha = 0.2, \beta = 1.75, \gamma = 0.5$$


The system is nonlinear because of the term $\beta S \cdot I$ in (1) and (2).

We simplify it to a linear system by removing (1) and replacing S by 1:

$$\begin{cases} E' = \beta I - \alpha E \\ I' = \alpha E - \gamma I \\ R' = \gamma I \end{cases}$$

This might be a good approximation when E, I, R are small and $S \approx 1$ (in the initial stage).

It does not work when a large percentage of the population is infected and will not give the curves you might have seen.



Now let us write this system in vector notation: $\vec{y}(t) = \begin{pmatrix} E(t) \\ I(t) \\ R(t) \end{pmatrix}$,

$$\begin{cases} E' = -\alpha E + \beta I + O \cdot R \\ I' = \alpha E - \gamma I + O \cdot R \\ R' = O \cdot E + \delta \cdot I + O \cdot R \end{cases}$$

Get $\vec{y}' = A\vec{y}$ with $A = \begin{pmatrix} -\alpha & \beta & 0 \\ \alpha & -\gamma & 0 \\ 0 & \delta & 0 \end{pmatrix}$

For COVID-19, $A = \begin{pmatrix} -0.2 & 1.75 & 0 \\ 0.2 & -0.5 & 0 \\ 0 & 0.5 & 0 \end{pmatrix}$

§ 8.1.2. Companion Systems

Similarly to 2nd order ODEs,
if we have an n th order ODE

$$y^{(n)} + a_{n-1}y^{(n-1)} + \dots + a_1y' + a_0y = 0$$

we can convert it to the $n \times n$ companion system

$$\vec{y}' = A\vec{y} \text{ where } \vec{y} = \begin{pmatrix} y \\ y' \\ \vdots \\ y^{(n-1)} \end{pmatrix}, A = \begin{pmatrix} 0 & 1 & & & \\ 0 & 0 & 1 & & \\ & & \ddots & & \\ & -a_0 & -a_1 & \dots & 0 \\ & & & & 1 \\ & & & & -a_{n-1} \end{pmatrix}$$

Example: Find a companion system

for $y''' + 2y'' + 3y' + 4y = 0$

TECHNIQUE
PRACTICE

Solution: Write $\vec{y} = \begin{pmatrix} y_1 \\ y_2 \\ y_3 \end{pmatrix}$ where

Then

$$y'_1 = y_2$$

$$y'_2 = y_3$$


$$y'_3 = y''' = -4y - 3y' - 2y'' = -4y_1 - 3y_2 - 2y_3$$

So $\vec{y}' = A\vec{y}$ where $A = \begin{pmatrix} y_1 & y_2 & y_3 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \\ -4 & -3 & -2 \end{pmatrix}$

One can also write 1st order companion systems for systems of higher order ODEs:

Example: 3 coupled Springs

MITx 3.5.12




$k_1, k_2, k_3 > 0$
Spring constants
 $m_1, m_2 > 0$
masses

We assume that there is a position of the masses in which all springs are in equilibrium and denote by $[x_1(t), x_2(t)]$ the displacements of the masses from that position.

Using Newton's 2nd law + Hooke's law we arrive to the 2nd order system

$$\begin{cases} m_1 \cdot x_1'' = -k_1 x_1 + k_2(x_2 - x_1) \\ m_2 \cdot x_2'' = -k_2(x_2 - x_1) - k_3 x_2 \end{cases}$$

For example, the right-hand side of the first equation describes the total force on the first mass:



$-k_1 x_1$ is the force of the first spring

$k_2(x_2 - x_1)$ is the force of the second spring
($x_2 - x_1$ is the displacement of the second spring from equilibrium length)

We can rewrite the above ODE system as

$$\begin{cases} x_1'' = -\frac{k_1+k_2}{m_1} x_1 + \frac{k_2}{m_1} x_2 \\ x_2'' = \frac{k_2}{m_2} x_1 - \frac{k_2+k_3}{m_2} x_2 \end{cases}$$

How to write the companion system?
We introduce the velocities

$$v_1 = x_1', \quad v_2 = x_2', \quad \text{then}$$

$$\begin{cases} x_1' = v_1 \\ x_2' = v_2 \\ v_1' = -\frac{k_1+k_2}{m_1} x_1 + \frac{k_2}{m_1} x_2 \\ v_2' = \frac{k_2}{m_2} x_1 - \frac{k_2+k_3}{m_2} x_2 \end{cases}$$

Now we got the companion system

where $\vec{y}(t) = \begin{pmatrix} x_1(t) \\ x_2(t) \\ v_1(t) \\ v_2(t) \end{pmatrix}$ and

$$\vec{y}' = A\vec{y}$$

$$A = \begin{pmatrix} x_1 & x_2 & v_1 & v_2 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \\ -\frac{k_1+k_2}{m_1} & \frac{k_2}{m_1} & 0 & 0 \\ \frac{k_2}{m_2} & -\frac{k_2+k_3}{m_2} & 0 & 0 \end{pmatrix}$$