

18.03 EXERCISES

1. First-order ODE's

1A. Introduction; Separation of Variables

1A-1. Verify that each of the following ODE's has the indicated solutions (c_i, a are constants):

a) $y'' - 2y' + y = 0, \quad y = c_1 e^x + c_2 x e^x$
b) $xy' + y = x \sin x, \quad y = \frac{\sin x + a}{x} - \cos x$

1A-2. On how many arbitrary constants (also called *parameters*) does each of the following families of functions depend? (There can be less than meets the eye...; a, b, c, d, k are constants.)

a) $c_1 e^{kx}$ b) $c_1 e^{x+a}$ c) $c_1 + c_2 \cos 2x + c_3 \cos^2 x$ d) $\ln(ax + b) + \ln(cx + d)$

1A-3. Write down an explicit solution (involving a definite integral) to the following initial-value problems (IVP's):

a) $y' = \frac{1}{y^2 \ln x}, \quad y(2) = 0$ b) $y' = \frac{y e^x}{x}, \quad y(1) = 1$

1A-4. Solve the IVP's (initial-value problems):

a) $y' = \frac{xy + x}{y}, \quad y(2) = 0$ b) $\frac{du}{dt} = \sin t \cos^2 u, \quad u(0) = 0$

1A-5. Find the general solution by separation of variables:

a) $(y^2 - 2y) dx + x^2 dy = 0$ b) $x \frac{dv}{dx} = \sqrt{1 - v^2}$
c) $y' = \left(\frac{y-1}{x+1}\right)^2$ d) $\frac{dx}{dt} = \frac{\sqrt{1+x}}{t^2+4}$

1B. Standard First-order Methods

1B-1. Test the following ODE's for exactness, and find the general solution for those which are exact.

a) $3x^2 y dx + (x^3 + y^3) dy = 0$ b) $(x^2 - y^2) dx + (y^2 - x^2) dy = 0$
c) $ve^{uv} du + ye^{uv} dv = 0$ d) $2xy dx - x^2 dy = 0$

1B-2. Find an integrating factor and solve:

a) $2x dx + \frac{x^2}{y} dy = 0$ b) $y dx - (x + y) dy = 0, \quad y(1) = 1$
c) $(t^2 + 4) dt + t dx = x dt$ d) $u(du - dv) + v(du + dv) = 0. \quad v(0) = 1$

1B-3. Solve the homogeneous equations

$$\text{a) } y' = \frac{2y-x}{y+4x} \quad \text{b) } \frac{dw}{du} = \frac{2uw}{u^2-w^2} \quad \text{c) } xy dy - y^2 dx = x\sqrt{x^2-y^2} dx$$

1B-4. Show that a change of variable of the form $u = \frac{y}{x^n}$ turns $y' = \frac{4+xy^2}{x^2y}$ into an equation whose variables are separable, and solve it.

(Hint: as for homogeneous equations, since you want to get rid of y and y' , begin by expressing them in terms of u and x .)

1B-5. Solve each of the following, finding the general solution, or the solution satisfying the given initial condition.

$$\begin{array}{ll} \text{a) } xy' + 2y = x & \text{b) } \frac{dx}{dt} - x \tan t = \frac{t}{\cos t}, \quad x(0) = 0 \\ \text{c) } (x^2 - 1)y' = 1 - 2xy & \text{d) } 3v dt = t(dt - dv), \quad v(1) = \frac{1}{4} \end{array}$$

1B-6. Consider the ODE $\frac{dx}{dt} + ax = r(t)$, where a is a positive constant, and $\lim_{t \rightarrow \infty} r(t) = 0$.

Show that if $x(t)$ is any solution, then $\lim_{t \rightarrow \infty} x(t) = 0$. (Hint: use L'Hospital's rule.)

1B-7. Solve $y' = \frac{y}{y^3+x}$. Hint: consider $\frac{dx}{dy}$.

1B-8. The **Bernoulli** equation. This is an ODE of the form $y' + p(x)y = q(x)y^n$, $n \neq 1$. Show it becomes linear if one makes the change of dependent variable $u = y^{1-n}$.

(Hint: begin by dividing both sides of the ODE by y^n .)

1B-9. Solve these Bernoulli equations using the method described in 1B-8:

$$\text{a) } y' + y = 2xy^2 \quad \text{b) } x^2y' - y^3 = xy$$

1B-10. The **Riccati** equation. After the linear equation $y' = A(x) + B(x)y$, in a sense the next simplest equation is the Riccati equation

$$y' = A(x) + B(x)y + C(x)y^2,$$

where the right-hand side is now a quadratic function of y instead of a linear function. In general the Riccati equation is not solvable by elementary means. However,

a) show that if $y_1(x)$ is a solution, then the general solution is

$$y = y_1 + u,$$

where u is the general solution of a certain Bernoulli equation (cf. 1B-8).

b) Solve the Riccati equation $y' = 1 - x^2 + y^2$ by the above method.

1B-11. Solve the following second-order autonomous equations (“autonomous” is an important word; it means that the independent variable does not appear explicitly in the equation — it does lurk in the derivatives, of course.)

$$\text{a) } y'' = a^2y \quad \text{b) } yy'' = y'^2 \quad \text{c) } y'' = y'(1+3y^2), \quad y(0) = 1, \quad y'(0) = 2$$

1B-12. For each of the following, tell what type of ODE it is — i.e., what method you would use to solve it. (Don't actually carry out the solution.) For some, there are several methods which could be used.

1. $(x^3 + y) dx + x dy = 0$
2. $\frac{dy}{dt} + 2ty - e^{-t} = 0$
3. $y' = \frac{x^2 - y^2}{5xy}$
4. $(1 + 2p) dq + (2 - q) dp = 0$
5. $\cos x dy = (y \sin x + e^x) dx$
6. $x(\tan y)y' = -1$
7. $y' = \frac{y}{x} + \frac{1}{y}$
8. $\frac{dv}{du} = e^{2u+3v}$
9. $xy' = y + xe^{y/x}$
10. $xy' - y = x^2 \sin x$
11. $y' = (x + e^y)^{-1}$
12. $y' + \frac{2y}{x} - \frac{y^2}{x} = 0$
13. $\frac{dx}{dy} = -x \left(\frac{2x^2y + \cos y}{3x^2y^2 + \sin y} \right)$
14. $y' + 3y = e^{-3t}$
15. $x \frac{dy}{dx} - y = \sqrt{x^2 + y^2}$
16. $\frac{y' - 1}{x^2} = 1$
17. $xy' - 2y + y^2 = x^4$
18. $y'' = \frac{y(y+1)}{y'}$
19. $t \frac{ds}{dt} = s(1 - \ln t + \ln s)$
20. $\frac{dy}{dx} = \frac{3 - 2y}{2x + y + 1}$
21. $x^2y' + xy + y^2 = 0$
22. $y' \tan(x + y) = 1 - \tan(x + y)$
23. $y ds - 3s dy = y^4 dy$
24. $du = -\frac{1 + u \cos^2 t}{t \cos^2 t} dt$
25. $y' + y^2 + (2x + 1)y + 1 + x + x^2 = 0$
26. $y'' + x^2y' + 3x^3 = \sin x$

1C. Graphical and Numerical Methods

1C-1. For each of the following ODE's, draw a direction field by using about five isoclines; the picture should be square, using the intervals between -2 and 2 on both axes. Then sketch in some integral curves, using the information provided by the direction field. Finally, do whatever else is asked.

a) $y' = -\frac{y}{x}$; solve the equation exactly and compare your integral curves with the correct ones.

b) $y' = 2x + y$; find a solution whose graph is also an isocline, and verify this fact analytically (i.e., by calculation, not from a picture).

c) $y' = x - y$; same as in (b).

d) $y' = x^2 + y^2 - 1$

e) $y' = \frac{1}{x + y}$; use the interval -3 to 3 on both axes; draw in the integral curves that pass respectively through $(0, 0)$, $(-1, 1)$, $(0, -2)$. Will these curves cross the line $y = -x - 1$? Explain by using the Intersection Principle (Notes G, (3)).

1C-2. Sketch a direction field, concentrating on the first quadrant, for the ODE

$$y' = \frac{-y}{x^2 + y^2} .$$

Explain, using it and the ODE itself how one can tell that the solution $y(x)$ satisfying the initial condition $y(0) = 1$

- a) is a decreasing function for $y > 0$;
- b) is always positive for $x > 0$.

1C-3. Let $y(x)$ be the solution to the IVP $y' = x - y$, $y(0) = 1$.

a) Use the Euler method and the step size $h = .1$ to find an approximate value of $y(x)$ for $x = .1, .2, .3$. (Make a table as in notes G).

Is your answer for $y(.3)$ too high or too low, and why?

b) Use the Modified Euler method (also called Improved Euler, or Heun's method) and the step size $h = .1$ to determine the approximate value of $y(.1)$. Is the value for $y(.1)$ you found in part (a) corrected in the right direction — e.g., if the previous value was too high, is the new one lower?

1C-4. Use the Euler method and the step size $.1$ on the IVP $y' = x + y^2$, $y(0) = 1$, to calculate an approximate value for the solution $y(x)$ when $x = .1, .2, .3$. (Make a table as in Notes G.) Is your answer for $y(.3)$ too high or too low?

1C-5. Prove that the Euler method converges to the exact value for $y(1)$ as the progressively smaller step sizes $h = 1/n$, $n = 1, 2, 3, \dots$ are used, for the IVP

$$y' = x - y, \quad y(0) = 1 .$$

(First show by mathematical induction that the approximation to $y(1)$ gotten by using the step size $1/n$ is

$$y_n = 2(1 - h)^n - 1 + nh .$$

The exact solution is easily found to be $y = 2e^{-x} + x - 1$.)

1C-6. Consider the IVP $y' = f(x)$, $y(0) = y_0$.

We want to calculate $y(2nh)$, where h is the step size, using n steps of the Runge-Kutta method.

The exact value, by Chapter D of the notes, is $y(2nh) = y_0 + \int_0^{2nh} f(x) dx$.

Show that the value for $y(2nh)$ produced by Runge-Kutta is the same as the value for $y(2nh)$ obtained by using Simpson's rule to evaluate the definite integral.

1C-7. According to the existence and uniqueness theorem, under what conditions on $a(x)$, $b(x)$, and $c(x)$ will the IVP

$$a(x)y' + b(x)y = c(x), \quad y(x_0) = y_0$$

have a unique solution in some interval $[x_0 - h, x_0 + h]$ centered around x_0 ?

1D. Geometric and Physical Applications

1D-1. Find all curves $y = y(x)$ whose graphs have the indicated geometric property. (Use the geometric property to find an ODE satisfied by $y(x)$, and then solve it.)

- a) For each tangent line to the curve, the segment of the tangent line lying in the first quadrant is bisected by the point of tangency.
- b) For each normal to the curve, the segment lying between the curve and the x -axis has constant length 1.
- c) For each normal to the curve, the segment lying between the curve and the x -axis is bisected by the y -axis.
- d) For a fixed a , the area under the curve between a and x is proportional to $y(x) - y(a)$.

1D-2. For each of the following families of curves,

- (i) find the ODE satisfied by the family (i.e., having these curves as its integral curves);
- (ii) find the orthogonal trajectories to the given family;
- (iii) sketch both the original family and the orthogonal trajectories.
 - a) all lines whose y -intercept is twice the slope
 - b) the exponential curves $y = ce^x$
 - c) the hyperbolas $x^2 - y^2 = c$
 - d) the family of circles centered on the y -axis and tangent to the x -axis.

1D-3. Mixing A container holds V liters of salt solution. At time $t = 0$, the salt concentration is c_0 g/liter. Salt solution having concentration c_1 is added at the rate of k liters/min, with instantaneous mixing, and the resulting mixture flows out of the container at the same rate. How does the salt concentration in the tank vary with time?

Let $x(t)$ be the *amount* of salt in the tank at time t . Then $c(t) = \frac{x(t)}{V}$ is the concentration of salt at time t .

- a) Write an ODE satisfied by $x(t)$, and give the initial condition.
- b) Solve it, assuming that it is pure water that is being added. (Lump the constants by setting $a = k/V$.)
- c) Solve it, assuming that c_1 is constant; determine $c(t)$ and find $\lim_{t \rightarrow \infty} c(t)$. Give an intuitive explanation for the value of this limit.
- d) Suppose now that c_1 is not constant, but is decreasing exponentially with time:

$$c_1 = c_0 e^{-\alpha t}, \quad \alpha > 0.$$

Assume that $a \neq \alpha$ (cf. part (b)), and determine $c(t)$, by solving the IVP. Check your answer by putting $\alpha = 0$ and comparing with your answer to (c).

1D-4. Radioactive decay A radioactive substance **A** decays into **B**, which then further decays to **C**.

- a) If the decay constants of **A** and **B** are respectively λ_1 and λ_2 (the decay constant is by definition $(\ln 2/\text{half-life})$), and the initial amounts are respectively A_0 and B_0 , set up an ODE for determining $B(t)$, the amount of **B** present at time t , and solve it. (Assume $\lambda_1 \neq \lambda_2$.)
- b) Assume $\lambda_1 = 1$ and $\lambda_2 = 2$. Tell when $B(t)$ reaches a maximum.

1D-5. Heat transfer According to Newton's Law of Cooling, the rate at which the temperature T of a body changes is proportional to the difference between T and the external temperature.

At time $t = 0$, a pot of boiling water is removed from the stove. After five minutes, the

water temperature is 80°C . If the room temperature is 20°C , when will the water have cooled to 60°C ? (Set up and solve an ODE for $T(t)$.)

1D-6. Motion A mass m falls through air under gravity. Find its velocity $v(t)$ and its terminal velocity (that is, $\lim_{t \rightarrow \infty} v(t)$) assuming that

- air resistance is kv (k constant; this is valid for small v);
- air resistance is kv^2 (k constant; this is valid for high v).

Call the gravitational constant g . In part (b), lump the constants by introducing a parameter $a = \sqrt{gm/k}$.

1D-7. A loaded cable is hanging from two points of support, with Q the lowest point on the cable. The portion QP is acted on by the total load W on it, the constant tension T_Q at Q , and the variable tension T at P . Both W and T vary with the point P .

Let s denote the length of arc QP .

a) Show that $\frac{dx}{T_Q} = \frac{dy}{W} = \frac{ds}{T}$.


b) Deduce that if the cable hangs under its own weight, and $y(x)$ is the function whose graph is the curve in which the cable hangs, then

(i) $y'' = k\sqrt{1 + y'^2}$, k constant

(ii) $y = \sqrt{s^2 + c^2} + c_1$, c, c_1 constants

c) Solve the suspension bridge problem: the cable is of negligible weight, and the loading is of constant horizontal density. (“Solve” means: find $y(x)$.)

d) Consider the “Marseilles curtain” problem: the cable is of negligible weight, and loaded with equally and closely spaced vertical rods whose bottoms lie on a horizontal line. (Take the x -axis as the line, and show $y(x)$ satisfies the ODE $y'' = k^2y$.)



1E. First-order autonomous ODE's

1E-1. For each of the following autonomous equations $dx/dt = f(x)$, obtain a qualitative picture of the solutions as follows:

(i) draw horizontally the axis of the dependent variable x , indicating the critical points of the equation; put arrows on the axis indicating the direction of motion between the critical points; label each critical point as stable, unstable, or semi-stable. Indicate where this information comes from by including in the same picture the graph of $f(x)$, drawn in dashed lines;

(ii) use the information in the first picture to make a second picture showing the tx -plane, with a set of typical solutions to the ODE: the sketch should show the main qualitative features (e.g., the constant solutions, asymptotic behavior of the non-constant solutions).

- $x' = x^2 + 2x$
- $x' = -(x - 1)^2$
- $x' = 2x - x^2$
- $x' = (2 - x)^3$

M.I.T. 18.03 Ordinary Differential Equations
18.03 Notes and Exercises

©Arthur Mattuck and M.I.T. 1988, 1992, 1996, 2003, 2007, 2011