

8. RELATIVE PROJ AND THE BLOW UP

We want to define a relative version of Proj, in pretty much the same way we defined a relative version of Spec. We start with a scheme X and a quasi-coherent sheaf \mathcal{S} sheaf of graded \mathcal{O}_X -algebras,

$$\mathcal{S} = \bigoplus_{d \in \mathbb{N}} \mathcal{S}_d,$$

where $\mathcal{S}_0 = \mathcal{O}_X$. It is convenient to make some simplifying assumptions:

(†) X is Noetherian, \mathcal{S}_1 is coherent, \mathcal{S} is locally generated by \mathcal{S}_1 .

To construct relative Proj, we cover X by open affines $U = \text{Spec } A$. $\mathcal{S}(U) = H^0(U, \mathcal{S})$ is a graded A -algebra, and we get $\pi_U: \text{Proj } \mathcal{S}(U) \rightarrow U$ a projective morphism. If $f \in A$ then we get a commutative diagram

$$\begin{array}{ccc} \text{Proj } \mathcal{S}(U_f) & \longrightarrow & \text{Proj } \mathcal{S}(U) \\ \pi_{U_f} \downarrow & & \downarrow \pi_U \\ U_f & \longrightarrow & U. \end{array}$$

It is not hard to glue π_U together to get $\pi: \mathbf{Proj } \mathcal{S} \rightarrow X$. We can also glue the invertible sheaves together to get an invertible sheaf $\mathcal{O}(1)$.

The relative construction is very similar to the old construction.

Example 8.1. *If X is Noetherian and*

$$\mathcal{S} = \mathcal{O}_X[T_0, T_1, \dots, T_n],$$

then satisfies (†) and $\mathbf{Proj } \mathcal{S} = \mathbb{P}_X^n$.

Given a sheaf \mathcal{S} satisfying (†), and an invertible sheaf \mathcal{L} , it is easy to construct a quasi-coherent sheaf $\mathcal{S}' = \mathcal{S} \star \mathcal{L}$, which satisfies (†). The graded pieces of \mathcal{S}' are $\mathcal{S}_d \otimes \mathcal{L}^d$ and the multiplication maps are the obvious ones. There is a natural isomorphism

$$\phi: P' = \mathbf{Proj } \mathcal{S}' \rightarrow P = \mathbf{Proj } \mathcal{S},$$

which makes the diagram commute

$$\begin{array}{ccc} P' & \xrightarrow{\phi} & P \\ & \searrow \pi' & \swarrow \pi \\ & X & \end{array}$$

and

$$\phi^* \mathcal{O}_P(1) \simeq \mathcal{O}_{P'}(1) \otimes \pi'^* \mathcal{L}.$$

Note that π is always proper; in fact π is projective over any open affine and properness is local on the base.

There are two very interesting family of examples of the construction of relative Proj. Suppose that we start with a locally free sheaf \mathcal{E} of rank $r \geq 2$. Note that

$$\mathcal{S} = \bigoplus \text{Sym}^d \mathcal{E},$$

satisfies (†). $\mathbb{P}(\mathcal{E}) = \mathbf{Proj} \mathcal{S}$ is the **projective bundle** over X associated to \mathcal{E} . The fibres of $\pi: \mathbb{P}(\mathcal{E}) \rightarrow X$ are copies of \mathbb{P}^{r-1} , where $n = r - 1$. We have

$$\bigoplus_{l=0}^{\infty} \pi_* \mathcal{O}_{\mathbb{P}(\mathcal{E})}(l) = \mathcal{S},$$

so that in particular

$$\pi_* \mathcal{O}_{\mathbb{P}(\mathcal{E})}(1) = \mathcal{E}.$$

Also there is a natural surjection

$$\pi^* \mathcal{E} \rightarrow \mathcal{O}_{\mathbb{P}(\mathcal{E})}(1).$$

Indeed, it suffices to check both statements locally, so that we may assume that X is affine. The first statement is then (4.21) and the second statement reduces to the statement that the sections x_0, x_1, \dots, x_n generate $\mathcal{O}_P(1)$. The most interesting result is:

Proposition 8.2. *Let $g: Y \rightarrow X$ be a morphism.*

Then a morphism $f: Y \rightarrow \mathbb{P}(\mathcal{E})$ over X is the same as giving an invertible sheaf \mathcal{L} on Y and a surjection $g^ \mathcal{E} \rightarrow \mathcal{L}$.*

Proof. One direction is clear; if $f: Y \rightarrow \mathbb{P}(\mathcal{E})$ is a morphism over X , then the surjective morphism of sheaves

$$\pi^* \mathcal{E} \rightarrow \mathcal{O}_{\mathbb{P}(\mathcal{E})}(1),$$

pullsback to a surjective morphism

$$g^* \mathcal{E} = f^*(\pi^* \mathcal{E}) \rightarrow \mathcal{L} = f^* \mathcal{O}_{\mathbb{P}(\mathcal{E})}(1).$$

Conversely suppose we are given an invertible sheaf \mathcal{L} and a surjective morphism of sheaves

$$g^* \mathcal{E} \rightarrow \mathcal{L}.$$

I claim that there is then a unique morphism $f: Y \rightarrow \mathbb{P}(\mathcal{E})$ over X , which induces the given surjection. By uniqueness, it suffices to prove this result locally. So we may assume that $X = \text{Spec } A$ is affine and

$$\mathcal{E} = \bigoplus_{i=0}^n \mathcal{O}_X,$$

is free. In this case surjectivity reduces to the statement that the images s_0, s_1, \dots, s_n of the standard sections generate \mathcal{L} , and the result reduces to one we have already proved. \square

Definition 8.3. Let X be a Noetherian scheme and let \mathcal{I} be a coherent sheaf of ideals on X . Let

$$\mathcal{S} = \bigoplus_{d=0}^{\infty} \mathcal{I}^d,$$

where $\mathcal{I}^0 = \mathcal{O}_X$ and \mathcal{I}^d is the d th power of \mathcal{I} . Then \mathcal{S} satisfies (\dagger) .

$\pi: \mathbf{Proj} \mathcal{S} \rightarrow X$ is called the **blow up** of \mathcal{I} (or Y , if Y is the subscheme of X associated to \mathcal{I}).

Example 8.4. Let $X = \mathbb{A}_k^n$ and let P be the origin. We check that we get the usual blow up. Let

$$A = k[x_1, x_2, \dots, x_n].$$

As $X = \text{Spec } A$ is affine and the ideal sheaf \mathcal{I} of P is the sheaf associated to $\langle x_1, x_2, \dots, x_n \rangle$,

$$Y = \mathbf{Proj} \mathcal{S} = \text{Proj } S,$$

where

$$S = \bigoplus_{d=0}^{\infty} I^d.$$

There is a surjective map

$$A[y_1, y_2, \dots, y_n] \rightarrow S,$$

of graded rings, where y_i is sent to x_i . $Y \subset \mathbb{P}_A^n$ is the closed subscheme corresponding to this morphism. The kernel of this morphism is

$$\langle y_i x_j - y_j x_i \rangle,$$

which are the usual equations of the blow up.

Definition 8.5. Let $f: X \rightarrow Y$ be a morphism of schemes. We are going to define the **inverse image ideal sheaf** $\mathcal{I}' \subset \mathcal{O}_Y$. First we take the inverse image of the sheaf $f^{-1}\mathcal{I}$, where we just think of f as being a continuous map. Then $f^{-1}\mathcal{I} \subset f^{-1}\mathcal{O}_Y$. Let $\mathcal{I}' = f^{-1}\mathcal{I} \cdot \mathcal{O}_Y$ be the ideal generated by the image of $f^{-1}\mathcal{I}$ under the natural morphism $f^{-1}\mathcal{O}_Y \rightarrow \mathcal{O}_X$.

Theorem 8.6 (Universal Property of the blow up). Let X be a Noetherian scheme and let \mathcal{I} be a coherent ideal sheaf.

If $\pi: Y \rightarrow X$ is the blow up of \mathcal{I} then $\pi^{-1}\mathcal{I} \cdot \mathcal{O}_Y$ is an invertible sheaf. Moreover π is universal amongst all such morphisms. If

$f: Z \rightarrow X$ is any morphism such that $f^{-1}\mathcal{I} \cdot \mathcal{O}_Z$ is invertible then there is a unique induced morphism $g: Z \rightarrow Y$ which makes the diagram commute

$$\begin{array}{ccc} Z & \xrightarrow{g} & Y \\ & \searrow f & \downarrow \pi \\ & & X. \end{array}$$

Proof. By uniqueness, we can check this locally. So we may assume that $X = \text{Spec } A$ is affine. As \mathcal{I} is coherent, it corresponds to an ideal $I \subset A$ and

$$X = \text{Proj} \bigoplus_{d=0}^{\infty} I^d.$$

Now $\mathcal{O}_Y(1)$ is an invertible sheaf on Y . It is not hard to check that $\pi^{-1}\mathcal{I} \cdot \mathcal{O}_Y = \mathcal{O}_Y(1)$.

Pick generators a_0, a_1, \dots, a_n for I . This gives rise to a surjective map of rings

$$\phi: A[x_0, x_1, \dots, x_n] \rightarrow I,$$

whence to a closed immersion $Y \subset \mathbb{P}_A^n$. The kernel of ϕ is generated by all homogeneous polynomials $F(x_0, x_1, \dots, x_n)$ such that $F(a_0, a_1, \dots, a_n) = 0$.

Now the elements a_0, a_1, \dots, a_n pullback to global sections s_0, s_1, \dots, s_n of the invertible sheaf $\mathcal{L} = f^{-1}\mathcal{I} \cdot \mathcal{O}_Y$ and s_0, s_1, \dots, s_n generate \mathcal{L} . So we get a morphism

$$g: Z \rightarrow \mathbb{P}_X^n,$$

over X , such that $g^*\mathcal{O}_{\mathbb{P}_X^n}(1) = \mathcal{L}$ and $g^{-1}x_i = s_i$. Suppose that $F(x_0, x_1, \dots, x_n)$ is a homogeneous polynomial in the kernel of ϕ . Then $F(a_0, a_1, \dots, a_n) = 0$ so that $F(s_0, s_1, \dots, s_n) = 0$ in $H^0(X, \mathcal{L}^d)$. It follows that g factors through Y .

Now suppose that $f: Z \rightarrow X$ factors through $g: Z \rightarrow Y$. Then

$$f^{-1}\mathcal{I} \cdot \mathcal{O}_Z = g^{-1}(\mathcal{I} \cdot \mathcal{O}_Y) \cdot \mathcal{O}_Z = g^{-1}\mathcal{O}_Y(1) \cdot \mathcal{O}_Z.$$

Therefore there is a surjective map

$$g^*\mathcal{O}_Y(1) \rightarrow \mathcal{L}.$$

But then this map must be an isomorphism and so $g^*\mathcal{O}_Y(1) = \mathcal{L}$. $s_i = g^*x_i$ and uniqueness follows. \square

Note that by the universal property, the morphism π is an isomorphism outside of the subscheme V defined by \mathcal{I} . We may put the universal property differently. The only subscheme with an invertible

ideal sheaf is a Cartier divisor (local generators of the ideal, give local equations for the Cartier divisor). So the blow up is the smallest morphism which turns a subscheme into a Cartier divisor. Perhaps surprisingly, therefore, blowing up a Weil divisor might give a non-trivial birational map.

If X is a variety it is not hard to see that π is a projective, birational morphism. In particular if X is quasi-projective or projective then so is Y . We note that there is a converse to this:

Theorem 8.7. *Let X be a quasi-projective variety and let $f: Z \rightarrow X$ be a birational projective morphism.*

Then there is an coherent ideal sheaf \mathcal{I} and a commutative diagram

$$\begin{array}{ccc} Z & \xrightarrow{\quad} & Y \\ & \searrow f & \swarrow \pi \\ & & X, \end{array}$$

where $\pi: Y \rightarrow X$ is the blow up of \mathcal{I} and the top row is an isomorphism.

It is interesting to figure out the geometry behind the example of a toric variety which is not projective. To warm up, suppose that we start with \mathbb{A}_k^3 . This is the toric variety associated to the fan spanned by e_1, e_2, e_3 . Imagine blowing up two of the axes. This corresponds to inserting two vectors, $e_1 + e_2$ and $e_1 + e_3$. However the order in which we blow up is significant. Let's introduce some notation. If we blow up the x -axis $\pi: Y \rightarrow X$ and then the y -axis, $\psi: Z \rightarrow Y$, let's call the exceptional divisors E_1 and E_2 , and let E'_1 denote the strict transform of E_1 on Z . E_1 is a \mathbb{P}^1 -bundle over the x -axis. The strict transform of the y -axis in Y intersects E_1 in a point p . When we blow up this curve, $E'_1 \rightarrow E_1$ blows up the point p . The fibre of E'_1 over the origin therefore consists of two copies Σ_1 and Σ_2 of \mathbb{P}^1 . Σ_1 is the strict transform of the fibre of E_1 over the origin and Σ_2 is the exceptional divisor. The fibre Σ of E_2 over the origin is a copy of \mathbb{P}^1 . Σ and Σ_2 are the same curve in Z .

The example of a toric variety which is not projective is obtained from \mathbb{P}^3 by blowing up three coordinate axes, which form a triangle. The twist is that we do something different at each of the three coordinate points. Suppose that $\pi: X \rightarrow \mathbb{P}^3$ is the birational morphism down to \mathbb{P}^3 , and let E_1, E_2 and E_3 be the three exceptional divisors. Over one point we extract E_1 first then E_2 , over the second point we extract first E_2 then E_3 and over the last point we extract first E_3 then E_1 .

To see what has gone wrong, we need to work in the homology and cohomology groups of X . Any curve C in X determines an element of $[C] \in H_2(X, \mathbb{Z})$. Any Cartier divisor D in X determines a class $[D] \in H^2(X, \mathbb{Z})$. We can pair these two classes to get an intersection number $D \cdot C \in \mathbb{Z}$. One way to compute this number is to consider the line bundle $\mathcal{L} = \mathcal{O}_X(D)$ associated to D . Then

$$D \cdot C = \deg \mathcal{L}|_C.$$

If D is ample then this intersection number is always positive. This implies that the class of every curve is non-trivial in homology.

Suppose the reducible fibres of E_1 , E_2 and E_3 over their images are $A_1 + A_2$, $B_1 + B_2$ and $C_1 + C_3$. Suppose that the general fibres are A , B and C . We suppose that A_1 is attached to B , B_1 is attached to C and C_1 is attached to A . We have

$$\begin{aligned} [A] &= [A_1] + [A_2] \\ &= [B] + [A_2] \\ &= [B_1] + [B_2] + [A_2] \\ &= [C] + [B_2] + [A_2] \\ &= [C_1] + [C_2] + [B_2] + [A_2] \\ &= [A] + [C_2] + [B_2] + [A_2], \end{aligned}$$

in $H_2(X, \mathbb{Z})$, so that

$$[A_2] + [B_2] + [C_2] = 0 \in H_2(X, \mathbb{Z}).$$

Suppose that D were an ample divisor on X . Then

$$0 = D \cdot ([A_2] + [B_2] + [C_2]) > D \cdot [A_2] + D \cdot [B_2] + D \cdot [C_2] > 0,$$

a contradiction.

There are a number of things to say about this way of looking at things, which lead in different directions. The first is that there is no particular reason to start with a triangle of curves. We could start with two conics intersecting transversally (so that they lie in different planes). We could even start with a nodal cubic, and just do something different over the two branches of the curve passing through the node. Neither of these examples are toric, of course. It is clear that in the first two examples, the morphism

$$\pi: X \longrightarrow \mathbb{P}^3,$$


is locally projective. It cannot be a projective morphism, since \mathbb{P}^3 is projective and the composition of projective is projective. It also follows that π is not the blow up of a coherent sheaf of ideals on \mathbb{P}^3 .

The third example is not even a variety. It is a complex manifold (and in fact it is something called an algebraic space). In particular the notion of the blow up in algebraic geometry is more delicate than it might first appear.

The second thing is to consider the difference between the order of blow ups of the two axes. Suppose we denote the composition of blowing up the x -axis and then the y -axis by $\pi_1: X_1 \rightarrow \mathbb{A}^3$ and the composition the other way by $\pi_2: X_2 \rightarrow \mathbb{A}^3$. Now X_1 and X_2 agree outside the origin. Let $\phi: X_1 \dashrightarrow X_2$ be the resulting birational map. If $A_1 + A_2$ is the fibre of π_1 over the origin and $B_1 + B_2$ is the fibre of π_2 over the origin, then ϕ is in fact an isomorphism outside A_2 and B_2 . So ϕ is a birational map which is an isomorphism in codimension one, in fact an isomorphism outside a curve, isomorphic to \mathbb{P}^1 . ϕ is an example of a *flop*. In terms of fans, we have four vectors v_1, v_2, v_3 and v_4 , such that

$$v_1 + v_3 = v_2 + v_4,$$

and any three vectors span the lattice. If σ is the cone spanned by these four vectors, then $Q = U_\sigma$ is the cone over a quadric. There are two ways to subdivide σ into two cones. Insert the edge connecting v_1 to v_3 or the edge corresponding to $v_2 + v_4$. The corresponding morphisms extract a copy of \mathbb{P}^1 and the resulting birational map between the two toric varieties is a (simple) flop. One can also insert the vector $w = v_1 + v_3$, to get a toric variety Y . The corresponding exceptional divisor is $\mathbb{P}^1 \times \mathbb{P}^1$. The toric varieties fit into a picture



The two maps $Y \rightarrow X_i$ correspond to the two projections of $\mathbb{P}^1 \times \mathbb{P}^1$ down to \mathbb{P}^1 . By (8.7) $\pi_i: X_i \rightarrow Q$ corresponds to blowing up a coherent ideal sheaf. In fact it corresponds to blowing up a Weil divisor (in fact this is given, as π_i does not extract any divisors), the plane determined by either ruling.

Finally, it is interesting to wonder more about the original examples of varieties which are not projective. Note that in the case when we blow up either a triangle or a conic if we make one flop then we get

a projective variety. Put differently, if we start with a projective variety then it is possible to get a non-projective variety by flopping a curve. When does flopping a curve mean that the variety is no longer projective? A variety is projective if it contains an ample divisor. Ample divisors intersect all curve positively. Note that any sum of ample divisors is ample.

Definition 8.8. *Let X be a proper variety. The **ample cone** is the cone in $H^2(X, \mathbb{R})$ spanned by the classes of the ample divisors.*

*The **Kleiman-Mori cone** $\overline{\text{NE}}(X)$ in $H_2(X, \mathbb{R})$ is the closure of the cone spanned by the classes of curves.*

The significance of all of this is the following:

Theorem 8.9 (Kleiman's Criteria). *Let X be a proper variety (or even algebraic space).*

A divisor D is ample if and only if the linear functional

$$\psi: H_2(X, \mathbb{R}) \longrightarrow \mathbb{R},$$

given by $\phi(\alpha) = [D] \cdot \alpha$ is strictly positive on $\overline{\text{NE}}(X) - \{0\}$.

Using Kleiman's criteria, it is not hard to show that if $\phi: X \dashrightarrow Y$ is a flop of the curve C and X is projective then Y is projective if and only if the class of $[C]$ generates a one dimensional face of $\overline{\text{NE}}(X)$.