

Lecture # 6

- Outline:
- 1) Review from last time
 - 2) Maps of S^1 to S^1
 - 3) Lifting closed curves to \mathbb{R}
 - 4) The degree of a map of S^1 to S^1
 - 5) Homotopy classes of curves
 - 6) Homotopy invariance of degree

Section 1 : Review

Definition :

• A closed curve in a surface Σ is a ^① continuous

^① map $\gamma : S^1 = \text{circle} \rightarrow \Sigma$.


① We send every pt in S^1 to a point in Σ .

② "Continuous" = we send points infinitesimally close


together in S^1 to points infinitesimally close together in Σ .

↔ We map S^1 into Σ w/ out ripping or cutting it


Example: 1)



2)



3)



Theorem :

Every compact orientable surface is homeomorphic to a connect sum $T^2 \# \dots \# T^2 \# S^2$ for some # of T^2 's.

Up Next :

- 1) Brouwer's Fixed Point Theorem
- 2) Fundamental Theorem of Algebra

Section 2: Maps of S^1 to S^1

Definition: • $S^1 = \{ (x, y) \text{ in } \mathbb{R}^2 \mid x^2 + y^2 = 1 \}$
= unit circle in the plane

Definition: A closed curve in S^1 = circle is a continuous
① map $\gamma: S^1 \rightarrow S^1$.

① We send every pt in S^1 to a point in S^1 .

② "Continuous" = we send points infinitesimally close together in S^1 to points infinitesimally close together in S^1 .

↳ We map S^1 into S^1 w/ out ripping or cutting it

Remark: Equivalently, a map $\gamma: S^1 \rightarrow S^1$ may be viewed as a continuous map

$$\gamma: [0, 2\pi] \rightarrow S^1$$

w/

$$\gamma(0) = \gamma(2\pi)$$

↳ i.e., a map of a circle is just a map of an interval that connects up at its end points.

↳ Intuitively, $\gamma: S^1 \rightarrow S^1$ gives a way of wrapping/laying a string onto a circle such that you can tie together its ends.

Example: 1) $\gamma_n: [0, 2\pi] \rightarrow S^1 \subseteq \mathbb{R}^2$ given by

$$\gamma_n(t) = (\cos(nt), \sin(nt))$$

↳ What is γ_0 ?

↳ What is γ_{-1} ?

↳ What is γ_n ?

* $\gamma_n(0) = \gamma_n(2\pi)$, ie, ends glue together

$$\gamma_n(0) = (1, 0) = (\cos(2\pi n), \sin(2\pi n)) = \gamma_n(2\pi).$$

2) Let $f: \mathbb{R} \rightarrow \mathbb{R}$ be any function st

$$f(0) = f(2\pi) + 2\pi \cdot n$$

for some n an integer.

$\gamma_f: [0, 2\pi] \rightarrow S^1$ given by

$$\gamma_f(t) = (\cos(f(t)), \sin(f(t)))$$

$$\hookrightarrow \text{Note } \gamma_f(0) = (\cos(f(0)), \sin(f(0)))$$



$$= (\cos(f(2\pi) + 2\pi n), \sin(f(2\pi) + 2\pi n))$$

$$= (\cos(f(2\pi)), \sin(f(2\pi)))$$



$$= \gamma_f(2\pi)$$

$\Rightarrow \gamma_f$ is a closed curve.

3) Crazier examples



Squash blue curve down onto circle to get actual map.



Section 3: Lifting closed curves to \mathbb{R}

Lemma: (Curve Lifting) Given a closed curve $\gamma: S^1 \rightarrow S^1$,

there exists a function $f: [0, 2\pi] \rightarrow \mathbb{R}$ st

1) $f(0) = f(2\pi) + 2\pi \cdot n$ for some integer n

2) $\gamma(t) = (\cos(f(t)), \sin(f(t)))$

$\hookrightarrow f$ is called a lift of γ to \mathbb{R} .

Remark:

1) f need not be unique.

2) If f is a lift of γ , then $f + 2\pi \cdot k$ is a lift of γ for every integer k .

3) These are the only other lifts of γ .

Proof :

1) Idea : "Unwind" the curve by noting the angle.

2) Notice that $\gamma(t) = (\cos(f(t)), \sin(f(t)))$

if and only if $f(t) =$ angle between $\gamma(t)$ and $(1,0)$

3) $f(t) =$ Accumulated angle of rotation of $\gamma(t)$

measured w/ respect to $(1,0)$

↳ rotate clockwise angle decreases

↳ rotate counter clockwise angle increases

↳ go around 5 times angle increases by

$$10\pi = 5 \cdot (2\pi).$$

4) By construction and (2), $\gamma(t) = (\cos(f(t)), \sin(f(t)))$

5) The only part in defining \tilde{f} where we have any choice is picking $\tilde{f}(0)$, that is, the starting angle from $(1, 0)$; any two choices differ by multiples of 2π .

So adding multiples of 2π to \tilde{f} gives all lifts

6) Notice that $\gamma(0) = \gamma(2\pi)$.

So total accumulated angle must be a multiple of 2π plus the starting angle.



So $\tilde{f}(0) = \tilde{f}(2\pi) + 2\pi \cdot n$ for some n . □

Example:

1) A lift of γ_n to \mathbb{R} is $\tilde{f}(t) = n \cdot t$.

2) A lift of γ_f to \mathbb{R} is \tilde{f} .



3)



$$\text{deg}(\gamma) = (f(2\pi) - f(0)) / 2\pi = (-2\pi - 0) / 2\pi = -1$$



4)

$$\deg(\gamma) = \left(\frac{5\pi}{2} - \frac{\pi}{2} \right) / 2\pi = 1$$



5)

$$\text{deg}(r) = 0.$$



Section 4: The degree of a map of S^1 to S^1

Definition: The degree of a closed curve $\gamma: S^1 \rightarrow S^1$ is

$$\deg(\gamma) = (f(2\pi) - f(0)) / 2\pi$$

where f is any lift of γ to \mathbb{R} .

Remark: 1) $\deg(\gamma)$ is independent of the choice of lift f .

↳ If g is another lift of γ , then

$$f = g + 2\pi \cdot k$$

$$\text{So } \frac{f(2\pi) - f(0)}{2\pi} = \frac{g(2\pi) - g(0)}{2\pi}.$$

Remark:

1) Intuitively, $\deg(\gamma) =$ signed # of times γ completely wraps around the circle

↳ signed: wraps clockwise = negative wrap

wraps counter clockwise = positive wrap

Example:

1) $\deg(\gamma_n) = n$

2) See above examples.

Section 5: Homotopy classes of curves

Definition: Two closed curves $\beta: S^1 \rightarrow S^1$ and $\gamma: S^1 \rightarrow S^1$ are homotopic if there is a continuous map $H: [0, 1] \times S^1 \rightarrow S^1$ satisfying

$$1) H(0, t) = \beta(t)$$

$$2) H(1, t) = \gamma(t)$$

Remark: Equivalently, $H: [0, 1] \times [0, 2\pi] \rightarrow S^1$ w/

$$1) H(0, t) = \beta(t)$$



$$2) H(1, t) = \gamma(t)$$

$$3) H(s, 0) = H(s, 2\pi)$$



Remark:

- 1) For each s_0 in $[0,1]$, $H(s_0, t)$ defines a closed curve in S' .
- 2) H parameterizes a family of curves that interpolates between β and γ .
- 3) Intuitively, H parameterizes how we can push, compress, deform the image of β in S' to the image of γ in S' .


Example :



* curve when
 s_0 is
 $H(s_0, t)$





Example 3




Remark 3

If we can rotate two curves to same curve, then they are homotopic.

Example 3



$\zeta = \zeta_{-1}$



Section 6: Homotopy invariance of degree

Theorem: Two closed curves $\beta: S^1 \rightarrow S^1$ and $\gamma: S^1 \rightarrow S^1$ are homotopic if and only if $\deg(\beta) = \deg(\gamma)$

Remark:

- 1) Notice that rotating the image of a curve in S^1 defines a continuous family of curves and thus a homotopy
 \Rightarrow Any curve is homotopic to a curve w/ $\gamma(0) = (1,0)$
- 2) If we rotate γ by θ , then the lift changes by $\frac{\theta}{2\pi} + \theta$
 $\Rightarrow \deg(\gamma) = \deg$ of rotated γ .

Claim 1:

If $\deg(\beta) = \deg(\gamma)$, then β is homotopic to γ .

Lemma:

Spse $\gamma(0) = (1, 0)$. If $\deg(\gamma) = n$, then γ is homotopic to $\gamma_n = (\cos(nt), \sin(nt))$.

Proof:

1) Let f be a lift of γ w/ $f(0) = 0$.

↳ we can do this since $\gamma(0) = (1, 0)$.

2) Define $\tilde{H}: [0, 1] \times [0, 2\pi] \rightarrow \mathbb{R}^2$ by

$$\tilde{H}(s, t) = (1-s) \cdot f(t) + s \cdot (n \cdot t)$$

3) Define $H: [0, 1] \times [0, 2\pi] \rightarrow S^1$ by

$$H(s, t) = (\cos(\tilde{H}(s, t)), \sin(\tilde{H}(s, t)))$$

4) We claim that H is a homotopy from γ to γ_n .

5) Check H glues up at ends

$$i) \quad \tilde{H}(s, 0) = (1-s) \cdot f(0) = 0$$

$$\begin{aligned} ii) \quad \tilde{H}(s, 2\pi) &= (1-s) \cdot f(2\pi n) + s \cdot 2\pi n \\ &= (1-s) \cdot 2\pi \deg(\gamma) + s \cdot 2\pi n \\ &= 2\pi n \end{aligned}$$

$$\begin{aligned} iii) \quad \Rightarrow H(s, 0) &= (\cos(0), \sin(0)) \\ &= (\cos(2\pi n), \sin(2\pi n)) \\ &= H(s, 2\pi) \end{aligned}$$

6) Check H is a homotopy from γ to γ_n

7) $\tilde{H}(0, t) = f(t)$

$$\Rightarrow H(0, t) = (\cos(f(t)), \sin(f(t))) = \gamma(t)$$

8) $\tilde{H}(1, t) = n \cdot t$

$$\Rightarrow H(1, t) = (\cos(nt), \sin(nt)) = \gamma_n(t)$$

* \cong = homotopic

Proof :

1) Spse $\deg(\gamma) = n = \deg(\beta)$

2) $n = \deg(\gamma) = \deg(\text{rotated } \gamma \text{ w/ starting pt } (1, 0))$

By lemma and fact that rotation is a homotopy,

$$\gamma \cong \text{rotated } \gamma \cong \gamma_n$$

3) Similarly, $\beta \cong \text{rotated } \beta \cong \gamma_n$

4) Chain together these homotopies to get $\beta \cong \gamma$.

Claim 2:

If β is homotopic to γ , then $\deg(\gamma) = \deg(\beta)$

Theorem:

(Homotopy Lifting) Let β and γ be closed curves in S^1 . Given a homotopy $H: [0,1] \times [0,2\pi] \rightarrow S^1$ w/ $H(0,t) = \beta(t)$ and $H(1,t) = \gamma(t)$.

There exists a continuous map

$$\tilde{H}: [0,1] \times [0,2\pi] \rightarrow \mathbb{R}$$

that satisfies

- 1) $H(s,t) = (\cos(\tilde{H}(s,t)), \sin(\tilde{H}(s,t)))$
- 2) For all s , $\tilde{H}(s,2\pi) - \tilde{H}(s,0) = 2\pi n$
- 3) \tilde{H} is unique up to adding a multiple of 2π .

Remark:

1) Prove the theorem by "unwrapping" in families.

$\tilde{H}(s,t)$ = lift of the curve $H(s,t)$ w/ s fixed.

↳ This essentially will imply (1), (3) in theorem

2) Since H parameterizes some family of curves, as we vary s , the accumulated amount of rotation can't jump (it is continuous).

3) For any lift, $f(2\pi) - f(0)$ is a multiple of 2π .

So $\tilde{H}(s, 2\pi) - \tilde{H}(s, 0)$ is a multiple of 2π for each s .

But it can't jump as s varies \Rightarrow must be constant.

Proof:

1) Let H be homotopy from β to γ .

2) Let \tilde{H} be a lift of H

3) $\tilde{H}(0, t)$ is a lift of β

$$\Leftrightarrow (\cos(\tilde{H}(0, t)), \sin(\tilde{H}(0, t))) = H(0, t) = \beta(t)$$

4) $\tilde{H}(1, t)$ is a lift of γ

$$\begin{aligned} 5) \text{ So } \deg(\beta) &= (\tilde{H}(0, 2\pi) - \tilde{H}(0, 0)) / 2\pi \\ &= (\tilde{H}(1, 2\pi) - \tilde{H}(1, 0)) / 2\pi \\ &= \deg(\gamma). \end{aligned}$$

□



