

# Complex Analysis and Riemann Surfaces

Note Title

9/2/2008

A graduate course taught by  
Prof. Duong H. Phong  
Columbia University, Fall 2008  
Notes taken by Qi You



## §1. Local Theory of Holomorphic Functions

Notation:  $f: \Omega \rightarrow \mathbb{C}$ ,  $\Omega$ : (connected) open domain in  $\mathbb{C}$

$$z = x + iy, \quad i^2 = -1$$

Def.  $f$ : holomorphic on  $\Omega \iff \forall z_0 \in \Omega, f'(z_0) \triangleq \lim_{h \rightarrow 0} \frac{f(z_0+h) - f(z_0)}{h}$  exists.

Example:  $f(z) = \bar{z}$ .

Although this function is linear, it's not holomorphic:

$z_0 = x_0 + iy_0$ ,  $h = h_1 + ih_2$ . Then  $\frac{f(z_0+h) - f(z_0)}{h} = \frac{x_0+h_1 - x_0}{h_1+ih_2} = \frac{h_1}{h_1+ih_2}$  has no limit as  $h \rightarrow 0$ .

### • Consequences of holomorphicity

Assume  $f$  holomorphic, i.e.  $f'(z) = \lim_{h \rightarrow 0} \frac{f(z+h) - f(z)}{h}$  exists,  $\forall z \in \Omega$ .

Take  $h = h_1 \rightarrow 0, \in \mathbb{R}$ , and view  $f(z) = f(x, y)$

$$\Rightarrow \lim_{h_1 \rightarrow 0} \frac{f(x+h_1, y) - f(x, y)}{h_1} = \frac{\partial f}{\partial x}(x, y)$$

On the other hand, take  $h = ih_2 \rightarrow 0, h_2 \in \mathbb{R}$ .

$$\Rightarrow \lim_{h_2 \rightarrow 0} \frac{f(x, y+h_2) - f(x, y)}{ih_2} = -i \frac{\partial f}{\partial y}(x, y)$$

$$\left. \begin{array}{l} \Rightarrow \frac{\partial f}{\partial x} = -i \frac{\partial f}{\partial y} \quad (= f'(z)) \end{array} \right\}$$

Moreover, if we write  $f = f(x, y) = u(x, y) + iv(x, y)$ ,  $u, v: \Omega \rightarrow \mathbb{R}$

$$\Rightarrow \frac{\partial u}{\partial x} = \frac{\partial v}{\partial y}, \quad \frac{\partial u}{\partial y} = -\frac{\partial v}{\partial x}$$

Notation:  $\frac{\partial f}{\partial \bar{z}} = \frac{1}{2} \left( \frac{\partial f}{\partial x} + i \frac{\partial f}{\partial y} \right)$ , then  $f$  holomorphic  $\Rightarrow \frac{\partial f}{\partial \bar{z}} = 0$ .

Thm. The following conditions are equivalent (in standard notations)

- (i)  $f$  holomorphic in  $\Omega$ .
- (ii)  $f$  is  $C^1$  in  $\Omega$  &  $\frac{\partial f}{\partial \bar{z}} \equiv 0$  in  $\Omega$
- (iii)  $f$  is  $C^1$  in  $\Omega$  &  $\forall$  region  $\bar{D} \subseteq \Omega$  with piecewise  $C^1$  boundary the line integral  $\oint_{\partial D} f(z) dz = 0$
- (iv)  $f$  is  $C^1$  in  $\Omega$  &  $\forall$  disc  $\bar{D}(z_0, r) \subseteq \Omega, z \in D(z_0, r)$ , we have 
$$f(z) = \frac{1}{2\pi i} \oint_{\partial D} \frac{f(w)}{w-z} dw \quad (\text{Cauchy integral formula})$$
- (v)  $\forall z_0 \in \Omega, \exists$  disc  $\bar{D}(z_0, r) \subseteq \Omega$  s.t.  $f(z) = \sum_{n=0}^{\infty} C_n (z-z_0)^n, \forall z \in D(z_0, r)$  (uniform convergence). In particular  $f \in C^{\omega}(\Omega) \subseteq C^{\infty}(\Omega)$ .

Observation:

Green's thm. in the plane: Let  $D$  be a region in  $\mathbb{R}^2$  with piecewise  $C^1$  boundary  $\partial D$ , then

$$\oint_{\partial D} P(x,y)dx + Q(x,y)dy = \iint_D \left( \frac{\partial Q}{\partial x} - \frac{\partial P}{\partial y} \right) dx dy$$

Apply this thm to the case of functions of a complex variable  $z$ ,  $f(z) = u + iv$

$$\oint_{\partial D} f(z) dz = \oint_{\partial D} (u+iv)(dx+idy) = \oint_{\partial D} u dx - v dy + i \oint_{\partial D} u dy + v dx$$

$$\stackrel{\text{Green}}{=} \iint_D \left( -\frac{\partial u}{\partial y} - \frac{\partial v}{\partial x} \right) dx dy + i \iint_D \left( \frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} \right) dx dy$$

$$= i \iint_D \left[ \left( \frac{\partial u}{\partial x} - \frac{\partial v}{\partial y} \right) + i \left( \frac{\partial u}{\partial y} + \frac{\partial v}{\partial x} \right) \right] dx dy$$

$$= 2i \iint_D \frac{\partial f}{\partial \bar{z}} dx dy$$

$$\Rightarrow \oint_{\partial D} f(z) dz = 0 \text{ if } f \text{ holomorphic } \left( \frac{\partial f}{\partial \bar{z}} \equiv 0 \right)$$

Proof of thm.

(ii)  $\Leftrightarrow$  (iii) :

" $\Rightarrow$ " is easy, since  $f \in C^1(\Omega)$ , we may apply Green's formula

" $\Leftarrow$ " We argue by contradiction. Assume  $\frac{\partial f}{\partial \bar{z}}(z_0) \neq 0$  for some  $z_0 \in \Omega$ .

Take an arbitrary disc  $D$  around  $z_0$ . By (iii)

$$0 = \left| \oint_{\partial D} f(z) dz \right| = \left| 2i \iint_D \frac{\partial f}{\partial \bar{z}} dx dy \right| = \left| 2i \iint_D \left( \frac{\partial f}{\partial \bar{z}}(z) - \frac{\partial f}{\partial \bar{z}}(z_0) \right) dx dy + 2i \frac{\partial f}{\partial \bar{z}}(z_0) \iint_D dx dy \right|$$

$$\geq 2 \left| \frac{\partial f}{\partial \bar{z}}(z_0) \right| \iint_D dx dy - 2 \left| \iint_D \left( \frac{\partial f}{\partial \bar{z}}(z) - \frac{\partial f}{\partial \bar{z}}(z_0) \right) dx dy \right|$$

$$\geq 2 \left| \frac{\partial f}{\partial \bar{z}}(z_0) \right| \text{Area}(D) - 2 \iint_D \left| \frac{\partial f}{\partial \bar{z}}(z) - \frac{\partial f}{\partial \bar{z}}(z_0) \right| dx dy$$

Since  $f$  is  $C^1$ ,  $\exists \delta > 0$  s.t.  $|z - z_0| < \delta \Rightarrow \left| \frac{\partial f}{\partial \bar{z}}(z) - \frac{\partial f}{\partial \bar{z}}(z_0) \right| < \frac{1}{2} \left| \frac{\partial f}{\partial \bar{z}}(z_0) \right|$

$$\Rightarrow 0 \geq \left| \frac{\partial f}{\partial \bar{z}}(z_0) \right| \pi \delta^2 > 0, \text{ contradiction.}$$

(iii)  $\Rightarrow$  (iv)

We fix  $z \in \Omega$ , and set  $g(w) = \frac{f(w) - f(z)}{w - z}$ ,  $w \in \Omega \setminus \{z\}$

Claim:  $\frac{\partial}{\partial \bar{w}} g(w) \equiv 0$  on  $\Omega \setminus \{z\}$ .

This is because  $\frac{\partial}{\partial \bar{w}} g(w) = \frac{\frac{\partial}{\partial \bar{w}}(f(w) - f(z))(w - z) - (f(w) - f(z)) \frac{\partial}{\partial \bar{w}}(w - z)}{(w - z)^2} = 0$  since  $f(w) - f(z)$ ,  $w - z$  are holomorphic.

Moreover,  $g$  is  $C^1$  on  $\Omega \setminus \{z\}$  since  $f(w) - f(z)$  and  $w - z$  are.

$\Rightarrow g$  satisfies (ii) on  $\Omega \setminus \{z\}$

$\Rightarrow g$  satisfies (iii) on  $\Omega \setminus \{z\}$

Apply (iii) to  $\varepsilon < |w - z| < \delta$  :



$$\Rightarrow 0 = \oint_{\partial\Omega} g(\omega) d\omega = \oint_{|\omega-z_1|=\delta} g(\omega) d\omega - \oint_{|\omega-z_1|=\varepsilon} g(\omega) d\omega \quad (*)$$

Claim:  $\oint_{|\omega-z_1|=\varepsilon} g(\omega) d\omega \rightarrow 0$  as  $\varepsilon \rightarrow 0$

This is because by Taylor's formula: (for  $C^1$ )

$$f(\omega) = f(z) + \frac{\partial f}{\partial \bar{\omega}}(\omega-z) - \frac{\partial f}{\partial \bar{\omega}}(\omega-z) + o(|\omega-z|)$$

$$\Rightarrow g(\omega) = \frac{\partial f}{\partial \bar{\omega}}(\omega-z) + o(|\omega-z|)$$

$\Rightarrow g$  can be extended as a continuous function on  $\Omega$ , thus bounded on  $|\omega-z_1| \leq \varepsilon$

$$\Rightarrow \left| \oint_{|\omega-z_1|=\varepsilon} g(\omega) d\omega \right| \leq \oint_{|\omega-z_1|=\varepsilon} \sup_{|\omega-z_1| \leq \varepsilon} |g(\omega)| = 2\pi\varepsilon \cdot \sup_{|\omega-z_1| \leq \varepsilon} |g(\omega)| \rightarrow 0 \text{ as } \varepsilon \rightarrow 0,$$

and we know that it's independent of  $\varepsilon$  by  $(*)$ .

$$\begin{aligned} \text{Thus } 0 &= \oint_{|\omega-z_1|=\delta} \frac{f(\omega)-f(z)}{\omega-z} d\omega = \oint_{|\omega-z_1|=\delta} \frac{f(\omega)}{\omega-z} d\omega - f(z) \oint_{|\omega-z_1|=\delta} \frac{d\omega}{\omega-z} \\ &= \oint_{|\omega-z_1|=\delta} \frac{f(\omega)}{\omega-z} d\omega - f(z) \int_{|\zeta|=\delta} \frac{d\zeta}{\zeta}, \text{ let } \zeta = \delta e^{i\theta} \\ &= \oint_{|\omega-z_1|=\delta} \frac{f(\omega)}{\omega-z} d\omega - f(z) \int_0^{2\pi} \frac{i\delta e^{i\theta}}{\delta e^{i\theta}} d\theta \\ &= \oint_{|\omega-z_1|=\delta} \frac{f(\omega)}{\omega-z} d\omega - 2\pi i f(z) \end{aligned}$$


$$\Rightarrow f(z) = \frac{1}{2\pi i} \oint_{|\omega-z_1|=\delta} \frac{f(\omega)}{\omega-z} d\omega$$

This is the case that  $z = z_0$  in (iv). More generally, for any point  $z$  in  $|z-z_0| < \delta$ , consider the integral of  $\frac{f(\omega)}{\omega-z}$  on the boundary of the region bounded by  $|\omega-z_0| < \delta$  and  $|\omega-z_1| < \varepsilon$ ,  $\varepsilon$  small enough so that it's contained in  $|\omega-z_0| < \delta$

we have:


$$\frac{1}{2\pi i} \oint_{|\omega-z_0|=\delta} \frac{f(\omega)}{\omega-z} d\omega - \frac{1}{2\pi i} \oint_{|\omega-z_1|=\varepsilon} \frac{f(\omega)}{\omega-z} d\omega = 0$$

$$\Rightarrow f(z) = \frac{1}{2\pi i} \oint_{|\omega-z_1|=\varepsilon} \frac{f(\omega)}{\omega-z} d\omega = \frac{1}{2\pi i} \oint_{|\omega-z_0|=\delta} \frac{f(\omega)}{\omega-z} d\omega$$



(iv)  $\Rightarrow$  (v)

$$\begin{aligned} \text{Now } f(z) &= \frac{1}{2\pi i} \oint_{|\omega-z_1|=\delta} \frac{f(\omega)}{(\omega-z_0) - (z-z_0)} d\omega \\ &= \frac{1}{2\pi i} \oint_{|\omega-z_1|=\delta} \frac{f(\omega)}{\omega-z_0} \frac{1}{1 - \frac{z-z_0}{\omega-z_0}} d\omega \\ &= \frac{1}{2\pi i} \oint_{|\omega-z_1|=\delta} \frac{f(\omega)}{\omega-z_0} \sum_{n=0}^{\infty} \left(\frac{z-z_0}{\omega-z_0}\right)^n d\omega \\ &= \frac{1}{2\pi i} \sum_{n=0}^{\infty} \left( \oint_{|\omega-z_1|=\delta} \frac{f(\omega)}{(\omega-z_0)^{n+1}} d\omega \right) (z-z_0)^n \end{aligned}$$



$$\left| \frac{z-z_0}{\omega-z_0} \right| = \frac{r}{\delta} < 1$$

Here the equalities are valid since  $\left| \frac{z-z_0}{\omega-z_0} \right| < 1$  and the convergence is uniform.

(v)  $\Rightarrow$  (i) trivial.


(i)  $\Rightarrow$  (ii) (Harder!)

We already know that  $\frac{\partial f}{\partial \bar{z}} = 0$  by analysis before the theorem. Thus our primary step would be to show that  $f$  holo.  $\Rightarrow f \in C^1$  (since we now don't know a priori that  $f$  is  $C^1$ , we can NOT apply Green's thm. to prove (i)  $\Rightarrow$  (iii))

It suffices to show that for any disc  $D \subseteq \Omega$ , there exists a  $C^1$  function  $F$  s.t.  $F'(z) = f(z)$  and  $\frac{\partial F}{\partial \bar{z}} = 0$ . Then  $F$  satisfies (ii) thus (v) by the previous proof. Then  $F$  would be  $C^\infty$  and so is  $f(z) = F'(z) \in C^\infty$ , in particular  $f(z)$  is  $C^1$ .

In the disc  $D$ , let  $F(z) = \int_{L_z} f(w) dw$ , where  $L_z$  is the line segment joining  $z_0$  and  $z$ . We compute  $F'(z) = \lim_{h \rightarrow 0} \frac{F(z+h) - F(z)}{h}$

$$\begin{aligned} &= \lim_{h \rightarrow 0} \frac{1}{h} (\int_{L_{z+h}} f(\cdot) - \int_{L_z} f(\cdot)) \\ \text{Claim} &= \lim_{h \rightarrow 0} \frac{1}{h} (\int_{L_z^{z+h}} f(\cdot)) \\ &= \lim_{h \rightarrow 0} \frac{1}{h} \int_0^1 f(z+th) h dt \\ &= f(z). \end{aligned}$$



Proof of the Claim:

The claim  $\Leftrightarrow \oint_{\text{Triangle}} f(z) dz = 0$

Let  $I = \oint_T f(z) dz$  and subdivide the triangle into the four equal pieces as in the picture on the right.

$$I = \sum_{i=1}^4 \oint_{T_i} f(z) dz$$

We can now pick one  $T_i$  ( $i=1, \dots, 4$ ) s.t. (say,  $T_1$ )

$$|\oint_{T_1} f(z) dz| \geq \frac{I}{4}$$

We further subdivide  $T_1$  into 4 equal pieces  $T_2^1, \dots, T_4^1$ , and pick  $T_2^1$  s.t.

$$|\oint_{T_2^1} f(z) dz| \geq \frac{1}{4} |\oint_{T_1} f(z) dz| \geq \frac{I}{4^2}$$

----- (iterate this process)



We obtain a sequence of "nested triangles":

$$T \supseteq T_1^1 \supseteq T_2^1 \supseteq \dots$$

$$\text{with } |\oint_{T_k^1} f(z) dz| \geq \frac{I}{4^k}$$

Moreover  $\bigcap_{k=1}^{\infty} T_k^1 = \{z_0\}$ . Since  $f$  is holomorphic at  $z_0$ ,  $f(z) = f(z_0) + f'(z_0)(z-z_0) + o(|z-z_0|)$  in a sufficiently small neighborhood of  $z_0$ . When  $k$  is large enough,  $T_k^1$  will be contained in this neighborhood. Thus

$$|\oint_{\partial T_k^1} f(z) dz| = |\oint_{\partial T_k^1} (f(z_0) + f'(z_0)(z-z_0) + o(|z-z_0|)) dz|$$




$$\begin{aligned}
&\leq \left| \oint_{\partial T_k^i} (f(z_0) + f'(z_0)(z-z_0)) dz \right| + \left| \oint_{\partial T_k^i} O(|z-z_0|) dz \right| \\
&\leq \oint_{\partial T_k^i} O(|z-z_0|) |dz| \\
&\leq 3 \cdot \text{diam } T_k^i \cdot \varepsilon \text{ diam } T_k^i \\
&= 3\varepsilon \cdot \frac{\text{diam } T}{2^k} \cdot \frac{\text{diam } T}{2^k} \\
&= 3 \cdot \frac{(\text{diam } T)^2}{4^k} \varepsilon \quad \text{where } \varepsilon \text{ is arbitrarily small, since } \lim_{z \rightarrow z_0} \frac{O(|z-z_0|)}{|z-z_0|} \rightarrow 0 \\
\Rightarrow \frac{I}{4^k} &\leq 3\varepsilon \frac{(\text{diam } T)^2}{4^k} \\
\Rightarrow I &\leq 3(\text{diam } T)^2 \cdot \varepsilon
\end{aligned}$$

Thus  $I = 0$  since  $\varepsilon$  is arbitrarily small. □

Examples of holomorphic functions

- $e^z \triangleq \sum_{n=0}^{\infty} \frac{z^n}{n!}$  converges for all  $z$ .  $\Rightarrow e^z$  is holomorphic on  $\mathbb{C}$  and  $\frac{d}{dz} e^z = e^z$ . and  $e^{z+w} = e^z e^w$
- We want to define  $\log z$  as the inverse of  $e^z$ , i.e.  $e^{\log z} = z$   
Set  $\log z = u + iv$   $z = e^{\log z} = e^u e^{iv} \Rightarrow u = |z|, v = \text{Arg } z$ , but until we introduce a cut on  $\mathbb{C}$ ,  $\text{Arg } z$  can't be a well-defined function on  $\mathbb{C}$ .



- Open mapping and maximal modulus theorems

Take  $f$  holomorphic in  $\Omega$

- 1).  $f \neq 0 \Rightarrow$  the zero's of  $f$  are isolated, i.e.  $f(z_0) = 0 \Rightarrow \exists V$ , neighborhood of  $z_0$  s.t.  $\forall z \in V \setminus \{z_0\}, f(z) \neq 0$

This is because, we can expand  $f$  as a power series near  $z_0$

$f(z) = \sum_{n=0}^{\infty} a_n (z-z_0)^n$  in a nhd of  $z_0$ .  $f \neq 0 \Rightarrow \exists n, a_n \neq 0$ . Take the smallest such  $n$ , say,  $a_n \neq 0$ .  $\Rightarrow f(z) = (z-z_0)^n (a_n + a_{n+1}(z-z_0) + \dots) = (z-z_0)^n g(z)$ , where  $g(z_0) = a_n \neq 0$ . Thus in a small enough neighborhood  $g(z) \neq 0$ .

Def: The order of vanishing of  $f$  at  $z_0 \triangleq N$ .

2). The class of meromorphic functions in  $\Omega$ .

Def:  $g$  meromorphic in  $\Omega \iff \forall z_0 \in \Omega, g(z_0) = \sum_{n=N}^{\infty} a_n (z-z_0)^n, N \in \mathbb{Z}$ .

Meromorphic functions satisfy the following version of Cauchy integral formula.

$$\frac{1}{2\pi i} \oint_{\partial D} g(z) dz = \sum_{z_i} \text{Res}(g)(z_i),$$

where  $z_i$  are poles of  $g$  inside  $D$  and there are no poles on  $\partial D$ . (The poles inside  $D$  are isolated thus finite, since  $\bar{D}$  compact). Here if  $z_i$  is a pole

$$g(z) = \sum_{n=N}^{\infty} a_n (z-z_i)^n \quad \text{Res}(g)(z_i) \triangleq a_{-1}$$

Pf: Consider  $g(z)$  as a holomorphic function on  $V \setminus \{z_1, \dots, z_N\}$ , where  $V$  is a neighborhood of  $\bar{D}$

$$\Rightarrow 0 = \oint_{\partial(D \setminus \bigcup_{i=1}^N \{z-z_i \leq \varepsilon\})} g(z) dz = \oint_{\partial D} g(z) dz - \sum_{i=1}^N \oint_{|z-z_i|=\varepsilon} g(z) dz$$

$$\Rightarrow \oint_{\partial D} g(z) dz = \sum_{i=1}^N \oint_{|z-z_i|=\varepsilon} g(z) dz$$

In each  $|z-z_i| \leq \varepsilon, g(z) = \sum_{n=N_i}^{\infty} a_n (z-z_i)^n$

$$\Rightarrow \oint_{|z-z_i|=\varepsilon} g(z) dz = \sum_{n=N_i}^{\infty} a_n \oint_{|z-z_i|=\varepsilon} (z-z_i)^n dz$$

$$\text{Now } \oint_{|z-z_i|=\varepsilon} (z-z_i)^n dz = \begin{cases} 0 & n \neq -1 \\ 2\pi i & n = -1 \end{cases}$$

$$\Rightarrow \oint_{\partial D} g(z) dz = \sum_{i=1}^N 2\pi i \text{Res}(g)(z_i) \quad \square$$

3). Simple consequences

$g(z)$  meromorphic in  $\Omega \Rightarrow g'(z)$  is meromorphic with poles at the poles of  $g(z)$

$\Rightarrow \frac{g'(z)}{g(z)}$  is meromorphic with poles at the poles and zeroes of  $g(z)$ ; and the residue of  $\frac{g'(z)}{g(z)}$  at a zero of  $g(z)$  is  $+1$  (to be counted with multiplicity), while is  $-1$  (to be counted with multiplicity) at a pole of  $g(z)$ .

Pf: Near a zero or pole of  $g(z)$ ,  $g(z) = \sum_{n=N_0}^{\infty} a_n (z-z_0)^n$  ( $a_{N_0} \neq 0$ )

$$\Rightarrow g(z) = (z-z_0)^{N_0} \sum_{n=N_0}^{\infty} a_n (z-z_0)^{n-N_0} = (z-z_0)^{N_0} u(z), u(z_0) \neq 0.$$

$$\Rightarrow \frac{g'(z)}{g(z)} = \frac{N_0 (z-z_0)^{N_0-1} u(z) + (z-z_0)^{N_0} u'(z)}{(z-z_0)^{N_0} u(z)} = \frac{N_0}{z-z_0} + \frac{u'(z)}{u(z)}$$

We have a pole of  $\frac{g'(z)}{g(z)}$  if  $N_0 \neq 0$ :

$$\begin{cases} N_0 > 0, & g(z) \text{ has a zero of order } N_0 \text{ at } z_0 \\ N_0 < 0, & g(z) \text{ has a pole of order } -N_0 \text{ at } z_0 \end{cases} \quad \square$$


In conclusion:

$$\frac{1}{2\pi i} \oint_{\partial D} \frac{g'(z)}{g(z)} dz = \sum_{\text{poles}} \text{Res}\left(\frac{g'(z)}{g(z)}\right)(z_i) = \sum_{\substack{\text{zeros} \\ \text{of } g}} 1 - \sum_{\substack{\text{poles} \\ \text{of } g}} 1$$



(Open Mapping Thm.)  $f: \Omega \rightarrow \mathbb{C}$  holomorphic, not constant. Assume  $f$  has order  $n > 0$  at  $z_0$  (i.e.  $f(z) = (z - z_0)^n u(z)$ ,  $u(z_0) \neq 0$ )  $\Rightarrow \exists U$  neighborhood of  $z_0$  and  $V$  neighborhood of  $0$  s.t.  $\forall v \in V, \exists$  exactly  $n$  points of  $z_1, \dots, z_n \in U$  s.t.  $f(z_1) = \dots = f(z_n) = v$

Pf: Since the zero's of  $f$  and  $f'$  are isolated, we may assume that  $S = \partial D$  is a small circle around  $z_0$  such that  $f(z) \neq 0, f'(z) \neq 0$  for  $z \in \bar{D} \setminus \{z_0\}$ .



Consider the following function of  $w$ :  $w \mapsto \frac{1}{2\pi i} \oint_S \frac{f'(z)}{f(z) - w} dz$

The R.H.S. = integer valued continuous function on  $\mathbb{C} \setminus f(S)$ , thus locally constant.

Let  $V$  be the connected component of  $\mathbb{C} \setminus f(S)$  containing  $0$ .

$\Rightarrow \forall w \in V \quad \frac{1}{2\pi i} \oint_S \frac{f'(z)}{f(z) - w} dz = n = \# \{ \text{zeros of } f(z) - w \text{ in } D \}$

Thus  $f(z) = w$  has exactly  $n$ -zeros for  $w \in V$  in  $f^{-1}(V) \cap D \cong U$ . The zero's are distinct since we have assumed that  $\forall z \neq z_0, f'(z) \neq 0$  □

Cor. If  $f$  is holomorphic and not a constant  $\Rightarrow$  the image of  $f$  is open.

Pf:  $\forall w \in \text{Im} f$ . Take  $V$  for  $f(z) - w$  as in the thm.  $V \subseteq \text{Im} f$ . □

Cor. (Maximum Modulus Principle)  $f$  holomorphic on  $\Omega$ . If  $\exists z_0 \in \Omega$  s.t.  $|f(z_0)| \geq |f(z)|, \forall z \in \Omega$ , then  $f$  is a constant.

Pf:



If  $f$  were not constant, then  $f(z_0)$  would be in an open set of images of  $f \Rightarrow$  there would be  $z' \in \Omega$  with  $|f(z')| > |f(z_0)|$ , contradiction. □

• Applications: method of residues

1). Calculate  $\int_0^{+\infty} \frac{1}{1+x^2} dx$  using residue formula.

$I = \frac{1}{2} \int_{-\infty}^{+\infty} \frac{1}{1+z^2} dz = \frac{1}{2} \lim_{R \rightarrow \infty} \int_{-R}^R \frac{1}{1+z^2} dz$ . Construct an integral contour as follows:

$$I = \frac{1}{2} \lim_{R \rightarrow \infty} \left( \int_{\Delta} \frac{dz}{1+z^2} - \int_{\Gamma} \frac{dz}{1+z^2} \right)$$


By the residue thm  $\int_{\Delta} \frac{dz}{1+z^2} = 2\pi i \operatorname{Res}_{(1+z^2)}(i)$ ,

$$\text{and } \frac{1}{1+z^2} = \frac{1}{z+i} \frac{1}{z-i} \Rightarrow \operatorname{Res}_{(1+z^2)}(i) = \frac{1}{2i}$$

$$\Rightarrow \int_{\Delta} \frac{dz}{1+z^2} = \pi$$

$$\text{Moreover } \left| \int_{\Gamma} \frac{dz}{1+z^2} \right| \leq \int_0^{\pi} \frac{|dz|}{|z^2-1|} = \int_0^{\pi} \frac{R d\theta}{R^2-1} = \frac{\pi R}{R^2-1} \rightarrow 0 \text{ as } R \rightarrow \infty.$$

$$\Rightarrow I = \frac{\pi}{2}.$$



2).  $\int_0^{\infty} \frac{\sin x}{x} dx$

First of all, the integral converges:  $x \rightarrow 0, \frac{\sin x}{x} \rightarrow 1$ , and the function is smooth (analytic)

thus integrable near 0;  $\left| \int_1^{\infty} \frac{\sin x}{x} dx \right| = \left| \frac{-\cos x}{x} \right|_1^{\infty} - \int_1^{\infty} \frac{\cos x}{x^2} dx \leq 1 + \int_0^{\infty} \frac{1}{x^2} dx$ , thus is

integrable near  $\infty$ .

We want to apply the method of residue. Consider  $e^{iz} = \cos z + i \sin z$

$$\int_{-\infty}^{\infty} \frac{\sin x}{x} dx = \operatorname{Im} \left( \int_{-\infty}^{\infty} \frac{e^{iz}}{z} dz \right)$$

To calculate  $\int_{-\infty}^{\infty} \frac{e^{iz}}{z} dz$  we take the following contour.

$$\oint \frac{e^{iz}}{z} dz = \operatorname{Res}_{(1/z)}(0) = 2\pi i$$

On I:  $\frac{e^{iz}}{z} = \frac{1}{z} + u(z)$  where  $u(z)$  is holomorphic

$$\int_I \frac{e^{iz}}{z} dz = \int_{\pi}^{2\pi} \frac{1}{z} dz + \int_{\pi}^{2\pi} u(r, \theta) i r e^{i\theta} d\theta$$

$$= \pi i + O(r) \rightarrow \pi i \quad (r \rightarrow 0)$$


$$\text{II and IV: } \left| \int_{\text{II}} \frac{e^{iz}}{z} dz \right| = \left| \int_0^R \frac{e^{i(R+iy)}}{R+iy} dy \right| = \left| \int_0^R \frac{e^{iR} e^{-y}}{R+iy} dy \right| \leq \int_0^R \left| \frac{e^{iR} e^{-y}}{R+iy} \right| dy$$

$$\leq \int_0^R \frac{e^{-y}}{R} dy \leq \frac{1}{R} \int_0^{\infty} e^{-y} dy \rightarrow 0 \quad (R \rightarrow \infty)$$

$$\text{III: } \left| \int_{\text{III}} \frac{e^{iz}}{z} dz \right| = \left| \int_R^{-R} \frac{e^{i(x-R)}}{x+iR} dx \right| \leq \int_R^{-R} \left| \frac{e^{i(x-R)}}{x+iR} \right| dx \leq e^{-R} \int_R^{-R} \frac{1}{R} dx = 2e^{-R} \rightarrow 0 \quad (R \rightarrow \infty)$$

Thus it follows that  $2\pi i = \int_{-\infty}^{\infty} \frac{e^{iz}}{z} dz + \pi i + 0 + 0$

$$\Rightarrow \int_0^{\infty} \frac{\sin x}{x} dx = \frac{1}{2} \int_{-\infty}^{\infty} \frac{\sin x}{x} dx = \frac{\pi}{2}.$$




3). Evaluate  $\int_0^{\infty} \frac{v^{z-1}}{1+v} dv$  ( $0 < z < 1$ , which ensures the convergence)

$v^{z-1} = e^{(z-1)\ln v}$ , but since "ln" function can only be well-defined after we introduce

a cut on the plane  $\mathbb{C}$ , we make the following contour on  $\mathbb{C} \setminus \{x \geq 0, y=0\}$

We pick the ordinary "ln" function on the upper half of the positive axis.

Now the function  $f(v) = \frac{v^{z-1}}{1+v}$  is holomorphic on  $\mathbb{C} \setminus \{-1\}$  and in particular has a simple pole inside our contour. Thus by the residue thm.



$$\oint f(v) dv = 2\pi i \operatorname{Res}(f, -1) = 2\pi i (e^{i\pi})^{z-1} = 2\pi i e^{i\pi(z-1)}$$

Moreover,  $|\int_{\text{I}} f(v) dv| = \left| \int_{2\pi}^0 \frac{(re^{i\theta})^{z-1}}{1+re^{i\theta}} r i e^{i\theta} d\theta \right| \leq \int_0^{2\pi} \frac{r^z}{1-r} dr = \frac{2\pi r^z}{1-r} \rightarrow 0 \quad (r \rightarrow 0)$

$|\int_{\text{III}} f(v) dv| = \left| \int_0^{2\pi} \frac{R^{z-1} e^{i(z-1)\theta}}{1+Re^{i\theta}} R i e^{i\theta} d\theta \right| \leq \int_0^{2\pi} \frac{R^z}{R-1} d\theta = \frac{2\pi R^z}{R-1} \rightarrow 0 \quad (R \rightarrow \infty)$

Now  $\int_{\text{II+IV}} f(v) dv = \int_0^R \frac{v^{z-1}}{1+v} dv - \int_0^R \frac{v^{z-1} e^{2\pi i(z-1)}}{1+v} dv$

$$= (1 - e^{2\pi i(z-1)}) \int_0^R \frac{v^{z-1}}{1+v} dv$$

$$\Rightarrow (1 - e^{2\pi i(z-1)}) \int_0^\infty \frac{v^{z-1}}{1+v} dv + 0 + 0 = 2\pi i e^{i\pi(z-1)}$$

$$\Rightarrow \int_0^\infty \frac{v^{z-1}}{1+v} dv = \frac{2\pi i e^{i\pi(z-1)}}{1 - e^{2\pi i(z-1)}} = \frac{\pi}{\sin \pi z}$$

□

### • Analytic Continuation

Basic Observation:

Let  $\Omega$  be a domain (connected),  $f$  a holomorphic function. If  $f=0$  on a non-empty open set  $\Omega' \subseteq \Omega$ , then  $f \equiv 0$  on  $\Omega$ .

Pf: Let  $\tilde{\Omega} = \{z_0 \in \Omega \mid f=0 \text{ in a neighborhood of } z_0\}$

(i).  $\tilde{\Omega} \supseteq \Omega'$  thus  $\neq \emptyset$ .

(ii).  $\tilde{\Omega}$  is open from definition


(iii).  $\tilde{\Omega}$  is closed

}  $\Rightarrow \tilde{\Omega} = \Omega$  since  $\Omega$  is connected.

•  $\forall z_1 \in \tilde{\Omega}^c$ , i.e.  $f$  is not identically 0 around  $z_1$ . Write  $f(z) = \sum_{n=0}^\infty C_n (z-z_1)^n$ , then some  $C_N$  must be non-zero. Take minimal such  $\Rightarrow f(z) = (z-z_1)^N \hat{f}(z)$  with  $\hat{f}(z_1) \neq 0$ .

Thus  $f(z) \neq 0$  on an open neighborhood where  $\hat{f}(z) \neq 0$ . □

A Basic Question: Given  $\Omega \subseteq \mathbb{C}$  and  $f$  holomorphic on  $\Omega$ . What is the largest  $\Omega'$  s.t.  $\exists g(z)$  holomorphic on  $\Omega'$  s.t.  $g|_\Omega = f$ . (Note that for smooth functions smooth extensions need not be unique:



$$y = e^{-\frac{1}{x^2}}$$

We may extend  $y$  by 0 on  $e^{-\frac{1}{x^2}}$  itself!

Model cases:

- Let  $\varphi \in C^\infty[0,1]$  and consider  $f(z) = \int_0^1 x^z \varphi(x) dx$ .

The integral converges for  $\operatorname{Re} z > -1$  and defines a holomorphic function on  $\{ \operatorname{Re} z > -1 \}$

Claim:  $f(z)$  admits a meromorphic extension with possibly poles at the negative integers.

Pf: Let  $N$  be any positive integer, we will write  $\varphi(x)$  in terms of its Taylor expansion at 0.  $\varphi(x) = \sum_{n=0}^N \frac{1}{n!} \varphi^{(n)}(0) x^n + E_N(x)$  where  $|E_N(x)| \leq C x^{N+1}$

Let  $\operatorname{Re} z > -1$ , then:


$$f(z) = \int_0^1 x^z \varphi(x) dx = \int_0^1 \sum_{n=0}^N \frac{1}{n!} \varphi^{(n)}(0) x^{n+z} dx + \int_0^1 E_N(x) x^z dx$$

Now, since  $|E_N(x)| \leq C x^{N+1}$   $\int_0^1 E_N(x) x^z dx$  converges on  $\operatorname{Re}(z+N+1) \geq -1$

or equivalently  $\operatorname{Re} z > -N-2$  and defines a holomorphic function there. Moreover

$$\int_0^1 \sum_{n=0}^N \frac{1}{n!} \varphi^{(n)}(0) x^{n+z} dx = \sum_{n=0}^N \frac{\varphi^{(n)}(0)}{n!} \frac{1}{n+1+z}$$

which is a meromorphic function with possibly poles at  $\{-1, -2, \dots, -N-1\}$ .



$$f(z) = \underbrace{\text{mero. fun}}_{\text{globally defined}} + \underbrace{\text{holo}}_{\text{defined only on } \operatorname{Re} z \geq -N-1}$$

This works for arbitrary  $N \in \mathbb{N} \Rightarrow f$  extends to a meromorphic function on  $\mathbb{C}$ , with residue at  $-N-1$  equal to  $\frac{\varphi^{(N)}(0)}{N!}$ .  $\square$

- The  $\Gamma(z)$ -function

$$\Gamma(z) = \int_0^\infty e^{-x} x^{z-1} dx \quad (\operatorname{Re} z > -1)$$

Claim:  $\Gamma(z)$  extends to a meromorphic function on  $\mathbb{C}$ , with simple poles in  $\{-n \mid n \in \mathbb{N}\}$ .

Pf:  $\Gamma(z) = \int_0^1 e^{-x} x^{z-1} dx + \int_1^\infty e^{-x} x^{z-1} dx$ , and  $\int_1^\infty e^{-x} x^{z-1} dx$  is well-defined and holomorphic in the whole plane  $\mathbb{C}$ .

Thus we may apply the previous result.  $\square$

- The  $\zeta(z)$  (Riemann zeta) function.

$$\zeta(s) \triangleq \sum_{n=1}^{\infty} \frac{1}{n^s}, \text{ well-defined for } \operatorname{Re}(s) > 1$$

Observe that  $\frac{1}{n^s}$  can be written as  $\int_0^{\infty} e^{-nt} t^{s-1} dt = \Gamma(s) n^{-s}$

$$\Rightarrow \zeta(s) = \sum_{n=1}^{\infty} \frac{1}{n^s} = \frac{1}{\Gamma(s)} \int_0^{\infty} \sum_{n=1}^{\infty} e^{-nt} t^{s-1} dt = \frac{1}{\Gamma(s)} \int_0^{\infty} \frac{1}{e^t - 1} t^{s-1} dt$$

Similar as above,  $\int_0^{\infty} \frac{1}{e^t - 1} t^{s-1} dt$  admits an extension to a meromorphic function on  $\mathbb{C}$ . (In fact holomorphic on  $\mathbb{C} \setminus \{1\}$ , with a simple pole at 1, since  $\Gamma(s)$  also has simple poles at  $-n$ .)

Ex. Compute to show that  $\zeta(0) = -\frac{1}{12}$ ,  $\zeta'(0) = -\frac{1}{2} \log 2\pi$

Now, take  $X$  to be a smooth compact manifold. Assume that  $\Delta: C^{\infty}(X) \rightarrow C^{\infty}(X)$  is an operator with eigenvalues  $\lambda_n$  ( $\lambda_n \geq 0$ )  $\Delta \varphi_n = \lambda_n \varphi_n$ .

We want to define  $\det \Delta = \prod_{n=1}^{\infty} \lambda_n$ , is this possible?

In interesting cases,  $\lambda_n \rightarrow \infty$ , there is no chance that it will converge.

E.g.  $X = S^1$ ,  $\Delta = -\frac{d^2}{d\theta^2}$   $\varphi = \sum_{n=-\infty}^{\infty} \varphi_n e^{in\theta}$  and  $\{e^{in\theta}\}$  is a basis

$$\Delta e^{in\theta} = -(in)^2 e^{in\theta} = n^2 e^{in\theta}$$

$$\prod n^2 = ?$$

In interesting cases,  $\det \Delta = \prod_{\lambda_n > 0} \lambda_n$ . We apply a zeta-function definition

Formally, define  $\zeta_{\Delta}(s) = \sum_{\lambda_n > 0} \frac{1}{\lambda_n^s}$ .

$$\Rightarrow \zeta'_{\Delta}(s) = \sum_{\lambda_n > 0} (e^{-s \ln \lambda_n})' = \sum_{\lambda_n > 0} -\ln \lambda_n \cdot \lambda_n^{-s}$$

$$\Rightarrow \zeta'_{\Delta}(0) = \sum_{\lambda_n > 0} -\ln \lambda_n = -\ln \prod_{\lambda_n > 0} \lambda_n$$

$$\Rightarrow \prod_{\lambda_n > 0} \lambda_n = e^{-\zeta'_{\Delta}(0)}$$

E.g.  $X = S^1$ ,  $\zeta_{\Delta}(s) = 2 \sum_{n=1}^{\infty} \frac{1}{n^{2s}} = 2\zeta(2s)$ .

Note that  $\lambda_n^{-s} = \frac{1}{\Gamma(s)} \int_0^{\infty} e^{-\lambda_n t} t^{s-1} dt$

$$\Rightarrow \sum_{\lambda_n > 0} \lambda_n^{-s} = \frac{1}{\Gamma(s)} \int_0^{\infty} \sum_{\lambda_n > 0} e^{-\lambda_n t} t^{s-1} dt. \text{ Define } K(t) = \sum_{\lambda_n > 0} e^{-\lambda_n t}$$

Using the same analysis as we did for  $\zeta(s)$ , we obtain a sufficient condition for  $\zeta_{\Delta}(s)$  to extend holomorphically for  $s$  near  $s=0$

Prop. Assume  $\lambda_n$  grows polynomially in  $n$  and  $K(t)$  is a smooth function for  $t > 0$ , furthermore:

(a).  $K(t) \leq C e^{-\mu t}$  for  $t \gg 0$

(b).  $K(t) \sim \sum_{i=1}^n C_i t^{-i} + \text{Smooth function for } t \in [0, 1)$

Then  $\zeta_\Delta(s)$  admits a meromorphic extension for  $s \in \mathbb{C}$ , which is holomorphic at 0.

Pf: Since  $|\int_0^\infty K(t) t^{s-1} dt| \leq \int_0^\infty |K(t)| t^{s-1} dt < \int_0^\infty C \cdot e^{-\mu t} t^{s-1} dt$  converges for  $\text{Re } s > -1$ , thus defines a holomorphic function there.

Now  $\int_0^\infty K(t) t^{s-1} dt = \int_0^1 K(t) t^{s-1} dt + \int_1^\infty K(t) t^{s-1} dt$ .

Since  $\int_1^\infty K(t) t^{s-1} dt$  is well-defined and holomorphic for all  $s \in \mathbb{C}$ , it suffices to show that  $\int_0^1 K(t) t^{s-1} dt$  can be so extended.

Indeed,  $\int_0^1 K(t) t^{s-1} dt = \sum_{m=0}^N C_m \int_0^1 t^{-m-1+s} ds + \int_0^1 E_N(t) t^{s-1} dt$   
 $= \sum_{m=1}^N \frac{C_m}{s-m} + \int_0^1 E_N(t) t^{s-1} dt$  for  $\text{Re}(s) > N+1$

Since  $\sum_{m=1}^N \frac{C_m}{s-m}$  is already meromorphic on  $\mathbb{C}$ , it suffices to show that  $\int_0^1 E_N(t) t^{s-1} dt$  extends to be a meromorphic function on  $\mathbb{C}$

Take the Taylor expansion of  $E_N(t) = \sum_{m=0}^M A_m t^m + F_M(t)$ ,  $A_m = \frac{E_N^{(m)}(0)}{m!}$ . Then  $|F_M(t)| \leq D \cdot t^{M+1}$  for  $t \in (0, 1)$ , and we have:

$$\int_0^1 (\sum_{m=0}^M A_m t^m + F_M(t)) t^{s-1} dt = \sum_{m=0}^M A_m \frac{1}{m+s} + \int_0^1 F_M(t) t^{s-1} dt$$

But  $|\int_0^1 F_M(t) t^{s-1} dt| < \int_0^1 |F_M(t)| t^{s-1} dt < \int_0^1 D \cdot t^{M+s} dt$ , converges for  $\text{Re } s > -M-1$  and defines a meromorphic function there. The result follows.

Finally  $\zeta_K(t)$  has at most simple poles at  $\{N, N-1, \dots, 1, 0, -1, \dots\}$  and  $\Gamma(s)^{-1}$  vanishes at  $s=0$ , the last statement follows. □


Observations: we shall later prove that if  $\Delta$  is an elliptic PDE of positive order, then  $\lambda_n \rightarrow \infty$  polynomially and  $\text{Tr} e^{-t\Delta}$  admits an expansion as in (b), where  $\text{Tr} e^{-t\Delta} = \dim \ker \Delta + \sum_{\lambda_n > 0} e^{-\lambda_n t}$ . We will show that  $\text{Tr} e^{-t\Delta}$  satisfies (b) by solving  $(-\frac{\partial}{\partial t} + \Delta) H(t) = 0$   $H(t)|_{t=0} = I$  and set  $H(t) = e^{-t\Delta}$ .

Basic example:

$$X = \mathbb{C} / \mathbb{Z} \oplus \mathbb{Z} \tau, \quad \tau = \tau_1 + \tau_2 i, \quad \tau_2 > 0$$

$$\Delta = -4 \frac{\partial^2}{\partial \bar{z} \partial z} \text{ acting on periodic functions}$$

$$\begin{cases} \varphi(z+1) = \varphi(z) \\ \varphi(z+\tau) = \varphi(z) \end{cases}$$



Ex. (1) Show that the eigenvalues of  $\Delta$  are given by  $\lambda_{mn} = \frac{4\pi^2}{\tau_2^2} |m+n\tau|^2$ ,  $m, n \in \mathbb{Z}$ .

(2)  $(\det \Delta)' = \frac{1}{(2\pi)^4} \tau_2^2 |\eta(\tau)|^4$  where  $\eta(\tau) = \tau^{1/2} \prod_{n=1}^{\infty} (1 - q^n)$ ,  $q = e^{2\pi\tau}$ .

## §2. Riemann Surfaces

- A Simple Model

Problem: We would like to define the function  $w = \sqrt{z}$  on a maximal domain, where it is holomorphic.

$\sqrt{z} = e^{\frac{1}{2}\ln z}$ , since there is  $\ln$  involved, we start by picking a branch of  $\ln z$  on  $\mathbb{C} \setminus \mathbb{R}_{>0}$ : if we choose the usual definition of  $\ln z$ , then  $z = re^{i\theta} \Rightarrow \sqrt{z} = \sqrt{r} e^{i\frac{\theta}{2}}$  and on upper and lower  $\mathbb{R}_{>0}$  axis,  $\sqrt{z}$  differ by  $e^{\frac{2\pi i}{2}} = e^{\pi i} = -1$ . Take 2 copies of  $\mathbb{C} \setminus \mathbb{R}_{>0}$ , then we can glue them together:




Glue  $I_+$  with  $II_-$  and glue  $I_-$  with  $II_+$ , we obtain  $X = (I) \cup (II)$  and define a function  $w$  on  $X$  as follows  $w = \sqrt{z}$  if  $z \in I$ ;  $w = -\sqrt{z}$  if  $z \in II$ .

Observations:


(1).  $w$  is continuous along the cut

(2).  $X \cong S^2 \setminus \{N, S\}$

For (2):



$\mathbb{C} \setminus \mathbb{R}_{>0} \cong$  upper hemi sphere






Claim: each point  $z_0$  of  $X$  admits a neighborhood  $W_{z_0}$  which is in 1-1 correspondence with a disc  $D$  in  $\mathbb{C}$ .

This is obvious for  $z_0 \notin \mathbb{R}_{>0}$  (the cut)  $W_{z_0} = \{z \mid |z - z_0| < \delta\}$

If  $z_0$  is on the cut, we can just take 2 half discs in I and II to glue together.




Thus we can give  $X = S^2 \setminus \{N, S\}$  coordinate charts by these  $W_{z_0}$ 's.  $W_{z_0} \rightarrow D$   
 $z \mapsto t = z - z_0$

Def. Let  $f$  be a function on  $S$ . We say  $f$  is holomorphic if  $\forall z_0 \in S$ ,  $f|_{W_{z_0}}(z(t))$  is holomorphic as a function of  $t$ .

In this sense,  $w(z) = \begin{cases} \sqrt{z}, & z \in I \\ -\sqrt{z}, & z \in II \end{cases}$  is holomorphic on  $X$ .

Claim: The point  $o$  (South pole) also admits a neighborhood  $V_o$  in 1-1 correspondence with a disk  $D$  in  $\mathbb{C}$ .

In fact:



and plug in  $o$ 's of  $V_I$  and  $V_{II}$  to map to  $o$  of  $D$


Now, let  $\hat{X} = X \cup \{S, N\}$ . A function  $f$  is said to be holomorphic near  $S$  if

$f|_{V_I \cup V_{II}}(z(t))$  is holomorphic in  $D$ .

Two natural examples are  $\omega(z(t))=t$  and  $z(t)=t^2$ , both holomorphic near  $S$ , and have zero's at  $S$  of order 1 and 2 respectively.

Near infinity of  $I$  and  $II$ , we introduce the coordinate  $u=\frac{1}{z}$  and  $u(\infty)=0$ .

Claim:  $\infty$  (north pole) also admits a neighborhood  $V_\infty$  in 1-1 correspondence with the unit disk  $D \subseteq \mathbb{C}$ , and the local coordinate is given by  $z \mapsto \frac{1}{\sqrt{z}}=t, z \in I$  and  $z \mapsto \frac{-1}{\sqrt{z}}=t, z \in II$ , and  $\infty$  (of both  $I$  and  $II$ )  $\mapsto 0$ .



$\omega|_{V_\infty}(z(t)) = \frac{1}{t}$ , simple pole at  $N$ , and  $z|_{V_\infty}(t) = \frac{1}{t^2}$ , double pole at  $t=0$ .

• In summary, the function  $\omega = \sqrt{z}$  defined on  $\mathbb{C} \setminus \mathbb{R}_+$  is extended to a meromorphic function on the space  $\hat{X} = I \sqcup II \sqcup \{S, N\} \cong S^2$ .

The function  $\omega$  has a simple 0 at  $0$ , and a simple pole at  $\infty$ .

The function  $z$  has a double 0 at  $0$ , and a double pole at  $\infty$ .

Under the natural involution (switching)  $I \leftrightarrow II$ , the function  $z$  is even and the function  $\omega$  is odd.

• Another Basic Example

$$w^2 = z(z-1)(z-\lambda) \quad (\lambda \neq 0, 1)$$

Problem: analytically continue  $w = \sqrt{z(z-1)(z-\lambda)}$  defined on some domain of  $\mathbb{C}$ .

For  $w$  to be defined, it suffices  $\sqrt{z}, \sqrt{z-1}, \sqrt{z-\lambda}$  be well-defined. Thus take the


domain  $\mathbb{C} \setminus L$ , where  $L$  passes through  $0, 1, \lambda$ .

Clearly,  $\sqrt{z}$ ,  $\sqrt{z-1}$  and  $\sqrt{z-\lambda}$  are all well defined and holomorphic on  $\mathbb{C} \setminus L$ .

Consider a point  $z \in [0, 1]$ :  $z \rightarrow e^{2\pi i} z$  on the circle  $C$ .

but  $\sqrt{z-1}$  and  $\sqrt{z-\lambda}$  are both well-defined, while  $\sqrt{z} \xrightarrow{C} -\sqrt{z}$

$$\Rightarrow W \rightarrow -W = -\sqrt{z(z-1)(z-\lambda)} \text{ under } C.$$



Next, consider  $z \in [1, \lambda]$

Similarly, we have

$$\sqrt{z} \xrightarrow{D} -\sqrt{z}$$

$$\sqrt{z-1} \xrightarrow{D} -\sqrt{z-1}$$

$$\sqrt{z-\lambda} \xrightarrow{D} \sqrt{z-\lambda}$$

$$\Rightarrow W \xrightarrow{D} W \text{ is invariant, continuous (holomorphic)}$$




Finally, consider  $z \in [\lambda, \infty)$ , and a similar loop.

$$\Rightarrow \sqrt{z} \rightarrow -\sqrt{z}, \sqrt{z-1} \rightarrow -\sqrt{z-1}, \sqrt{z-\lambda} \rightarrow -\sqrt{z-\lambda}$$


and  $W \rightarrow -W$  under the loop.

In summary, singularities occur only on  $[0, 1] \cup [\lambda, \infty)$ .



Thus  $W = \pm \sqrt{z(z-1)(z-\lambda)}$  are holomorphic on  $\mathbb{C} \setminus ([0, 1] \cup [\lambda, \infty))$ . Take two copies and glue similarly as in the previous case.




SII




SII





We can construct explicitly local coordinates near the south pole  $S$ .



Claim:  $w$  is holomorphic near  $0$  ( $S$ ). (similarly for  $1$  and  $\lambda$ )

$$w = \sqrt{t^2(t^2-1)(t^2-\lambda)} = t \sqrt{(t^2-1)(t^2-\lambda)}; \text{ for } |t-0| \ll 0, \sqrt{(t^2-1)(t^2-\lambda)} \text{ is holomorphic}$$

$\Rightarrow w$  has a simple  $0$  at  $t=0$

Similarly, near  $\infty$  ( $N$ ), introduce the holomorphic coordinate  $t = \frac{1}{\sqrt{z}}$  on  $I$  and  $t = -\frac{1}{\sqrt{z}}$  on  $II$ .  $w = \sqrt{\frac{1}{t^2}(\frac{1}{t^2}-1)(\frac{1}{t^2}-\lambda)} = \frac{1}{t^3} \sqrt{(1-t^2)(1-\lambda t^2)}$ ; for  $|t-0| \ll 0$   $\sqrt{(1-t^2)(1-\lambda t^2)}$  is holomorphic


$\Rightarrow w$  is meromorphic with a pole of order  $3$ .

In conclusion,  $w$  has been extended to a meromorphic function on  $T$  and  $w$  has  $3$  simple  $0$ 's and a pole of order  $3$ .

The function  $z$  is also extended meromorphically, with an order  $2$   $0$  at  $S$  and an order  $2$  pole at  $N$ .

- Question: There are many tori with a complex structure (i.e. a notion of holomorphic or meromorphic function applies), for instance:

$$T_\tau \triangleq \mathbb{C} / \mathbb{Z} \oplus \mathbb{Z}\tau, \quad \tau \in \mathbb{C}, \operatorname{Im}\tau > 0$$



A function  $\varphi$  on  $T_\tau$  can be identified with a function on  $\mathbb{C}$  satisfying  $\varphi(z+m+n\tau) = \varphi(z)$ ,  $\forall m, n \in \mathbb{Z}$ , and  $\varphi$  on  $T_\tau$  is holomorphic (meromorphic) if the corresponding  $\varphi$  on  $\mathbb{C}$  is holomorphic (meromorphic).

Thus we need to see if  $X$  can be identified with  $T_\tau$  for some  $\tau$ , and if yes how do we determine such a  $\tau$ ?

- Key strategy: There are very few (in fact, only constants, by maximal modulus theorem) holomorphic functions on  $\hat{X}$ . Rather we work instead with

(1). holomorphic forms

and/or

(2). meromorphic functions

And then proceed in the following 3 steps.

Step 1: Construct these objects (← explicit forms)

Step 2: Develop techniques for manipulating them (← Riemann bilinear relations and abelian integrals)

Step 3: Use them to construct a holomorphic map  $\hat{X} \rightarrow \mathbb{C} / \mathbb{Z} + \mathbb{Z}\tau$  (← The Abel map: Jacobi inversion thm, based on Abel's thm.)

### I. Construction of holomorphic differentials.

Recall that  $\hat{X}$  has two meromorphic functions on  $\hat{X}$ , namely  $z: \hat{X} \rightarrow \mathbb{C} \cup \{\infty\}$ ,

$$p \mapsto z(p) \quad \text{and} \quad w(p) = \begin{cases} \sqrt{z(z-1)(z-\lambda)} & \text{on I} \\ -\sqrt{z(z-1)(z-\lambda)} & \text{on II} \end{cases}$$

Consider the differential " $\frac{dz}{w}$ " on  $\hat{X}$

Key observation: " $\frac{dz}{w}$ " is actually a holomorphic differential form on all of  $\hat{X}$

To see this, express  $\frac{dz}{w}$  in a local coordinate system.

$$\frac{dz}{w} = \frac{dz}{\sqrt{z(z-1)(z-\lambda)}}. \quad \text{Near } z=0, \text{ recall that the holomorphic coordinate is given by}$$

$$t = \sqrt{z} \Rightarrow z = t^2, \text{ thus } dz = 2t dt$$

$$\Rightarrow \frac{dz}{w} = \frac{2t dt}{\sqrt{t^2(t-1)(t-\lambda)}} = \frac{2 dt}{\sqrt{(t-1)(t-\lambda)}}, \text{ which is holomorphic for } |t| \ll 1.$$

The same reasoning applies to  $z_0 = 1, \lambda$ . Near  $z_0 = \infty$ , the holomorphic coordinate is given by  $t = \frac{1}{\sqrt{z}} \Rightarrow z = \frac{1}{t^2}, dz = -2 \frac{dt}{t^3}$

Thus  $\frac{dz}{w} = \frac{-2dt/t^3}{\sqrt{\frac{1}{t^2}(t^2-1)(t^2-\lambda^2)}} = \frac{-2dt/t^3}{\frac{1}{t^2}\sqrt{(1-t^2)(1-\lambda^2 t^2)}} = \frac{-2dt}{\sqrt{(1-t^2)(1-\lambda^2 t^2)}}$ , which is again holomorphic for  $|t| < 1$ .

Using this form  $\omega = \frac{dz}{w}$ , we can construct a map from  $\hat{X}$  to  $\mathbb{C}/\text{lattice}$  in the following way:

(The Abel map) Fix  $P_0 \in \hat{X}, \forall P \in \hat{X}$ .

Consider  $\int_{P_0}^P \omega$ , integration along  $\gamma$ . If  $\gamma$  can be continuously deformed to  $\gamma'$ , then  $\int_{\gamma} \omega = \int_{\gamma'} \omega$ .

Pick  $A, B$  two cycles on  $\hat{X}$

(A topological fact): Any two  $\gamma, \gamma'$  connecting  $P_0$  and  $P$  have  $\gamma - \gamma' \simeq nA + mB, n, m \in \mathbb{Z}$   
 $\Rightarrow \int_{\gamma} \omega = \int_{\gamma'} \omega + n \oint_A \omega + m \oint_B \omega, n, m \in \mathbb{Z}$ .

Thus we may view  $\int_{\gamma} \omega$  as an equivalence class in  $\mathbb{C}/\mathbb{Z}\alpha + \mathbb{Z}\beta$  rather than a complex number, where  $\alpha = \oint_A \omega, \beta = \oint_B \omega$





We shall later show that  $\alpha \neq 0, \beta \neq 0$ , and then we can normalize  $\omega$  and define  $\hat{\omega} = \frac{\omega}{\oint_A \omega} \Rightarrow \oint_A \hat{\omega} = 1$ , and  $\tau \triangleq \frac{\beta}{\alpha} = \frac{\oint_B \hat{\omega}}{\oint_A \hat{\omega}}$ . The Abel map is then defined as:


$$\hat{X} \longrightarrow \mathbb{C}/\mathbb{Z} + \mathbb{Z}\tau. \quad P \mapsto A(P) \triangleq \left[ \int_{P_0}^P \hat{\omega} \right]$$

Furthermore, we will also show that  $\text{Im}\tau > 0$ , so that  $\mathbb{C}/\mathbb{Z} + \mathbb{Z}\tau \cong \mathbb{T}^2$

Thm. (Jacobi Inversion Thm.) The Abel map is holomorphic from  $\hat{X}$  to  $\mathbb{C}/\mathbb{Z} + \mathbb{Z}\tau$ , and is 1-1 and onto.

Pf: We may choose the cycles  $A$  and  $B$  as follows:





$\Rightarrow \alpha = 2 \int_0^1 \frac{dx}{\sqrt{x(x-1)(x-\lambda)}}$ . We will show later that  $\alpha \in \mathbb{R}$ .

The Abel map is holomorphic since  $dA(p) = \omega(p)$ . Furthermore  $\omega$  is never 0  $\Rightarrow$  the Abel map is locally an isomorphism. The open mapping thm  $\Rightarrow$  the image of  $A$  is open; while  $\hat{X}$  compact  $\Rightarrow$  the image is closed. Thus we conclude that  $A$  maps  $\hat{X}$  holomorphically onto  $\mathbb{C}/\mathbb{Z} + \mathbb{Z}z$ .

To show that it is 1-1, we shall use Abel's thm, to be proved below, which states that  $\forall P, Q \in \hat{X}$ ,  $A(P) = A(Q)$  iff there exists a meromorphic function  $f$ , with a simple zero at  $P$  and a simple pole at  $Q$ .


Thus if  $P \neq Q$  but  $A(P) = A(Q)$  we would obtain a meromorphic function  $f$  with a simple pole at  $Q$ . Consider the meromorphic differential  $f \frac{dz}{\omega}$ , which has a simple pole at  $Q$ , since  $\frac{dz}{\omega}$  is globally holomorphic and non-vanishing. Thus the residue of  $f \frac{dz}{\omega}$  is then non-zero at  $Q$ , which violates the following lemma.  $\square$

Lemma. A meromorphic differential  $\omega$  on a compact Riemann surface has

$$\sum_{P: \text{poles}} (\text{Res } \omega)(P) = 0$$

Pf: Let  $P_1, \dots, P_k$  be all the poles of  $\omega$ . Then

$$\begin{aligned} \sum_{i=1}^k (\text{Res } \omega)(P_i) &= \sum_{i=1}^k \oint_{\partial D_i} \omega = - \int_{X \setminus \bigcup_{i=1}^k D_i} \omega \\ &= - \int_{X \setminus \bigcup_{i=1}^k D_i} \omega \\ &= 0 \end{aligned}$$



The last equality holds since  $\omega$  is a holomorphic differential form on  $X \setminus \bigcup_{i=1}^k D_i$ .  $\square$

Thm. (Abel) Let  $P_1, \dots, P_M, Q_1, \dots, Q_N$  be points on  $\hat{X}$ , counted with multiplicity. Then  $\exists f$  on  $\hat{X}$ ,  $f$  meromorphic with 0's at  $P_i$  and poles at  $Q_j \iff M=N$  and  $\sum_{i=1}^M A(P_i) = \sum_{j=1}^N A(Q_j)$ , the addition being induced from the group addition of  $\mathbb{C}$ .

Sketch of proof of Abel's theorem.

- Idea: try to construct / identify  $f$  from  $\frac{df}{f}$ , which is easier to deal with since its residues are always  $\pm 1$ :  $+1$  for a pole of  $f$ ,  $-1$  for a zero of  $f$ , both to be counted with multiplicity.

The basic building block is the so called Abelian differential of the 3<sup>rd</sup> kind. Lemma:  $\forall Q_1 \neq Q_2$  on  $\hat{X}$ ,  $\exists$  a meromorphic form  $\omega_{Q_1, Q_2}(P)$  with simple poles at exactly  $Q_1$  and  $Q_2$  and residues  $-1, +1$  respectively.

Assuming this lemma, a candidate of  $\frac{df}{f}$  would be

$$\frac{df}{f} \stackrel{?}{=} \underbrace{\sum_{i=1}^N \omega_{P_0, P_i}}_{\substack{\text{with a simple} \\ \text{pole at each } P_i \\ \text{whose residue is } 1}} - \underbrace{\sum_{i=1}^N \omega_{P_0, Q_i}}_{\substack{\text{with a simple} \\ \text{pole at each } Q_i \\ \text{whose residue is } -1}} \quad (\text{Fixing } P_0)$$

no poles at  $P_0$  since they all got canceled

Then  $\frac{df}{f} - \sum_{i=1}^N \omega_{P_0, P_i} + \sum_{i=1}^N \omega_{P_0, Q_i} = c \cdot \frac{dz}{z}$  for some constant  $c$ .

In case we know  $\frac{df}{f}$ , we may reconstruct  $f$  by formally:

$$f = \exp\left(\int^z \frac{df}{f}\right)$$


where we have to make sure that the right hand side is well-defined. This will be the part where  $\sum_{i=1}^M A(P_i) = \sum_{j=1}^N A(Q_j)$  becomes a necessary and sufficient condition.

Existence of  $\omega_{Q_1, Q_2}$  (proof of lemma)

Now we come back to the realization of  $\hat{X}$  as  $\omega^2 = z(z-1)(z-\lambda)$ , i.e. a ramified double cover of the Riemann sphere  $S^2 \cong \mathbb{P}^1 = \mathbb{C} \cup \{\infty\}$







Now given  $Q_1, Q_2$  on  $\hat{X}$ , we try to construct  $\omega_{Q_1, Q_2}$ . Firstly we assume that  $z(Q_1) \neq z(Q_2)$ . Consider:

$$\omega_{Q_1, Q_2} \stackrel{?}{=} \frac{1}{(z - z(Q_1))(z - z(Q_2))} \frac{dz}{w}$$

But this actually doesn't work since  $z(\tilde{Q}_1) = z(Q_1)$ ,  $z(\tilde{Q}_2) = z(Q_2)$ , there will be 2 more poles than we actually need. Hence we can try to multiply a function to kill these poles off. Take:

$$\omega_{Q_1, Q_2} = \frac{\alpha z + \beta + w}{(z - z(Q_1))(z - z(Q_2))} \frac{dz}{w}$$

( $w$  must be involved since  $z$  is an "even" function, wherever  $Q_i$  occurs,  $\tilde{Q}_i$  would be of the same value for  $z$ ).

Thus we want:  $\alpha z + \beta + w = 0$ ,  $z = z(Q_1)$ ,  $w = w(\tilde{Q}_1)$

$$\alpha z + \beta + w = 0, \quad z = z(Q_2), \quad w = w(\tilde{Q}_2)$$

For this equation to have a solution, it suffices that  $\begin{vmatrix} z(Q_1) & 1 \\ z(Q_2) & 1 \end{vmatrix} \neq 0$ , which is automatic since we assumed  $z(Q_1) \neq z(Q_2)$  from the outset. Moreover, this function doesn't create new poles at  $\infty$  since  $|z| = O(|z_1|)$ ,  $|w| = O(|z_1|^{\frac{3}{2}})$  as  $|z| \rightarrow \infty$  and the denominator is  $O(|z|^2)$  as  $z \rightarrow \infty$

If  $z(Q_1) = z(Q_2)$ , it suffices to consider  $\frac{1}{z - z(Q_1)} \frac{dz}{w}$

If one of  $Q_i$ , say  $Q_1$ , is  $\infty$ , then it suffices to take  $\frac{w - w(\tilde{Q}_2)}{z - z(Q_2)} \frac{dz}{w}$

Furthermore, the above expressions even make sense if one of  $Q_1, Q_2$ , or both are the ramification points. In fact, the numerator is solved to be  $\frac{w_1 - w_2}{z_2 - z_1} z + \frac{z_2 w_1 - z_1 w_2}{z_2 - z_1} + w$  ( $w_1 = w(\tilde{Q}_1)$ ,  $z_1 = z(Q_1)$ ,  $w_2 = w(\tilde{Q}_2)$ ,  $z_2 = z(Q_2)$ ). If  $w_1 = 0$ , this just gives  $\frac{w_2(z - z_1)}{z_1 - z_2} + w$ , but  $(z - z_1)$  is a zero of order 2 at the ramification point while  $w$  is a 0 of order 1 at  $z_1$ . Similarly if both are ramification points, then the numerator just gives  $w$ , which vanishes to order 1 at  $Q_1$  and  $Q_2$ .

We may also construct meromorphic forms with a double pole at any  $Q \in \hat{X}$ .

If  $Q \neq 0, 1, \lambda$  or  $\infty$ , then  $w(\tilde{Q}) = -w(Q) \neq 0$  we may try  $\frac{\alpha z + \beta w + \gamma}{(z - z(Q))^2} \frac{dz}{w}$  such that  $\alpha z + \beta w + \gamma$  vanishes to order 2 at  $\tilde{Q}$ , where  $\tilde{Q}, Q$  are the two distinct points on  $\hat{X}$  lying over  $z(Q)$ . Consider:

$$\alpha z + \beta w + \gamma = ((z(Q_1)-1)(z(Q_1)-\lambda) + z(Q_1)(z(Q_1)-1) + z(Q_1)(z(Q_1)-\lambda))(z-z(Q_1)) - 2w(\tilde{Q})(w-w(\tilde{Q})))$$

We claim that the above expression vanishes to order  $\geq 2$  at  $\tilde{Q}$ .

$$\begin{aligned} \text{Indeed } (w-w(\tilde{Q})+w(\tilde{Q}))^2 &= z(z-1)(z-\lambda) = (z-z(Q_1)+z(Q_1))(z-z(Q_1)+z(Q_1)-1)(z-z(Q_1)+z(Q_1)-\lambda) \\ (w-w(\tilde{Q}))^2 + 2w(\tilde{Q})(w-w(\tilde{Q})) + w(\tilde{Q})^2 &= z(Q_1)(z(Q_1)-1)(z(Q_1)-\lambda) \\ &\quad + ((z(Q_1)-1)(z(Q_1)-\lambda) + z(Q_1)(z(Q_1)-1) + z(Q_1)(z(Q_1)-\lambda))(z-z(Q_1)) \\ &\quad + (z(Q_1)+z(Q_1)-1+z(Q_1)-\lambda)(z-z(Q_1))^2 \\ &\quad + (z-z(Q_1))^3 \end{aligned}$$

(note that  $z(Q_1) = z(\tilde{Q})$ ,  $w(\tilde{Q})^2 = z(\tilde{Q})(z(\tilde{Q})-1)(z(\tilde{Q})-\lambda)$ )

$$\begin{aligned} \Rightarrow ((z(Q_1)-1)(z(Q_1)-\lambda) + z(Q_1)(z(Q_1)-1) + z(Q_1)(z(Q_1)-\lambda))(z-z(Q_1)) - 2w(\tilde{Q})w \\ = (w-w(\tilde{Q}))^2 - (z(Q_1)+z(Q_1)-1+z(Q_1)-\lambda)(z-z(Q_1))^2 - (z-z(Q_1))^3 \\ = O(|z-z(\tilde{Q})|^2) \end{aligned}$$

Since  $w-w(\tilde{Q})$  is locally a holomorphic function in  $z-z(Q)$ , i.e.  $w-w(\tilde{Q}) = (z-z(Q)) \cdot h(z-z(Q))$ .

Moreover, since in our case  $w(\tilde{Q}) = -w(Q) \neq 0$ ,

$$\alpha z + \beta w + \gamma \Big|_{(z(Q), w(Q))} = -4w(\tilde{Q})^2 \neq 0$$



and  $O(|\alpha z + \beta w + \gamma|) = O(|z|^{\frac{3}{2}})$  ( $|z| \rightarrow \infty$ ), while  $O(|z-z(Q)|^2) = O(|z|^2)$  ( $|z| \rightarrow \infty$ ).

It follows that  $\frac{\alpha z + \beta w + \gamma}{(z-z(Q))^2} \frac{dz}{w}$  satisfies the required conditions.

If  $z(Q) = 0, 1$ , or  $\lambda$ , it suffices to take  $\frac{1}{z-z(Q)} \frac{dz}{w}$ , since  $(z-z(Q))$  vanishes to order 2 at these ramification points.

If  $z(Q) = \infty$ , it suffices to take  $z \frac{dz}{w}$ , since  $z$  has an order 2 pole at  $\infty$ .

In general, the above constructions can be visualized by considering the following picture of tori (as elliptic curves)




For the next step, we need to deal with  $\int^z \omega_{P_0, P_1}$  and  $\int^z \omega$ , and some computation needs to be done.


Method of abelian integral (Riemann)

Pick some fixed  $A, B$  cycles on  $\hat{X}$ ,  $P_0 \in \hat{X}$

$$f(z) \triangleq \int_{P_0}^z \omega \quad (I)$$



On the right hand side picture,  $f(z)$  is well-defined on  $\hat{X}_{cut}$  since  $\hat{X}_{cut}$  is simply connected. However, it may not be doubly periodic.



i.e.  $f(z) \neq f(z+A)$  and  $f(z) \neq f(z+B)$ . Thus it may not correspond to a function on  $\hat{X}$ . However, the differences are under control:

$$f(z+A) - f(z) = \int_z^{z+A} \omega = \oint_A \omega$$

$$f(z'+B) - f(z') = \int_{z'}^{z'+B} \omega = \oint_B \omega$$


Example of using abelian integral

Let  $f$  be the abelian integral (I). Consider  $\frac{i}{2} \int_{\hat{X}} \omega \wedge \bar{\omega}$ . In local coordinate systems we may write  $\omega = \psi(t) dt$ . Then  $\omega \wedge \bar{\omega} = |\psi(t)|^2 dt \wedge d\bar{t} = |\psi(t)|^2 (-2i dt_1 \wedge dt_2)$ ,  $t = t_1 + it_2$   
 $\Rightarrow \frac{i}{2} \omega \wedge \bar{\omega} = |\psi(t)|^2 dt_1 \wedge dt_2$ , thus  $\frac{i}{2} \int_{\hat{X}} \omega \wedge \bar{\omega} > 0$ .

Moreover, on  $\hat{X}_{cut}$ ,  $0 < \frac{i}{2} \int_{\hat{X}} \omega \wedge \bar{\omega} = \frac{i}{2} \int_{\hat{X}_{cut}} d(f(t) \cdot \bar{\omega})$  (The last equality holds because  $d(f(z) \bar{\omega}) = (df(z)) \cdot \bar{\omega} + f(z) d\bar{\omega} = (\frac{\partial f}{\partial t} dt + \frac{\partial f}{\partial \bar{t}} d\bar{t}) \psi(t) dt + f(t) (\frac{\partial \bar{\psi}}{\partial t} dt + \frac{\partial \bar{\psi}}{\partial \bar{t}} d\bar{t})$  and  $\frac{\partial f}{\partial \bar{t}} = \frac{\partial \bar{f}}{\partial t} = 0$  since  $f$  is holomorphic; or it can be seen topologically since  $d\omega = 0$  and  $\omega|_{\hat{X}_{cut}} = df$  since  $\hat{X}_{cut}$  is contractible)


By Stokes' thm:

$$\begin{aligned}
 0 < \int_{\hat{X}} \omega \wedge \bar{\omega} &= \frac{i}{2} \int_{\hat{X}_{\text{int}}} d(f(t) \bar{\omega}) = \frac{i}{2} \int_{\partial \hat{X}_{\text{int}}} f(t) \bar{\omega} \\
 &= \frac{i}{2} \left\{ \int_A f(z) \bar{\omega}(z) - \int_A f(z+B) \bar{\omega}(z+B) \right. \\
 &\quad \left. + \int_B f(z+A) \bar{\omega}(z+A) - \int_B f(z) \bar{\omega}(z) \right\} \\
 &= \frac{i}{2} \left\{ -\oint_A \bar{\omega} \oint_B \omega + \oint_B \bar{\omega} \oint_A \omega \right\} \\
 &= \text{Im} \left( \oint_A \bar{\omega} \oint_B \omega \right)
 \end{aligned}$$



The second last equality holds because  $\omega, \bar{\omega}$  are both defined on  $\hat{X}$ , thus must agree on boundaries of  $\hat{X}_{\text{int}}$ , and  $f(z+A) - f(z) = \oint_A \omega$ ,  $f(z+B) - f(z) = \oint_B \omega$

If we normalize  $\oint_A \omega = 1 \Rightarrow \text{Im} \oint_B \omega > 0$ , proving that the image of the Abel map is a genuine torus.



Example of method of abelian differential

Let  $\omega_{Q_1, Q_2}(z)$  be a meromorphic differential with simple poles at  $Q_1, Q_2$ , whose residues are  $+1$  at  $Q_2$  and  $-1$  at  $Q_1$ . Fix cycles  $A, B$  ( $\neq Q_1, Q_2$ ) and normalize  $\omega_{Q_1, Q_2}$  so that  $\oint_A \omega_{Q_1, Q_2} = 0$ . This can be done since we have assumed  $\oint_A \omega = 1$ , and subtracting  $\oint_A \omega_{Q_1, Q_2} \cdot \omega$  from  $\omega_{Q_1, Q_2}$  won't affect its poles.

Consider the integral  $\oint_C f \cdot \omega_{Q_1, Q_2}$ :


Deforming  $C$  towards  $Q_1$  and  $Q_2$  gives:

$$\begin{aligned}
 \oint_C f \omega_{Q_1, Q_2} &= 2\pi i \sum_{Q_i} \text{Res}(f \cdot \omega_{Q_1, Q_2})(Q_i) = 2\pi i (f(Q_2) - f(Q_1)) \\
 &(\equiv 2\pi i (A(Q_2) - A(Q_1)) \pmod{\mathbb{Z} + \mathbb{Z}z})
 \end{aligned}$$

Deforming  $C$  towards  $\partial \hat{X}_{\text{int}}$  gives:

$$\begin{aligned}
 \oint_C f \omega_{Q_1, Q_2} &= \int_A (f(z) - f(z+B)) \omega_{Q_1, Q_2} - \int_B (f(z) - f(z+A)) \omega_{Q_1, Q_2} \\
 &= -\oint_A \omega_{Q_1, Q_2} \oint_B \omega + \oint_B \omega_{Q_1, Q_2} \oint_A \omega \\
 &= \oint_B \omega_{Q_1, Q_2} \quad \text{since we normalized } \oint_A \omega_{Q_1, Q_2} = 0, \oint_A \omega = 1.
 \end{aligned}$$

$$\Rightarrow \frac{1}{2\pi i} \oint_B \omega_{Q_1, Q_2} = f(Q_2) - f(Q_1)$$



Proof of Abel's thm.

" $\Rightarrow$ " Assume a meromorphic function  $\varphi$  with the desired property exists. Then  $\frac{d\varphi}{\varphi}$  is a meromorphic form  $\Rightarrow 0 = \sum \text{Res}(\frac{d\varphi}{\varphi}) = M - N$ .

To see the other property, observe that  $\frac{d\varphi}{\varphi} = \sum_{i=1}^N \omega_{P_i} - \sum_{j=1}^N \omega_{Q_j} + \lambda \omega$  for some  $\lambda \in \mathbb{C}$ . This is because:

Lemma:  $\tilde{\omega}$  is holomorphic on  $\hat{X} \Rightarrow \tilde{\omega} = \lambda \omega$ .

Pf: First of all, we may subtract  $(\oint_A \tilde{\omega}) \cdot \omega$  from  $\tilde{\omega}$  and assume that  $\oint_A \tilde{\omega} = 0$

Consider the abelian integral of  $\tilde{\omega}$ , we obtain:

$$0 \leq \frac{i}{2} \int_{\hat{X}_{\text{out}}} \tilde{\omega} \wedge \bar{\tilde{\omega}} = \text{Im}(\oint_A \tilde{\omega} \cdot \oint_B \bar{\tilde{\omega}})$$

But  $\oint_A \tilde{\omega} = 0 \Rightarrow \frac{i}{2} \int_{\hat{X}_{\text{out}}} \tilde{\omega} \wedge \bar{\tilde{\omega}} = 0$ . Locally,  $\tilde{\omega} = \psi(t) dt \Rightarrow \frac{i}{2} \tilde{\omega} \wedge \bar{\tilde{\omega}} = |\psi(t)|^2 dt_1 \wedge dt_2$

Thus  $\psi(t) \equiv 0 \Rightarrow \tilde{\omega} \equiv 0$ .

□ of lemma.

Now we can evaluate the periods of  $\oint_C \frac{d\varphi}{\varphi}$ . This must be  $2\pi i \cdot n$  for some  $n \in \mathbb{Z}$  since anyway it's to calculate the residues within (or outside, which doesn't matter)  $C$

$$2\pi i n = \oint_A \frac{d\varphi}{\varphi} = \oint_A \sum_{i=1}^N (\omega_{P_i} - \omega_{Q_i}) + \oint_A c \cdot \omega = c \cdot \oint_A \omega = c$$

since we have assumed that  $\oint_A \omega_{P_i} = 0$ . On the other hand:

$$2\pi i m = \oint_B \frac{d\varphi}{\varphi} = \oint_B \sum_{i=1}^N (\omega_{P_i} - \omega_{Q_i}) + \oint_B c \cdot \omega$$

$$= 2\pi i \cdot \sum_{i=1}^N (f(P_i) - f(Q_i)) + cZ$$

$$\Rightarrow \sum_{i=1}^N f(Q_i) = \sum_{i=1}^N f(P_i) + (n+mZ) \equiv \sum_{i=1}^N f(Q_i) \pmod{\mathbb{Z} + \mathbb{Z}Z}$$


" $\Leftarrow$ " Assume Abel's relation holds, i.e.  $\sum_{i=1}^N f(P_i) = \sum_{j=1}^N f(Q_j) + n + mZ$ .

Define  $\varphi(z) = \exp(\int_M^z \sum_{i=1}^N (\omega_{P_i} - \omega_{Q_j}) + c\omega)$ , where  $c$  is to be chosen.

For  $\varphi$  to be a well-defined meromorphic function on

$\hat{X}$ , it's sufficient and necessary that:

- (1).  $\varphi(z)$  is independent of choices of path.
- (2).  $\varphi(z+A) = \varphi(z)$
- (3).  $\varphi(z+B) = \varphi(z)$



(1) is easy since choosing different path amounts to adding  $2\pi i \cdot$  (residues of  $P_i, Q_j$ 's) which is an integral multiple of  $2\pi i$  and doesn't affect the value of  $\varphi$ .

(2): we have assumed that  $\oint_A \omega_{P_i} = 0$  and  $\oint_A \omega = 1$  thus it sufficient if  $c = 2\pi i \cdot k, k \in \mathbb{Z}$

(3):  $\sum_{j=1}^N \oint_B (\omega_{P_0 P_i} - \omega_{P_0 Q_i}) = 2\pi i \sum_{j=1}^N (f(P_j) - f(Q_j))$ ,  $\oint_B \omega = z$ . By our assumption  $2\pi i \sum_{j=1}^N (f(P_j) - f(Q_j)) = 2\pi i n + 2\pi i m z$ . Choose  $c = 2\pi i m$  and condition (3) and (2) are both satisfied.  $\square$

### §3. Function Theory on Tori

We have seen that on the R.S.  $\hat{X} : w^2 = z(z-1)(z-\lambda)$ , there are

(1) A holomorphic form  $\omega = \frac{dz}{\sqrt{z(z-1)(z-\lambda)}}$

(2)  $\forall P, Q \in \hat{X}, P \neq Q, \exists$  a meromorphic form  $\omega_{PQ}$  with poles at  $P, Q$ .

(3)  $\forall P \in \hat{X}, \exists$  a meromorphic form  $\omega_P$  with a double pole at  $P$ .

We have also shown that  $\forall P \in \hat{X}, P \mapsto A(P) = \int_{P_0}^P \omega \in \mathbb{C}/\mathbb{Z} + \mathbb{Z}\tau$ , where

$$\tau = \left( \int_0^\lambda \frac{dz}{\sqrt{z(z-1)(z-\lambda)}} \right) / \left( \int_0^1 \frac{dz}{\sqrt{z(z-1)(z-\lambda)}} \right)$$

#### Several View Points

- 1). Abelian Integral (done in §2 already)
- 2). Weierstrass function  $\wp(z), \zeta(z), \sigma(z)$
- 3). Jacobi theta function
- 4).  $\bar{\partial}$ -construction (P.D.E.)

#### • Function Theory According To Weierstrass

Let  $\mathbb{C}/\mathbb{Z} + \mathbb{Z}\tau$  be a given complex torus,  $\text{Im}\tau > 0$ . A function  $\varphi$  on  $\mathbb{C}/\mathbb{Z} + \mathbb{Z}\tau$  is simply a function on  $\mathbb{C}$  satisfying a doubly periodic condition:

$$\begin{cases} \varphi(z+1) = \varphi(z) \\ \varphi(z+\tau) = \varphi(z) \end{cases}$$

A holomorphic 1-form :  $z \mapsto z+m+n\tau$  by  $\mathbb{Z} \oplus \mathbb{Z}$  action. But  $dz$  is invariant under this action  $\Rightarrow dz$  is a holomorphic 1-form on  $\hat{X}$ .

Suppose  $P=0$ , we want to construct a meromorphic form with a double pole at  $P$ . In view of the form  $dz$ , we can identify forms and functions, which is possible because  $dz$  has neither 0's nor poles,  $f(z) \leftrightarrow f(z)dz$ . Thus we need to construct a function with exactly a double pole at 0 and which is doubly periodic:

Our first try would be to average out  $\frac{1}{z^2}$  over the lattice  $\mathcal{L} \cong \mathbb{Z} + \mathbb{Z}\tau$

$\sum_{w \in \mathcal{L}} \frac{1}{(z+w)^2}$ , but unfortunately it doesn't converge. ( $\iint_{\mathbb{R}^n} \frac{dx}{(1+|x|)^p} < \infty$  if  $p > n$ )

Define  $\beta(z) \triangleq \frac{1}{z^2} + \sum_{w \in \mathcal{L}^*} \left\{ \frac{1}{(z+w)^2} - \frac{1}{w^2} \right\}$   $\mathcal{L}^* = \mathcal{L} \setminus \{0\}$ . Note that when  $|w| \rightarrow \infty$ ,  $\frac{1}{(z+w)^2} - \frac{1}{w^2} = \frac{1}{w^2} \left( \frac{1}{(1+\frac{z}{w})^2} - 1 \right) = O\left(\frac{1}{|w|}\right)$ ,  $\forall z$  fixed. Thus the sum converges.

From the discussion above, the series converges for any  $z$  and defines a meromorphic function with poles on  $\mathcal{L}$ .

Claim:  $\beta(z)$  is doubly periodic.

Indeed, we can compute that  $\beta'(z) = -\frac{2}{z^3} - \sum_{w \in \mathcal{L}^*} \frac{2}{(z+w)^3} = -2 \sum_{w \in \mathcal{L}} \frac{1}{(z+w)^3}$   
 $\Rightarrow \beta'(z)$  is doubly periodic:  $\beta'(z+1) = \beta'(z)$ ,  $\beta'(z+z) = \beta'(z)$ .

But this implies  $\frac{d}{dz}(\beta(z+1) - \beta(z)) = 0$ ,  $\frac{d}{dz}(\beta(z+z) - \beta(z)) = 0$ .  
 $\Rightarrow \beta(z+1) = \beta(z) + C_1$ ,  $\beta(z+z) = \beta(z) + C_2$ .

Moreover, since  $\beta(z)$  is even (the lattice is symmetric w.r.t. 0), taking  $z$  to be  $-\frac{1}{2}$ ,  $-\frac{z}{2}$  respectively gives  $\beta(\frac{1}{2}) = \beta(-\frac{1}{2}) + C_1$ ,  $\beta(\frac{z}{2}) = \beta(-\frac{z}{2}) + C_2 \Rightarrow C_1 = C_2 = 0$ .

Observe that on  $\mathbb{C}$  we have the function  $z$  which is holomorphic with a simple zero. Thus functions with given zero's and poles can immediately be written as  $f(z) = \frac{\pi(z - p_i)}{\pi(z - q_j)}$ .

Is there an analogue for such a function on the torus? i.e. a function with a single 0? Not true by maximal modulus principle. But there is an adequate replacement.

Idea: 1). Integrate  $\beta(z)$  twice to get a  $\log z$  and take exponential

2). Integrals give rise to Abelian integrals, so we need to keep track of the periods:  $(\varphi(z+1) - \varphi(z))$ ,  $(\varphi(z+z) - \varphi(z))$ .

Integral of  $\beta(z)$ .

Consider  $\frac{1}{z} + \sum_{w \in \mathcal{L}^*} \left\{ \frac{1}{z+w} + \frac{z}{w^2} \right\}$ , which is the formal integration of  $-\beta(z)$ . But again this has convergence problems. How to correct this?



$$\frac{1}{z+w} = \frac{1}{w} \cdot \frac{1}{(1+\frac{z}{w})} = \frac{1}{w} (1 - \frac{z}{w} + \frac{z^2}{w^2} - \dots) \Rightarrow \frac{1}{z+w} + \frac{z}{w^2} = \frac{1}{w} + \underbrace{O(\frac{1}{|w|^3})}_{\text{converges}}$$

Define  $\zeta(z) = \frac{1}{z} + \sum_{w \in \mathbb{Z}^*} \left\{ \frac{1}{z+w} - \frac{1}{w} + \frac{z}{w^2} \right\}$ , which is meromorphic on the whole plane and with simple poles at  $\mathbb{Z}$ .

Clearly,  $\zeta'(z) = -\beta(z)$  (we are just subtracting constants from the formal anti-derivative of  $\beta(z)$ ). We need to determine  $\zeta(z+1) - \zeta(z)$ ,  $\zeta(z+z) - \zeta(z)$

$$\text{But } \frac{d}{dz} (\zeta(z+1) - \zeta(z)) = -\beta(z+1) + \beta(z) = 0 \Rightarrow \zeta(z+1) = \zeta(z) + \eta_1$$


$$\text{Similarly } \zeta(z+z) = \zeta(z) + \eta_2$$

Lemma:  $z\eta_1 - \eta_2 = 2\pi i$

$$\text{Pf: } \int_{\partial \text{out}} \zeta(z) dz = 2\pi i \cdot \text{Res} \zeta(z) = 2\pi i$$

On the other hand:

$$\begin{aligned} \int_{\partial \text{out}} \zeta(z) dz &= \int_z^{z+1} \zeta(z) dz + \int_{z+1}^{z+1+z} \zeta(z) dz \\ &\quad - \int_{z+z}^{z+1+z} \zeta(z) dz - \int_z^{z+z} \zeta(z) dz \\ &= \int_z^{z+1} (\zeta(z) - \zeta(z+z)) dz + \int_z^{z+z} (\zeta(z+1) - \zeta(z)) dz \\ &= -\eta_2 + z\eta_1 \end{aligned}$$



□

Next, we integrate  $\zeta(z)$  and take exponential, thus the resulting  $2\pi i$ 's from integrating  $\zeta(z)$  won't affect the result:

$$e^{\int \zeta(z) dz} = z \prod_{w \in \mathbb{Z}^*} (z+w) e^{(-\frac{z}{w} + \frac{z^2}{2w^2})}$$

which has convergence problem again. Instead:

$$\text{Define } \sigma(z) \triangleq z \prod_{w \in \mathbb{Z}^*} (1 + \frac{z}{w}) e^{(-\frac{z}{w} + \frac{z^2}{2w^2})}$$

Here we factored out  $\prod_{w \in \mathbb{Z}^*} w$ , a "constant", which made the whole thing diverge.

The function converges now since if we take a cut on  $\mathbb{C}$ , take log and obtain

$$\begin{aligned} \log z + \sum_{w \in \mathbb{Z}^*} (\log(1 + \frac{z}{w}) - \frac{z}{w} + \frac{z^2}{2w^2}) &= \log z + \sum_{w \in \mathbb{Z}^*} (\frac{z}{w} - \frac{z^2}{2w^2} + O(\frac{1}{|w|^3}) - \frac{z}{w} + \frac{z^2}{2w^2}) \\ &= \log z + \sum_{w \in \mathbb{Z}^*} \underbrace{O(\frac{1}{|w|^3})}_{\text{converges!}} \end{aligned}$$

Clearly  $\sigma(z)$  is holomorphic on  $\mathbb{C}$ , with simple 0's at  $\mathbb{Z}$ . Observe that

$\sigma(z+1)/\sigma(z) = \zeta(z)$ , thus

$$\frac{\sigma(z+1)}{\sigma(z+1)} - \frac{\sigma(z)}{\sigma(z)} = \zeta(z+1) - \zeta(z) = \eta_1$$

$$\Rightarrow \log \sigma(z+1) - \log \sigma(z) = \eta_1 z + C_1 \pmod{2\pi i \mathbb{Z}}$$

$$\Rightarrow \sigma(z+1) = \sigma(z) e^{\eta_1 z + C_1}$$

Taking  $z = -\frac{1}{2}$ , and observing that  $\sigma(z)$  is odd, we have:

$$\sigma\left(\frac{1}{2}\right) = \sigma\left(-\frac{1}{2}\right) e^{-\frac{1}{2}\eta_1 + C_1} = -\sigma\left(\frac{1}{2}\right) e^{-\frac{1}{2}\eta_1 + C_1}$$

$$(\sigma\left(\frac{1}{2}\right) \neq 0) \Rightarrow \sigma(z+1) = -\sigma(z) e^{\eta_1(z+\frac{1}{2})}$$

Similarly, we have  $\sigma(z+2) = -\sigma(z) e^{\eta_1(z+\frac{3}{2})}$ .

Proof of Abel's Theorem (second proof)

"Given  $\sum_{i=1}^N A(P_i) = \sum_{j=1}^N A(Q_j) \Leftrightarrow \exists f$  meromorphic with zero's at  $P_i$  and poles at  $Q_j$ ".

With the  $\sigma(z)$  function, we may try  $f(z) = \frac{\prod_{i=1}^N (\sigma(z-P_i))}{\prod_{j=1}^N (\sigma(z-Q_j))}$  and see if they descend to a function on  $\hat{X}$ , i.e. if it's doubly periodic on  $\mathbb{C}$ :

$$? \begin{cases} f(z+1) = f(z) \\ f(z+2) = f(z) \end{cases}$$

$$\begin{aligned} \text{But } f(z+1) &= \frac{\prod_{i=1}^N (\sigma(z+1-P_i))}{\prod_{j=1}^N (\sigma(z+1-Q_j))} \\ &= \frac{[(-1)^N \prod_{i=1}^N \sigma(z-P_i) e^{\eta_1(z-P_i+\frac{1}{2})}]}{[(-1)^N \prod_{j=1}^N \sigma(z-Q_j) e^{\eta_1(z-Q_j+\frac{1}{2})}]} \\ &= \left( \frac{\prod_{i=1}^N \sigma(z-P_i)}{\prod_{j=1}^N \sigma(z-Q_j)} \right) e^{\eta_1(\sum Q_j - \sum P_i)} \end{aligned}$$

The Abel map in this case is  $A(P) = \int_0^P dz = P$ , thus  $\sum_{i=1}^N A(P_i) = \sum_{j=1}^N A(Q_j)$

$\Rightarrow \sum_{j=1}^N Q_j - \sum_{i=1}^N P_i = n + mz$ . It is not a priori true that  $\eta_1(n+mz) = 0$ .

The correct solution is thus change different  $P_i, Q_j$ 's, for instance we may take  $Q'_1 = Q_1 - n - mz$ ,  $Q'_i = Q_i, i=2, \dots, N$  and take


$$\bullet f(z) \triangleq \frac{\prod_{i=1}^N (\sigma(z-P_i))}{\prod_{j=1}^N (\sigma(z-Q'_j))}$$

which would then descend to  $\hat{X}$ .

Similar as in the previous section, we may produce:

- A form with a double pole at  $P$ :  $\omega_P(z) \triangleq \beta(z-P)dz$ . Since  $\beta(z)$  is already a meromorphic function on  $\mathbb{C}$  with a double pole at 0 which descends to  $\hat{X}$ .
- A form with 2 simple poles at  $P, Q$ :  $\omega_{PQ} \triangleq (\zeta(z-P) - \zeta(z-Q))dz$ . Although  $\zeta(z)$  is not a function on  $\hat{X}$ , the difference is. Since  $\zeta(z+1) - \zeta(z) = \eta_1$ ;  $\zeta(z+\tau) - \zeta(z) = \eta_2$ .

Recall that the Riemann surface was originally defined by  $w^2 = z(z-1)(z-\lambda)$



First: the lattice  $\mathcal{L}$  is given, we may express  $\beta(z)$  as a Laurent series:

$$\begin{aligned}
 \beta(z) &= \frac{1}{z^2} + \sum_{w \in \mathcal{L}^*} \left( \frac{1}{(w+z)^2} - \frac{1}{w^2} \right) \\
 &= \frac{1}{z^2} + \sum_{w \in \mathcal{L}^*} \left( \frac{1}{w^2} \left( \frac{1}{1+\frac{z}{w}} \right)^2 - \frac{1}{w^2} \right) \\
 &= \frac{1}{z^2} + \sum_{w \in \mathcal{L}^*} \left( \frac{1}{w^2} \sum_{k=0}^{\infty} (-1)^k (k+1) \left( \frac{z}{w} \right)^k - \frac{1}{w^2} \right) \\
 &= \frac{1}{z^2} + \sum_{w \in \mathcal{L}^*} \left( \sum_{\ell=1}^{\infty} (2\ell+1) \frac{z^{2\ell}}{w^{2\ell+2}} \right) \quad (\text{odd order terms get cancelled since}) \\
 &= \frac{1}{z^2} + \sum_{k=1}^{\infty} (2k+1) z^{2k} \sum_{w \in \mathcal{L}^*} \frac{1}{w^{2k+2}} \quad \mathcal{L}^* \text{ is symmetric: } w \in \mathcal{L}^* \Rightarrow -w \in \mathcal{L}^*
 \end{aligned}$$

Define  $G_k(\mathcal{L}) \triangleq \sum_{w \in \mathcal{L}^*} \frac{1}{w^{2k}}$ , the Eisenstein series, then:

$$\beta(z) = \frac{1}{z^2} + \sum_{k=1}^{\infty} (2k+1) G_{k+1} z^{2k}$$

Differentiate, we obtain:

$$\begin{aligned}
 \beta'(z) &= -\frac{2}{z^3} + \sum_{k=1}^{\infty} (2k+1) 2k G_{k+1} z^{2k-1} \\
 &= -\frac{2}{z^3} + 6G_2 z + 20G_3 z^3 + O(z^5)
 \end{aligned}$$

$$\Rightarrow (\beta'(z))^2 = \frac{4}{z^6} - \frac{24G_2}{z^2} - 80G_3 + O(z^2)$$

And we have:

$$\begin{aligned}
 \beta(z)^3 &= \left( \frac{1}{z^2} + 3G_2 z^2 + 5G_3 z^4 + O(z^6) \right)^3 \\
 &= \left( \frac{1}{z^2} + 6G_2 + 10G_3 z^2 + O(z^4) \right) \left( \frac{1}{z^2} + 3G_2 z^2 + 5G_3 z^4 + O(z^6) \right) \\
 &= \frac{1}{z^6} + 9G_2 \frac{1}{z^2} + 15G_3 + O(z^2)
 \end{aligned}$$

$$\Rightarrow (\beta'(z))^2 - 4(\beta(z))^3 = -60G_2 \frac{1}{z^2} - 140G_3 + O(z^2)$$

$$\Rightarrow (\beta'(z))^2 - 4(\beta(z))^3 + 60G_2 \beta(z) + 140G_3 = O(z^2) = 0 \quad (\text{Liouville's theorem})$$

Define  $g_2 = 60G_2$ ,  $g_3 = 140G_3$ , then we conclude that:

$$(\beta'(z))^2 = 4\beta(z)^3 - g_2\beta(z) - g_3$$

Compared with  $w^2 = z(z-1)(z-\lambda)$ , which may be changed into the form

$$w^2 = 4z^3 - g_2z - g_3$$

by a linear transformation. Hence the inverse map is given by:

$$z = \beta(s) \quad w = \beta'(s)$$

Remark: the Weierstrass  $\beta(z)$  function answers an old question from calculus:

What is an elliptic integral?

$$z = \int^z \frac{\beta'}{\sqrt{4\beta^3 - g_2\beta - g_3}} dz = \int^{\beta(z)} \frac{du}{\sqrt{4u^3 - g_2u - g_3}} = E(\beta(z))$$

i.e.  $E$  (elliptic integral) is the inverse of Weierstrass function. Compare with the more familiar:

$$\int^u \frac{du}{\sqrt{1-u^2}} = \arcsin u, \quad \text{and} \quad \arcsin(\sin z) = z$$

### • Function Theory According To Jacobi

- Weierstrass: easy to follow, not easy to generalize to other R.S.
- Jacobi: generalize well to the other R.S.'s. The key notion is the "theta functions"
- Similar as above, we need to construct:
  - 1). A holomorphic form  $\omega = dz$
  - 2). Meromorphic forms with simple poles at  $P \neq Q$ :  $\omega_{PQ}$ .
  - 3). Meromorphic forms with a double pole at  $P$ :  $\omega_P$

Fix  $\mathbb{C}/\mathbb{Z} + \mathbb{Z}z$ : ( $\text{Im}z > 0$ )

- Define  $\Theta(z|z) = \sum_{n=-\infty}^{\infty} e^{\pi i n^2 z + 2\pi i n z}$ ,  $z \in \mathbb{C}$

This series converges for any  $z$ , because  $|e^{\pi i n^2 z + 2\pi i n z}| = e^{-\pi n^2 \text{Im}z - 2\pi n \text{Im}z}$  decays like a Gaussian w.r.t.  $|n|$ , and  $\Theta(z|z)$  is holomorphic in  $z$  (and  $z$ ).

Key Transformations:

(I).  $\Theta(z+1|z) = \Theta(z|z)$ .

(II).  $\Theta(z+z|z) = e^{-2\pi i z - \pi i z} \Theta(z|z)$ . Note in particular that  $-2\pi i z - \pi i z$  is linear in  $z$ .

$$\begin{aligned}
 \text{Pf: } \theta(z+z) &= \sum_{n=-\infty}^{\infty} e^{i\pi n^2 z + 2\pi i n(z+z)} = \sum_{n=-\infty}^{\infty} e^{i\pi z(n^2+2n+1) + 2\pi i n z} \\
 &= \sum_{n=-\infty}^{\infty} e^{i\pi z(n+1)^2 - i\pi z - 2\pi i z + 2\pi i(n+1)z} \\
 &= e^{-2\pi i z - \pi i z} \sum_{n=-\infty}^{\infty} e^{i\pi z(n+1)^2 + 2\pi i(n+1)z} \\
 &= e^{-2\pi i z - \pi i z} \theta(z)
 \end{aligned}$$


Thm.  $\theta(z|z) = 0 \Leftrightarrow z = \frac{1+z}{2} \pmod{\mathcal{L}}$ , where  $\mathcal{L} = \mathbb{Z} + \mathbb{Z}z$

Pf: We count the number of zero's inside a fundamental parallelogram.

i.e. we compute the integral  $\frac{1}{2\pi i} \oint_{\partial \mathcal{R}_{out}} \frac{\theta'(z)}{\theta(z)} dz = \# \{ \text{zero's} \}$

inside a fundamental parallelogram.

$$\begin{aligned}
 \frac{1}{2\pi i} \oint_{\partial \mathcal{R}_{out}} \frac{\theta'(z)}{\theta(z)} dz &= \frac{1}{2\pi i} \left\{ \int_0^1 \frac{\theta'}{\theta} dz + \int_1^{1+z} \frac{\theta'}{\theta} dz - \int_z^{z+1} \frac{\theta'}{\theta} dz - \int_0^z \frac{\theta'}{\theta} dz \right\} \\
 &= \frac{1}{2\pi i} \left\{ \int_0^1 \left( \frac{\theta'}{\theta}(z) - \frac{\theta'}{\theta}(z+z) \right) dz - \int_0^z \left( \frac{\theta'}{\theta}(z) - \frac{\theta'}{\theta}(z+1) \right) dz \right\}
 \end{aligned}$$



Now by (I) and (II).  $\theta'/\theta(z) - \theta'/\theta(z+z) = (\ln \theta(z))' - (\ln \theta(z+z))' = 2\pi i$

$$\frac{\theta'}{\theta}(z) - \frac{\theta'}{\theta}(z+1) = (\ln \theta(z))' - (\ln \theta(z+1))' = 0$$

$$\Rightarrow \frac{1}{2\pi i} \oint_{\partial \mathcal{R}_{out}} \frac{\theta'}{\theta} dz = 1$$

To find where the zero is, consider  $\tilde{\theta}(z|z) \triangleq \theta(z + \frac{1+z}{2} | z)$ .

$$\begin{aligned}
 \tilde{\theta}(z|z) &= \sum_{n=-\infty}^{\infty} e^{i\pi n^2 z + 2\pi i n(z + \frac{1+z}{2})} = \sum_{n=-\infty}^{\infty} e^{i\pi n^2 z + i\pi n z + \pi i n + 2\pi i n z} \\
 &= \sum_{n=-\infty}^{\infty} e^{i\pi z(n^2+n+\frac{1}{4}) - \frac{1}{4}i\pi z + 2\pi i(n+\frac{1}{2})z - \pi i z + \pi i n} \\
 &= e^{-\frac{1}{4}i\pi z - \pi i z} \sum_{n=-\infty}^{\infty} e^{i\pi z(n+\frac{1}{2})^2 + 2\pi i(n+\frac{1}{2})z} \cdot (-1)^n
 \end{aligned}$$

$$\text{Evaluated at } 0, \text{ we have: } \tilde{\theta}(0|z) = e^{-\frac{1}{4}i\pi z} \sum_{n=-\infty}^{\infty} e^{\frac{i\pi z}{4}(2n+1)^2} (-1)^n$$

But  $(2n+1)^2 = (2k+1)^2$  ( $k \geq 0$ )  $\Rightarrow 2n+1 = \pm(2k+1) \Rightarrow n=k$  or  $-k-1$ , which are of different parity.  $\Rightarrow \tilde{\theta}(0|z) = 0$ . The result follows.  $\square$

Rmk: The factor of  $\theta(z+z|z) = e^{-2\pi i z - \pi i z} \cdot \theta(z|z)$  shows that  $\theta(z|z)$  is a section of a line bundle  $\mathcal{L}$  on  $\hat{X}$ , and the above computation shows that  $C_1(\mathcal{L}) = 1$ .

Rmk: Define  $\theta_1(z|z) \triangleq e^{\pi i \frac{z}{4} + \pi i(z+\frac{1}{2})} \theta(z + \frac{1+z}{2} | z)$ , then  $\theta_1$  is an odd function.

This again explains that  $\theta(z + \frac{1+z}{2} | z)$  has a zero at 0. Moreover  $\theta_1$  satisfies the following transformation relations:

$$(III). \quad \Theta_1(z+1|z) = -\Theta_1(z|z)$$

$$(IV). \quad \Theta_1(z+z|z) = -e^{-i\pi z - 2\pi iz} \Theta_1(z|z)$$

Pf: By definition,  $\Theta_1(z) = e^{\frac{\pi i}{2}} \sum_{n=-\infty}^{\infty} e^{i\pi z(n+\frac{1}{2})^2 + 2\pi i(n+\frac{1}{2})z + n\pi i}$   
 $= \sum_{n=-\infty}^{\infty} e^{i\pi z(n+\frac{1}{2})^2 + 2\pi i(n+\frac{1}{2})z + \pi i(n+\frac{1}{2})}$

Thus  $\Theta_1(-z) = \sum_{n=-\infty}^{\infty} e^{i\pi z(-(n+\frac{1}{2}))^2 + 2\pi i(-(n+\frac{1}{2}))z + \pi i(-(n+\frac{1}{2})) + (2n+1)\pi i}$   
 $= (-1) \sum_{k=-\infty}^{\infty} e^{i\pi z(k+\frac{1}{2})^2 + 2\pi i(k+\frac{1}{2})z + \pi i(k+\frac{1}{2})}$   
 $= -\Theta_1(z)$

For the transformation relations, we note that

$$\Theta_1(z) = e^{\frac{1}{4}\pi z + \pi i(z+\frac{1}{2})} \tilde{\Theta}(z|z) = e^{\frac{\pi i}{2}} \sum_{n=-\infty}^{\infty} e^{i\pi z(n+\frac{1}{2})^2 + 2\pi i(n+\frac{1}{2})z} (-1)^n$$

Hence:  $\Theta_1(z+1) = e^{\frac{\pi i}{2}} \sum_{n=-\infty}^{\infty} e^{i\pi z(n+\frac{1}{2})^2 + 2\pi i(n+\frac{1}{2})z + 2\pi i(n+\frac{1}{2})} (-1)^n$   
 $= -e^{\frac{\pi i}{2}} \sum_{n=-\infty}^{\infty} e^{i\pi z(n+\frac{1}{2})^2 + 2\pi i(n+\frac{1}{2})z} (-1)^n = -\Theta_1(z)$

$\Theta_1(z+z) = e^{\frac{\pi i}{2}} \sum_{n=-\infty}^{\infty} e^{i\pi z(n+\frac{1}{2})^2 + 2\pi i(n+\frac{1}{2})(z+z)} (-1)^n$   
 $= e^{\frac{\pi i}{2}} \sum_{n=-\infty}^{\infty} e^{i\pi z(n+\frac{1}{2})^2 + 2\pi i z(n+\frac{1}{2}) + \pi i z - \pi i z + 2\pi i(n+\frac{1}{2}+1)z - 2\pi i z} (-1)(-1)^{n+1}$   
 $= -e^{\frac{\pi i}{2}} \left( \sum_{n=-\infty}^{\infty} e^{i\pi z(n+\frac{1}{2})^2 + 2\pi i(n+\frac{1}{2}+1)z} (-1)^{n+1} \right) \cdot e^{-\pi i z - 2\pi i z}$   
 $= -e^{-\pi i z - 2\pi i z} \Theta_1(z)$

□

Proof of Abel's Theorem (third proof)

"Given  $\sum_{i=1}^N A(P_i) = \sum_{j=1}^N A(Q_j) \Leftrightarrow \exists f$  meromorphic with zero's at  $P_i$  and poles at  $Q_j$ ".

Now given  $\sum P_i = \sum Q_j + n + mz$ , and to construct  $f$ , we now use  $\Theta$ -function:

For a first attempt, try  $f = \left( \prod_{i=1}^N \Theta_1(z - P_i) \right) / \left( \prod_{j=1}^N \Theta_1(z - Q_j) \right)$

$$f(z+1) = \frac{\prod_{i=1}^N \Theta_1(z+1 + \frac{1+z}{2} - P_i)}{\prod_{j=1}^N \Theta_1(z+1 + \frac{1+z}{2} - Q_j)} = \frac{(-1)^N \prod_{i=1}^N \Theta_1(z + \frac{1+z}{2} - P_i)}{(-1)^N \prod_{j=1}^N \Theta_1(z + \frac{1+z}{2} - Q_j)} = f(z)$$

$$f(z+z) = \frac{\prod_{i=1}^N \Theta_1(z+z + \frac{1+z}{2} - P_i)}{\prod_{j=1}^N \Theta_1(z+z + \frac{1+z}{2} - Q_j)}$$

$$= \frac{(-1)^N \prod_{i=1}^N \Theta_1(z + \frac{1+z}{2} - P_i) \cdot e^{-2\pi i(z-P_i) - \pi iz}}{(-1)^N \prod_{j=1}^N \Theta_1(z + \frac{1+z}{2} - Q_j) \cdot e^{-2\pi i(z-Q_j) - \pi iz}}$$

$$= e^{2\pi i(\sum Q_j - \sum P_i)} f(z)$$

So again we may replace one of the points, say,  $Q_1$  by  $\tilde{Q}_1 + n + mz$ , so that  $\sum P_i = \sum Q_j$ , and the result follows.  $\square$

Meromorphic Forms:

- We can also construct  $\omega_{pq}(z)$ , a meromorphic form with simple poles at  $p \neq q$  as follows:

Since  $\theta_1$  satisfies the transformation rules III and IV,  $(\ln \theta_1)' = \theta_1' / \theta_1$  will be a meromorphic function with a simple pole at the origin, and furthermore it satisfies:

$$\theta_1'(z+1) / \theta_1(z+1) = \theta_1'(z) / \theta_1(z)$$

$$\theta_1'(z+\tau) / \theta_1(z+\tau) = -2\pi i + \theta_1'(z) / \theta_1(z)$$

$$\Rightarrow \frac{\theta_1'(z-P)}{\theta_1(z-P)} - \frac{\theta_1'(z-Q)}{\theta_1(z-Q)}$$

is doubly periodic with simple poles at  $P$  and  $Q$ , and thus defines the required function on  $\hat{X}$ .

- To construct  $\omega_p$  with a double pole at  $P$ , note that:

$(\ln \theta_1)''(z) = (\frac{\theta_1''}{\theta_1})'(z)$  has a double pole at  $P$  and is doubly periodic on  $\mathbb{C}$ , thus it suffices to consider  $-(\frac{d^2}{dz^2} \ln \theta_1)(z-P)$ .

- We can also connect Jacobi's theory with Weierstrass's theory, and we have the following identities.

$$\beta(z) = -\frac{d^2}{dz^2} \log \theta_1(z) + c(z)$$

Indeed,  $\beta(z) - (-\frac{d^2}{dz^2} \log \theta_1(z))$  is holomorphic on  $\mathbb{C}$  and doubly periodic.

$$\sigma(z) = e^{\frac{1}{2} \eta_1 z^2} \theta_1(z|z) / \theta_1(0|z)$$

Pf: We first show that:  $\frac{\sigma'(z)}{\sigma(z)} = \eta_1 z + \frac{\theta_1'(z)}{\theta_1(z)}$  ..

Indeed, we have  $\frac{\sigma'(z)}{\sigma(z)} = \zeta(z)$  and  $\zeta(z+1) = \zeta(z) + \eta_1$ ,  $\zeta(z+\tau) = \zeta(z) + \eta_2$ ,  $2\eta_1 - \eta_2 = 2\pi i$

as proved before, similarly we have:

$$\begin{cases} \frac{\theta_1'}{\theta_1}(z+1) + \eta_1(z+1) - \frac{\theta_1'}{\theta_1}(z) - \eta_1 z = \eta_1 \\ \frac{\theta_1'}{\theta_1}(z+\tau) + \eta_1(z+\tau) - \frac{\theta_1'}{\theta_1}(z) - \eta_1 z = -2\pi i + \eta_1 \tau = \eta_2 \end{cases}$$

$\Rightarrow (\eta_1 z - \frac{\theta_1'}{\theta_1}(z)) - \zeta(z)$  is doubly periodic and holomorphic, thus constant

But we also know that  $\zeta(z) \rightarrow \frac{1}{z}$  and  $\eta_1 z - \frac{\theta_1'}{\theta_1}(z) \rightarrow \frac{1}{z}$  where  $z \rightarrow 0$  and both have no constant terms.

$$\Rightarrow \eta_1 z - \frac{\theta_1'}{\theta_1}(z) = \zeta(z).$$

It follows that  $C \cdot \sigma(z) = e^{\frac{1}{2}\eta_1 z^2} \cdot \theta_1(z|z)$ . To specify  $C$ , we note that  $\sigma'(0) = 1$  and  $\theta_1(z|z)(0) = 0 \Rightarrow C = \lim_{z \rightarrow 0} e^{\frac{1}{2}\eta_1 z^2} \cdot \theta_1(z|z) / \sigma(z) = \theta_1'(0|z)$ .

The formula follows. □

### Special Properties of $\theta$ -Functions (two important formulae)

- Product representation: set  $q = e^{\pi i z}$

$$\theta(z|z) = \prod_{n=1}^{\infty} (1 - q^{2n}) \prod_{m=0}^{\infty} (1 + q^{2m+1} e^{2\pi i z}) (1 + q^{2m+1} e^{-2\pi i z}), \quad (1)$$

- Observation: since  $z, -\frac{1}{z}$  are parameters for the same lattice (differ by an  $SL(2, \mathbb{Z})$ -action:  $\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix}$ ), there should be some relation between  $\theta(z|z)$  and  $\theta(z|-\frac{1}{z})$ . Indeed, there is the **modular transformation law**:

$$\theta(z|-\frac{1}{z}) = \sqrt{\frac{z}{i}} e^{i\pi z z^2} \theta(zz|z) \quad (2)$$

where  $\sqrt{t}$  is the main branch which is positive for  $t > 0$ . In particular:

$$\theta(0|-\frac{1}{z}) = \sqrt{\frac{z}{i}} \theta(0|z) \quad (\text{duality})$$

Proof of the product presentation.

Step 1: Denote the R.H.S. of (1) by  $T(z|z)$ , we will first show that

$$\theta(z|z) = C(q) T(z|z) \text{ for some constant } C(q). \quad (1)'$$

First of all,  $T$  vanishes when  $1 + q e^{-2\pi i z} = 0$ , or  $e^{\pi i - 2\pi i z} = e^{\pi i}$ . In particular it vanishes when  $z = \frac{1+z}{2}$ .

Secondly,  $T$  transforms similarly as  $\theta$  does under translations by  $\mathbb{Z}$ , i.e.

$$\begin{cases} T(z+1|z) = T(z|z) & (3) \\ T(z+z|z) = e^{-2\pi i z - \pi i z} T(z|z) & (4) \end{cases}$$

Then it follows that  $T$  vanishes at every  $\frac{1+z}{2} + \mathbb{Z}$  and  $\frac{T}{\theta}$  is a holomorphic, doubly periodic function on  $\mathbb{C}$ , thus must be constant.

Now let's check (3) and (4):

(3) follows easily since it's invariant termwise.



$$\begin{aligned}
(4): T(z+z|z) &= \prod_{n=1}^{\infty} (1-q^{2n})(1+q^{2n-1}e^{2\pi i(z+z)})(1+q^{2n-1}e^{-2\pi i(z+z)}), \\
&= \prod_{n=1}^{\infty} (1-q^{2n})(1+q^{2n+1}e^{2\pi iz})(1+q^{2n-3}e^{-2\pi iz}), \\
&= \left( \prod_{n=1}^{\infty} (1-q^{2n})(1+q^{2n-1}e^{2\pi iz})(1+q^{2n-1}e^{-2\pi iz}) \right) (1+q^{-1}e^{-2\pi iz}) / (1+qe^{2\pi iz}), \\
&= T(z|z)(qe^{2\pi iz})^{-1} \left( \frac{1+q^{-1}e^{-2\pi iz}}{1+qe^{2\pi iz}} \right) \\
&= e^{-2\pi iz - \pi iz} T(z|z)
\end{aligned}$$

Step 2. Firstly we shall show that  $c(q) = 1$ .

Claim:  $c(q) = c(q^4)$ . (5)

Then it follows that  $c(q) = c(q^4) = c(q^{16}) = \dots = c(q^{n^2}) = \dots$ . Since  $\text{Im}z > 0$ ,  $|q^{n^2}| = e^{n^2\pi \text{Im}z} < 1$   
 $\Rightarrow c(q) = \lim_{q \rightarrow 0} c(q)$ . But in both cases,  $\Theta(0|z) \rightarrow 1, T(0|z) \rightarrow 1$ , when  $n^2 \text{Im}z \rightarrow \infty$   
 $\Rightarrow c(q) = \lim_{q \rightarrow 0} c(q) = 1$ .

Now we prove (5). Apply (1)' at  $z = \frac{1}{2}$ , we obtain that:

$$\begin{aligned}
\Theta\left(\frac{1}{2}|z\right) &= \sum_{n \in \mathbb{Z}} q^{n^2} (-1)^n, \\
\text{and } T\left(\frac{1}{2}|z\right) &= \prod_{n \geq 1} (1-q^{2n})(1-q^{2n-1})^2 = \prod_{n \geq 1} (1-q^n)(1-q^{2n-1}) \\
\Rightarrow c(q) &= \left( \sum_{n \in \mathbb{Z}} q^{n^2} (-1)^n \right) / \prod_{n \geq 1} (1-q^n)(1-q^{2n-1}). \quad (6)
\end{aligned}$$

Again apply (1)' at  $z = \frac{1}{4}$  we obtain that:

$$\Theta\left(\frac{1}{4}|z\right) = \sum_{n \in \mathbb{Z}} q^{n^2} i^n$$

But observe that  $n = \pm(2k+1), k \in \mathbb{N} \Rightarrow q^{(2k+1)^2} i^{2k+1} + q^{-(2k+1)^2} i^{-(2k+1)} = q^{2k+1} (-1)^k (i + i^{-1}) = 0$

$$\Rightarrow \Theta\left(\frac{1}{4}|z\right) = \sum_{n \text{ even}} q^{n^2} i^n = \sum_{n \geq 1} q^{4n^2} (-1)^n$$

$$\begin{aligned}
T\left(\frac{1}{4}|z\right) &= \prod_{n \geq 1} (1-q^{2n})(1+q^{2n-1}i)(1-q^{2n-1}i) \\
&= \prod_{n \geq 1} (1-q^{2n})(1+q^{4n-2}) \\
&= \prod_{n \geq 1} (1-q^{4n})(1-q^{4n-2})(1+q^{4n-2}) \\
&= \prod_{n \geq 1} (1-q^{4n})(1-q^{8n-4}) \\
&= \prod_{n \geq 1} (1-(q^4)^n)(1-(q^4)^{2n-1})
\end{aligned}$$

$$\Rightarrow c(q) = \left( \sum_{n \geq 1} (q^4)^n (-1)^n \right) / \prod_{n \geq 1} (1-(q^4)^n)(1-(q^4)^{2n-1}) \quad (7)$$

Compare (6) and (7), we obtain that  $c(q) = c(q^4)$  as asserted, and finishes the proof of the product presentation.  $\square$

Proof of the modular transformation.

Note that  $\Theta(z|\tau)$  is holomorphic in both  $z$  and  $\tau$ , it suffices to prove for  $z \in \mathbb{R}$  and  $\tau = i\tau_2, \tau_2 > 0$ .

Now the R.H.S. of (2) reads:

$$\begin{aligned} \sqrt{\tau_2} e^{-\pi\tau_2 z^2} \Theta(iz\tau_2 | i\tau_2) &= \sqrt{\tau_2} e^{-\pi\tau_2 z^2} \sum_{n \in \mathbb{Z}} e^{-\pi\tau_2 n^2 + 2\pi i(iz\tau_2)n} \\ &= \sqrt{\tau_2} e^{-\pi\tau_2 z^2} \sum_{n \in \mathbb{Z}} e^{-\pi\tau_2 n^2 - 2\pi\tau_2 z n} \\ &= \sqrt{\tau_2} \sum_{n \in \mathbb{Z}} e^{-\pi\tau_2 (z+n)^2} \end{aligned}$$

Thus (2)  $\Leftrightarrow \sum_{n \in \mathbb{Z}} e^{-\frac{\pi}{\tau_2} n^2 + 2\pi i n z} = \sqrt{\tau_2} \sum_{n \in \mathbb{Z}} e^{-\pi\tau_2 (z+n)^2}$ , which is actually a special case of the Poisson summation formula:

Thm. (**Poisson Summation Formula**) Let  $f$  be a smooth, rapidly decaying function. Define the Fourier transform  $\hat{f}(\xi) = \int_{\mathbb{R}} e^{-2\pi i x \xi} f(x) dx$ . Then we have:

$$\sum_{n \in \mathbb{Z}} f(\theta + n) = \sum_{n \in \mathbb{Z}} e^{2\pi i n \theta} \hat{f}(n) \quad \forall \theta \in \mathbb{R}.$$

In particular, let  $\theta = 0$ , then we have:

$$\sum_{n \in \mathbb{Z}} f(n) = \sum_{n \in \mathbb{Z}} \hat{f}(n)$$

"Rapidly decaying" guarantees that  $\sum_{n \in \mathbb{Z}} f(\theta + n)$  converges for all  $\theta \in \mathbb{R}$ . For example take the Gaussian:  $f = e^{-\frac{\pi}{\tau_2} x^2}$ ,  $\hat{f}(\xi) = \sqrt{\tau_2} e^{-2\pi \tau_2 \xi^2}$ . Thus apply this formula to our problem:  $f = e^{-\pi\tau_2 z^2}$ ,  $z \in \mathbb{R} \Rightarrow \hat{f}(\xi) = \frac{1}{\sqrt{\tau_2}} e^{-\frac{\pi \xi^2}{\tau_2}}$

$$\begin{aligned} \Rightarrow \sum_{n \in \mathbb{Z}} e^{-\pi\tau_2 (z+n)^2} &= \sum_{n \in \mathbb{Z}} e^{2\pi i n z} \frac{1}{\sqrt{\tau_2}} e^{-\frac{\pi n^2}{\tau_2}} \\ \Rightarrow \sqrt{\tau_2} \sum_{n \in \mathbb{Z}} e^{-\pi\tau_2 (z+n)^2} &= \sum_{n \in \mathbb{Z}} e^{-\frac{\pi n^2}{\tau_2} + 2\pi i n z}, \text{ as desired.} \end{aligned}$$

Proof of Poisson summation formula.

For a rapidly decaying function  $f$ , we may define  $\psi(\theta) = \sum_{n \in \mathbb{Z}} f(n+\theta)$ . Moreover  $\psi(\theta+1) = \psi(\theta)$ . Thus  $\psi$  is a smooth periodic function, which can be expanded into Fourier series:  $\psi(\theta) = \sum_n c_n e^{2\pi i n \theta}$ , where  $c_n = \int_0^1 \psi(\theta) e^{-2\pi i n \theta} d\theta$

$$\begin{aligned} \text{But } c_n &= \int_0^1 e^{-2\pi i n \theta} \left( \sum_{k \in \mathbb{Z}} f(k+\theta) \right) d\theta = \sum_{k \in \mathbb{Z}} \int_0^1 e^{-2\pi i n \theta} f(k+\theta) d\theta \\ &= \sum_{k \in \mathbb{Z}} \int_k^{k+1} e^{-2\pi i n \theta} f(\theta) d\theta = \int_{\mathbb{R}} e^{-2\pi i n \theta} f(\theta) d\theta = \hat{f}(n) \end{aligned}$$


It follows that:  $\psi(\theta) = \sum_{n \in \mathbb{Z}} \hat{f}(n) e^{2\pi i n \theta}$ . Hence by definition of  $\psi$  we have:

$$\sum_{n \in \mathbb{Z}} f(n+\theta) = \sum_{n \in \mathbb{Z}} \hat{f}(n) e^{2\pi i n \theta}.$$

□

• Function Theory Using PDE

Goal: To construct holomorphic and meromorphic forms on a surface  $\hat{X}$ , on which every point has locally holomorphic coordinates.




Idea: Suppose we want to construct a form  $\omega_p$  meromorphic on  $\hat{X}$  with a double pole at  $p$ . Using local coordinate charts, we have:

$$\begin{array}{ccc} \text{neighborhood } V \text{ of } p & \longleftrightarrow & \text{Disk } D \text{ in } \mathbb{C} \\ p & \longleftrightarrow & 0 \\ z & \longleftrightarrow & t(z) \end{array}$$

(1). Take  $\frac{dt}{t^2}$  to obtain a form  $\tilde{\omega}$  on  $V$ .

(2). Extend  $\tilde{\omega}$  to  $\hat{X}$  by considering  $\chi \cdot \tilde{\omega}$ , where  $\chi \in C_0(V)$ ,  $\chi \equiv 1$  in a smaller neighborhood of  $p$  in  $V$ . But  $\chi \cdot \tilde{\omega}$  is no longer meromorphic on  $\hat{X}$ . We shall correct  $\chi \tilde{\omega}$  by subtracting a form  $\psi$ , so that  $\bar{\partial}\psi = \bar{\partial}(\chi \tilde{\omega})$ , where  $\bar{\partial} = \frac{\partial}{\partial \bar{z}}$ .



More precisely, we proceed as follows:

Thm. (Hodge Decomposition - a simple version). Let  $\Phi$  be any smooth  $(0,1)$ -form on  $\hat{X}$ , locally  $\Phi = \Phi(z) d\bar{z}$ . Then there exists a smooth function  $f \in C^\infty(\hat{X})$  and a  $(0,1)$ -form  $\Phi_0$  s.t.

$$\Phi = \frac{\partial f}{\partial \bar{z}} d\bar{z} + \Phi_0 \quad \text{and} \quad \frac{\partial \Phi_0}{\partial \bar{z}} = 0$$

Observation:  $\bar{\Phi}_0$  is a holomorphic  $(1,0)$ -form and we shall show that the space of holomorphic  $(1,0)$ -forms is finite dimensional.

Assuming the Hodge decomposition thm, consider the following form

$$\Phi \triangleq \frac{\partial}{\partial \bar{z}} (\chi(z) \frac{1}{z}) d\bar{z} \in C^\infty(\hat{X} \setminus \{p\}).$$

on  $\hat{X}$ . Since  $\frac{\partial}{\partial \bar{z}} (\frac{1}{z}) = 0$  for  $z \neq p$ , we may just extend it by 0 at  $p$ , and 0 outside  $V$ , then it's a smooth  $(0,1)$ -form on  $\hat{X}$ .

By Hodge decomposition, we may write it as  $\bar{\Phi} = \partial_{\bar{z}} f \cdot d\bar{z} + \bar{\Phi}_0$ , with  $\bar{\Phi}_0$  an anti-holomorphic  $(0,1)$ -form. Now we try the form  $\omega \triangleq \partial_{\bar{z}}(\chi \cdot \frac{1}{z} - f(z)) dz$ .


Claim:  $\omega$  is holomorphic.

This is because  $\frac{\partial}{\partial \bar{z}} \omega = \partial_{\bar{z}}(\partial_{\bar{z}}(\chi \cdot \frac{1}{z} - f(z))) dz = \partial_{\bar{z}}(\bar{\Phi} - \partial_{\bar{z}} f \cdot d\bar{z}) dz = \partial_{\bar{z}} \bar{\Phi}_0 dz = 0$ .

Moreover we may construct a form with simple poles at  $P \neq Q$ .

Observation: how can one construct a form with a single pole?


Naively, one may try  $\partial_{\bar{z}}(\chi \cdot \ln z)$ . But  $\ln z$  has singularities along a "long" cut, which is not compact, and the above "cut off" trick doesn't apply. (supp  $\chi$  is compact!)



However, if  $P \neq Q$  are inside a same coordinate chart, we may introduce a cut by connecting  $P$  and  $Q$ , which is compact and the "cut off" technique applies:

$$\bar{\Phi}_{PQ} \triangleq \partial_{\bar{z}}(\chi(z) \cdot \ln(\frac{z-P}{z-Q}))$$

If  $P \neq Q$  are not in the same chart, we may just take a sequence  $\{P_i\}_{i=0}^n$ ,  $P_0 = P$ ,  $P_n = Q$ , and each neighboring ones are in a same chart. Then apply the above construction to get  $\bar{\Phi}_{P_i P_{i+1}}$  and define  $\bar{\Phi}_{PQ} = \sum_{i=0}^{n-1} \bar{\Phi}_{P_i P_{i+1}}$



Proof of Hodge decomposition.

More general statements will be proven latter, which holds for any dimension. At the present time, we give an explicit proof for  $\mathbb{C}/\mathbb{Z} + \mathbb{Z}z$ , using explicit formula.

Observation: on  $\mathbb{C}/\mathbb{Z} + \mathbb{Z}z$ , forms can be identified with doubly periodic functions

on  $\mathbb{C}$  via the correspondence:  $\bar{\Phi}(z) d\bar{z} \leftrightarrow \bar{\Phi}(z)$ . Thus the problem becomes:

Given a function  $\bar{\Phi}$ ,  $?\exists f, \bar{\Phi} = \frac{\partial}{\partial \bar{z}} f$ .

1). Solving  $\bar{\partial}$ -equation in  $\mathbb{C}$

Let  $\Phi$  be a smooth function with compact support. Define  $f(z) \triangleq \frac{1}{2\pi i} \iint_{\mathbb{C}} \frac{\Phi(\xi)}{\xi - z} d\xi \wedge d\bar{\xi}$ .

Then  $\frac{\partial f}{\partial \bar{z}} = \Phi$ .

$$\begin{aligned}
 \text{pf: } \frac{\partial f}{\partial \bar{z}} &= \frac{1}{2\pi i} \frac{\partial}{\partial \bar{z}} \iint_{\mathbb{C}} \frac{\Phi(\xi)}{\xi - z} d\xi \wedge d\bar{\xi} \\
 &= \frac{1}{2\pi i} \frac{\partial}{\partial \bar{z}} \iint_{\mathbb{C}} \frac{\Phi(\omega+z)}{\omega} d\omega \wedge d\bar{\omega} \quad (\omega = \xi - z) \\
 &= \frac{1}{2\pi i} \iint_{\mathbb{C}} \frac{1}{\omega} \frac{\partial}{\partial \bar{z}} (\Phi(\omega+z)) d\omega \wedge d\bar{\omega} \\
 &= \frac{1}{2\pi i} \iint_{\mathbb{C}} \frac{1}{\omega} \frac{\partial}{\partial \bar{\omega}} (\Phi(\omega+z)) d\omega \wedge d\bar{\omega} \quad \left( \frac{\partial}{\partial \bar{z}} \Phi(z+\omega) = \frac{\partial}{\partial \bar{(z+\omega)}} \Phi(z+\omega) = \frac{\partial}{\partial \bar{\omega}} \Phi(z+\omega) \right) \\
 &= \frac{1}{2\pi i} \lim_{\epsilon \rightarrow 0} \iint_{|\omega| \geq \epsilon} \frac{1}{\omega} \frac{\partial \Phi}{\partial \bar{\omega}}(\omega+z) d\omega \wedge d\bar{\omega} \\
 &= \frac{1}{2\pi i} \lim_{\epsilon \rightarrow 0} \iint_{|\omega| \geq \epsilon} -d\left(\frac{1}{\omega} \Phi(\omega+z) d\omega\right) \\
 &= \frac{1}{2\pi i} \lim_{\epsilon \rightarrow 0} \oint_{|\omega|=\epsilon} \frac{1}{\omega} \Phi(\omega+z) d\omega \\
 &= \frac{1}{2\pi i} 2\pi i \Phi(z) \\
 &= \Phi(z). \quad \square
 \end{aligned}$$

Using this result, we may also solve the equation  $\Delta g = \Phi$ ,  $\Delta = \partial_z \partial_{\bar{z}}$ , where  $\Phi$  is a smooth function with compact support. Indeed, just take

$$g(z) = \frac{1}{2\pi i} \iint_{\mathbb{C}} \log|z-\omega|^2 \Phi(\omega) d\omega \wedge d\bar{\omega}$$

This is because:

$$\begin{aligned}
 \partial_z g &= \frac{1}{2\pi i} \iint_{\mathbb{C}} \frac{1}{z-\omega} \Phi(\omega) d\omega \wedge d\bar{\omega} \\
 \text{and } \partial_{\bar{z}} \partial_z g &= \Phi(z)
 \end{aligned}$$

2). The torus case:  $\mathbb{C}/\mathbb{Z} + \mathbb{Z}\tau$

• Question: Given a  $C^\infty$ -function  $\varphi$  on  $\mathbb{C}/\mathbb{Z} + \mathbb{Z}\tau$ , when can we solve:

- $\frac{\partial f}{\partial \bar{z}} = \varphi$  ( $f$  doubly periodic w.r.t.  $\mathbb{Z} + \mathbb{Z}\tau$ )
- $\Delta g = \varphi$  ( $\Delta = \partial_z \partial_{\bar{z}}$ , again  $g$  doubly periodic)

Since on  $\mathbb{C}$ ,  $\varphi \in C_0^\infty(\mathbb{C})$ , we can solve these equations by explicit formula:

- $f(z) = \frac{1}{2\pi i} \iint_{\mathbb{C}} \frac{\varphi(\omega)}{z-\omega} d\omega \wedge d\bar{\omega}$
- $g(z) = \frac{1}{2\pi i} \iint_{\mathbb{C}} (\log|z-\omega|^2) \varphi(\omega) d\omega \wedge d\bar{\omega}$


Returning to the torus case  $\mathbb{C}/\mathbb{Z} + \mathbb{Z}\tau$ , we try a formula of similar kind:

$$g(z) = \frac{1}{2\pi i} \iint_{\mathbb{C}/\mathbb{Z} + \mathbb{Z}\tau} G(z-\omega) \varphi(\omega) d\omega \wedge d\bar{\omega}.$$

Observations:

By  $\iint_{\mathbb{C}/\mathbb{Z}+\mathbb{Z}\tau}$ , we mean an integral over a fundamental domain  $D$ . Furthermore, since:

$$\iint_{D+m+n\tau} G(z-w)\varphi(w)dw\wedge d\bar{w} = \iint_D G(z-m-n\tau-w)\varphi(w'+m+n\tau)dw'\wedge d\bar{w}' \\ = \iint_D G(z-m-n\tau-w)\varphi(w')dw'\wedge d\bar{w}' ,$$



when  $G(z-w)$  is doubly periodic, the integral is doubly periodic and  $g(z)$  is well-defined. Thus we look for  $G(z)$  doubly periodic and  $G(z) \sim \log|z|^2$  for  $z \sim 0$ .

Try  $\log|\Theta_1(z|\tau)|^2$ . Note that  $\Theta_1(z|\tau) = \Theta_1(0|\tau) + z\Theta_1'(0|\tau) + z^2E(z)$ ,  $E(z)$  holo. or  $\Theta_1 = z(\Theta_1'(0|\tau) + zE(z))$  as  $\Theta_1(0|\tau) = 0$ . Thus we try  $\log\left|\frac{\Theta_1(z|\tau)}{\Theta_1'(0|\tau)}\right|^2$

Recall that:  $\Theta_1(z+1|\tau) = -\Theta_1(z|\tau)$

$$\Theta_1(z+\tau|\tau) = -e^{-\pi iz - 2\pi i z^2} \Theta_1(z|\tau)$$

Clearly:  $\begin{cases} \frac{|\Theta_1(z+1|\tau)|^2}{|\Theta_1'(0|\tau)|^2} = \frac{|\Theta_1(z|\tau)|^2}{|\Theta_1'(0|\tau)|^2} \\ \frac{|\Theta_1(z+\tau|\tau)|^2}{|\Theta_1'(0|\tau)|^2} = |e^{-\pi iz - 2\pi i z^2}|^2 \frac{|\Theta_1(z|\tau)|^2}{|\Theta_1'(0|\tau)|^2} = |e^{\pi z_2 + 2\pi y}|^2 \frac{|\Theta_1(z|\tau)|^2}{|\Theta_1'(0|\tau)|^2} , z = z_1 + iz_2, z = x + iy. \end{cases}$

$$\Rightarrow \begin{cases} \log \frac{|\Theta_1(z+1|\tau)|^2}{|\Theta_1'(0|\tau)|^2} = \log \frac{|\Theta_1(z|\tau)|^2}{|\Theta_1'(0|\tau)|^2} \\ \log \frac{|\Theta_1(z+\tau|\tau)|^2}{|\Theta_1'(0|\tau)|^2} = 2\pi z_2 + 4\pi y + \log \frac{|\Theta_1(z|\tau)|^2}{|\Theta_1'(0|\tau)|^2} \end{cases}$$

Thus we may try instead  $\log \frac{|\Theta_1(z|\tau)|^2}{|\Theta_1'(0|\tau)|^2} - \frac{2\pi}{\tau_2} y^2$ . Under the transformation  $z \mapsto z+\tau$  (or  $x+yi \mapsto x+z_1+(y+z_2)i$ ,  $y \mapsto y+z_2$ ):

$$\frac{2\pi}{\tau_2} y^2 \mapsto \frac{2\pi}{\tau_2} (y^2 + 2yz_2 + \tau_2^2) = \frac{2\pi}{\tau_2} y^2 + 4\pi y + 2\pi z_2$$


which cancels the extra factor.

Def: (Green's function on torus).  $G(z) = \log \frac{|\Theta_1(z|\tau)|^2}{|\Theta_1'(0|\tau)|^2} - \frac{2\pi}{\tau_2} (\text{Im } z)^2$ .

Set  $g(z) = \frac{1}{2\pi i} \iint_{\mathbb{C}/\mathbb{Z}+\mathbb{Z}\tau} G(z-w)\varphi(w)dw\wedge d\bar{w}$  (\*). We compute  $\Delta(g(z)) = \partial\bar{z}\partial z g(z)$

We shall bring ourselves back to the complex plane case as follows:

Fix an arbitrary  $z \in \mathbb{C}/\mathbb{Z}+\mathbb{Z}\tau$ , and let  $\chi_w \in C^\infty(\mathbb{C}/\mathbb{Z}+\mathbb{Z}\tau)$ ,  $\chi(w) \equiv 1$  in a neighborhood of  $z$ ;  $\chi(w) = 0$  outside a neighborhood of  $z$ :



$$\iint_{\mathbb{C}/\mathbb{Z}+\mathbb{Z}\tau} G(z-w)\varphi(w)dw\wedge d\bar{w} = \iint G(z-w)(\chi(w)\varphi(w))dw\wedge d\bar{w} + \iint G(z-w)(1-\chi(w))\varphi(w)dw\wedge d\bar{w}$$

$G(z-w)(1-\chi(w))\varphi(w)$  is  $C^\infty$  on  $\mathbb{C}/\mathbb{Z}+\mathbb{Z}\tau$  since singularity of  $G$  occurs only when  $w=z$   
 Thus:  $\Delta_z \iint_{\mathbb{C}/\mathbb{Z}+\mathbb{Z}\tau} G(z-w)(1-\chi)\varphi(w)dw\wedge d\bar{w} = \iint (\Delta_z G(z-w))(1-\chi)\varphi(w)dw\wedge d\bar{w}$   
 $= \iint -\frac{\pi}{z_2} (1-\chi(w))\varphi(w)dw\wedge d\bar{w}$

(In the last equality, we used the fact that:

$$\partial_z \partial_{\bar{z}} (G(z-w)) = \partial_z \partial_{\bar{z}} \left( \log \frac{|\Theta(z-w|z)|^2}{|\Theta'(0|z)|^2} - \frac{2\pi}{z_2} (\text{Im}(z-w))^2 \right) = -\frac{\pi}{z_2}.$$

To calculate  $\Delta_z \iint G(z-w)(\chi(w)\varphi(w))dw\wedge d\bar{w}$ , observe that for  $w \in \text{Supp } \chi$  and hence near  $z$ , we have  $|\frac{\Theta(z-w|z)}{\Theta'(0|z)}|^2 = |z-w|^2 |1+(z-w)h(z-w)|^2$ ,  $h$  holo.

$$\Rightarrow \log \left| \frac{\Theta(z-w|z)}{\Theta'(0|z)} \right|^2 = \log |z-w|^2 + \log |1+(z-w)h(z-w)|^2$$

$$\Rightarrow G(z-w) = \log |z-w|^2 + \log |1+(z-w)h(z-w)|^2 - \frac{2\pi}{z_2} \text{Im}(z-w)^2$$

$$\Rightarrow \Delta_z \iint G(z-w)(\chi(w)\varphi(w))dw\wedge d\bar{w} = \Delta_z \iint \log |z-w|^2 \chi \cdot \varphi(w)dw\wedge d\bar{w} + 0$$

$$+ \iint \Delta_z \left( -\frac{2\pi}{z_2} \text{Im}(z-w)^2 \right) \chi \varphi(w)dw\wedge d\bar{w}$$

and by the  $\mathbb{C}$ -case  $\Delta_z \iint \log |z-w|^2 \chi \cdot \varphi(w)dw\wedge d\bar{w} = \varphi(z)$

Altogether:

$$\Delta g = \varphi(z) - \frac{1}{z_2 i} \iint_{\mathbb{C}/\mathbb{Z}+\mathbb{Z}\tau} \varphi(w)dw\wedge d\bar{w} \quad (**)$$

and we make the following observations:

- If  $\iint_{\mathbb{C}/\mathbb{Z}+\mathbb{Z}\tau} \varphi(w)dw\wedge d\bar{w} = 0$ ,  $\Delta g = \varphi$  admits a solution and is given explicitly by the formula (\*).
- Furthermore, if the equation  $\Delta g = \varphi$  admits a solution,  $\iint_{\mathbb{C}/\mathbb{Z}+\mathbb{Z}\tau} \varphi dw\wedge d\bar{w} = 0$   
 Indeed  $\iint \varphi dz\wedge d\bar{z} = \iint \partial_z \partial_{\bar{z}} g(z) dz\wedge d\bar{z} = \iint d(\partial_{\bar{z}} g(z) d\bar{z}) = 0$ , by Stokes thm.
- The space  $\{\varphi \mid \Delta g = \varphi \text{ is solvable}\}$  has codimension 1 in  $C^\infty(\mathbb{C}/\mathbb{Z}+\mathbb{Z}\tau)$
- Furthermore,  $(**) \Rightarrow \varphi(z) = \partial_{\bar{z}} \partial_z g + \varphi_0 = \partial_{\bar{z}} f + \varphi_0$ ,  $f = \partial_z g$  and  $\varphi_0$  a constant.  
 This is exactly the Hodge decomposition thm on  $\mathbb{C}/\mathbb{Z}+\mathbb{Z}\tau$ .


## §4. General Theory

Def:  $X$  is called a Riemann surface (R.S.) if  $X = \cup_{\alpha} X_{\alpha}$  with the property

$\Phi_{\alpha}: X_{\alpha} \rightarrow D \subseteq \mathbb{C}$  ( $D$ : unit disk) and  $\forall \alpha, \beta$

$$\Phi_{\beta} \circ \Phi_{\alpha}^{-1}: \Phi_{\alpha}(X_{\alpha} \cap X_{\beta}) \rightarrow \Phi_{\beta}(X_{\alpha} \cap X_{\beta})$$

is holomorphic,  $1-1$ ,  $(\Phi_{\beta} \circ \Phi_{\alpha}^{-1})'(z) \neq 0$  and with holomorphic inverse.



For notational sanity, we will use  $X_{\alpha} \ni z \xrightarrow{\Phi_{\alpha}} z_{\alpha} \in D$ .

Def. of line bundles (generalization of differential forms)

A line bundle  $L$  is an assignment:  $L \leftrightarrow \{t_{\alpha\beta}\}$ , where  $t_{\alpha\beta}$  are functions on  $X_{\alpha} \cap X_{\beta}$ , satisfying the following cocycle conditions.

$$\begin{cases} t_{\alpha\beta} \cdot t_{\beta\gamma} \equiv t_{\alpha\gamma} & \text{on } X_{\alpha} \cap X_{\beta} \cap X_{\gamma} \\ t_{\alpha\alpha} = 1 \end{cases}$$

$L$  is said to be  $C^{\infty}$  if each  $t_{\alpha\beta}$  is smooth

$L$  is said to be holomorphic if each  $t_{\alpha\beta}$  is holomorphic

$L$  is said to be antiholomorphic if each  $t_{\alpha\beta}$  is antiholomorphic.

To a line bundle corresponds its space of smooth sections.  $L \leftrightarrow \Gamma(X, L)$ .

$$\Gamma(X, L) = \{(\varphi_{\alpha}) \mid \varphi_{\alpha}: X_{\alpha} \rightarrow \mathbb{C}, C^{\infty}\text{-function on } X_{\alpha} \text{ and } \varphi_{\alpha} = t_{\alpha\beta} \varphi_{\beta} \text{ on } X_{\alpha} \cap X_{\beta}\}$$

To a holomorphic line bundle there is also the space of holomorphic sections:

$$H^0(X, L) = \{(\varphi_{\alpha}) \mid \varphi_{\alpha} \text{ holomorphic on } X_{\alpha} \text{ and } \varphi_{\alpha} = t_{\alpha\beta} \varphi_{\beta} \text{ on } X_{\alpha} \cap X_{\beta}\}$$

Examples:

(1).  $L =$  the trivial bundle  $\leftrightarrow \{t_{\alpha\beta} \equiv 1, \forall \alpha, \beta\}$



Then  $\Gamma(X, L) = \{(\varphi_\alpha), \varphi_\alpha \text{ functions on } X_\alpha; \varphi_\alpha = \varphi_\beta \text{ on } X_\alpha \cap X_\beta\}$   
 $= \{ \text{all functions on } X \}$

(2). The canonical bundle  $\Lambda^{1,0}(X)$  (or  $K_X$ )

$\Lambda^{1,0}(X) = \{t_{\alpha\beta} = \frac{\partial z_\beta}{\partial z_\alpha} \text{ on } X_\alpha \cap X_\beta\}$ . The chain rule  $\Rightarrow \{t_{\alpha\beta}\}$  satisfy the cocycle condition  
 $\Gamma(X, L) = \{(\varphi_\alpha): \varphi_\alpha \text{ defined on } X_\alpha \text{ and } \varphi_\alpha = \frac{\partial z_\beta}{\partial z_\alpha} \varphi_\beta\}$   
 $= \{(\varphi_\alpha dz_\alpha), \varphi_\alpha dz_\alpha = \varphi_\beta \frac{\partial z_\beta}{\partial z_\alpha} dz_\alpha = \varphi_\beta dz_\beta \text{ on } X_\alpha \cap X_\beta\}$   
 $= \{ \text{Smooth } (1,0)\text{-forms on } X \}$

In general, for arbitrary complex manifold of dim  $n$ ,  $X = \cup_\alpha X_\alpha$  and  $\Phi_\alpha: X_\alpha \rightarrow D \subseteq \mathbb{C}^n$   
 $z \mapsto z_\alpha = (z_\alpha^j)_{j=1}^n \in \mathbb{C}^n$ , we may similarly define a rank  $n$  vector bundle  $\Lambda^{1,0}(X)$ :  
 $\Lambda^{1,0}(X) = \{t_{\alpha\beta} = (\frac{\partial z_\beta^j}{\partial z_\alpha^i})_{n \times n}$ , holomorphic functions taking value in  $GL_n(\mathbb{C})\}$ .

In  $\dim_{\mathbb{C}} X = 1$ , we will use the notation  $K_X$  synonymously.

(3). The holo. tangent bundle

$L \leftrightarrow \{t_{\alpha\beta} = (\frac{\partial z_\alpha}{\partial z_\beta})^{-1} = \frac{\partial z_\alpha}{\partial z_\beta} \text{ on } X_\alpha \cap X_\beta\}$   
 $\Rightarrow \Gamma(X, L) = \{(\varphi_\alpha): \varphi_\alpha = \frac{\partial z_\alpha}{\partial z_\beta} \varphi_\beta \text{ on } X_\alpha \cap X_\beta\}$   
 $= \{(\varphi_\alpha \frac{\partial}{\partial z_\alpha}), \varphi_\alpha \frac{\partial}{\partial z_\alpha} = \varphi_\beta \frac{\partial z_\alpha}{\partial z_\beta} \frac{\partial}{\partial z_\alpha} = \varphi_\beta \frac{\partial}{\partial z_\beta}\}$

More generally, given a bundle  $L$  on  $X$ ,  $L \leftrightarrow \{t_{\alpha\beta}\}$ , we can define another bundle  $L^k \leftrightarrow \{(t_{\alpha\beta})^k\}$ ,  $k \in \mathbb{Z}$ .

E.g.  $L = K_X$ ,  $L^3 \leftrightarrow \{(\frac{\partial z_\alpha}{\partial z_\beta})^3\}$  and  $\Gamma(X, L) = \{(\varphi_\alpha): \varphi_\alpha \text{ defined on } X_\alpha \text{ with } \varphi_\alpha (dz_\alpha)^3 = \varphi_\beta (dz_\beta)^3\}$ .

Covariant derivatives, metrics, and curvature on line bundles

Question: Can we differentiate sections of  $L$ ?

$\varphi \in \Gamma(X, L) \leftrightarrow \{(\varphi_\alpha(z)): \varphi_\alpha: X_\alpha \rightarrow \mathbb{C}, \varphi_\alpha = t_{\alpha\beta} \varphi_\beta\}$

A naive attempt: Just take  $\{\frac{\partial \varphi_\alpha}{\partial z_\alpha}\}$ , can this be glued back to a section of some line bundle?

$$\frac{\partial \varphi_\alpha}{\partial \bar{z}_\alpha} = \frac{\partial (t_{\alpha\beta} \varphi_\beta)}{\partial \bar{z}_\alpha} = \frac{\partial t_{\alpha\beta}}{\partial \bar{z}_\alpha} \varphi_\beta + \frac{\partial \varphi_\beta}{\partial \bar{z}_\alpha} t_{\alpha\beta} = \frac{\partial t_{\alpha\beta}}{\partial \bar{z}_\alpha} \varphi_\beta + \frac{\partial \varphi_\beta}{\partial \bar{z}_\beta} \frac{\partial \bar{z}_\beta}{\partial \bar{z}_\alpha} t_{\alpha\beta}$$

Observe that if  $L$  holomorphic, then  $\frac{\partial t_{\alpha\beta}}{\partial \bar{z}_\alpha} = 0$ , thus

$$\frac{\partial \varphi_\alpha}{\partial \bar{z}_\alpha} = t_{\alpha\beta} \frac{\partial \bar{z}_\beta}{\partial \bar{z}_\alpha} \frac{\partial \varphi_\beta}{\partial \bar{z}_\beta}$$

i.e.  $(\frac{\partial \varphi_\alpha}{\partial \bar{z}_\alpha})$  is a section of  $L \otimes \Lambda^{1,0}(X) = L \otimes \Lambda^{0,1}(X)$ . This is the so called Chern connection.

Def. The **Chern connection** is defined by:

$$\bar{\partial}: \Gamma(X, L) \rightarrow \Gamma(X, L \otimes \Lambda^{0,1}), \quad \varphi = \{\varphi_\alpha\} \mapsto \bar{\partial}\varphi = \left\{ \frac{\partial \varphi_\alpha}{\partial \bar{z}_\alpha} \right\}$$

Def. A **metric** on  $L = \{t_{\alpha\beta}\}$  is an assignment  $h$ :

$$h = \{h_\alpha\}: h_\alpha: X_\alpha \rightarrow \mathbb{R}^{>0}, \text{ smooth, strictly positive on } X_\alpha \text{ satisfying } h_\alpha = |t_{\alpha\beta}|^{-2} h_\beta \text{ on } X_\alpha \cap X_\beta$$

Equivalently,  $h$  is a strictly positive section in  $L^{-1} \otimes \bar{L}^{-1}$ . There are plenty of such sections by glueing local sections.

Def: Given  $\varphi \in \Gamma(X, L)$ ,  $\varphi = \{\varphi_\alpha\}$ , we can define its **norm** by:

$$\|\varphi\| \triangleq \varphi_\alpha \bar{\varphi}_\alpha h_\alpha$$

Well-defined since on  $X_\alpha \cap X_\beta$ ,  $\varphi_\alpha \bar{\varphi}_\alpha h_\alpha = t_{\alpha\beta} \varphi_\beta \cdot \bar{t}_{\alpha\beta} \bar{\varphi}_\beta |t_{\alpha\beta}|^{-2} h_\beta = \varphi_\beta \bar{\varphi}_\beta h_\beta$ .

Given a metric  $h$  on  $L$ , we can define the covariant derivative  $\nabla$  as follows:

$$\begin{aligned} \nabla_{\bar{z}}: \Gamma(X, L) &\longrightarrow \Gamma(X, L \otimes \Lambda^{1,0}), \\ \varphi = \{\varphi_\alpha\} &\mapsto h_\alpha^{-1} \frac{\partial}{\partial \bar{z}_\alpha} (h_\alpha \varphi_\alpha) \end{aligned}$$

Indeed,  $\{h_\alpha \varphi_\alpha\} \in \Gamma(X, L^{-1} \otimes \bar{L}^{-1} \otimes L) \cong \Gamma(X, \bar{L}^{-1})$  is a section of an antiholomorphic bundle and  $\frac{\partial}{\partial \bar{z}_\alpha} (h_\alpha \varphi_\alpha)$  is thus a well-defined section of  $\Gamma(X, \bar{L}^{-1} \otimes \Lambda^{1,0})$ . Hence  $\{h_\alpha^{-1} \frac{\partial}{\partial \bar{z}_\alpha} (h_\alpha \varphi_\alpha)\}$  is a well-defined section of  $\Gamma(X, L \otimes \bar{L} \otimes \bar{L}^{-1} \otimes \Lambda^{1,0}) = \Gamma(X, L \otimes \Lambda^{1,0})$ .

Another way of writing  $\nabla_{\bar{z}}$  would be useful:  $\nabla_{\bar{z}} \varphi = \{\partial_{\bar{z}_\alpha} \varphi_\alpha + h_\alpha^{-1} \partial_{\bar{z}_\alpha} h_\alpha\} \varphi_\alpha$   
 $= \{\partial_{\bar{z}_\alpha} \varphi_\alpha + \partial_{\bar{z}_\alpha} (\log h_\alpha)\} \varphi_\alpha$ . Define  $\Gamma_\alpha = \partial_{\bar{z}_\alpha} (\log h_\alpha)$ , then  $\nabla_{\bar{z}} \varphi = \{\partial_{\bar{z}_\alpha} \varphi_\alpha + \Gamma_\alpha \varphi_\alpha\}$

• Commutation rules:

Fix  $L \rightarrow X$  a holomorphic line bundle  $L = \{t_{\alpha\beta}\}$ , metric  $h = \{h_\alpha\}$ ,  $\varphi = \{\varphi_\alpha\} \in \Gamma(X, L)$

$$\begin{aligned}
\nabla_{\bar{z}} \nabla_{\bar{z}} \varphi - \nabla_{\bar{z}} \nabla_z \varphi &= \nabla_{\bar{z}}(\partial_{\bar{z}} \varphi) - \nabla_{\bar{z}}(h^{-1} \partial_z(h\varphi)) \\
&= h^{-1} \partial_{\bar{z}}(h(\partial_{\bar{z}} \varphi)) - \partial_{\bar{z}}(h^{-1} \partial_z(h\varphi)) \\
&= \partial_z \partial_{\bar{z}} \varphi + \Gamma \cdot \partial_{\bar{z}} \varphi - \partial_{\bar{z}}(\Gamma \varphi + \partial_z \varphi) \\
&= -(\partial_{\bar{z}} \Gamma) \varphi
\end{aligned}$$

Note that  $\nabla_{\bar{z}} \varphi \in \Gamma(X, L \otimes \Lambda^{1,0})$ , a section of a holomorphic line bundle, to which  $\partial_{\bar{z}}$  applies;  $\partial_{\bar{z}} \varphi \in \Gamma(X, L \otimes \Lambda^{0,1})$ ,  $h \partial_{\bar{z}} \varphi \in \Gamma(X, L^{-1} \otimes \bar{L}^{-1} \otimes L \otimes \Lambda^{0,1}) \cong \Gamma(X, \bar{L}^{-1} \otimes \Lambda^{0,1})$ , i.e. a section of an anti-holomorphic bundle, to which  $\partial_z$  applies.

Hence we obtain:

$$[\nabla_{\bar{z}}, \nabla_z] \varphi = F_{\bar{z}z} \cdot \varphi$$

where  $F_{\bar{z}z} = -\partial_{\bar{z}} \Gamma = -\partial_{\bar{z}} \partial_z \log h$ , which is called the *curvature* of the line bundle  $L$  w.r.t.  $h$ .

Observation:  $F_{\bar{z}z} = \{F_{\bar{z}_\alpha z_\alpha}\}$  is a section of  $\Lambda^{1,1} \cong \Lambda^{1,0} \otimes \Lambda^{0,1}$ .  
 Explicitly,  $\partial_{\bar{z}_\alpha} \partial_{z_\alpha} \log h_\alpha = \frac{\partial \bar{z}_\beta}{\partial \bar{z}_\alpha} \cdot \frac{\partial z_\beta}{\partial z_\alpha} \partial_{\bar{z}_\beta} \partial_{z_\beta} (\log h_\beta - \log t_{\alpha\beta} - \log \bar{t}_{\alpha\beta})$   
 $= \frac{\partial \bar{z}_\beta}{\partial \bar{z}_\alpha} \cdot \frac{\partial z_\beta}{\partial z_\alpha} \partial_{\bar{z}_\beta} \partial_{z_\beta} \log h_\beta$

It's not cohomologically trivial since in general  $\{(\partial_{z_\alpha} \log h_\alpha) dz_\alpha\}$  is not a global form on  $X$ .

• Basic residue formula for holomorphic line bundles.

For any  $L \rightarrow X$ , holomorphic line bundle, any metric  $h$  and any meromorphic section  $\varphi = \{\varphi_\alpha\}$ , i.e.  $\varphi_\alpha$  is a meromorphic function on  $X_\alpha$ :

$$\#\{\text{zeroes of } \varphi\} - \#\{\text{poles of } \varphi\} = \frac{i}{2\pi} \int_X F_{\bar{z}z} dz \wedge d\bar{z}$$

Observations:

- (1). L.H.S. doesn't depend on the metric chosen.
- (2). R.H.S. doesn't depend on the meromorphic section chosen.
- (3). Meromorphic sections always exist (to be proven). We want to know if holomorphic sections exist. If  $\frac{i}{2\pi} \int_X F_{\bar{z}z} dz \wedge d\bar{z} < 0$ , then for any  $\varphi$ ,  $\#\{\text{poles of } \varphi\} > 0 \Rightarrow \nexists$  holomorphic sections. Thus the sign of curvature is of great importance.

Def. (The 1<sup>st</sup> Chern class)  $C_1(L) \triangleq \frac{i}{2\pi} \int_X F_{\bar{z}z} dz \wedge d\bar{z}$ .

Proof of the formula.

We compute the R.H.S., Locally:

$$F_{\bar{z}z} dz \wedge d\bar{z} = -\partial_z \partial_{\bar{z}} \log h dz \wedge d\bar{z} = -d(\partial_{\bar{z}} \log h) d\bar{z}$$


Notice that  $\partial_{\bar{z}} \log h d\bar{z}$  is not a section of  $\Lambda^{1,0}$ . Note that:

$$F_{\bar{z}z} dz \wedge d\bar{z} = -\partial_z \partial_{\bar{z}} (\log \|\varphi\|^2) \cdot dz \wedge d\bar{z},$$

away from the zeroes and poles of  $\varphi$ , which are denoted as  $\{P_1, \dots, P_N\}$ . Take  $\varepsilon$  small enough, so that each disk  $D_\varepsilon(P_i, \varepsilon) (\cong D_i)$  contains only one  $P_i$ . Thus

$$\begin{aligned} \int_X F_{\bar{z}z} dz \wedge d\bar{z} &= \lim_{\varepsilon \rightarrow 0} \int_{X \setminus \bigcup_{i=1}^N D_i} -\partial_z \partial_{\bar{z}} (\log \|\varphi\|^2) \cdot dz \wedge d\bar{z} \\ &= \lim_{\varepsilon \rightarrow 0} \sum_{i=1}^N \oint_{\partial D_i} \partial_{\bar{z}} (\log \|\varphi\|^2) d\bar{z} \end{aligned}$$

On each  $D_i$  (contained in a coordinate chart),  $\varphi = z^M \Phi(z)$ ,  $\Phi$ : holo,  $\Phi(0) \neq 0$ .



$$\begin{aligned} \oint_{\partial D_i} \partial_{\bar{z}} (\log \|\varphi\|^2) d\bar{z} &= \oint_{\partial D_i} \partial_{\bar{z}} (\log |z|^{2M} + \log |h|\Phi(z)|^2) d\bar{z} \\ &= \oint_{\partial D_i} \left( \frac{M}{z} + \partial_{\bar{z}} \log |h|\Phi(z)|^2 \right) d\bar{z} \\ &= -2\pi i M + O(\varepsilon) \rightarrow -2\pi i M \quad (\varepsilon \rightarrow 0) \\ \Rightarrow \frac{i}{2\pi} \int_X F_{\bar{z}z} dz \wedge d\bar{z} &= \sum_{i=1}^N M_i = \#\{\text{zeroes of } \varphi\} - \#\{\text{poles of } \varphi\} \quad \square \end{aligned}$$

### • The Riemann-Roch Theorem

Notation:  $H^0(X, L) \triangleq \{\text{holomorphic sections of } L\} = \{\varphi = (\varphi_\alpha) \mid \partial_{\bar{z}_\alpha} \varphi_\alpha = 0\}$

For any line bundle  $L \rightarrow X$ , we have the following formula (Riemann-Roch):

$$\dim H^0(X, L) - \dim H^0(X, L^{-1} \otimes K_X) = C_1(L) + \frac{1}{2} \chi(X)$$

where  $K_X = \Lambda^{1,0}(X)$ ,  $\chi(X) \triangleq C_1(K_X^{-1}) =$  the Euler characteristic of  $X$ .

Furthermore, we shall prove the following (Gauss-Bonnet):

If  $X$  is a surface with  $g$  holes, then  $\chi(X) = 2 - 2g$

Corollaries:

1). Let  $X$  be a Riemann surface with  $g$  holes. Take  $L = K_X$ , then we have  $K_X^{-1} \otimes K_X$  is trivial, thus by R.R. Thm:

$$\dim H^0(X, L) - 1 = -\chi(X) + \frac{1}{2}\chi(X) = g - 1$$

$$\Rightarrow \dim H^0(X, L) = g$$

E.g. On the torus, we exhibited  $\omega = \frac{dz}{\sqrt{z(z-1)(z-\lambda)}}$ , which is a basis of  $H^0(X, K_X)$ .

2). Use of point bundles


Fix a R.S.  $X$  and  $p \in X$ . We construct a line bundle in the following way:

Pick a coordinate system in a neighborhood  $X_0$  of  $p$ , and set  $X_\infty = X \setminus \{p\}$

Let  $L$  be  $\{t_{\infty 0}(z) = z \text{ on } X_0 \cap X_\infty\}$ . This defines a holomorphic line bundle which admits a holomorphic section  $1_p = 1$  on  $X_\infty$ ;  $z$  on  $X_0$ , and  $1_p|_{X_0} = z = z \cdot 1 = t_{\infty 0} \cdot 1|_{X_\infty}$ . Thus  $1_p$  is a section of  $L \cong [nP]$ , which is

holomorphic, and has exactly 1 zero at  $p$ . In particular

$$C_1(L) = \#(\text{zeros of } 1_p) - \#(\text{poles of } 1_p) = 1.$$



Similarly, given any integer  $n \in \mathbb{Z}$ , we may define  $[nP]$  by the transition function  $\{z^n\}$ , and a holomorphic / meromorphic section  $1_{nP}$  depending on  $n \geq 0$  /  $n < 0$ . It follows that  $C_1([nP]) = n$ .

More generally, given  $P_1, \dots, P_M$  on  $X$  and integers  $n_1, \dots, n_M$ , we may construct  $[n_1 P_1 + \dots + n_M P_M] = L \cong [n_1 P_1] \otimes \dots \otimes [n_M P_M]$ , and  $C_1(L) = \sum_{i=1}^M n_i$

Proof that any holomorphic line bundle admits a non-trivial meromorphic section.

Pick a point  $P$ , consider  $L \otimes [nP]$ .  $C_1(L \otimes [nP]) = C_1(L) + n$ . By R.R. Thm.

$$\dim H^0(X, L \otimes [nP]) - \dim H^0(X, L^{-1} \otimes [nP] \otimes K) = C_1(L) + n - \frac{1}{2}\chi(X) > 0 \quad (n \gg 0)$$

$$\Rightarrow \dim H^0(X, L \otimes [nP]) > 0 \quad (n \gg 0)$$

$\Rightarrow \exists \varphi \neq 0$ ,  $\varphi$  a section of  $L \otimes [nP]$ ,  $\varphi$  holomorphic.

$\Rightarrow \psi = \varphi \cdot 1_{-nP}$  is a non-zero meromorphic section of  $L$ .

3). Given any two points  $P_1 \neq P_2$ , we show that R.R.  $\Rightarrow \exists$  meromorphic form  $\varphi$  with simple poles  $P_1$  and  $P_2$ .

Consider the bundle  $L = [-P_1 - P_2] \Rightarrow C_1(L) = -2$  and  $L$  can't admit any holomorphic section.

$$\begin{aligned} R.R \Rightarrow 0 - \dim H^0([P_1 + P_2] \otimes K_X) &= -2 + 1 - g \\ \Rightarrow \dim H^0([P_1 + P_2] \otimes K_X) &= g + 1. \end{aligned}$$

Let  $\psi_1, \dots, \psi_g$  be a basis of  $H^0(X, K_X)$ , then  $1_{[P_1 + P_2]} \cdot \psi_1, \dots, 1_{[P_1 + P_2]} \cdot \psi_g$  are linearly independent sections of  $[P_1 + P_2] \otimes K_X$ . Let  $\Phi$  be a holomorphic section of the bundle which is linearly independent of  $1_{[P_1 + P_2]} \cdot \psi_1, \dots, 1_{[P_1 + P_2]} \cdot \psi_g$ , and set  $\varphi = 1_{[-P_1 - P_2]} \cdot \Phi$  is a section we need. (Potentially  $\varphi$  may not have 2 poles, then it would have no pole at all since  $\sum_{p=P_i} \text{Res}(\varphi)(p) = 0$ . In that case  $\varphi$  is a holomorphic section and  $1_{[P_1 + P_2]} \cdot \varphi = \Phi$  would be holomorphic thus a linear combination of  $\psi_1, \dots, \psi_g$ , which contradicts the fact that  $\Phi$  being linearly independent from  $1_{[P_1 + P_2]} \cdot \psi_1, \dots, 1_{[P_1 + P_2]} \cdot \psi_g$ .)

## Proof of Riemann-Roch Theorem

- Basic idea behind index theorems - heuristic point of view

Assume  $D$  is a linear operator between two vector spaces:  $D: H_1 \rightarrow H_2$  where  $H$  and  $H_1$  are Hilbert spaces (in general,  $H_1 \neq H_2$ , for instance,  $\bar{\partial}: \Gamma(X, L) \rightarrow \Gamma(X, L \otimes \Lambda^{0,1})$ ,  $\varphi \mapsto \bar{\partial}\varphi$ ). Since the spectrum of  $D$  is not available. Consider  $\Delta_- = D^*D$ , where  $D^*$  is the adjoint of  $D$  (which is defined by  $\langle Du, v \rangle = \langle u, D^*v \rangle$ ,  $\forall u \in H_1, v \in H_2$ ),  $\Delta_-: H_1 \rightarrow H_1$ , self-adjoint. Hence we can talk about the spectrum of  $\Delta_-$ . Spectrum  $\Delta_- = \{ \lambda_{\bar{n}} : \exists 0 \neq \varphi_n \in H_1, \Delta_- \varphi_n = \lambda_{\bar{n}} \varphi_n \}$ . Furthermore,  $\ker \Delta_- = \{ \varphi : D^*D\varphi = 0 \} = \{ \varphi : \langle \varphi, D^*D\varphi \rangle = 0 \} = \{ \varphi : \|D\varphi\| = 0 \} = \{ \varphi : D\varphi = 0 \} = \ker D$ .

Another natural operator:  $\Delta_+ = DD^*: H_2 \rightarrow H_2$  and again, we define the spectrum  $\Delta_+ = \{ \lambda_n^+ : \exists \psi \in H_2, \Delta_+ \psi = \lambda_n^+ \psi \}$ , and similarly  $\ker \Delta_+ = \ker D^*$ .

A simple observation: the spectra of  $\Delta_-$  and  $\Delta_+$  are the same (counted with multiplicity) are the same except for the kernels. Indeed,  $D^*D\varphi = \lambda_{\bar{n}} \varphi \Rightarrow \Delta_+(D\varphi) = DD^*D\varphi = \lambda_{\bar{n}} D\varphi$ . i.e.  $\forall \lambda_{\bar{n}}$  a non-zero eigen-value of  $\Delta_-$ ,  $\lambda_{\bar{n}}$  is also an eigenvalue of  $\Delta_+$ .

Consider,  $\forall t > 0$ ,  $\text{Tr}(e^{-t\Delta_-}) - \text{Tr}(e^{-t\Delta_+})$ . Here we assume that  $\Delta_-$ ,  $\Delta_+$  can be diagonalized, i.e.  $\exists$  an o.n.b.  $\{\varphi_n^-\}$  of  $H_1$  with  $\Delta_- \varphi_n^- = \lambda_n^- \varphi_n^-$ , an o.n.b.  $\{\varphi_n^+\}$  of  $H_2$  with  $\Delta_+ \varphi_n^+ = \lambda_n^+ \varphi_n^+$ . We may define  $f(\Delta_-) \varphi_n^- = f(\lambda_n^-) \varphi_n^-$ .

e.g.  $e^{-t\Delta_-} = \sum \frac{(-t)^n}{n!} (\Delta_-)^n$ , then  $e^{-t\Delta_-} \varphi_n^- = e^{-t\lambda_n^-} \varphi_n^-$ .

In general,  $\varphi = \sum C_n \varphi_n^-$ ,  $f(\Delta_-) \varphi = \sum C_n f(\Delta_-) \varphi_n^- = \sum C_n f(\lambda_n^-) \varphi_n^-$ . Note that  $\varphi \in H \Rightarrow \sum |C_n|^2 < \infty$ , for  $f(\Delta_-) \varphi \in H$ , we need  $\sum |C_n|^2 |f(\lambda_n^-)|^2 < \infty$ . In our case it's good since  $\lambda_n > 0$ ,  $e^{-t\lambda_n} < 1$ .

$\text{Tr}(f(\Delta_-)) \triangleq \sum_n f(\lambda_n^-)$  if it converges. Then

$$\text{Tr}(e^{-t\Delta_-}) - \text{Tr}(e^{-t\Delta_+}) = \sum e^{-t\lambda_n^-} - \sum e^{-t\lambda_n^+} = \dim \ker D - \dim \ker D^{\dagger}$$

Our next observation is that, the LHS can be computed using perturbation theory. More precisely, set  $P = e^{-t\Delta}$ , then

$$\begin{cases} (\frac{\partial}{\partial t} + \Delta)P = 0 & \text{heat equation} \\ P|_{t=0} = \text{Id} \end{cases}$$

and solutions of the heat equation can be computed asymptotically for small  $t$ .

### Details

Let  $L \rightarrow X$  be a holomorphic line bundle. We want to set up:

$$\bar{\partial}: \Gamma(X, L) \rightarrow \Gamma(X, L \otimes \Lambda^{0,1})$$

- Inner products on  $\Gamma(X, L)$  and  $\Gamma(X, L \otimes \Lambda^{0,1})$

Pick a metric  $h$  on  $L$ , and a metric  $g$  on  $(\Lambda^{1,0})^{-1} = \{f_{\alpha} \frac{\partial}{\partial z_{\alpha}}\}$  (the holomorphic tangent bundle). Recall that for a holo. line bundle  $L$ , a metric is a positive section of  $L^{-1} \otimes \bar{L}^{-1}$ . Thus  $g$  is a positive section of  $\Lambda^{1,0} \otimes \bar{\Lambda}^{1,0} \cong \Lambda^{1,1}$ , a positive (1,1)-form; we shall write  $g_{\bar{z}z}$  to stress it's a (1,1)-form.

Def.  $\forall \varphi, \psi \in \Gamma(X, L)$ ,  $\Phi, \bar{\Psi} \in \Gamma(X, L \otimes \Lambda^{0,1})$ , we define inner products

$$\langle \varphi, \psi \rangle \triangleq \int_X h \varphi \bar{\psi} g_{\bar{z}z}$$

$$\langle \Phi, \bar{\Psi} \rangle \triangleq \int_X h \Phi \bar{\Psi}$$

Here note that  $h \varphi \bar{\psi} g_{\bar{z}z} \in \Gamma(X, L^{-1} \otimes \bar{L}^{-1} \otimes L \otimes \bar{L} \otimes \Lambda^{1,1}) = \Gamma(X, \Lambda^{1,1})$

$$h \Phi \bar{\Psi} \in \Gamma(X, L^{-1} \otimes \bar{L}^{-1} \otimes L \otimes \Lambda^{0,1} \otimes \bar{L} \otimes \Lambda^{1,0}) = \Gamma(X, \Lambda^{1,1}),$$

hence both are (1,1) forms, and thus  $\int_X$  makes sense.

- Formal adjoint  $\bar{\partial}^\dagger$  of  $\bar{\partial}$

$\bar{\partial}^\dagger: \Gamma(X, L \otimes \Lambda^{0,1}) \rightarrow \Gamma(X, L)$  w.r.t. the inner products above is defined by

$$\langle \bar{\partial}\varphi, \Phi \rangle = \langle \varphi, \bar{\partial}^\dagger\Phi \rangle$$

Explicitly:  $\int_X h \bar{\partial}\bar{z}\varphi \cdot \bar{\Phi} = \int_X h \cdot \varphi \cdot \bar{\partial}^\dagger\bar{\Phi} g_{\bar{z}\bar{z}}$

The L.H.S. =  $\int_X \bar{\partial}\bar{z}\varphi \cdot \overline{h\bar{\Phi}} = -\int_X \varphi \bar{\partial}\bar{z}(h\bar{\Phi})$  (integration by parts)

$$= -\int_X h\varphi \overline{g^{\bar{z}\bar{z}} h^{-1} \bar{\partial}\bar{z}(h\bar{\Phi})} g_{\bar{z}\bar{z}} \quad (g^{\bar{z}\bar{z}} \cdot g_{\bar{z}\bar{z}} = 1)$$

$$\Rightarrow \bar{\partial}^\dagger\bar{\Phi} = -g^{\bar{z}\bar{z}} \nabla_{\bar{z}}\bar{\Phi}.$$

(Indeed,  $g^{\bar{z}\bar{z}} \nabla_{\bar{z}}\bar{\Phi} \in \Gamma(X, (\Lambda^{1,1})^{-1} \otimes L \otimes \Lambda^{0,1} \otimes \Lambda^{1,0}) = \Gamma(X, L)$  so this is a valid def.)

Then we define  $\Delta_- = \bar{\partial}^\dagger\bar{\partial}: \Gamma(X, L) \rightarrow \Gamma(X, L)$

$$\Delta_+ = \bar{\partial}\bar{\partial}^\dagger: \Gamma(X, L \otimes \Lambda^{0,1}) \rightarrow \Gamma(X, L \otimes \Lambda^{0,1})$$

Thm. (Spectral Decomposition of Laplacian  $\Delta_-$ )

(1).  $\exists \{ \varphi_n \in \Gamma(X, L), n \in \mathbb{Z}_+ \}$  an o.n.b. of  $H_1 = \overline{\Gamma(X, L)}$  which consists only of eigenfunctions for  $\Delta_-$ , i.e.  $\Delta_- \varphi_n = \lambda_n \varphi_n$ .

(2).  $\lambda_n \geq 0$  and  $\lambda_n \rightarrow \infty$  at least at polynomial growth rate as  $n \rightarrow \infty$ , i.e.  $C \cdot n^\nu \geq \lambda_n \geq D \cdot n^\delta$  for some  $C, D, \nu, \delta > 0$ .

(3). For any  $k \in \mathbb{Z}_+$ ,  $\exists \delta_k$  s.t.  $\|\varphi_n\|_{C^k} \leq A_k |\lambda_n|^{-\delta_k}$  for some constant  $A_k$ .

Here  $\|\varphi\|_{C^0} \triangleq \sup_x |\varphi|^2 h$ ;  $\|\varphi\|_{C^1} \triangleq \|\varphi\|_{C^0} + \sup_x |\nabla_{\bar{z}}\varphi|^2 h g^{\bar{z}\bar{z}} + \sup_x |\nabla_z\varphi|^2 h g^{\bar{z}\bar{z}}$ , etc.

(4). Each eigenvalue  $\lambda_n$  occurs with finite multiplicity. ( $\Rightarrow \ker \Delta_-$  is finite dimensional).

Assuming this thm, we define  $e^{-t\Delta_-}$  and its trace as follows:

Def.  $u \in \overline{\Gamma(X, L)}$ , write  $u = \sum_{n=0}^\infty u_n \varphi_n$ ,  $u_n = \langle u, \varphi_n \rangle$ , with the series converging in the  $L^2$ -sense w.r.t.  $\langle, \rangle$  on  $H_1$ . Define:

$$e^{-t\Delta_-} u \triangleq \sum_{n=0}^\infty e^{-t\lambda_n} u_n \varphi_n \quad (*).$$

The R.H.S is a well-defined element of  $H_1$ , in view of the

Thm. (Riesz - Fisher)  $\sum_{n=0}^\infty u_n \varphi_n$  converges in  $H_1$  iff  $\sum_{n=0}^\infty |u_n|^2 < \infty$ .

Pf:  $\{ \sum_{n=0}^N u_n \varphi_n \}$  converges iff it's Cauchy iff  $\| \sum_{n=N}^M u_n \varphi_n \|^2 < \epsilon$  for  $M, N \gg 0$  iff  $\sum_{n=0}^\infty |u_n|^2 < +\infty$ .

□



Observe that  $\{\varphi_n\}$  is also an o.n.b. of eigen-functions for  $e^{-t\Delta^-}$ . Indeed

$$e^{-t\Delta^-} \varphi_n = (e^{-t\lambda_n^-}) \varphi_n$$

with eigenvalues  $\{e^{-t\lambda_n^-}\}$ . It follows from our thm that

$$\text{Tr}(e^{-t\Delta^-}) = \sum_n e^{-t\lambda_n^-} < \infty, \quad \forall t > 0$$

since  $\lambda_n^- \geq c \cdot n^\delta$  for  $n \gg 0$ .

$$\begin{aligned} \text{It follows that } \dim \ker \bar{\partial} - \dim \ker \bar{\partial}^+ &= \dim \ker \Delta^- - \dim \ker \Delta^+ \\ &= \text{Tr}(e^{-t\Delta^-}) - \text{Tr}(e^{-t\Delta^+}) \end{aligned}$$

Evaluating  $\text{Tr}(e^{-t\Delta^-})$

1). Expressing it as a kernel.

$$\begin{aligned} \forall u \in \Gamma(X, L). \quad (e^{-t\Delta^-} u)(z) &= \sum_n e^{-t\lambda_n^-} u_n \varphi_n(z) \\ &= \sum_n e^{-t\lambda_n^-} \left( \int_X h u \bar{\varphi}_n g_{\bar{w}w} \frac{i}{2} d\omega \wedge d\bar{\omega} \right) \varphi_n(z) \\ &= \int_X \left( \sum_n e^{-t\lambda_n^-} \varphi_n(z) \overline{\varphi_n(w)} h(w) g_{\bar{w}w}(w) \right) u(w) \frac{i}{2} d\omega \wedge d\bar{\omega} \end{aligned}$$

Note that we may interchange sum and integral since  $\|\varphi_n\|_{C^k} \leq A_k |\lambda_n|^{-\delta_k}$  and thus the sum converges.

$$\Rightarrow (e^{-t\Delta^-} u)(z) = \int_X K_t(z, w) u(w) \frac{i}{2} d\omega \wedge d\bar{\omega}, \quad \text{where}$$

$$K_t(z, w) = \sum_n e^{-t\lambda_n^-} \underbrace{\varphi_n(z)}_{\in L_z} \underbrace{\overline{\varphi_n(w)}}_{\bar{L}_w} \underbrace{h(w)}_{L_w \otimes \bar{L}_w} \underbrace{g_{\bar{w}w}(w)}_{\Lambda_w^1} \quad (\text{heat kernel})$$

$$\Rightarrow K_t(z, z) \in \Gamma(X, \Lambda^{1,1})$$

$$\begin{aligned} \Rightarrow \int_X K_t(z, z) \frac{i}{2} dz \wedge d\bar{z} &= \int_X \sum_n e^{-t\lambda_n^-} |\varphi_n(z)|^2 h(z) g_{\bar{z}z} \frac{i}{2} dz \wedge d\bar{z} \\ &= \sum_n e^{-t\lambda_n^-} \int_X |\varphi_n(z)|^2 h(z) g_{\bar{z}z} \frac{i}{2} dz \wedge d\bar{z} \\ &= \sum_n e^{-t\lambda_n^-} \|\varphi_n\|^2 \\ &= \sum_n e^{-t\lambda_n^-} \\ &= \text{Tr} e^{-t\Delta^-} \end{aligned}$$

2). Determine  $K_t(z, z)$  for small  $t$ .

Claim: (a) There exists an asymptotic expansion of  $K_t(z, z)$ , in the sense that

$$K_t(z, z) = \frac{U_{\bar{z}z}}{t} + V_{\bar{z}z} + O(t)$$

$$\text{and } \|K_t(z, z) - \frac{U_{\bar{z}z}}{t} - V_{\bar{z}z}\|_{C^m} \leq A_m t \quad (t \ll 1)$$

Clearly  $U_{\bar{z}z} \in \Gamma(X, \Lambda^{1,1})$  and so is  $V_{\bar{z}z}$ .

(b).  $U_{\bar{z}z}$ ,  $V_{\bar{z}z}$  depend only on  $h$  of  $L$  and  $g$  of  $\Lambda^{-1,0}$ .

Key observation:  $U_{\bar{z}z}$  doesn't depend on the derivatives of  $h$  and  $g$ , while  $V_{\bar{z}z}$  depends on derivatives of  $h, g$  up to order 2. (This will follow from heat kernel equation theory to be explained below).

$$\begin{cases} U_{\bar{z}z} = a g_{\bar{z}z} \\ V_{\bar{z}z} = b F_{\bar{z}z}^h + c F_{\bar{z}z}^g \end{cases}$$

(This can be guessed from power counting)

We write specifically for  $\Delta^-$  and  $\Delta^+$

$$\text{kernel of } \Delta^-: K_{\bar{z}z}^-(z, \bar{z}) = a^- \frac{g_{\bar{z}z}}{t} + b^- F_{\bar{z}z}^h + c^- F_{\bar{z}z}^g$$

$$\text{kernel of } \Delta^+: K_{\bar{z}z}^+(z, \bar{z}) = a^+ \frac{g_{\bar{z}z}}{t} + b^+ F_{\bar{z}z}^h + c^+ F_{\bar{z}z}^g$$

$$\Rightarrow \text{Tr}(e^{-t\Delta^-}) - \text{Tr}(e^{-t\Delta^+}) = (a^- - a^+) \frac{\text{Vol}(X)}{t} + (b^- - b^+) C_1(L) + (c^- - c^+) C_1(K_X^{-1})$$

Clearly  $a^- = a^+$  otherwise the L.H.S. would blow-up.

$$\Rightarrow \text{Tr}(e^{-t\Delta^-}) - \text{Tr}(e^{-t\Delta^+}) = \beta C_1(L) + \gamma C_1(K_X^{-1})$$

We can even determine the coefficients  $\beta$  and  $\gamma$ .

- **Serre duality**: We know that  $\text{Tr}(e^{-t\Delta^-}) - \text{Tr}(e^{-t\Delta^+}) = \dim \ker \bar{\partial} - \dim \ker \bar{\partial}^+$  but how does  $\ker \bar{\partial}^+ \sim H^0(L^{-1} \otimes K_X)$ ?

Claim:  $\Phi \mapsto h\bar{\Phi}$  gives an isomorphism:  $\Gamma(X, L \otimes \Lambda^{0,1}) \rightarrow \Gamma(X, L^{-1} \otimes K_X)$

Indeed  $\Gamma(X, L \otimes \Lambda^{0,1}) \ni \Phi \mapsto h\bar{\Phi} \in \Gamma(X, \bar{L} \otimes \Lambda^{1,0} \otimes L^{-1} \otimes \bar{L}^{-1}) = \Gamma(X, L^{-1} \otimes \Lambda^{1,0})$ .

Moreover,  $\Phi \in \ker \bar{\partial}^+ \Leftrightarrow \bar{\partial}^+ \Phi = 0 \Leftrightarrow g^{\bar{z}z} h^{-1} (\partial_{\bar{z}} h \bar{\Phi}) = 0$

$$\Leftrightarrow \partial_{\bar{z}} h \bar{\Phi} = 0 \Leftrightarrow \partial_{\bar{z}} h \bar{\Phi} = 0 \quad (h \text{ is positive, thus real})$$

$$\Leftrightarrow h \bar{\Phi} \in H^0(X, L^{-1} \otimes K_X).$$

Thus  $\dim \ker \bar{\partial}^+ = \dim H^0(X, L^{-1} \otimes K_X)$ .

Now  $\dim \ker \bar{\partial} - \dim \ker \bar{\partial}^+ = \dim H^0(X, L) - \dim H^0(X, L^{-1} \otimes K_X)$

$$= \beta C_1(L) + \gamma C_1(K_X^{-1})$$

Later we will see that  $\beta$  and  $\gamma$  are universal constants.

- Special cases

1).  $L = \text{trivial} \Rightarrow 1 - \dim H^0(X, K_X) = 0 + \chi C_1(K_X^{-1})$

2).  $L = K_X \Rightarrow \dim H^0(X, K_X) - 1 = \beta C_1(K_X) + \chi \cdot C_1(K_X^{-1})$

Combining 1), 2)  $\Rightarrow \chi = \frac{\beta}{2}$ . To determine the over all value (assuming Gauss-Bonnet thm.), consider  $X = S^2$ ,  $C_1(K_X^{-1}) = 2 - 2g = 2 \Rightarrow C_1(K_X) = -2$  and  $H^0(X, K_X) = 0$ .

$\Rightarrow 1 - 0 = \chi C_1(K_X^{-1}) = 2\chi \Rightarrow \chi = \frac{1}{2}$ . Hence Riemann-Roch follows.

• Road map and summary: three main ingredients.

(1). Spectral theory of  $\Delta_- = \bar{\partial}^+ \bar{\partial}$ ,  $\Delta_+ = \bar{\partial} \bar{\partial}^+$

(2). Small time expansion of  $K_{\pm}^{\pm}(z, \bar{z})$

(3). Gauss-Bonnet  $\chi(X) = \frac{i}{2\pi} \int_X F_{\bar{z}z} dz \wedge d\bar{z} = 2 - 2g$ ,  $g = \# \text{ holes}$ .

$F_{\bar{z}z}$ : curvature of  $g_{\bar{z}z}$ .

Proof of Gauss-Bonnet

• Elements of Riemannian Geometry

$X$ : (real) manifold of dim  $n$ .  $X = \cup X_{\alpha}$ ,  $\Phi_{\alpha}: X_{\alpha} \rightarrow B \subseteq \mathbb{R}^n$ ,  $\Phi_{\alpha} \circ \Phi_{\beta}^{-1}$  smooth with smooth metric, Jacobian  $\neq 0$ .

Locally, on each chart,  $\{x^i, i=1, \dots, n\}$  be local charts. A metric is

$$ds^2 = \sum_{ij} g_{ij} dx^i dx^j$$

with  $g_{ij}(x)$  a symmetric, positive definite metric, transforming in such a way

as to make  $ds^2$  invariant. i.e. on  $X_{\alpha} \cap X_{\beta}$   $g_{ij}(x_{\alpha}) dx_{\alpha}^i dx_{\alpha}^j = g_{kl}(x_{\beta}) dx_{\beta}^k dx_{\beta}^l$

$$\Leftrightarrow g_{kl}^{\beta}(x_{\beta}) \frac{\partial x_{\beta}^k}{\partial x_{\alpha}^i} \frac{\partial x_{\beta}^l}{\partial x_{\alpha}^j} = g_{ij}^{\alpha}(x_{\alpha}).$$

Levi-Civita Connection

A vector field  $V$  on  $X$  corresponds to  $(V^i(x_{\alpha}) | i=1, \dots, n)$ , i.e. a vector valued function on  $X_{\alpha}$ , transforming in such a way that  $\{V^i \frac{\partial}{\partial x_{\alpha}^i}\}$  is invariant.

i.e. on  $X_{\alpha} \cap X_{\beta}$ ,  $V_{\alpha}^i(x_{\alpha}) \frac{\partial}{\partial x_{\alpha}^i} = V_{\beta}^j(x_{\beta}) \frac{\partial}{\partial x_{\beta}^j} \Leftrightarrow V_{\alpha}^i(x_{\alpha}) = V_{\beta}^j(x_{\beta}) \frac{\partial x_{\beta}^j}{\partial x_{\alpha}^i}$

Fix a metric  $ds^2 = g_{ij} dx^i dx^j$

Claim:  $\exists ! \{ \Gamma_{ij}^k | i, j, k=1, \dots, n \}$  so that  $((\nabla_m V)^i \triangleq \partial_m V^i + \Gamma_{mk}^i V^k)$  is a vector field

and: 1).  $\partial_m \langle V, W \rangle = \langle \nabla_m V, W \rangle + \langle V, \nabla_m W \rangle$

2).  $\Gamma_{ij}^k = \Gamma_{ji}^k$  (torsion free)

Indeed  $\Gamma_{ij}^k$  satisfying these two conditions are computed to be

$$\Gamma_{ij}^k = \frac{1}{2} g^{kl} (\partial_i g_{ej} + \partial_j g_{ie} - \partial_l g_{ij}).$$

Proof. 1)  $\Rightarrow \partial_m (g_{kl} V^k W^l) = (\partial_m V^k) g_{kl} W^l + g_{kl} \Gamma_{mp}^k V^p W^l + V^k g_{kl} (\partial_m W^l) + g_{kl} V^k \Gamma_{mp}^l W^p$   
 $\Rightarrow (\partial_m g_{kl}) V^k W^l = g_{kl} \Gamma_{mp}^k V^p W^l + g_{kl} \Gamma_{mp}^l V^k W^p = g_{pl} \Gamma_{mk}^p V^k W^l + g_{kp} \Gamma_{ml}^p V^k W^l$   
 $\Rightarrow \partial_m g_{kl} = g_{pl} \Gamma_{mk}^p + g_{kp} \Gamma_{ml}^p$ . (a)

Permuting m, k, l. we obtain:

$$\partial_k g_{lm} = g_{pm} \Gamma_{kl}^p + g_{lp} \Gamma_{km}^p \quad (b)$$

$$\partial_l g_{mk} = g_{pk} \Gamma_{lm}^p + g_{mp} \Gamma_{lk}^p \quad (c)$$

(b)+(c)-(a), using 2), we obtain:

$$2g_{pm} \Gamma_{kl}^p = \partial_l g_{mk} + \partial_k g_{lm} - \partial_m g_{kl}$$

Multiplying both sides by  $\frac{1}{2} g^{qm}$  and summing over p, we obtain:

$$\Gamma_{kl}^q = \frac{1}{2} g^{qm} (\partial_l g_{mk} + \partial_k g_{lm} - \partial_m g_{kl})$$

This computation also establishes the uniqueness of  $\{\Gamma_{ij}^k\}$  once the metric is fixed.  $\square$

The  $\nabla_m$ 's don't commute, and we have:

$$[\nabla_l, \nabla_m] V^i = R_{lm}^i{}^p V^p.$$

Def. (Curvatures)

1).  $R_{lm}^i{}^p = \partial_l \Gamma_{mp}^i - \partial_m \Gamma_{lp}^i + \Gamma_{lq}^i \Gamma_{mp}^q - \Gamma_{mq}^i \Gamma_{lp}^q$  is the Riemannian curvature

2).  $R_{mp} = R_{lm}^l{}^p$  is the Ricci curvature.

3).  $R = g^{pm} R_{mp}$  is the scalar curvature.

Note that our convention for raising and lowering indices:  $V^i \rightarrow V_j \triangleq V^i g_{ij}$ .  
 for example,  $R = R^p{}_p = g^{pm} R_{mp} = R_m{}^m$ .

**Example:**  $X$  a Riemann surface, with a metric  $g_{z\bar{z}}$  on  $\Lambda^{-1,0}$  gives rise to a Riemannian metric on  $X$ . We set up a dictionary.

- Dictionary. On a holomorphic chart  $z = x^1 + ix^2 \leftrightarrow (x^1, x^2)$

1).  $V \in \Gamma(X, \Lambda^{-1,0})$ ,  $V = V \partial_{\bar{z}}$ .  $\leftrightarrow$   $v$ : vector field  $V + \bar{V}$

Explicitly,  $V \partial_{\bar{z}} = (V^1 + iV^2) \cdot \frac{1}{2}(\partial_{x^1} - i\partial_{x^2}) = \frac{1}{2}(V^1 \partial_{x^1} + V^2 \partial_{x^2}) + \frac{i}{2}(V^2 \partial_{x^1} - V^1 \partial_{x^2})$

$\Rightarrow v = V^1 \partial_{x^1} + V^2 \partial_{x^2}$ . Conversely, define  $J: V^1 \partial_{x^1} + V^2 \partial_{x^2} \mapsto V^2 \partial_{x^1} - V^1 \partial_{x^2}$ , then  $V = \frac{1}{2}(v + iJv)$ .

2). A metric  $g_{\bar{z}z}$  on  $\Lambda^{-1,0}$ :  $\|V\| = V \bar{V} g_{\bar{z}z}$ . Define  $\|v\|^2 \triangleq \|V\|^2 = ((V^1)^2 + (V^2)^2) g_{\bar{z}z}$

Compared with  $\|v\|^2 = \sum g_{ij} v^i v^j \Rightarrow ds^2 = g_{ij} dx^i dx^j = g_{\bar{z}z}((dx^1)^2 + (dx^2)^2)$

Thus  $(g_{\bar{z}z}) \leftrightarrow ds^2 = g_{\bar{z}z}((dx^1)^2 + (dx^2)^2)$

3). Connection

$$\begin{array}{ccc} \nabla_{\bar{z}} V & \leftrightarrow & \nabla_{\bar{z}} v = (\nabla_{\bar{z}} + \nabla_{\bar{z}})(V + \bar{V}) \\ \text{(Chern connection)} & & \text{Levi-Civita connection} \\ \text{defined by } g_{\bar{z}z} & & \text{w.r.t. } ds^2 = g_{\bar{z}z}((dx^1)^2 + (dx^2)^2) \end{array}$$

Dependence of  $\Gamma_{ij}^k$  and  $R_{lm}{}^ip$  on the metric  $g_{ij}$ .

We determine the variations of  $\Gamma_{ij}^k$  and  $R_{lm}{}^ip$  under variations of  $g$ .

(The derivative (linear approximation) is simpler than the function itself). Since

$$g_{ij} \rightsquigarrow \Gamma_{ij}^k, \quad \delta g_{ij} \rightsquigarrow \delta \Gamma_{ij}^k.$$

Key variation formulae: (to be proven later)

$$1). \delta \Gamma_{ij}^k = \frac{1}{2} g^{kl} (\nabla_j \delta g_{il} + \nabla_i \delta g_{lj} - \nabla_l \delta g_{ij})$$

$$2). \delta R_{jm}{}^l{}_k = \nabla_j (\delta \Gamma_{mk}^l) - \nabla_m (\delta \Gamma_{jk}^l)$$

$$3). \delta R_{ml} = \nabla_j (\delta \Gamma_{ml}^j) - \frac{1}{2} \nabla_m \nabla_l (g^{jp} \delta g_{pj})$$

$$4). \delta R = -\delta g^{lm} R_{ml} + \nabla^j \nabla^l \delta g_{lj} - \Delta (g^{ml} \delta g_{lm})$$

where  $\delta g^{ij} \triangleq (\delta g_{kl}) g^{ki} g^{lj}$ .

Using these formula, we claim:

Define the functional on the space of all Riemannian metrics on  $X$ .

$$g_{ij} \mapsto I(g) = \frac{1}{2\pi} \int_X (R \sqrt{g}) dx,$$

then  $\delta I(g) = 0$  for any  $\delta g_{ij}$ .

Pf of claim:

$$\delta I(g) = \frac{1}{2\pi} \int_X \delta(R\sqrt{g}) dx = \frac{1}{2\pi} \int_X (\delta R \cdot \sqrt{g} + \frac{1}{2} \delta g / \sqrt{g} \cdot R) dx$$

where  $g = \det(g_{ij})$ . Note that  $\delta \log(\det(g_{ij})) = g^{ij} \delta g_{ij}$ .

(Indeed, we may assume that  $(g_{ij}) = \text{diag}(\lambda_1, \dots, \lambda_n) \Rightarrow \log(\det(g_{ij})) = \sum \log \lambda_i \Rightarrow \delta \log \det(g_{ij}) = \sum \frac{\delta \lambda_i}{\lambda_i} = \text{tr}((g_{ij})^{-1} \delta(g_{ij})) = g^{ij} \delta g_{ij}$ .)

$$\begin{aligned} \Rightarrow \delta I(g) &= \frac{1}{2\pi} \int_X (\delta R + \frac{1}{2} (\delta \log g) \cdot R) \cdot \sqrt{g} dx \\ &= \frac{1}{2\pi} \int_X (-\delta g^{lm} R_{ml} + \nabla^i \nabla^l \delta g_{ij} - \Delta(g^{ml} \delta g_{em}) + \frac{1}{2} g^{ij} \delta g_{ij} R) \sqrt{g} dx \\ &= \frac{1}{2\pi} \int_X (-\delta g^{lm} (R_{ml} - \frac{1}{2} g_{ml} R) + \nabla^i \nabla^l \delta g_{ij} - \Delta(g^{ml} \delta g_{ml})) \sqrt{g} dx \end{aligned}$$

Observation: In case  $\dim X = 2$   $-R_{ml} + \frac{1}{2} R g_{ml} = 0$  and  $\nabla^i \nabla^l \delta g_{ij} - \Delta(g^{ml} \delta g_{ml})$  is exact thus the integral vanishes.

The first identity is a consequence of the symmetries of the Riemannian curvature tensor:

Recall that  $[\nabla_j, \nabla_k] V^l = R_{jk}{}^l{}_m V^m$ , and

- $R_{jkpm} = -R_{kjpm} = -R_{jkmp}$
- $R_{jkpm} = R_{pmjk}$ .

In dimension 2, we calculate the Ricci tensor  $R_{ij} = R_{ilj}{}^l = g^{lp} R_{iljp}$ :

$$R_{11} = g^{lp} R_{1l1p} = g^{22} R_{1212} \quad ; \quad R_{12} = g^{lp} R_{1l2p} = g^{21} R_{1221} = -g^{21} R_{1212} = R_{21}$$

$$R_{22} = g^{lp} R_{2l2p} = g^{11} R_{2121} = g^{11} R_{1212}$$

$$\Rightarrow \begin{pmatrix} R_{11} & R_{12} \\ R_{21} & R_{22} \end{pmatrix} = R_{1212} \begin{pmatrix} g^{22} & -g^{21} \\ -g^{12} & g^{11} \end{pmatrix} = \frac{1}{g} R_{1212} \begin{pmatrix} g_{11} & g_{12} \\ g_{21} & g_{22} \end{pmatrix}$$


$$\Rightarrow R_{ml} - \frac{1}{2} R g_{ml} = \frac{1}{g} R_{1212} (g_{ml} - \frac{1}{2} g^{ij} g_{ij} g_{ml}) = 0$$

Now the Gauss-Bonnet formula follows easily from this result:


$$\frac{1}{2\pi} \int_X R \sqrt{g} dx = 2 - 2h$$

Here we temporarily write  $h = \# \text{ holes of } X$ ,  $g = \det(g_{ij})$

Decompose  $X$  as follows: (surgery)




Now  $\frac{1}{2\pi} \int_X R \sqrt{g} dx = \frac{1}{2\pi} \int_{\text{Body}} R \sqrt{g} dx + \sum \frac{1}{2\pi} \int_{\text{Handle}} R \sqrt{g} dx$ . We calculate each integral:



$$\frac{1}{2\pi} \int_{\text{Body}} R \sqrt{g} dx = \frac{1}{2\pi} \int_{\text{Body}+2h \text{ caps}} R \sqrt{g} dx - \frac{1}{2\pi} \int_{2h \text{ caps}} R \sqrt{g} dx.$$

By the previous claim,  $\frac{1}{2\pi} \int_{\text{Body}+2h \text{ caps}} R \sqrt{g} dx = \frac{1}{2\pi} \int_{S^2} R' \sqrt{g'} dx$ , where  $(g'_{ij})$ ,  $g'$ ,  $R'$  are derived from the standard metric on  $S^2$  as the unit sphere, and a direct calculation shows that  $R' = 1$ , thus  $\frac{1}{2\pi} \int_{S^2} R' \sqrt{g'} dx = 4\pi/2\pi = 2$ .

Moreover, for each cap  $\frac{1}{2\pi} \int_{\text{cap}} R \sqrt{g} dx = \frac{1}{2} \left( \frac{1}{2\pi} \int_{\text{sphere}} R \sqrt{g} \right) = \frac{1}{2} \cdot 2 = 1$



Lastly, for each handle,



$$\begin{aligned} \frac{1}{2\pi} \int_{\text{handle}} R \sqrt{g} dx &= \frac{1}{2\pi} \int_{\text{handle}+2 \text{ caps}} R \sqrt{g} dx - \frac{1}{2\pi} \int_{2 \text{ caps}} R \sqrt{g} dx \\ &= 2 - 2 \cdot 1 = 0 \end{aligned}$$

$$\begin{aligned} \text{Altogether: } \frac{1}{2\pi} \int_X R \sqrt{g} dx &= \frac{1}{2\pi} \int_{\text{Body}} R \sqrt{g} dx + \sum \frac{1}{2\pi} \int_{\text{Handle}} R \sqrt{g} dx \\ &= 2 - 2h \end{aligned}$$

Main valuation formulae.

Now let's derive our key variational formulae. First of all, assuming 1) and 2)

$$1). \delta \Gamma_{ij}^k = \frac{1}{2} g^{kl} (\nabla_j \delta g_{il} + \nabla_i \delta g_{lj} - \nabla_l \delta g_{ij})$$

$$2). \delta R_{jm}^l{}_k = \nabla_j (\delta \Gamma_{mk}^l) - \nabla_m (\delta \Gamma_{jk}^l)$$

we shall evaluate  $\delta R_{mk}$  and  $\delta R$ :

$$\delta R_{mk} = \delta (R_{jm}{}^j{}_k) = \delta (\nabla_j \delta \Gamma_{mk}^j - \nabla_m \delta \Gamma_{jk}^j)$$

$$\begin{aligned} \text{However, note that } \delta \Gamma_{jk}^j &= \frac{1}{2} g^{jp} (\nabla_k \delta g_{jp} + \nabla_j \delta g_{pk} - \nabla_p \delta g_{jk}) \\ &= \frac{1}{2} g^{jp} \nabla_k \delta g_{jp} \\ &= \frac{1}{2} \nabla_k (g^{jp} \delta g_{jp}) \end{aligned}$$

The second equality holds because  $\nabla_j \delta g_{pk} - \nabla_p \delta g_{jk}$  is antisymmetric in  $p$  and  $j$  while  $g^{jp}$  is symmetric. The last equality holds since  $\nabla_j g_{pk} = 0$ , which follows from

$$\partial_j g_{pk} = g_{pl} \Gamma_{jk}^l + g_{lk} \Gamma_{jp}^l$$

(known as Ricci's lemma, see the general remark on covariant differentiation of tensors.)

$$\text{Thus } \delta R_{mk} = \nabla_j (\delta \Gamma_{mk}^j) - \frac{1}{2} \nabla_m \nabla_k (g^{jp} \delta g_{jp}), \text{ as claimed in } \Rightarrow.$$

Next.  $\delta R = \delta (g^{km} R_{mk})$

$$= (\delta g^{km}) \cdot R_{mk} + g^{km} \delta R_{mk}$$

$$= -g^{kp} (\delta g_{pq}) g^{qm} R_{mk} + g^{km} (\nabla_j (\delta \Gamma_{mk}^j) - \frac{1}{2} \nabla_m \nabla_k (g^{jp} \delta g_{jp})).$$

$$\text{(Here we used that } G \cdot G^{-1} = \text{Id} \Rightarrow (\delta G) \cdot G^{-1} + G \cdot \delta(G^{-1}) = 0 \Rightarrow \delta G^{-1} = -G^{-1} (\delta G) G^{-1}.)$$

$$= -\delta g^{km} R_{mk} + \nabla_j (g^{km} \delta \Gamma_{mk}^j) - \frac{1}{2} \Delta (g^{jp} \delta g_{jp})$$

$$\text{(where by definition, } \Delta f \triangleq g^{km} \nabla_m \nabla_k f = g^{km} \nabla_m (\partial_k f), \text{ for any smooth } f).$$

We compute  $g^{km} \delta \Gamma_{mk}^j$ :

$$g^{km} \delta \Gamma_{mk}^j = g^{km} \cdot \frac{1}{2} g^{jp} (\nabla_k \delta g_{mp} + \nabla_m \delta g_{pk} - \nabla_p \delta g_{mk})$$

$$= g^{km} g^{jp} \nabla_k \delta g_{mp} - \frac{1}{2} g^{jp} \nabla_p (g^{km} \delta g_{mk})$$

$$\text{(since } \nabla_k \delta g_{mp} + \nabla_m \delta g_{pk} \text{ is symmetric in } k, p \text{ and so is } g^{km}; \nabla_p g^{km} = 0.)$$

$$= \nabla_k (\delta g^{kj}) - \frac{1}{2} g^{jp} \nabla_p (g^{km} \delta g_{mk})$$

Plugging in, we obtain:

$$\delta R = -\delta g^{km} R_{mk} + \nabla_j \nabla_k (\delta g^{kj}) - \Delta (g^{jp} \delta g_{pj})$$

which is the key variational formula 4), with the term  $\nabla_j \nabla_k (\delta g^{kj}) - \Delta (g^{jp} \delta g_{pj})$  being an exact term.



Thirdly, we calculate  $\delta R_{jm}{}^l{}_k$ . One way of doing this is via the formula

$$R_{jm}{}^l{}_k = \partial_j \Gamma_{mk}^l - \partial_m \Gamma_{jk}^l + \Gamma_{jp}^l \Gamma_{mk}^p - \Gamma_{mp}^l \Gamma_{jk}^p.$$

Instead, we shall use the defining equation below to simplify computations:

$$[\nabla_j, \nabla_m] V^l = R_{jm}{}^l{}_k V^k$$

$\Rightarrow \delta([\nabla_j, \nabla_m] V^l) = \delta(R_{jm}{}^l{}_k V^k) = (\delta R_{jm}{}^l{}_k) V^k$ , since  $\{V^k\}$  is a fixed vector field thus independent of variations. On the other hand:

$$\begin{aligned} \delta([\nabla_j, \nabla_m] V^l) &= \delta(\nabla_j \nabla_m V^l - \nabla_m \nabla_j V^l) \\ &= (\delta \nabla_j) \nabla_m V^l + \nabla_j (\delta \nabla_m) V^l - (\delta \nabla_m) \nabla_j V^l - \nabla_m (\delta \nabla_j) V^l \\ &= ((\delta \nabla_j) \nabla_m V^l - \nabla_m (\delta \nabla_j) V^l) + (\nabla_j (\delta \nabla_m) V^l - (\delta \nabla_m) \nabla_j V^l) \\ &= [\delta \nabla_j, \nabla_m] V^l + [\nabla_j, \delta \nabla_m] V^l \end{aligned}$$

Observation: (to be shown in the remark after the verification of these formulae)

If  $W_m^l$  is a (1,1)-tensor, then  $\nabla_j (W_m^l) = \partial_j W_m^l + \Gamma_{jk}^l W_m^k - \Gamma_{jm}^k W_k^l$ . Hence viewing  $\delta \nabla_j$  as an (infinitesimal) difference of two connections, which is a tensor, we have:

$$(\delta \nabla_j) \nabla_m V^l = \underline{\delta \Gamma_{jk}^l} \nabla_m V^k - \delta \Gamma_{jm}^k \nabla_k V^l.$$

while  $\nabla_m (\delta \nabla_j) V^l = \nabla_m (\delta \Gamma_{jk}^l V^k) = (\nabla_m \delta \Gamma_{jk}^l) \cdot V^k + \underline{\delta \Gamma_{jk}^l} \cdot \nabla_m V^k$ . Note the underlined terms cancel when subtracting. Furthermore, the expression

$$[\delta \nabla_j, \nabla_m] V^l + [\nabla_j, \delta \nabla_m] V^l = [\delta \nabla_j, \nabla_m] V^l - [\delta \nabla_m, \nabla_j] V^l$$

is anti-symmetric in  $j, m$ . It follows that  $-\delta \Gamma_{jm}^k \nabla_k V^l$ , which is symmetric in  $j, m$ , will eventually get cancelled in the final expression. Hence

$$\delta R_{jm}{}^l{}_k V^k = -(\nabla_m \delta \Gamma_{jk}^l) + \nabla_j \delta \Gamma_{mk}^l V^k$$

Since this is true for arbitrary vector fields, we obtain 2).

$$\delta R_{jm}{}^l{}_k = -(\nabla_m \delta \Gamma_{jk}^l) + \nabla_j \delta \Gamma_{mk}^l$$

Finally, it remains to show 1). the variation of  $\Gamma_{ij}^k$ .

Recall that  $\Gamma_{ij}^k = \frac{1}{2} g^{kl} (\partial_i g_{ej} + \partial_j g_{ie} - \partial_e g_{ij})$ , consequently:

$$\delta \Gamma_{ij}^k = \frac{1}{2} g^{kl} (\partial_i \delta g_{ej} + \partial_j \delta g_{ie} - \partial_e \delta g_{ij}) - \frac{1}{2} g^{ks} \delta g_{sr} g^{rl} (\partial_i g_{ej} + \partial_j g_{ie} - \partial_e g_{ij})$$

Since we know that  $\delta \Gamma_{ij}^k$  should be an (infinitesimal) tensor, the terms as  $\partial_i \delta g_{ej}$  will have to be replaced by tensors like  $\nabla_j \delta g_{ej}$ , and formula 1) follows. More explicitly

$$\frac{1}{2} g^{ks} \delta g_{sr} g^{rl} (\partial_i g_{ej} + \partial_j g_{ie} - \partial_e g_{ij}) = g^{ks} \delta g_{sr} \Gamma_{ij}^r$$

and from taking covariant derivative of (2,0)-tensor  $\delta g_{ij}$ , we have:

$$\begin{aligned}\partial_i(\delta g_{lj}) &= \nabla_i(\delta g_{lj}) + \Gamma_{il}^s \delta g_{sj} + \Gamma_{ij}^s \delta g_{ls} \\ \partial_j(\delta g_{il}) &= \nabla_j(\delta g_{il}) + \Gamma_{ji}^s \delta g_{sl} + \Gamma_{jl}^s \delta g_{is} \\ -\partial_l(\delta g_{ij}) &= -\nabla_l(\delta g_{ij}) - \Gamma_{li}^s \delta g_{sj} - \Gamma_{lj}^s \delta g_{is}\end{aligned}$$

Summing up, it gives:

$$\begin{aligned}\delta \Gamma_{ij}^k &= \frac{1}{2} g^{kl} (\nabla_i \delta g_{lj} + \nabla_j \delta g_{il} - \nabla_l \delta g_{ij}) + g^{kl} \delta g_{ls} \Gamma_{ij}^s - g^{ks} \delta g_{sr} \Gamma_{ij}^r \\ &= \frac{1}{2} g^{kl} (\nabla_i \delta g_{lj} + \nabla_j \delta g_{il} - \nabla_l \delta g_{ij})\end{aligned}$$

and finishes the proof of 1).

Rmk: Throughout, we used the covariant derivative of tensor fields, which is uniquely extended from covariant derivative of vector fields by the rule:

- $\nabla_i T_{i_1 \dots i_s}^{j_1 \dots j_r}$  is a tensor of type  $(r, s+1)$
- $\nabla_i f = \partial_i f$ ,  $\nabla_i v^j = \partial_i v^j + \Gamma_{ik}^j v^k$
- $\nabla_i (T_{i_1 \dots i_s}^{j_1 \dots j_r} \cdot R_{l_1 \dots l_u}^{k_1 \dots k_u}) = (\nabla_i T_{i_1 \dots i_s}^{j_1 \dots j_r}) R_{l_1 \dots l_u}^{k_1 \dots k_u} + T_{i_1 \dots i_s}^{j_1 \dots j_r} \cdot \nabla_i R_{l_1 \dots l_u}^{k_1 \dots k_u}$
- $\nabla_i$  commutes with contraction of indices.

For example:  $(\partial_i v^j) w_j + v^j (\partial_i w_j) = \partial_i (v^j w_j)$

$$= \nabla_i (v^j w_j)$$

by b)

by c) and d)

$$= (\nabla_i v^j) w_j + v^j (\nabla_i w_j)$$

$$= (\partial_i v^j) w_j + \Gamma_{ik}^j v^k w_j + v^j (\nabla_i w_j)$$

$$\Rightarrow v^j (\nabla_i w_j) = v^j \partial_i w_j - v^j \Gamma_{ij}^k w_k$$

$$\Rightarrow \nabla_i w_j = \partial_i w_j - \Gamma_{ij}^k w_k.$$

Sketch of spectral decomposition of Laplacian.

Recall our setting:  $L \rightarrow X$ , holomorphic line bundle with a metric  $h$ .  $g_{\bar{z}z}$  is a metric on  $\Lambda^{1,0}$ , we have  $\bar{\partial}$  and its formal adjoint:

$$\Gamma(X, L) \xrightleftharpoons[\bar{\partial}^*]{\bar{\partial}} \Gamma(X, L \otimes \Lambda^{0,1})$$

adjoint in the sense that  $\langle \bar{\partial} \varphi, \psi \rangle = \langle \varphi, \bar{\partial}^* \psi \rangle$ .

• Key fact: A Priori Estimate:  $\forall \varphi \in \Gamma(X, L)$  a smooth section

$$\|\varphi\|_{(2)} \leq C \cdot (\|\Delta \varphi\|_{(0)} + \|\varphi\|_{(1)})$$

where  $\|\varphi\|_{(s)}$  denotes the Sobolev norm defined below:

Def: (Sobolev norm  $\|\varphi\|_{(s)}$ )

$$\|\varphi\|_{(0)}^2 \triangleq \|\varphi\|^2, \text{ the } L^2\text{-norm of } \varphi.$$

$$\|\varphi\|_{(1)}^2 \triangleq \|\varphi\|_{(0)}^2 + \|\nabla_{\bar{z}}\varphi\|^2 + \|\nabla_z\varphi\|^2$$

$$\|\varphi\|_{(2)}^2 \triangleq \|\varphi\|_{(1)}^2 + \|\nabla_z\nabla_{\bar{z}}\varphi\|^2 + \|\nabla_{\bar{z}}\nabla_z\varphi\|^2 + \|\nabla_z\nabla_z\varphi\|^2 + \|\nabla_{\bar{z}}\nabla_{\bar{z}}\varphi\|^2$$

...

Here  $\|\cdot\|$  are various  $L^2$ -norms in different line bundles. For instance

$$\nabla_{\bar{z}}\varphi \in \Gamma(X, L \otimes \Lambda^{1,0}), \quad \|\nabla_{\bar{z}}\varphi\|^2 \triangleq \int_X \nabla_{\bar{z}}\varphi \overline{\nabla_{\bar{z}}\varphi} h$$

$$\nabla_z\nabla_{\bar{z}}\varphi \in \Gamma(X, L \otimes \Lambda^{0,1} \otimes \Lambda^{1,0}), \quad \|\nabla_z\nabla_{\bar{z}}\varphi\|^2 \triangleq \int_X \nabla_z\nabla_{\bar{z}}\varphi \cdot \overline{\nabla_z\nabla_{\bar{z}}\varphi} h g^{z\bar{z}} dzd\bar{z}$$

In general, in  $n$ -dimensional case  $\|\varphi\|_{(s)} \triangleq \sum_{0 \leq p \leq s} \int_X g^{i_1\bar{i}_1} \dots g^{i_p\bar{i}_p} \nabla_{i_1} \dots \nabla_{i_p}\varphi \cdot \overline{\nabla_{i_1} \dots \nabla_{i_p}\varphi} h \, dv$ .

where  $dv$  is the volume element,  $i_1, \dots, i_p \in \{1, \dots, n, \bar{1}, \dots, \bar{n}\}$ .

w. r. t.  $\|\cdot\|_{(s)}$ , we define  $H_{(s)} \triangleq \overline{\Gamma(X, L)}$ , the completion, the Sobolev space. (This is a stronger sense of convergence by giving restrictions on all  $\leq s$ th derivatives).

A trivial observation:  $\|\Delta\varphi\|_{(0)} \leq C\|\varphi\|_{(2)}$ .

Indeed, recall that  $\bar{\partial}^+\varphi = -g^{z\bar{z}}\nabla_{\bar{z}}\varphi$ ,  $\Delta\varphi = -g^{z\bar{z}}\nabla_z\nabla_{\bar{z}}\varphi$ , which is just a rescaling of  $\nabla_z\nabla_{\bar{z}}\varphi$ .

General a priori estimate:  $\forall s \in \mathbb{N} \cup \{0\}$ ,  $\exists C_s > 0$  s.t.  $\forall \varphi \in \Gamma(X, L)$ .

$$\|\varphi\|_{(s+2)}^2 \leq C_s (\|\Delta\varphi\|_{(s)}^2 + \|\varphi\|_{(s+1)}^2)$$

Two basic lemmas about Sobolev spaces.

• Sobolev lemma:  $X$ : a manifold of real dimension  $n$ . Then  $\forall \varphi \in \Gamma(X, L)$ .

(a).  $\|\varphi\|_{C^k} \leq C \cdot \|\varphi\|_{(k+s)}$ ,  $\forall s > \frac{n}{2}$

(b).  $H_{(k+s)} \subseteq C^k(X, L)$

(c). As a consequence of (b),  $\bigcap_{s \geq 0} H_{(s)} = \Gamma(X, L)$

• Rellich's lemma.  $X$ : compact manifold. Let  $t > s$ . Then a sequence  $\{\varphi_j\}$  in  $H_{(t)}$  with  $\|\varphi_j\| \leq C$  (bounded) has a subsequence converging in  $H_{(s)}$ . (AKA:  $H_{(s)} \hookrightarrow H_{(t)}$  being compact).

Using A Priori Estimate:

Basic observations:

(a). We can extend  $\Delta: \Gamma(X, L) \rightarrow \Gamma(X, L)$  to  $\Delta: H_{(2)} \rightarrow H_{(0)}$ , a continuous map.

Namely,  $\forall \varphi \in H_{(2)}$ ,  $\varphi = \lim \varphi_n$  w.r.t.  $\|\cdot\|_{(2)}$ , then by the trivial observation made above:

$$\|\Delta\varphi_i - \Delta\varphi_j\|_{(0)} \leq C \cdot \|\varphi_i - \varphi_j\|_{(2)} \rightarrow 0 \text{ as } (i, j) \rightarrow \infty.$$

(b). Define the range of  $\Delta$ :  $\text{Range}\Delta = \{\Delta\varphi \mid \varphi \in H_{(2)}\}$ . Then a priori estimate implies that  $\text{Range}\Delta$  is closed.

Claim: A priori estimate  $\Rightarrow \exists C > 0$  s.t.  $\forall \varphi \in \Gamma(X, L)$ ,  $\varphi \perp \text{Ker}\Delta$ , we have

$$\|\varphi\|_{(2)} \leq C \cdot \|\Delta\varphi\|_{(0)}. \quad (*)$$

Then  $\forall \Phi \in \text{Range}\Delta$ ,  $\Phi_n = \Delta\varphi_n$ ,  $\varphi_n \perp \text{Ker}\Delta$ , and  $\Phi_n \rightarrow \Phi$  in  $H_{(0)}$ . Now by (\*)  $\|\varphi_n - \varphi_m\|_{(2)} \leq C \cdot \|\Delta\varphi_n - \Delta\varphi_m\|_{(0)} = C \|\Phi_n - \Phi_m\|_{(0)} \rightarrow 0 \Rightarrow \{\varphi_n\}$  converges in  $H_2$ , say,  $\varphi_n \rightarrow \varphi$ . Then  $\Delta\varphi = \lim \Delta\varphi_n = \Phi$ .

Pf of claim:

Otherwise,  $\forall n \in \mathbb{N}$ ,  $\exists \varphi_n \in \Gamma(X, L)$ ,  $\varphi_n \perp \text{Ker}\Delta$  with  $\|\varphi_n\|_{(2)} \geq n \|\Delta\varphi_n\|_{(0)}$ . Define:

$\psi_n \triangleq \varphi_n / \|\varphi_n\|_{(2)}$ . Then  $\|\Delta\psi_n\|_{(0)} \leq \frac{1}{n}$ ,  $\|\psi_n\|_{(2)} = 1$ .

By a priori estimate,  $\|\psi_n - \psi_m\|_{(2)} \leq D \cdot (\|\Delta\psi_n - \Delta\psi_m\|_{(0)} + \|\psi_n - \psi_m\|_{(1)})$  for some  $D > 0$ .

Rellich's lemma  $\Rightarrow \exists$  subsequence, which we may assume to be  $\{\psi_n\}$  to start with,

s.t.  $\|\psi_n - \psi_m\|_{(1)} \rightarrow 0$  as  $n, m \rightarrow \infty$ .  $\Rightarrow \psi_n \rightarrow \psi$  in  $H_{(2)}$ , with  $\|\psi\|_{(2)} = \lim \|\psi_n\|_{(2)} = 1$ .

and  $\Delta\psi = \lim_n \Delta\psi_n = 0 \Rightarrow \psi \in \text{Ker}\Delta \subseteq H_{(2)}$ . On the other hand,  $\psi_n \in (\text{Ker}\Delta)^\perp \Rightarrow$

$\psi \in (\text{Ker}\Delta)^\perp \subseteq H_{(2)}$ . It follows that  $\psi = 0$ , contradiction to  $\|\psi\|_{(2)} = 1$ .

Now  $\text{Range}\Delta$  being closed allows us to construct an "inverse" of  $\Delta$  as follows. By basic Hilbert space theory,  $H_{(0)} = \text{Range}\Delta \oplus (\text{Range}\Delta)^\perp$ . (In general, if we don't know that  $\text{Range}\Delta$  is closed, we can only say that  $H_{(0)} = \overline{\text{Range}\Delta} \oplus (\text{Range}\Delta)^\perp$ .)

Given  $\psi \in H_{(0)}(X, L)$ ,  $\psi = \Delta\varphi + \psi^\perp$ , and  $\varphi$  is unique by requiring that  $\varphi \perp \text{Ker}\Delta$  (this will be explained in more detail in the next chapter). Then define  $G\psi \triangleq \varphi$ . i.e.

$$\Delta G = I - \text{Pr}_{(\text{Range}\Delta)^\perp}.$$

Finally,  $G: H_{(0)} \rightarrow H_{(2)} \hookrightarrow H_{(0)}$  is a self-adjoint operator on a Hilbert space

and we may talk about eigenvalues of  $G$ . Riesz-Schauder thm  $\Rightarrow \exists$  orthonormal basis of eigen-functions  $\{\varphi_n\}$

In general, we can construct extensions of  $\Delta$  to

$H_{(s+2)} \rightarrow H_{(s)}$  and  $G: H_{(s)} \rightarrow H_{(s+2)}$ , then the general a priori estimate shows that  $\text{Range } \Delta$  is closed and  $G$  is similarly constructed. Now:

$$G\varphi_n = \lambda_n \varphi_n \in H_{(0)} \supseteq H_{(2)}$$

$\Rightarrow \varphi_n \in H_{(2)} \Rightarrow \varphi_n \in H_{(4)} \Rightarrow \dots \Rightarrow \varphi_n \in \bigcap_n H_{(n)} = \Gamma'(X, L)$ , by Sobolev's lemma.

A sketch of A Priori Estimate on Riemann surfaces.

$$\begin{aligned} \forall \varphi \in \Gamma'(X, L), \|\Delta\varphi\|_{(0)} &= \int_X g^{\bar{z}\bar{z}} \nabla_{\bar{z}} \nabla_{\bar{z}} \varphi \overline{\nabla_{\bar{z}} \nabla_{\bar{z}} \varphi} h g^{\bar{z}\bar{z}} g_{\bar{z}\bar{z}} dz d\bar{z} \\ &= \int_X \nabla_{\bar{z}} \nabla_{\bar{z}} \varphi \cdot \overline{\nabla_{\bar{z}} \nabla_{\bar{z}} \varphi} h g^{\bar{z}\bar{z}} dz d\bar{z} \\ &= \|\nabla_{\bar{z}} \nabla_{\bar{z}} \varphi\|^2 \end{aligned}$$

The remaining terms  $\|\nabla_{\bar{z}} \nabla_{\bar{z}} \varphi\|^2$  can also be bounded by  $\|\Delta\varphi\|_{(0)}$ . For example:

$$\begin{aligned} \|\nabla_{\bar{z}} \nabla_{\bar{z}} \varphi\|^2 &= \int_X \nabla_{\bar{z}} \nabla_{\bar{z}} \varphi \overline{\nabla_{\bar{z}} \nabla_{\bar{z}} \varphi} (g^{\bar{z}\bar{z}})^2 h g_{\bar{z}\bar{z}} dz d\bar{z} \\ &= - \int_X \nabla_{\bar{z}} \varphi \overline{\nabla_{\bar{z}} \nabla_{\bar{z}} \nabla_{\bar{z}} \varphi} g^{\bar{z}\bar{z}} h dz d\bar{z} \text{ by integration by parts} \\ &= - \int_X \nabla_{\bar{z}} \varphi \overline{(\nabla_{\bar{z}} \nabla_{\bar{z}} \nabla_{\bar{z}} \varphi + R_{\bar{z}\bar{z}} \nabla_{\bar{z}} \varphi)} g^{\bar{z}\bar{z}} h dz d\bar{z} \\ &= \int_X \nabla_{\bar{z}} \nabla_{\bar{z}} \varphi \overline{\nabla_{\bar{z}} \nabla_{\bar{z}} \varphi} g^{\bar{z}\bar{z}} h dz d\bar{z} + \text{terms in } \|\nabla_{\bar{z}} \varphi\|^2 \\ &= \int_X \nabla_{\bar{z}} \nabla_{\bar{z}} \varphi \cdot \overline{\nabla_{\bar{z}} \nabla_{\bar{z}} \varphi} g^{\bar{z}\bar{z}} h dz d\bar{z} + \text{terms in } \|\nabla_{\bar{z}} \varphi\|^2 \text{ or } \|\nabla_{\bar{z}} \varphi\|^2 \\ &\leq \|\Delta\varphi\|_{(0)} + C \|\varphi\|_{(1)} \end{aligned}$$

$\Rightarrow \|\varphi\|_{(2)} \leq C(\|\Delta\varphi\|_{(0)} + \|\varphi\|_{(1)})$ , as asserted.

## §5. Curvatures on Vector Bundles

Def. of Complex Manifolds

$X = \cup_{\mu} X_{\mu}$ ,  $\Phi_{\mu}: X_{\mu} \rightarrow \mathbb{C}^n$ : local coordinate charts, with  $\Phi_{\mu} \circ \Phi_{\nu}^{-1}|_{\Phi_{\nu}(X_{\mu} \cap X_{\nu})}$  holomorphic, 1-1 and has invertible Jacobian.

Observation: Let  $f: \Omega \rightarrow \mathbb{C}$ ,  $(z_1, \dots, z_n) \mapsto f(z_1, \dots, z_n)$ ,  $\Omega \subseteq \mathbb{C}^n$ . We say that  $f$  is holomorphic if  $f$  is holomorphic in each variable  $z_1, \dots, z_n$ . Such  $f$  is characterized by:

Thm. (Hartog)  $f$  is holomorphic (in the above sense) iff  $f$  can be expanded as a power series near any point  $\zeta \in \Omega$ :

$$f(z) = \sum_{|\alpha|=0}^{\infty} C_{\alpha} (z - \zeta)^{\alpha}$$

for  $|z - \zeta| < \varepsilon$ , some  $\varepsilon > 0$ . □

Notation:  $\alpha$  is the multi-index:  $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$ .  $\zeta^{\alpha} \triangleq \zeta_1^{\alpha_1} \dots \zeta_n^{\alpha_n}$ .  $|\alpha| \triangleq \alpha_1 + \dots + \alpha_n$ .

• There is no such characterization in the smooth category.

Notation:  $F: \Omega \rightarrow \mathbb{C}^n$ ,  $F: (z_1, \dots, z_n) \mapsto (f_1(z), \dots, f_n(z))$ .  $F$  is said to be holomorphic if each of its component  $f_i$  is holomorphic. The Jacobian of  $F$  is defined as:

$$\text{Jac}(F) = \left( \frac{\partial f_i}{\partial z_j} \right)_{n \times n}$$

Holomorphic Vector Bundle

$E \xrightarrow{\pi} X$  is a holomorphic vector bundle of rank  $r$

$\Leftrightarrow X = \cup_{\mu} X_{\mu}$ , with  $\{(t_{\mu\nu}^{\alpha\beta})(z) \text{ on } X_{\mu} \cap X_{\nu}, 1 \leq \alpha, \beta \leq r\}$  with

- $(t_{\mu\nu}^{\alpha\beta})_{r \times r}$  holomorphic and invertible.
- $t_{\mu\nu}^{\alpha\beta} t_{\nu\rho}^{\beta\gamma} = t_{\mu\rho}^{\alpha\gamma}$  on  $X_{\mu} \cap X_{\nu} \cap X_{\rho}$

(Recall that for holomorphic line bundles over a Riemann surface:  $L \rightarrow X$

$\Leftrightarrow \{t_{\mu\nu}(z): X_{\mu} \cap X_{\nu} \rightarrow \mathbb{C}^*$ , holomorphic;  $t_{\mu\nu}(z)t_{\nu\rho}(z) = t_{\mu\rho}(z)$  on  $X_{\mu} \cap X_{\nu} \cap X_{\rho}\}$ .)

(Smooth) Sections of  $E \rightarrow X$ .

$\Gamma(X, E) \ni \varphi \Leftrightarrow \{\varphi_{\mu}^{\alpha}(z): \text{smooth functions defined on } X_{\mu}, 1 \leq \alpha \leq r, \text{ with } (\varphi_{\mu}^1, \dots, \varphi_{\mu}^r)^t \text{ the}$

$\mathbb{C}^r$ -valued function satisfying  $\varphi_{\mu}^{\alpha}(z) = t_{\mu\nu}^{\alpha\beta} \varphi_{\nu}^{\beta}(z)$  on  $X_{\mu} \cap X_{\nu}\}$ .

## Covariant Derivatives of Sections of Holomorphic Vector Bundles

$E \rightarrow X$ : holomorphic vector bundle.  $\Gamma(X, E) \ni \varphi = \{\varphi_\mu\}$

$$\begin{aligned} \bar{\partial}\varphi &\triangleq \left\{ \frac{\partial}{\partial \bar{z}^i} \varphi d\bar{z}^j \right\} \\ &= \left\{ \frac{\partial}{\partial \bar{z}^\mu} \varphi_\mu^\alpha d\bar{z}^\mu \right\} \text{ on } X_\mu \end{aligned}$$

• How do they glue together?

Recall:  $\varphi_\mu^\alpha = t_{\mu\nu}^\alpha{}_\beta \varphi_\nu^\beta$  on  $X_\mu \cap X_\nu$ .

$$\Rightarrow \frac{\partial}{\partial \bar{z}^\mu} \varphi_\mu^\alpha = \frac{\partial}{\partial \bar{z}^\mu} (t_{\mu\nu}^\alpha{}_\beta \varphi_\nu^\beta) = t_{\mu\nu}^\alpha{}_\beta \frac{\partial}{\partial \bar{z}^\mu} \varphi_\nu^\beta = t_{\mu\nu}^\alpha{}_\beta \frac{\partial \bar{z}^\nu}{\partial \bar{z}^\mu} \frac{\partial}{\partial \bar{z}^\nu} \varphi_\nu^\beta.$$

$$\Rightarrow \frac{\partial}{\partial \bar{z}^\mu} \varphi_\mu^\alpha = t_{\mu\nu}^\alpha{}_\beta \left( \frac{\partial}{\partial \bar{z}^\nu} \varphi_\nu^\beta \right) \frac{\partial \bar{z}^\nu}{\partial \bar{z}^\mu}$$

i.e. it's a section of  $E \otimes \Lambda^{0,1}$ , where  $\Lambda^{0,1}$  is the vector bundle on  $X$  with transition functions  $\tilde{t}_{\mu\nu}{}^k{}_j = \frac{\partial \bar{z}^\nu}{\partial \bar{z}^\mu}$  (transforms as row vectors, so that  $\left\{ \frac{\partial}{\partial \bar{z}^\mu} \varphi_\mu^\alpha d\bar{z}^\mu \right\}$  is invariant).

To differentiate in the  $z^j$  direction, we need a connection. An important connection is a unitary connection:

• Let  $\{H_{\bar{\mu}\alpha}\}$  be a metric on  $E \rightarrow X$ , i.e. given  $\varphi \in \Gamma(X, E)$ , we associate it its length

$$\|\varphi\|^2 = \left\{ \|\varphi\|_\mu^2 \triangleq (H_\mu)_{\bar{\mu}\alpha}(z) \varphi_\mu^\alpha \overline{\varphi_\mu^\beta} \text{ s.t. } \|\varphi\|_\mu^2 = \|\varphi\|_\nu^2 \text{ on } X_\mu \cap X_\nu \right\}$$

$$\Rightarrow (H_\mu)_{\bar{\mu}\alpha} \varphi_\mu^\alpha \overline{\varphi_\mu^\beta} = (H_\nu)_{\bar{\nu}\delta} \varphi_\nu^\delta \overline{\varphi_\nu^\gamma} \text{ on } X_\mu \cap X_\nu$$

$$\Rightarrow (H_\mu)_{\bar{\mu}\alpha} t_{\mu\nu}^\alpha{}_\delta \varphi_\nu^\delta \overline{t_{\mu\nu}^\beta{}_\gamma \varphi_\nu^\gamma} = (H_\nu)_{\bar{\nu}\delta} \varphi_\nu^\delta \overline{\varphi_\nu^\gamma} \text{ on } X_\mu \cap X_\nu$$

$$\text{Hence } (H_\mu)_{\bar{\mu}\alpha} t_{\mu\nu}^\alpha{}_\delta t_{\mu\nu}^\beta{}_\gamma = (H_\nu)_{\bar{\nu}\delta} \quad (*)$$

Def. A metric on  $E \rightarrow X$  is an assignment of  $(H_\mu)_{\bar{\mu}\alpha}$  on  $X_\mu$  s.t.  $(*)$  holds and  $\|\varphi\|^2 \geq 0$ , " $=$ " iff  $\varphi = 0$ .

Observation: a short hand for a metric is that:

$$\|\varphi\|_\mu^2 = (\dots \overline{\varphi_\mu^\alpha} \dots) \left( (H_\mu)_{\bar{\mu}\beta} \right) \begin{pmatrix} \varphi_\mu^\beta \\ \vdots \\ \varphi_\mu^\alpha \end{pmatrix}$$

so that if  $\begin{pmatrix} \varphi_\mu^\beta \\ \vdots \\ \varphi_\mu^\alpha \end{pmatrix} = (t_{\mu\nu}^\beta{}_\gamma) \begin{pmatrix} \varphi_\nu^\gamma \\ \vdots \\ \varphi_\nu^\alpha \end{pmatrix}$ , then

$$(H_\nu)_{\bar{\nu}\delta} = \overline{t_{\mu\nu}^\beta{}_\gamma} (H_\mu)_{\bar{\mu}\alpha} t_{\mu\nu}^\alpha{}_\delta$$

i.e. if we define  $(t_{\mu\nu}^\dagger)_{\bar{\beta}}{}^\gamma = \overline{(t_{\mu\nu}^\beta{}_\gamma)}$ , then  $H_\nu = t_{\mu\nu}^\dagger H_\mu t_{\mu\nu}$ .

Now given a metric  $H_{\bar{\mu}\beta}$  on  $E$ , we can define the corresponding covariant derivative on  $\Gamma(X, E)$ :

Def: (Connection, unitary).  $\varphi \in \Gamma(X, E)$ ,  $\nabla\varphi \triangleq \{ \nabla_j \varphi dz^j \}$ , where  $\nabla_j \varphi = H^{\alpha\bar{\beta}} \frac{\partial}{\partial z^i} (H_{\bar{\beta}\gamma} \varphi^\gamma)$ .

Here  $H^{\alpha\bar{\beta}} H_{\bar{\beta}\gamma} = \delta^\alpha{}_\gamma$ .  $\nabla: \Gamma(X, E) \rightarrow \Gamma(X, E \otimes \Lambda^{1,0})$  is the unitary connection on  $E$ .

This definition makes sense since  $\{H_{\bar{\nu}\nu}\varphi^\nu\}$  is a section of the bundle  $E^+$ , which is anti-holomorphic and thus  $\{\partial_j H_{\bar{\nu}\nu}\varphi^\nu\} \in \Gamma(X, E^+ \otimes \Lambda^{1,0})$ , tensoring with  $H^{\alpha\bar{\beta}}$  gives us a section of  $E \otimes \Lambda^{1,0}$ .

$$\begin{aligned} \text{Now, } \nabla_j \varphi^\alpha &= H^{\alpha\bar{\beta}} \{H_{\bar{\nu}\nu} \partial_j \varphi^\nu + (\partial_j H_{\bar{\nu}\nu}) \varphi^\nu\} \\ &= \delta^\alpha_\nu \partial_j \varphi^\nu + (H^{\alpha\bar{\beta}} \partial_j H_{\bar{\beta}\nu}) \varphi^\nu \\ &= \partial_j \varphi^\alpha + (H^{\alpha\bar{\beta}} \partial_j H_{\bar{\beta}\nu}) \varphi^\nu \quad (*) \end{aligned}$$

Denote  $A_{j\nu}^\alpha = H^{\alpha\bar{\beta}} \partial_j H_{\bar{\beta}\nu}$ ,  $1 \leq \alpha, \beta \leq r$ ,  $1 \leq j \leq n$ .

Def. (Connection, general form). A connection on  $E$  is an assignment  $\{A_{j\nu}^\alpha\}$  satisfying the requirement that  $(*)$  defines a section in  $\Gamma(X, E \otimes \Lambda^{1,0})$ .

Thus, a unitary connection is a connection with  $A_{j\nu}^\alpha = H^{\alpha\bar{\beta}} \partial_j H_{\bar{\beta}\nu}$ . View  $A_{j\nu}^\alpha$  as entries of the matrix  $A_j = H^{-1} \partial_j H$ .

It's also convenient to introduce the connection form  $A \triangleq A_j dz^j$ . However,  $A$  is not globally defined, i.e.  $A_j$  doesn't transform as a global  $(1,0)$ -form under change of coordinates. This is readily seen since by  $(*)$ ,  $\nabla_j \varphi^\alpha$  transforms like a tensor while  $\partial_j \varphi^\alpha$  doesn't.

In the special case of line bundles,  $\text{rank } E = 1$ ,  $H = h$  is a complex scalar function, thus we have our former  $A_j = h^{-1} \partial_j h = \partial_j (\log h)$ .

Commutation Rules for Covariant Derivative.

$$\begin{aligned} \text{We compute } \nabla_{\bar{k}} \nabla_j \varphi^\alpha - \nabla_j \nabla_{\bar{k}} \varphi^\alpha &= \nabla_{\bar{k}} (\partial_j \varphi^\alpha + A_{j\nu}^\alpha \varphi^\nu) - \nabla_j (\partial_{\bar{k}} \varphi^\alpha) \\ &= \partial_{\bar{k}} \partial_j \varphi^\alpha + \partial_{\bar{k}} (A_{j\nu}^\alpha \varphi^\nu) - \partial_j \partial_{\bar{k}} \varphi^\alpha - A_{j\nu}^\alpha \partial_{\bar{k}} \varphi^\nu \\ &= (\partial_{\bar{k}} A_{j\nu}^\alpha) \varphi^\nu \end{aligned}$$

Def. (Curvature of  $E$  w.r.t.  $H_{\bar{\alpha}\beta}$ )

$F_{\bar{k}j}^\alpha \triangleq -\partial_{\bar{k}} A_{j\nu}^\alpha = -\partial_{\bar{k}} (H^{\alpha\bar{\beta}} \partial_j H_{\bar{\beta}\nu}) = [-\partial_{\bar{k}} (H^{-1} \partial_j H)]^\alpha_\nu$  is the curvature associated with metric  $H_{\bar{\alpha}\beta}$ . Thus in summary, we have the key formula:

$$\bullet [\nabla_{\bar{k}}, \nabla_j] \varphi^\alpha = -F_{\bar{k}j}^\alpha \varphi^\nu$$



Invariant Point of View.

Given a bundle  $E \rightarrow X$ , we can construct the bundle  $\text{End}(E) \rightarrow X$  as follows:

$\Gamma(X, \text{End}E) \ni T \Leftrightarrow \{T_{\mu\beta}^{\alpha}(z)\}$ , smooth matrix-valued function defined on  $X_{\mu}$ , satisfying the condition that  $\{T_{\mu\beta}^{\alpha} \varphi_{\mu}^{\beta}\}$  is a globally defined section of  $\Gamma(X, E)$  whenever  $\varphi = \{\varphi_{\mu}^{\alpha}\} \in \Gamma(X, E)$

More explicitly,  $T_{\mu\beta}^{\alpha} \varphi_{\mu}^{\beta} = t_{\mu\nu}^{\alpha\gamma} T_{\nu\delta}^{\gamma} \varphi_{\nu}^{\delta}$  on  $X_{\mu} \cap X_{\nu}$ . But since  $\varphi_{\mu}^{\beta} = t_{\mu\nu}^{\beta\delta} \varphi_{\nu}^{\delta}$ , we have

$$T_{\mu\beta}^{\alpha} \varphi_{\mu}^{\beta} = T_{\mu\beta}^{\alpha} t_{\mu\nu}^{\beta\delta} \varphi_{\nu}^{\delta} = t_{\mu\nu}^{\alpha\gamma} T_{\nu\delta}^{\gamma} \varphi_{\nu}^{\delta}$$

Hence:  $T_{\mu\beta}^{\alpha} t_{\mu\nu}^{\beta\delta} = t_{\mu\nu}^{\alpha\gamma} T_{\nu\delta}^{\gamma}$

Or more compactly, in matrix form:  $T_{\mu} = t_{\mu\nu} T_{\nu} t_{\mu\nu}^{-1}$

Observation:  $F_{\bar{k}j}$  is a section of  $\Gamma(X, \text{End}E)$ . This follows by def. of  $\text{End}E$  and the key formula above.

We may also introduce the curvature form  $F = F_{\bar{k}j} dz^i \wedge d\bar{z}^k$ , which is a section of  $\Gamma(X, \text{End}E \otimes \Lambda^{1,1})$ . First let's review:

Formalism of Differential Forms

Digression: de Rham complex

Recall that if  $X$  is a smooth manifold, then we have the notions:

- $\Gamma(X, \Lambda^p) = \{ \frac{1}{p!} \varphi_{i_1 \dots i_p} dx^{i_1} \wedge \dots \wedge dx^{i_p} \}$
- The exterior derivative:  $d: \Gamma(X, \Lambda^p) \rightarrow \Gamma(X, \Lambda^{p+1})$ .

Also recall that  $d$  is defined by:

(1). On functions,  $df \triangleq \sum_i \frac{\partial f}{\partial x^i} dx^i$  in a local coordinate chart.

(2). On higher forms, it's extended by:  $d(\frac{1}{p!} \sum_I \varphi_I dx^I) = \frac{1}{p!} \sum_I \frac{\partial \varphi_I}{\partial x^i} dx^i \wedge dx^I$

This local definition is well-defined globally. For instance, on different coordinate neighborhoods,  $\sum \varphi_i dx^i = \sum \tilde{\varphi}_j dy^j \Rightarrow \varphi_i = \tilde{\varphi}_j \frac{\partial y^j}{\partial x^i}$ . Thus, by def.

$$\begin{aligned} d(\sum \varphi_i dx^i) &= \sum_{ij} \frac{\partial \varphi_i}{\partial x^j} dx^j \wedge dx^i \\ &= \frac{1}{2} \sum_{ij} (\frac{\partial \varphi_i}{\partial x^j} - \frac{\partial \varphi_j}{\partial x^i}) dx^j \wedge dx^i \\ &= \frac{1}{2} \sum_{ijk} (\frac{\partial \varphi_k}{\partial x^j} \frac{\partial y^k}{\partial x^i} + \tilde{\varphi}_k \frac{\partial^2 y^k}{\partial x^i \partial x^j} - \frac{\partial \tilde{\varphi}_k}{\partial x^i} \frac{\partial y^k}{\partial x^j} - \tilde{\varphi}_k \frac{\partial^2 y^k}{\partial x^j \partial x^i}) dx^i \wedge dx^j \\ &= \frac{1}{2} \sum_{ijk\ell} (\frac{\partial \tilde{\varphi}_k}{\partial y^{\ell}} \frac{\partial y^{\ell}}{\partial x^j} \frac{\partial y^k}{\partial x^i} - \frac{\partial \tilde{\varphi}_k}{\partial y^{\ell}} \frac{\partial y^{\ell}}{\partial x^i} \frac{\partial y^k}{\partial x^j}) dx^i \wedge dx^j \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2} \sum_{k,l} \left( \frac{\partial \hat{\varphi}_k}{\partial y^l} - \frac{\partial \hat{\varphi}_l}{\partial y^k} \right) dy^l \wedge dy^k \\
&= d \left( \sum_k \hat{\varphi}_k dy^k \right).
\end{aligned}$$

Now in our case,  $E \rightarrow X$  is a vector bundle, we can similarly define, with a fixed connection  $A$  (associated with a Hermitian metric)

- $\Gamma(X, E) \ni \varphi \xrightarrow{d_A} d_A \varphi \triangleq \nabla_j \varphi dz^j + \nabla_{\bar{j}} \varphi d\bar{z}^{\bar{j}} \in \Gamma(X, E \otimes \Lambda^1)$ .
- $d_A \left( \frac{1}{p!q!} \sum \varphi_{\bar{j}_1 \dots \bar{j}_q} dz^i d\bar{z}^{\bar{j}} \right) \triangleq \frac{1}{p!q!} (d_A \varphi_{\bar{j}_1 \dots \bar{j}_q}) dz^i d\bar{z}^{\bar{j}} \quad I = (i_1, \dots, i_p), \quad \bar{J} = (\bar{j}_1, \dots, \bar{j}_q)$

In this notation,  $\varphi_{\bar{j}_1 \dots \bar{j}_q} dz^i d\bar{z}^{\bar{j}} = \varphi_{\bar{j}_2 \dots \bar{j}_q, i_1 \dots i_p} dz^{i_1} \dots dz^{i_p} d\bar{z}^{\bar{j}_1} \dots d\bar{z}^{\bar{j}_q}$ .

Now, in this formalism, we have the following basic identities.

- $d_A^2 \varphi = F \wedge \varphi$
- $F = dA + A \wedge A$
- $d_A F = 0$  (the 2<sup>nd</sup>- Bianchi's identity)

Proof of Identities.

Take  $\varphi \in \Gamma(X, E)$ , we compute:

$$d_A \varphi = \nabla_{\bar{k}} \varphi d\bar{z}^{\bar{k}} + \nabla_k \varphi dz^k$$

$$d_A^2 \varphi = (d_A \nabla_{\bar{k}} \varphi) d\bar{z}^{\bar{k}} + (d_A \nabla_k \varphi) dz^k$$

$$= (\nabla_{\bar{j}} \nabla_{\bar{k}} \varphi dz^j + \nabla_{\bar{j}} \nabla_{\bar{k}} \varphi d\bar{z}^{\bar{j}}) d\bar{z}^{\bar{k}} + (\nabla_j \nabla_k \varphi dz^j + \nabla_j \nabla_k \varphi d\bar{z}^{\bar{j}}) dz^k$$

$$= \underbrace{\nabla_{\bar{j}} \nabla_{\bar{k}} \varphi d\bar{z}^{\bar{j}} d\bar{z}^{\bar{k}}}_{\textcircled{1}} + \underbrace{\nabla_{\bar{j}} \nabla_k \varphi dz^j d\bar{z}^{\bar{k}} + \nabla_j \nabla_k \varphi d\bar{z}^{\bar{j}} dz^k}_{\textcircled{2}} + \underbrace{\nabla_j \nabla_k \varphi dz^j dz^k}_{\textcircled{3}}$$

$$\begin{aligned}
\textcircled{1} &= \frac{1}{2} (\nabla_{\bar{j}} \nabla_{\bar{k}} \varphi d\bar{z}^{\bar{j}} d\bar{z}^{\bar{k}} - \nabla_{\bar{k}} \nabla_{\bar{j}} \varphi d\bar{z}^{\bar{j}} d\bar{z}^{\bar{k}}) \\
&= \frac{1}{2} (\partial_{\bar{j}} \partial_{\bar{k}} \varphi d\bar{z}^{\bar{j}} d\bar{z}^{\bar{k}} - \partial_{\bar{k}} \partial_{\bar{j}} \varphi d\bar{z}^{\bar{j}} d\bar{z}^{\bar{k}}) \\
&= 0
\end{aligned}$$

$$\begin{aligned}
\textcircled{2} &= \frac{1}{2} (\nabla_{\bar{j}} \nabla_k \varphi - \nabla_k \nabla_{\bar{j}} \varphi) dz^j d\bar{z}^{\bar{k}} \\
&= \frac{1}{2} (\partial_{\bar{j}} (\partial_k \varphi + A_k \varphi) + A_j (\partial_k \varphi + A_k \varphi) - \text{terms } (j \leftrightarrow k)) dz^j d\bar{z}^{\bar{k}} \\
&= \frac{1}{2} (\underbrace{\partial_{\bar{j}} \partial_k \varphi}_{\text{sym. in } j, k} + (\partial_{\bar{j}} A_k) \varphi + \underbrace{A_k \partial_{\bar{j}} \varphi}_{\text{sym. in } j, k} + A_j \partial_k \varphi + A_j A_k \varphi - \text{terms } (j \leftrightarrow k)) dz^j d\bar{z}^{\bar{k}}
\end{aligned}$$

$$= \frac{1}{2} [(\partial_j A_k - \partial_k A_j) + (A_j A_k - A_k A_j)] \varphi dz^j dz^k$$

*Claim*  
 $= 0$  since  $A_j$  is induced from a Hermitian metric.

Proof of claim:

$$\partial_j A_k = \partial_j (H^{-1} \partial_k H) = -H^{-1} (\partial_j H) H^{-1} \partial_k H$$

(Recall that  $H^{-1} H = I \Rightarrow (\partial_j H^{-1}) H + H^{-1} \partial_j H = 0 \Rightarrow \partial_j H^{-1} = -H^{-1} \partial_j H H^{-1}$ )

Hence  $(\partial_j A_k - \partial_k A_j) + (A_j A_k - A_k A_j)$

$$= -H^{-1} (\partial_j H) H^{-1} (\partial_k H) + H^{-1} (\partial_k H) H^{-1} (\partial_j H) + (H^{-1} \partial_j H) (H^{-1} \partial_k H) - (H^{-1} \partial_k H) (H^{-1} \partial_j H)$$

$$= 0$$

$$\begin{aligned} \textcircled{3} &= \nabla_j \nabla_{\bar{k}} \varphi dz^j d\bar{z}^k + \nabla_{\bar{j}} \nabla_k \varphi d\bar{z}^j dz^k \\ &= \nabla_j \nabla_{\bar{k}} \varphi dz^j d\bar{z}^k - \nabla_{\bar{k}} \nabla_j \varphi d\bar{z}^j dz^k \\ &= [\nabla_j \nabla_{\bar{k}}] \varphi dz^j d\bar{z}^k \\ &= F_{\bar{k}j} \varphi dz^j d\bar{z}^k \end{aligned}$$

Summarizing, we have:  $d_A^2 \varphi = F_{\bar{k}j} \varphi dz^j d\bar{z}^k = F \wedge \varphi$

Next, we observe that:

$$\begin{aligned} dA &= d(\sum A_j dz^j) \\ &= \sum dA_j dz^j \\ &= \sum \partial_{\bar{k}} A_j d\bar{z}^k dz^j + \partial_k A_j dz^k dz^j \\ &= \frac{1}{2} \sum (\partial_k A_j - \partial_j A_k) dz^k dz^j + \sum \partial_{\bar{k}} A_j d\bar{z}^k dz^j \\ &= \frac{1}{2} (A_j A_k - A_k A_j) dz^k dz^j + F_{\bar{k}j} d\bar{z}^k dz^j \quad (\text{by the claim above}) \\ &= A_j A_k dz^k dz^j + F_{\bar{k}j} d\bar{z}^k dz^j \\ &= -A_j dz^j A_k dz^k + F_{\bar{k}j} d\bar{z}^k dz^j \\ &= -A \wedge A + F \end{aligned}$$

$\Rightarrow F = dA + A \wedge A$ , as asserted.

Thirdly,  $dF = d(dA + A \wedge A)$

$$\begin{aligned}
&= dA \wedge A - A \wedge dA \\
&= (F - A \wedge A) \wedge A - A \wedge (F - A \wedge A) \\
&= F \wedge A - A \wedge F
\end{aligned}$$

Or equivalently,  $dF + A \wedge F - F \wedge A = 0$ .

Observation:  $d_A F = dF + A \wedge F - F \wedge A$ .

Recall that  $F \in \Gamma(X, \text{End} E \otimes \wedge^2 T^* X)$ ,  $F = F_{\bar{r}j} dz^j d\bar{z}^k$ , then  $d_A F = (d_A F_{\bar{r}j}) dz^j d\bar{z}^k$ . The observation comes from the simple fact a connection on  $E$  determines a connection on  $\text{End} E \cong E \otimes E^*$  (In fact, over the tensor algebra of  $E$  and  $E^*$ ). Let  $T \in \Gamma(X, \text{End} E)$ ,  $\varphi \in \Gamma(X, E)$ , then  $T\varphi \in \Gamma(X, E)$ , and:

$$\begin{aligned}
\nabla_j (T\varphi) &= (\nabla_j T)\varphi + T\nabla_j \varphi \\
\Rightarrow (\nabla_j T)\varphi &= \nabla_j (T\varphi) - T\nabla_j \varphi
\end{aligned}$$

Explicitly:

$$\begin{aligned}
(\nabla_j T)\varphi &= \partial_j (T\varphi) + A_j T\varphi - T\partial_j \varphi - T A_j \varphi \\
&= (\partial_j T)\varphi + T\partial_j \varphi + A_j T\varphi - T\partial_j \varphi - T A_j \varphi \\
&= (\partial_j T + A_j T - T A_j)\varphi \\
\Rightarrow \nabla_j T &= \partial_j T + A_j T - T A_j \\
\Rightarrow d_A T &= dT + A \wedge T - T \wedge A.
\end{aligned}$$

Componentwise:  $\nabla_j T^\alpha_\beta = \partial_j T^\alpha_\beta + A_j^\alpha_\nu T^\nu_\beta - T^\alpha_\nu A_j^\nu_\beta$ . Notice that  $-A_j^\nu_\beta$  comes from the connection of  $E^*$ . This finishes the proof of the basic equalities.

Special case of  $E = T^{1,0} X$ .

$\Gamma(X, T^{1,0} X) = \{V^j \partial_j \mid \text{holomorphic vector fields}\}$

$\Gamma(X, T^{1,0} X) \ni V \leftrightarrow V^j_\mu(z)$  on  $X_\mu$  s.t.  $V^j_\mu(z) = \frac{\partial z^j_\mu}{\partial z^k} V^k_\nu(z)$ , i.e.  $t_{\mu\nu}^j k = \frac{\partial z^j_\mu}{\partial z^k}$ .

In this case, the connection takes the form  $A_j^\alpha_\beta = A_j^l_k$  ( $j, k, l \in \{1, \dots, n\}$ ), and the curvature takes the form  $F_{\bar{r}j}^\alpha_\beta = F_{\bar{r}j}^{lm}$  ( $j, k, l, m \in \{1, \dots, n\}$ ).

Def. A metric  $g_{\bar{r}j}$  on  $T^{1,0} X$  is said to be Kähler if  $\partial_l g_{\bar{r}j} = \partial_j g_{\bar{r}l}$ .

This condition is invariant under change of coordinates because it's equivalent to the global condition  $0 = d\omega = d(\sum g_{\bar{r}j} dz^j d\bar{z}^k)$ .

For a Kähler metric  $g_{\bar{k}j}$  (we denote the curvature for tangent bundles by  $R$  instead of  $K$ ), we have:

$$R_{\bar{k}j\bar{p}m} = R_{\bar{p}m\bar{k}j} = R_{\bar{k}m\bar{p}j} \quad (\text{the 1st Bianchi identity})$$

where  $R_{\bar{k}j\bar{p}m} = g_{\bar{p}l} R_{\bar{k}j}{}^l{}_m$ .

Proof.  $R_{\bar{k}j\bar{p}m} = g_{\bar{p}l} R_{\bar{k}j}{}^l{}_m$

$$= g_{\bar{p}l} (-\partial_{\bar{k}} (g^{l\bar{q}} \partial_j g_{\bar{q}m}))$$

$$= g_{\bar{p}l} (-\partial_{\bar{k}} g^{l\bar{q}} \partial_j g_{\bar{q}m} - g^{l\bar{q}} \partial_{\bar{k}} \partial_j g_{\bar{q}m})$$

$$= g_{\bar{p}l} (g^{l\bar{s}} (\partial_{\bar{k}} g_{\bar{s}r}) g^{r\bar{q}} \partial_j g_{\bar{q}m} - g^{l\bar{q}} \partial_{\bar{k}} \partial_j g_{\bar{q}m})$$

$$= \partial_{\bar{k}} g_{\bar{p}r} g^{r\bar{q}} \partial_j g_{\bar{q}m} - \partial_{\bar{k}} \partial_j g_{\bar{p}m}$$

By def. of Kähler metrics,  $(j \leftrightarrow m)$  ( $\bar{k} \leftrightarrow \bar{p}$ ) doesn't change  $R_{\bar{k}j\bar{p}m}$ .

All of the above formalism extends trivially to the case of smooth bundles. Consider  $E \rightarrow X$  a smooth complex vector bundle over a smooth manifold. By def., a smooth vector bundle is defined by smooth transition functions  $\{t_{\mu\nu} : \text{smooth invertible matrix valued functions on } X_\mu \cap X_\nu\}$ .

Def: A connection on  $E \rightarrow X = \cup_\mu X_\mu$  is given by  $A_\mu = A_j^\alpha{}_\beta dx^j$   $1 \leq \alpha, \beta \leq r$  on  $X_\mu$  satisfying:  $\Gamma(X, E) \ni \varphi = \{\varphi^\alpha\} \mapsto \nabla_j \varphi = \partial_j \varphi^\alpha + A_j^\alpha{}_\beta \varphi^\beta \in \Gamma(X, E \otimes \Lambda^1)$ .

Def: The curvature tensor is defined by:

$$[\nabla_i, \nabla_j] \varphi^\alpha = -F_{ij}{}^\alpha{}_\beta \varphi^\beta$$

It follows from a simple computation that

$$F_{ij}{}^\alpha{}_\beta = -(\partial_i A_j^\alpha{}_\beta - \partial_j A_i^\alpha{}_\beta + A_i^\alpha{}_\gamma A_j^\gamma{}_\beta - A_j^\alpha{}_\gamma A_i^\gamma{}_\beta).$$

If  $X$  has more structure, say,  $X$  is a complex manifold,  $x \mapsto (z, \bar{z})$ , we are interested in a more special class of connections, which respect the complex structure of  $X$  as much as possible.

Def.  $A = A_j^\alpha{}_\beta dx^j$  is called a Chern connection if  $A_j^\alpha{}_\beta d\bar{z}^j = 0$ , i.e.  $\nabla_j \varphi^\alpha = \partial_j \varphi^\alpha$ .

Assume that  $A$  is a Chern connection, i.e.  $A_{\bar{j}}$  is 0, we have

$$\begin{aligned} F_{\bar{j}} &= -(\partial_{\bar{k}} A_j - \underbrace{\partial_j A_{\bar{k}}}_{0} + \underbrace{A_{\bar{k}} A_j}_{0} - \underbrace{A_j A_{\bar{k}}}_{0}) \\ &= -\partial_{\bar{k}} A_j. \end{aligned}$$

However,  $F_{ij}$  may not be 0. However, we have the following characterization:

Thm. (Newlander & Nirenberg) If  $A$  is a Chern connection with  $F_{ij}=0$  ( $F_{\bar{i}\bar{j}}=0$  by def. of Chern connection), then  $E \rightarrow X$  admits a holomorphic structure.

For a proof, c.f. Donaldson-Kronheimer, The Geometry of Four Manifolds)

### Characteristic Classes

$E \rightarrow X$ : smooth complex vector bundle over compact  $X$ . Let  $A$  be any connection on  $E$

Def:  $\tilde{C}_m(A) = \text{tr}(\wedge^m F) = \text{tr}(\underbrace{F \wedge F \wedge \dots \wedge F}_m) \in \Gamma(X, \wedge^{2m})$

where  $F = F_{ij} dx^i dx^j \in \Gamma(X, \text{End} E \otimes \wedge^2)$ .

Basic observation:  $d\tilde{C}_m(A) = 0$

$$\begin{aligned} \text{Indeed, } d\tilde{C}_m(A) &= \text{Tr}(d(F \wedge \dots \wedge F)) \\ &= \text{Tr}(dF \wedge F \wedge \dots \wedge F + F \wedge dF \wedge \dots \wedge F + \dots + F \wedge F \wedge \dots \wedge dF) \\ &= \text{Tr}(m dF \wedge F \wedge \dots \wedge F) \quad (\text{since } \text{Tr}(AB) = \text{Tr}(BA)) \\ &= \text{Tr}(m(F \wedge A - A \wedge F) \wedge F \wedge \dots \wedge F) \quad (2^{\text{nd}} \text{ Bianchi's identity}) \\ &= \text{Tr}(m(A \wedge F \wedge \dots \wedge F - A \wedge F \wedge \dots \wedge F)) \\ &= 0. \end{aligned}$$

Def. We define the Chern classes of  $E$  as  $[\tilde{C}_m(A)]$ .

Claim:  $[\tilde{C}_m(A)]$  is independent of choices of  $A$ . More precisely, if  $A'$  is any other connection, we can write:

$$\tilde{C}_m(A') - \tilde{C}_m(A) = d(m \int_0^1 \text{Tr}(B \wedge F_t^{m-1}) dt)$$

where  $B = A' - A$ ,  $A_t = A + tB$  and  $F_t$  is the curvature of the connection  $A_t$ . Note that  $B$  is an  $\text{End} E$ -valued 1-form. ( $\because \forall \varphi \in \Gamma(X, E)$ ,  $\partial_j \varphi + A_j \varphi$  and  $\partial_j \varphi + A'_j \varphi$  are

both global  $\Rightarrow A_j \varphi - A'_j \varphi$  is global  $\Rightarrow A_j - A'_j \in \Gamma(X, \text{End} E)$ .

Proof.

$$\begin{aligned} \tilde{C}_m(A') - \tilde{C}_m(A) &= \int_0^1 \frac{d\tilde{C}_m(A_t)}{dt} dt \\ &= \int_0^1 \frac{d}{dt} (\text{Tr}(F_t \wedge \dots \wedge F_t)) dt \\ &= \int_0^1 \text{Tr}(\dot{F}_t \wedge F_t \wedge \dots \wedge F_t + F_t \wedge \dot{F}_t \wedge \dots \wedge F_t + \dots + F_t \wedge F_t \wedge \dots \wedge \dot{F}_t) dt \end{aligned}$$

$$\text{Since } F_t = dA_t + A_t \wedge A_t \Rightarrow \dot{F}_t = d\dot{A}_t + \dot{A}_t \wedge A_t + A_t \wedge \dot{A}_t.$$

$$\Rightarrow \dot{F}_t = dB + B \wedge A_t + A_t \wedge B$$

$$\Rightarrow \text{Tr}(\dot{F}_t \wedge F_t \wedge \dots \wedge F_t + F_t \wedge \dot{F}_t \wedge \dots \wedge F_t + \dots + F_t \wedge F_t \wedge \dots \wedge \dot{F}_t)$$

$$= \text{Tr}(m \dot{F}_t \wedge F_t \wedge \dots \wedge F_t)$$

$$= \text{Tr}(m(dB + B \wedge A_t + A_t \wedge B) \wedge F_t \wedge \dots \wedge F_t)$$

$$= m \text{Tr}(d(B \wedge F_t \wedge \dots \wedge F_t) + B \wedge (\sum_{i=1}^{m-1} \Lambda^i F_t \wedge dF_t \wedge \Lambda^{m-i} F_t) + B \wedge A_t \wedge \Lambda^{m-1} F_t + A_t \wedge B \wedge \Lambda^{m-1} F_t)$$

$$= m \text{Tr}(d(B \wedge \Lambda^{m-1} F_t) + B \wedge (\sum_{i=1}^{m-1} \Lambda^i F_t \wedge (F_t \wedge A_t - A_t \wedge F_t) \wedge \Lambda^{m-i-1} F_t) + B \wedge A_t \wedge \Lambda^{m-1} F_t - B \wedge \Lambda^{m-1} F_t \wedge A_t)$$

$$= m \text{Tr}(d(B \wedge \Lambda^{m-1} F_t) + B \wedge (\sum_{i=1}^{m-1} \Lambda^i F_t \wedge A_t \wedge \Lambda^{m-i-1} F_t - \sum_{i=0}^{m-2} \Lambda^i F_t \wedge A_t \wedge \Lambda^{m-i} F_t + A_t \wedge \Lambda^{m-1} F_t - \Lambda^{m-1} F_t \wedge A_t))$$

$$= m \text{Tr}(d(B \wedge \Lambda^{m-1} F_t) + B \wedge (\Lambda^{m-1} F_t \wedge A_t - A_t \wedge \Lambda^{m-1} F_t + A_t \wedge \Lambda^{m-1} F_t - \Lambda^{m-1} F_t \wedge A_t))$$

$$= m \text{Tr}(d(B \wedge \Lambda^{m-1} F_t))$$

$$= d(m \text{Tr}(B \wedge \Lambda^{m-1} F_t)).$$

Note that in the computation we used the fact that  $\text{Tr}(ABC) = \text{Tr}(BCA)$  and differential forms form a  $\mathbb{Z}/2$ -graded ring. □

Def. The ordinary Chern classes  $C_m(E)$  are obtained from  $\tilde{C}_m(E)$  as follows:

The two basis of symmetric functions are related by polynomial relations:

i. e. if

$$\begin{cases} \sigma_1 = \chi_1 + \dots + \chi_r \\ \sigma_2 = \sum_{i \neq j} \chi_i \chi_j \\ \dots \\ \sigma_r = \chi_1 \dots \chi_r \end{cases} \quad \begin{cases} S_1 = \chi_1 + \dots + \chi_r \\ S_2 = \chi_1^2 + \dots + \chi_r^2 \\ \dots \\ S_r = \chi_1^r + \dots + \chi_r^r \end{cases}$$

then  $S_i = P_i(\sigma_1, \dots, \sigma_r)$ ,  $i=1, \dots, r$ . For instance:  $S_1 = \sigma_1$ ,  $S_2 = \sigma_1^2 - \sigma_2$ , .. ... Then:

$$C_i(E) \cong P_i(\tilde{C}_1(E), \dots, \tilde{C}_r(E)).$$

Note that  $C_1(E)$  is well-defined by the ring structure of de Rham cohomology.

Interlude: Maxwell equations, geometric interpretation.

$$X = \mathbb{R}^{1,3}, \quad ds^2 = -(dt)^2 + dx^2 + dy^2 + dz^2, \quad L = \mathbb{R}^{1,3} \times \mathbb{C}.$$

Let the connection be given by  $A = A_j dx^j = \underbrace{-\varphi dx^0}_{\text{potential}} + \underbrace{A_x dx + A_y dy + A_z dz}_{\text{vector potential}}.$

$$\begin{aligned} F &= dA + A \wedge A \\ &= dA \quad (\text{since } A \text{ is just a 1-form (U(1)-connection).}) \\ &= \sum_{\mu} A_{\mu} dx^{\mu} \\ &= \frac{1}{2} \sum \left( \frac{\partial A_{\mu}}{\partial x^{\nu}} - \frac{\partial A_{\nu}}{\partial x^{\mu}} \right) dx^{\nu} \wedge dx^{\mu} \\ &= \frac{1}{2} F_{\mu\nu} dx^{\nu} \wedge dx^{\mu}, \end{aligned}$$

where we define the curvature  $F_{\mu\nu} = \frac{1}{2} \left( \frac{\partial A_{\mu}}{\partial x^{\nu}} - \frac{\partial A_{\nu}}{\partial x^{\mu}} \right)$ , which is also referred to as the field strength in physics literature.

Def. The electric field  $\vec{E} = (E_1, E_2, E_3)$  is defined by  $E_j = F_{j0}$ .

The magnetic field  $\vec{B} = (B_1, B_2, B_3)$  is defined by  $B_x = F_{yz}, B_y = F_{zx}, B_z = F_{xy}$ .

We also write  $F_{\mu\nu}$  in a matrix form:

$$F_{\mu\nu} = \begin{pmatrix} 0 & -E_x & -E_y & -E_z \\ E_x & 0 & -F_{yx} & -F_{zx} \\ E_y & F_{yx} & 0 & -F_{zy} \\ E_z & F_{zx} & F_{zy} & 0 \end{pmatrix}$$

Recall that  $F$  satisfies Bianchi's identity:  $d_A F = 0$

$$\Rightarrow 0 = d_A F = dF + A \wedge F - F \wedge A = dF$$

$$\Rightarrow 0 = \frac{1}{2} \sum dF_{\mu\nu} dx^{\nu} \wedge dx^{\mu} = \frac{1}{2} \frac{\partial F_{\mu\nu}}{\partial x^{\rho}} dx^{\rho} \wedge dx^{\nu} \wedge dx^{\mu}.$$

$$\Rightarrow \partial_{\rho} F_{\mu\nu} + \partial_{\mu} F_{\nu\rho} + \partial_{\nu} F_{\rho\mu} = 0, \quad \forall \rho, \mu, \nu.$$

In terms of the electric and magnetic fields:

(1). All  $\rho, \mu, \nu$  are space indices: ( $\rho = x, \mu = y, \nu = z$ )

$$\Rightarrow \partial_x F_{yz} + \partial_y F_{zx} + \partial_z F_{xy} = 0$$

$$\text{i.e. } \partial_x B_x + \partial_y B_y + \partial_z B_z = 0$$

$$\text{or } \nabla \cdot \vec{B} = 0$$



(2). One index is the time index

$$\begin{cases} \partial_t F_{xy} + \partial_x F_{y0} + \partial_y F_{0x} = 0 & \text{i.e. } \partial_t B_z + \partial_x E_y - \partial_y E_x = 0 \\ \partial_t F_{yz} + \partial_y F_{z0} + \partial_z F_{0y} = 0 & \text{i.e. } \partial_t B_x + \partial_y E_z - \partial_z E_y = 0 \\ \partial_t F_{zx} + \partial_z F_{x0} + \partial_x F_{0z} = 0 & \text{i.e. } \partial_t B_y + \partial_z E_x - \partial_x E_z = 0 \end{cases}$$

or  $\partial_t \vec{B} + \vec{\nabla} \times \vec{E} = 0.$

Thus the Bianchi's identity accounts for 2 of Maxwell's equations. The other 2 equations arise from a variational principle, i.e. given a connection  $A$ , we can associate with it its action  $I(A) = \int_{\mathbb{R}^4} |F|^2$ . The electric magnetic fields are the ones that minimize this action. Namely:

$$\delta I(A) = \frac{\delta I}{\delta A} \cdot \delta A$$

and the critical points are the ones satisfying  $\delta I / \delta A = 0$ .

In the case of connections, the critical point equation is:

$$\nabla^\mu F_{\mu\nu} = 0 \quad (\text{Yang-Mills Equation})$$

This will be shown later. Explicitly in the present case,  $\nabla^\mu F_{\mu\nu} = \partial_\rho (g^{\rho\mu} F_{\mu\nu})$ , where  $g^{\mu\nu} = \text{diag}(-1, 1, 1, 1)$ , and thus:

$$g^{\rho\mu} F_{\mu\nu} = \begin{cases} -F_{0\nu} & \rho=0 \\ F_{\rho\nu} & \rho=x, y, z. \end{cases}$$

(3). Take  $\nu=0$ , then:

$$0 = \partial_\rho (g^{\rho\mu} F_{\mu 0}) = \partial_0 (-F_{00}) + \partial_x F_{x0} + \partial_y F_{y0} + \partial_z F_{z0}.$$

$$\text{i.e. } \partial_x E_x + \partial_y E_y + \partial_z E_z = 0$$

$$\text{or } \nabla \cdot \vec{E} = 0$$

(4). Take  $\nu = x$  (or  $y, z$ ), then

$$\begin{cases} 0 = \partial_\rho (g^{\rho\mu} F_{\mu x}) = \partial_0 (-F_{0x}) + \partial_x F_{xx} + \partial_y F_{yx} + \partial_z F_{zx} \\ 0 = \partial_\rho (g^{\rho\mu} F_{\mu y}) = \partial_0 (-F_{0y}) + \partial_x F_{xy} + \partial_y F_{yy} + \partial_z F_{zy} \\ 0 = \partial_\rho (g^{\rho\mu} F_{\mu z}) = \partial_0 (-F_{0z}) + \partial_x F_{xz} + \partial_y F_{yz} + \partial_z F_{zz} \end{cases}$$

$$\text{i.e. } \begin{cases} \partial_t E_x - (\partial_y B_z - \partial_z B_y) = 0 \\ \partial_t E_y - (\partial_z B_x - \partial_x B_z) = 0 \\ \partial_t E_z - (\partial_x B_y - \partial_y B_x) = 0 \end{cases}$$

$$\text{or } \partial_t \vec{E} - \nabla \times \vec{B} = 0$$

Variational formula.

Given a variation  $A \rightsquigarrow A + \delta A$  (note that  $\delta A$  is a 1-form). Recall that

$$F_{jk} = -(\partial_j A_k - \partial_k A_j + A_j A_k - A_k A_j)$$

$$\begin{aligned} \Rightarrow \delta F_{jk} &= -(\partial_j \delta A_k - \partial_k \delta A_j + \delta A_j \cdot A_k + A_j \delta A_k - \delta A_k \cdot A_j - A_k \delta A_j) \\ &= -(\partial_j \delta A_k + A_j \delta A_k - \delta A_k \cdot A_j - (\partial_k \delta A_j + A_k \delta A_j - \delta A_j \cdot A_k)) \\ &= -(\nabla_j \delta A_k - \nabla_k \delta A_j) \end{aligned}$$

$$\text{Thus, } \delta I = \delta \int_X \langle F, F \rangle$$

$$= 2 \int_X \langle \delta F, F \rangle$$

$$= 2 \int_X (\nabla_j \delta A_k - \nabla_k \delta A_j) g^{j\ell} g^{km} F_{m\ell}$$

$$= 2 \int_X -\delta A_k \cdot g^{km} \nabla^\ell F_{m\ell} + \delta A_j g^{j\ell} \nabla^m F_{m\ell} \quad (\text{Integration by parts})$$

$$= 4 \int_X \delta A_k \cdot g^{km} \nabla^\ell F_{m\ell}$$

The right hand side is linear in  $\delta A_k$  now, thus

$$\delta I / \delta A = 0 \Leftrightarrow \nabla^\ell F_{m\ell} = 0.$$

A basic example of complex manifolds:  $\mathbb{C}P^n$ .

Def.  $\mathbb{C}P^n = \mathbb{C}^{n+1} \setminus \{0\} / \sim$ , where  $(\zeta^0, \dots, \zeta^n) \sim \lambda(\zeta^0, \dots, \zeta^n)$ ,  $\forall \lambda \in \mathbb{C}^*$ .

We use the following coordinate system,  $\forall j=0, \dots, n$ .

$$X_j \triangleq \{[\zeta^0 : \dots : \zeta^n] \mid \zeta_j \neq 0\}$$

and we identify  $X^j$  with  $\mathbb{C}^n$  via:

$$X_j \ni (\zeta^0, \dots, \zeta^n) \mapsto (\frac{\zeta^0}{\zeta_j}, \dots, \frac{\zeta^j}{\zeta_j}, \dots, \frac{\zeta^n}{\zeta_j}) \triangleq z \in \mathbb{C}^n$$

The transition functions, for instance, on  $X_0 \cap X_n$ , is given by:

$$\begin{array}{c} [\zeta^0 : \dots : \zeta^n] \\ \swarrow \quad \searrow \\ z \triangleq (\frac{\zeta^1}{\zeta^0}, \dots, \frac{\zeta^n}{\zeta^0}) \quad (\frac{\zeta^0}{\zeta^n}, \dots, \frac{\zeta^{n-1}}{\zeta^n}) \triangleq w \end{array}$$

$$\text{then } w^0 = \frac{1}{z^n}, w^1 = \frac{z^1}{z^n}, w^2 = \frac{z^2}{z^n}, \dots$$

Def. The universal line bundle  $L^{-1}$  assigns each point of  $\mathbb{C}P^n$  the line it represents.

Note that the total space of  $L^{-1} \setminus \{0\}$ -sections  $\cong \mathbb{C}^{n+1}$ . We trivialize  $L^{-1}$  in the following way:

$$\text{On } X_0, (\zeta^0, \dots, \zeta^n) \mapsto ([\zeta^0 : \dots : \zeta^n], \zeta^0) \in \mathbb{C}P^n \times \mathbb{C}$$

$$\text{On } X_n, (\zeta^0, \dots, \zeta^n) \mapsto ([\zeta^0 : \dots : \zeta^n], \zeta^n) \in \mathbb{C}P^n \times \mathbb{C}$$

Since  $\zeta^0 = (\zeta^j / \zeta^n) \cdot \zeta^n$ , we define  $L^{-1}$  as the holomorphic line bundle with transition functions  $t_{jk} = \zeta^j / \zeta^k$  on  $X_j \cap X_k$ , i.e.

$$\Gamma(X, L^{-1}) \ni \varphi \iff \{ \varphi_j = \zeta^j / \zeta^k \cdot \varphi_k \text{ on } X_j \cap X_k \}.$$

An example of a meromorphic section is given as follows:

$$\text{On } X_0, \text{ set } \varphi_0 = 1. \text{ This determines } \varphi_j = \frac{\zeta^j}{\zeta^0} \cdot 1 = \frac{\zeta^j}{\zeta^0} \text{ on } X_j \cap X_0.$$

This is meromorphic section of  $L^{-1}$  with poles along the codimension 1 subvariety  $\{\zeta_0 = 0\}$ .

The universal bundle  $L^{-1}$  admits a natural metric:

The invariant way of defining it is if  $\varphi = (\zeta^0, \dots, \zeta^n)$ , set

$$\|\varphi\|^2 = \sum_{i=0}^n |\zeta^i|^2$$

Locally, say, on  $X_0$ :

$$\begin{aligned} \|\varphi\|^2 &= |\zeta^0|^2 \left( 1 + \left| \frac{\zeta^1}{\zeta^0} \right|^2 + \dots + \left| \frac{\zeta^n}{\zeta^0} \right|^2 \right) \\ &= |\zeta^0|^2 h(z). \end{aligned}$$

Thus  $h(z) = (1 + \|z\|^2)$ , where  $\|z\|^2 = |z_1|^2 + \dots + |z_n|^2$  on each coordinate chart.

Curvature of  $L^{-1}$ .

By our previous formula:

$$\begin{aligned} F_{\bar{k}j} &= -\partial_{\bar{k}} \partial_j \log h \\ &= -\partial_{\bar{k}} \partial_j (\log(1 + \|z\|^2)) \\ &= -\partial_{\bar{k}} \left( \frac{\bar{z}^j}{1 + \|z\|^2} \right) \\ &= - \left\{ \frac{\delta_{kj}}{1 + \|z\|^2} - \frac{\bar{z}^j z^k}{(1 + \|z\|^2)^2} \right\} \end{aligned}$$

Observe that if  $v = (v^1, \dots, v^n)$  is a vector

$$F_{\bar{k}j} v^j \bar{v}^k = - \left\{ \frac{\|v\|^2}{1 + \|z\|^2} - \frac{\langle v, z \rangle \langle z, v \rangle}{(1 + \|z\|^2)^2} \right\}$$

$$= - \left\{ \frac{\|v\|^2 + \|v\|^2 \|z\|^2 - |\langle v, z \rangle|^2}{(1 + \|z\|^2)^2} \right\}$$

$$< - \frac{\|v\|^2}{(1 + \|z\|^2)^2}$$

Thus the curvature is a negative definite (1,1) form.

Def: Let the hyperplane bundle  $L$  be the inverse of  $L^{-1}$ , i.e.  $L$  is defined by the transition functions  $\{\frac{\zeta^k}{\zeta^j}$  on  $X_j \cap X_k\}$ . i.e.

$$\Gamma(X, L) \ni \varphi \longleftrightarrow \{\varphi_j = \zeta^k / \zeta^j \varphi_k \text{ on } X_j \cap X_k\}$$

Similarly,  $L^m = \{(\zeta^k / \zeta^j)^m \text{ on } X_j \cap X_k\}$ .

$L$  admits a natural metric  $h_L(z) \triangleq \frac{1}{1 + \|z\|^2}$ , the Fubini-Study metric. The curvature of this metric is:

$$(F_L)_{\bar{k}j} = \left\{ \frac{\delta_{jk}}{1 + \|z\|^2} - \frac{z_k \bar{z}_j}{(1 + \|z\|^2)^2} \right\}$$

which is strictly positive:

$$(F_L)_{\bar{k}j} \cup^j \bar{v}^k = \left\{ \frac{\|v\|^2}{1 + \|z\|^2} - \frac{\langle v, z \rangle \langle z, v \rangle}{(1 + \|z\|^2)^2} \right\}$$

$$= \left\{ \frac{\|v\|^2 + \|v\|^2 \|z\|^2 - |\langle v, z \rangle|^2}{(1 + \|z\|^2)^2} \right\}$$

$$> \frac{\|v\|^2}{(1 + \|z\|^2)^2}$$

The holomorphic sections of  $L^m$  can be identified with homogeneous polynomials  $p(\zeta^0, \zeta^1, \dots, \zeta^n)$  of order  $m$ . (Homogeneous means  $p(\lambda \zeta^0, \lambda \zeta^1, \dots, \lambda \zeta^n) = \lambda^m p(\zeta^0, \zeta^1, \dots, \zeta^n)$ .)

Given  $p(\zeta)$ , we may define, for instance on  $X_0$ ,  $p_0(z) = p(1, z^1, \dots, z^n)$ . Then, on  $X_0 \cap X_n$ :

$$p_0(1, z_1, \dots, z_n) = p\left(1, \frac{\zeta^1}{\zeta^0}, \dots, \frac{\zeta^n}{\zeta^0}\right)$$

$$= \left(\frac{1}{\zeta^0}\right)^m p(\zeta^0, \zeta^1, \dots, \zeta^n)$$

$$= \left(\frac{\zeta^n}{\zeta^0}\right)^m p\left(\frac{\zeta^0}{\zeta^n}, \frac{\zeta^1}{\zeta^n}, \dots, 1\right)$$

$$= \left(\frac{\zeta^n}{\zeta^0}\right)^m p_n(\omega^0, \omega^1, \dots, 1)$$

In summary, we have the following set-up:  $L \rightarrow \mathbb{C}P^n$  admits a metric  $h_L$  with strictly positive curvature  $F_L$ . Then we may pick  $F_L$  to be a metric for  $\mathbb{C}P^n$  since after all it's just a (1,1)-form. Thus in the future, the Fubini-

Study metric will mean 2 things:

- 1). The metric  $h_L$  on  $L$
- 2). Curvature  $F_L$  of  $(L, h_L)$ , regarded as a metric  $g_{\bar{k}j} = (F_L)_{\bar{k}j}$  on  $\mathbb{C}P^n$ .

Exercise: Compute  $g^{\bar{k}l}$ ,  $\det g_{\bar{k}j}$  and  $R_{\bar{k}j}$  ( $= (n+1)g_{\bar{k}j}$ ).

Lemma. (Linear algebra). If  $u \in \mathbb{C}^n$  is a unit vector,  $\lambda \in \mathbb{C} \setminus \{0, 1\}$ , then

$$(\lambda \text{Id} - u \bar{u}^t)^{-1} = \lambda^{-1} \text{Id} + ((\lambda - 1)^{-1} - \lambda^{-1}) u \bar{u}^t.$$

Pf:  $(\lambda \text{Id} - u \bar{u}^t)(\lambda^{-1} \text{Id} + ((\lambda - 1)^{-1} - \lambda^{-1}) u \bar{u}^t)$   
 $= \text{Id} + \lambda((\lambda - 1)^{-1} - \lambda^{-1}) u \bar{u}^t - \lambda^{-1} u \bar{u}^t - ((\lambda - 1)^{-1} - \lambda^{-1}) u \bar{u}^t$   
 $= \text{Id} + \frac{1}{\lambda - 1} u \bar{u}^t - \lambda^{-1} u \bar{u}^t - \frac{1}{\lambda - 1} u \bar{u}^t + \lambda^{-1} u \bar{u}^t$   
 $= \text{Id}$

□

Using this lemma, we can calculate  $(g_{\bar{k}j})^{-1}$ :

$$(g_{\bar{k}j}) = \left( \frac{\delta_{jk}}{1 + \|z\|^2} - \frac{z_k \bar{z}_j}{(1 + \|z\|^2)^2} \right) = \frac{\|z\|^2}{(1 + \|z\|^2)^2} \left( \frac{1 + \|z\|^2}{\|z\|^2} \text{Id} - \frac{z \bar{z}^t}{\|z\|^2} \right)$$

$$\Rightarrow (g_{\bar{k}j})^{-1} = \frac{(1 + \|z\|^2)^2}{\|z\|^2} \left( \frac{\|z\|^2}{1 + \|z\|^2} \text{Id} + \left( \|z\|^2 - \frac{\|z\|^2}{1 + \|z\|^2} \right) \frac{z \bar{z}^t}{\|z\|^2} \right)$$

$$= (1 + \|z\|^2) \text{Id} + (1 + \|z\|^2) \|z\|^2 \frac{z \bar{z}^t}{\|z\|^2}$$

$$\Rightarrow g^{\bar{k}l} = (1 + \|z\|^2) \delta_{\bar{k}l} + (1 + \|z\|^2) \|z\|^2 z^{\bar{l}} \bar{z}^{\bar{k}}$$

To calculate the determinant, we use:

Lemma. (Linear algebra). If  $u \in \mathbb{C}^n$  is a unit vector,  $\lambda \in \mathbb{C} \setminus \{0, 1\}$ , then

$$\det(\lambda \text{Id} - u \bar{u}^t) = \lambda^{n-1} (\lambda - 1)$$

Pf: Complete  $u$  to an orthonormal basis  $\{u, u_1, \dots, u_{n-1}\}$ , then

$$\begin{cases} (\lambda \text{Id} - u \bar{u}^t) u = (\lambda - 1) u, \\ (\lambda \text{Id} - u \bar{u}^t) u_i = \lambda u_i \end{cases}$$

$\Rightarrow \{u, u_1, \dots, u_{n-1}\}$  diagonalizes  $\lambda \text{Id} - u \bar{u}^t \Rightarrow \det(\lambda \text{Id} - u \bar{u}^t) = \lambda^{n-1} (\lambda - 1)$ .

□

It follows  $\det(g_{\bar{k}j}) = \frac{\|z\|^{2n}}{(1 + \|z\|^2)^{2n}} \cdot \frac{(1 + \|z\|^2)^{n-1}}{\|z\|^{2(n-1)}} \left( \frac{1 + \|z\|^2}{\|z\|^2} - 1 \right) = \frac{1}{(1 + \|z\|^2)^{n+1}}$ .

Finally, the Ricci curvature:

Lemma. If  $A(t)$  is an  $n \times n$  matrix valued function,  $A(0)$  invertible, then:

$$(\log \det A(t))'(0) = \operatorname{tr} A^{-1}(0) A'(0).$$

Pf: Over  $\mathbb{C}$ , let  $J = B^{-1} A B$  be the Jordan canonical form,  $\lambda_i$  the eigen-values.

Then  $\log \det A(t) = \log \det J = \sum \log \lambda_i$

$$\Rightarrow (\log \det A(t))'(0) = \sum \lambda_i^{-1} \lambda_i'(0)$$

$$= \operatorname{tr} J^{-1} J'(0)$$

$$= \operatorname{tr} (B^{-1} A B)^{-1} (-B^{-1} B' B^{-1} A B + B^{-1} A B + B^{-1} A B')$$

$$= \operatorname{tr} (-B^{-1} A^{-1} B' B^{-1} A B + B^{-1} A^{-1} B B^{-1} A' B + B^{-1} A^{-1} B B^{-1} A B')$$

$$= \operatorname{tr} (-B^{-1} B^{-1} + A^{-1} A' + B^{-1} B')$$

$$= \operatorname{tr} (A^{-1} A')$$

□

$$\text{Now: } R_{\bar{k}j} = R_{\bar{k}j}^{\ell\ell} = -\partial_{\bar{k}} (g^{\ell\bar{q}} \partial_j g_{\bar{q}\ell})$$

$$= -\partial_{\bar{k}} \partial_j (\log \det g_{\bar{q}\ell})$$

$$= -\partial_{\bar{k}} \partial_j (-(n+1) \log(1 + \|z\|^2))$$

$$= (n+1) \left( \frac{\delta_{jk}}{1 + \|z\|^2} - \frac{\bar{z}^j z^k}{(1 + \|z\|^2)^2} \right)$$

$$= (n+1) g_{\bar{k}j}$$

**Exercise:** Our metric  $g_{\bar{k}j} = \partial_{\bar{k}} \partial_j \log h$  is always Kähler since  $\partial_{\bar{q}} g_{\bar{k}j} = \partial_j g_{\bar{q}k}$ . Now show that if  $X$  is Kähler,  $Y$  is a complex submanifold, then  $Y$  is Kähler. In particular any complex submanifold of  $\mathbb{C}P^n$  is Kähler.

Pf: The question is local, thus we may assume that  $Y$  is locally the zero set of some holomorphic functions  $f_{r+1}, \dots, f_n$ , with linearly independent differentials.

Complete  $f_{r+1}, \dots, f_n$  to a local coordinate chart  $\{z_1, \dots, z_n\}$ , say,  $z_{r+1} = f_{r+1}, \dots, z_n = f_n$ . Then at  $p \in Y$ ,  $T_p^0 M = \mathbb{C} \langle \partial_1, \dots, \partial_r \rangle$ , and  $\{g_{\bar{k}j}\}|_Y$  satisfies  $\partial_{\bar{q}} g_{\bar{k}j}|_Y = \partial_j g_{\bar{q}k}|_Y$ . The result follows.

## §6. Kodaira Vanishing Theorem

### Road Map:

Our first goal is the Kodaira Embedding Thm:

(Kodaira Embedding Thm) Let  $L \rightarrow X$  be a positive line bundle over a compact complex manifold. Then for  $m$  large enough, the map

$$\begin{aligned} X &\longrightarrow \mathbb{C}P^{Nm} \\ z &\longmapsto [S_0(z), \dots, S_{Nm}(z)] \end{aligned}$$

is an embedding. Here  $\{S_\alpha(z), \alpha=0, \dots, Nm\}$  is a basis of the space  $H^0(X, L^m)$  and  $\dim H^0(X, L^m) = Nm+1$

Main Ingredients of the proof.

Need: many holomorphic sections of  $L^m$

- Vanishing thms: If  $E$  is a holomorphic bundle, when is  $\ker \square|_{E \otimes \Lambda^{p,q}} = ?$

$$\dots \xrightleftharpoons[\bar{\partial}^+]{\bar{\partial}} E \otimes \Lambda^{p,q-1} \xrightleftharpoons[\bar{\partial}^+]{\bar{\partial}} E \otimes \Lambda^{p,q} \xrightleftharpoons[\bar{\partial}^+]{\bar{\partial}} E \otimes \Lambda^{p,q+1} \xrightleftharpoons[\bar{\partial}^+]{\bar{\partial}} \dots$$

$$\square = \bar{\partial} \bar{\partial}^+ + \bar{\partial}^+ \bar{\partial}$$

Observation: Naively,  $\ker \square|_{E \otimes \Lambda^{p,q}}$  depends on metrics because  $\bar{\partial}^+$  does. It's however only dependent on the complex structure:

- Hodge decomposition thm.

$$\ker \square|_{E \otimes \Lambda^{p,q}} \equiv H_{\bar{\partial}}^{p,q}(X, E) \quad : \text{ Dolbeault cohomology.}$$

which depends only on the complex structure.

- Sheaf cohomology:  $H_{\bar{\partial}}^{p,q}(X, E) = H^q(X, \Omega^p(E))$ , where  $\Omega^p(E)$  is the sheaf of  $E$ -valued  $(p,0)$  forms.

Advantage of sheaf cohomology:

$$\begin{aligned} 0 &\longrightarrow \mathcal{F} \longrightarrow \mathcal{G} \longrightarrow \mathcal{H} \longrightarrow 0 \\ \Rightarrow 0 &\longrightarrow H^0(X, \mathcal{F}) \longrightarrow H^0(X, \mathcal{G}) \longrightarrow H^0(X, \mathcal{H}) \\ &\longrightarrow H^1(X, \mathcal{F}) \longrightarrow \dots \end{aligned}$$

Then once we know  $H^1(X, \mathcal{F})$  vanishes and  $H^0(X, \mathcal{H})$  is big enough, it would imply  $H^0(X, \mathcal{G})$  is big enough.

Bochner - Kodaira Formulas.

• Characteristic feature: In geometry, there are many "Laplacians". A Bochner-Kodaira formula is a formula of the type:

$$\Delta = \tilde{\Delta} + \text{Curvature terms.}$$

where  $\Delta, \tilde{\Delta}$  are different Laplacians.

Base case:  $E \rightarrow X$ : holomorphic line bundle.  $H_{\bar{\alpha}\beta}$ : metric on  $E$ .  $g_{\bar{k}j}$ : metric on  $X$ . (Hermitian metric  $\Rightarrow \overline{H_{\bar{\alpha}\beta}} = H_{\beta\alpha}$ ). We have:

$$\dots \xrightarrow{\bar{\partial}} E \otimes \Lambda^{p,q-1} \xrightarrow{\bar{\partial}} E \otimes \Lambda^{p,q} \xrightarrow{\bar{\partial}} E \otimes \Lambda^{p,q+1} \xrightarrow{\bar{\partial}} \dots$$

In presence of metrics  $H_{\bar{\alpha}\beta}, g_{\bar{k}j}$ , there is a norm on  $\Gamma(X, E \otimes \Lambda^{p,q})$ , given by:

$\forall \varphi \in \Gamma(X, E \otimes \Lambda^{p,q}), \varphi = \frac{1}{p!q!} \sum \varphi_{\bar{J}I}^\alpha dz^I d\bar{z}^{\bar{J}}$ , where

$$\begin{cases} dz^I = dz^{i_1} \wedge \dots \wedge dz^{i_p} \\ d\bar{z}^{\bar{J}} = d\bar{z}^{\bar{j}_1} \wedge \dots \wedge d\bar{z}^{\bar{j}_q} \\ \varphi_{\bar{J}I}^\alpha = \varphi_{\bar{j}_1 \dots \bar{j}_q, i_1 \dots i_p}^\alpha \end{cases}$$

$$\|\varphi\|^2 \triangleq \frac{1}{p!q!} \int_X \varphi_{\bar{J}I}^\alpha \overline{\varphi_{\bar{K}L}^\beta} H_{\bar{\beta}\alpha} g^{k\bar{j}} g^{i\bar{l}} \cdot \frac{\omega^n}{n!}$$

$$= \frac{1}{p!q!} \int_X \varphi_{\bar{j}_1 \dots \bar{j}_q, i_1 \dots i_p}^\alpha \overline{\varphi_{\bar{k}_1 \dots \bar{k}_q, l_1 \dots l_p}^\beta} H_{\bar{\beta}\alpha} g^{k_1 \bar{j}_1} \dots g^{k_q \bar{j}_q} g^{i_1 \bar{l}_1} \dots g^{i_p \bar{l}_p} \frac{\omega^n}{n!}$$

Here  $\omega$  denotes the symplectic form  $\omega = \frac{i}{2} g_{\bar{k}j} dz^j \wedge d\bar{z}^{\bar{k}}$ , and  $\frac{\omega^n}{n!}$  is the volume form on  $X$ :  $\frac{\omega^n}{n!} = \det(g_{\bar{k}j}) \prod_{j=1}^n (dz^j \wedge d\bar{z}^{\bar{j}})$

Note that we can also define the inner product similarly.

Now, we define the formal adjoint  $\bar{\partial}^\dagger$  by:

$$\langle \bar{\partial}\varphi, \psi \rangle = \langle \varphi, \bar{\partial}^\dagger\psi \rangle$$

for any  $\varphi \in \Gamma(X, E \otimes \Lambda^{p,q}), \psi \in \Gamma(X, E \otimes \Lambda^{p,q+1})$ , with compact support. Then, there is a natural Laplacian  $\square = \bar{\partial}\bar{\partial}^\dagger + \bar{\partial}^\dagger\bar{\partial}$  on  $\Gamma(X, E \otimes \Lambda^{p,q})$ . However, there is another natural Laplacian (metric):

$$\Gamma(X, E \otimes \Lambda^{p,q}) \ni \varphi \mapsto -g^{j\bar{k}} \nabla_{\bar{j}} \nabla_{\bar{k}} \varphi_{\bar{J}K}^\alpha \in \Gamma(X, E \otimes \Lambda^{p,q}).$$

Question: Compare " $\square$ " and " $-g^{j\bar{k}} \nabla_{\bar{j}} \nabla_{\bar{k}}$ ".



A simple example:  $\square$  on  $\Gamma(\mathbb{C}^n, \Lambda^{0,1})$ , with flat metric  $g_{\bar{k}j} = \delta_{kj}$ . ( $E$  trivial)

We compute  $\bar{\partial}$  on  $\Lambda^{0,0}$  and  $\Lambda^{0,1}$ . Take  $f \in \Lambda^{0,0}$ ,  $\varphi \in \Lambda^{0,1}$ .

$$\bar{\partial}f = \partial_{\bar{j}} \int d\bar{z}^j$$

$$\bar{\partial}\varphi = \bar{\partial}(\varphi_{\bar{j}} d\bar{z}^j) = \bar{\partial}\varphi_{\bar{j}} d\bar{z}^j = \partial_{\bar{k}}\varphi_{\bar{j}} d\bar{z}^k \wedge d\bar{z}^j = \frac{1}{2}(\partial_{\bar{k}}\varphi_{\bar{j}} - \partial_{\bar{j}}\varphi_{\bar{k}}) d\bar{z}^k \wedge d\bar{z}^j$$

i.e.  $(\bar{\partial}\varphi)_{\bar{j}\bar{k}} = \partial_{\bar{k}}\varphi_{\bar{j}} - \partial_{\bar{j}}\varphi_{\bar{k}}$  in components.

Next we compute  $\bar{\partial}^+$  on  $\Lambda^{0,2}$  and  $\Lambda^{0,1}$ , by the defining equation:

$$\langle \bar{\partial}\varphi, \psi \rangle = \langle \varphi, \bar{\partial}^+\psi \rangle, \quad \forall \varphi \in \Gamma_c(\mathbb{C}^n, \Lambda^{0,1}), \quad \psi \in \Gamma(\mathbb{C}^n, \Lambda^{0,2})$$

$$\text{l.h.s.} = \frac{1}{2} \int_X (\bar{\partial}\varphi)_{\bar{j}\bar{k}} \overline{\psi_{\bar{l}\bar{m}}} g^{\bar{l}j} g^{m\bar{k}} \cdot \text{vol}$$

$$= \int_X (\partial_{\bar{k}}\varphi_{\bar{j}}) \overline{\psi_{\bar{l}\bar{m}}} g^{\bar{l}j} g^{m\bar{k}} \cdot \text{vol} \quad (\text{note that } \partial_{\bar{k}}\varphi_{\bar{j}} - \partial_{\bar{j}}\varphi_{\bar{k}} \text{ anti-sym in } \bar{k}, \bar{j} \text{ but}$$

$$= - \int_X \varphi_{\bar{j}} \overline{\nabla_{\bar{k}}\psi_{\bar{l}\bar{m}}} g^{k\bar{m}} \cdot \text{vol} \quad \overline{\psi_{\bar{l}\bar{m}}} g^{\bar{l}j} g^{m\bar{k}} \text{ is also anti-sym in } \bar{k}, \bar{j}, \text{ giving } 2)$$

The last step using integration by parts and the reason we use  $\nabla_{\bar{k}}$  will be clarified in detail later.

Thus, on  $\Lambda^{0,2}$ ,  $(\bar{\partial}^+\psi)_{\bar{l}} = -\nabla_{\bar{k}}\psi_{\bar{l}\bar{m}} g^{k\bar{m}} = -\nabla_{\bar{k}}\psi_{\bar{l}\bar{k}}$  (flat metric).

On  $\Lambda^{0,1}$ , take  $\varphi \in \Gamma_c(\mathbb{C}^n, \Lambda^{0,0})$ ,  $\psi \in \Gamma(\mathbb{C}^n, \Lambda^{0,1})$ :

$$\text{l.h.s.} = \int_X (\partial_{\bar{j}}\varphi) \overline{\psi_{\bar{l}}} g^{\bar{l}j} \cdot \text{vol}$$

$$= \int_X \varphi (-\nabla_{\bar{j}}\overline{\psi_{\bar{l}}}) g^{j\bar{l}} \cdot \text{vol}$$

Thus on  $\Lambda^{0,1}$ ,  $\bar{\partial}^+\psi = -\nabla_{\bar{j}}\psi_{\bar{l}} g^{j\bar{l}} = -\nabla_{\bar{l}}\psi_{\bar{l}} = -\partial_{\bar{l}}\psi_{\bar{l}}$

Hence on  $\Lambda^{0,1}$ , the Laplacian  $\square$  is given by ( $\varphi = \varphi_{\bar{j}} d\bar{z}^j$ )

$$(\square\varphi)_{\bar{j}} = \{ \bar{\partial}\bar{\partial}^+\varphi \}_{\bar{j}} + \{ \bar{\partial}^+\bar{\partial}\varphi \}_{\bar{j}}$$

$$= \partial_{\bar{j}}(-\partial_{\bar{l}}\varphi_{\bar{l}}) + (-\partial_{\bar{k}}(\bar{\partial}\varphi)_{\bar{j}\bar{k}})$$

$$= -\partial_{\bar{j}}\partial_{\bar{l}}\varphi_{\bar{l}} - (\partial_{\bar{k}}(\partial_{\bar{k}}\varphi_{\bar{j}} - \partial_{\bar{j}}\varphi_{\bar{k}}))$$

$$= -\partial_{\bar{j}}\partial_{\bar{l}}\varphi_{\bar{l}} + \partial_{\bar{k}}\partial_{\bar{j}}\varphi_{\bar{k}} - \partial_{\bar{k}}\partial_{\bar{k}}\varphi_{\bar{j}}$$

$$= -\partial_{\bar{k}}\partial_{\bar{k}}\varphi_{\bar{j}}$$

Note that in the last step, we used the fact that on flat spaces,  $\partial_{\bar{j}}\partial_{\bar{k}} = \partial_{\bar{k}}\partial_{\bar{j}}$  (or more precisely,  $\nabla_{\bar{j}}\nabla_{\bar{k}} - \nabla_{\bar{k}}\nabla_{\bar{j}} = 0$ ). In general, this results in curvature terms. In this case, two Laplacians agree.

General case:  $\square = \bar{\partial}\bar{\partial}^+ + \bar{\partial}^+\bar{\partial}$  on  $E \otimes \Lambda^{p,q}$ .

(1). Computation of  $\bar{\partial}$  on  $E \otimes \Lambda^{p,q}$ .

$$\bar{\partial} \left( \frac{1}{p!q!} \sum \varphi_{\bar{j}_1}^\alpha dz^1 \wedge \dots \wedge dz^p \wedge d\bar{z}^1 \wedge \dots \wedge d\bar{z}^q \right) = \frac{1}{p!q!} \sum \partial_{\bar{k}} \varphi_{\bar{j}_1}^\alpha dz^1 \wedge \dots \wedge dz^p \wedge d\bar{z}^1 \wedge \dots \wedge d\bar{z}^q$$

Here we need some explanation: this means without anti-symmetrization,

$$(\bar{\partial} \varphi)_{\bar{j}_1 \bar{k}}^\alpha = (q+1) \partial_{\bar{k}} \varphi_{\bar{j}_1}^\alpha.$$

However, with antisymmetrization, we can also write:

$$\begin{aligned} (\bar{\partial} \varphi)_{\bar{j}_1 \bar{k}}^\alpha &= (\bar{\partial} \varphi)_{\bar{j}_2 \dots \bar{j}_1 \bar{k}} \\ &= \partial_{\bar{k}} \varphi_{\bar{j}_2 \dots \bar{j}_1} - \partial_{\bar{j}_1} \varphi_{\bar{j}_2 \dots \bar{j}_2 \bar{k}} - \partial_{\bar{j}_2} \varphi_{\bar{j}_2 \dots \bar{k} \bar{j}_1} - \dots - \partial_{\bar{j}_q} \varphi_{\bar{k} \bar{j}_2 \dots \bar{j}_1}. \end{aligned}$$

Note also that, even  $E \otimes \Lambda^{p,q}$  is no longer a holomorphic bundle.  $\bar{\partial}$  makes sense since the process of antisymmetrizing kills higher order differentiations, just as in de Rham case.

(2). Computation of  $\bar{\partial}^\dagger$  on  $E \otimes \Lambda^{p,q}$

Again, take  $\varphi \in \Gamma(X, E \otimes \Lambda^{p,q})$   $\psi \in \Gamma(X, E \otimes \Lambda^{p,q+1})$ . By definition, we have:

$$\langle \bar{\partial} \varphi, \psi \rangle = \langle \varphi, \bar{\partial}^\dagger \psi \rangle.$$

$$\text{l.h.s.} = \frac{1}{p!(q+1)!} \int_X (\bar{\partial} \varphi)_{\bar{j}_1 \bar{k}}^\alpha \overline{\psi_{\bar{k} \bar{m} \bar{\ell}}^\beta} H_{\bar{\beta} \alpha} g^{k\bar{j}} g^{i\bar{m}} g^{\ell \bar{k}} \frac{\omega^n}{n!}$$

Observation:

(a). We can write  $(\bar{\partial} \varphi)_{\bar{j}_1 \bar{k}}^\alpha$  in terms of genuine covariant derivative.

$$\begin{aligned} \text{e.g. } \varphi \in \Gamma(X, E \otimes \Lambda^{p,1}), (\bar{\partial} \varphi)_{\bar{j}_1 \bar{k}}^\alpha &= \partial_{\bar{k}} \varphi_{\bar{j}_1}^\alpha - \partial_{\bar{j}_1} \varphi_{\bar{k}}^\alpha \\ &= \nabla_{\bar{k}} \varphi_{\bar{j}_1}^\alpha - \Gamma_{\bar{k} \bar{j}}^{\bar{m}} \varphi_{\bar{m} 1}^\alpha - (\nabla_{\bar{j}} \varphi_{\bar{k} 1}^\alpha - \Gamma_{\bar{j} \bar{k}}^{\bar{m}} \varphi_{\bar{m} 1}^\alpha) \end{aligned}$$

(Here recall that on a (1,0)-form  $\nabla_{\bar{k}} \varphi_{\bar{j}} = \partial_{\bar{k}} \varphi_{\bar{j}} - \Gamma_{\bar{k} \bar{j}}^{\bar{m}} \varphi_{\bar{m}}$   $\Rightarrow \nabla_{\bar{k}} \varphi_{\bar{j}} = \partial_{\bar{k}} \varphi_{\bar{j}} - \Gamma_{\bar{k} \bar{j}}^{\bar{m}} \varphi_{\bar{m}}$ ).

$$\text{i.e. } (\bar{\partial} \varphi)_{\bar{j}_1 \bar{k}}^\alpha = (\nabla_{\bar{k}} \varphi_{\bar{j}_1}^\alpha - \nabla_{\bar{j}_1} \varphi_{\bar{k}}^\alpha) - (\Gamma_{\bar{k} \bar{j}}^{\bar{m}} - \Gamma_{\bar{j} \bar{k}}^{\bar{m}}) \varphi_{\bar{m} 1}^\alpha.$$

The last term, being the difference of two connections, is a tensor, called the torsion tensor.

(b).  $(g_{\bar{k} \bar{j}})$  Kähler  $\iff$  Torsion tensor = 0

Recall that,  $\Gamma_{\bar{k} \bar{j}}^{\bar{m}} = g^{m\bar{p}} \partial_{\bar{k}} g_{\bar{p} \bar{j}}$ , then  $\Gamma_{\bar{k} \bar{j}}^{\bar{m}} - \Gamma_{\bar{j} \bar{k}}^{\bar{m}} = g^{m\bar{p}} (\partial_{\bar{k}} g_{\bar{p} \bar{j}} - \partial_{\bar{j}} g_{\bar{p} \bar{k}})$ . Since  $(g^{m\bar{p}})$  is invertible, the last term vanishes iff  $\partial_{\bar{k}} g_{\bar{p} \bar{j}} - \partial_{\bar{j}} g_{\bar{p} \bar{k}} = 0$ , iff  $(g_{\bar{k} \bar{j}})$  Kähler.

• Henceforth, we will assume that  $(g_{\bar{k} \bar{j}})$  is Kähler.

Now we have:

$$\begin{aligned} &\frac{1}{p!(q+1)!} \int_X (\bar{\partial} \varphi)_{\bar{j}_1 \bar{k}}^\alpha \overline{\psi_{\bar{k} \bar{m} \bar{\ell}}^\beta} H_{\bar{\beta} \alpha} g^{k\bar{j}} g^{i\bar{m}} g^{\ell \bar{k}} \frac{\omega^n}{n!} \\ &= \frac{1}{p!q!} \int_X \nabla_{\bar{k}} \varphi_{\bar{j}_1}^\alpha \overline{\psi_{\bar{k} \bar{m} \bar{\ell}}^\beta} H_{\bar{\beta} \alpha} g^{k\bar{j}} g^{i\bar{m}} g^{\ell \bar{k}} \frac{\omega^n}{n!} \end{aligned}$$

Here we introduce the following trick:

Lemma. (Ricci).  $(\nabla_{\bar{k}} \varphi_{j\bar{i}}^\alpha) g^{k\bar{j}} = \nabla_{\bar{k}} (g^{k\bar{j}} \varphi_{j\bar{i}}^\alpha)$ .

Pf: The Chern connection preserves both the complex structure and metric structure:

$$\begin{aligned} \nabla_{\bar{k}} g_{i\bar{j}} &= \partial_{\bar{k}} g_{i\bar{j}} - \Gamma_{\bar{k}\bar{i}}^{\bar{j}} g_{\bar{j}} \\ &= \partial_{\bar{k}} g_{i\bar{j}} - g^{p\bar{l}} \partial_{\bar{k}} g_{i\bar{p}} \cdot g_{\bar{l}\bar{j}} \\ &= \partial_{\bar{k}} g_{i\bar{j}} - \partial_{\bar{k}} g_{i\bar{j}} \\ &= 0. \end{aligned}$$

□

Notice that  $g^{k\bar{j}} \varphi_{j\bar{i}}^\alpha$  is a section of a holomorphic bundle, thus:

$$\begin{aligned} & \frac{1}{p!q!} \int_X \nabla_{\bar{k}} \varphi_{j\bar{i}}^\alpha \overline{\psi_{\bar{k}m\bar{l}}^\beta} H_{\bar{\beta}\alpha} g^{k\bar{j}} g^{i\bar{m}} g^{l\bar{n}} \frac{\omega^n}{n!} \\ &= \frac{1}{p!q!} \int_X \nabla_{\bar{k}} (g^{k\bar{j}} \varphi_{j\bar{i}}^\alpha) \overline{\psi_{\bar{k}m\bar{l}}^\beta} g^{m\bar{i}} g^{k\bar{l}} H_{\bar{\beta}\alpha} \cdot \frac{\omega^n}{n!} \\ &= \frac{1}{p!q!} \int_X \partial_{\bar{k}} (g^{k\bar{j}} \varphi_{j\bar{i}}^\alpha) \overline{W_{k\alpha}^{i\bar{l}}} \frac{\omega^n}{n!} \\ &= - \frac{1}{p!q!} \int_X (g^{k\bar{j}} \varphi_{j\bar{i}}^\alpha) \partial_{\bar{k}} (\overline{W_{k\alpha}^{i\bar{l}}}) \frac{\omega^n}{n!}. \end{aligned}$$

where we denote for short  $\overline{\psi_{\bar{k}m\bar{l}}^\beta} g^{m\bar{i}} g^{k\bar{l}} H_{\bar{\beta}\alpha} = \overline{W_{k\alpha}^{i\bar{l}}}$ ; the last step used integration by parts.

Now, locally,  $\omega^n = \det(g_{\bar{q}p}) \prod_{i=1}^n dz^i \wedge d\bar{z}^i$ . thus

$$\begin{aligned} \partial_{\bar{k}} (W^{\bar{k}} \det g_{\bar{q}p}) &= \partial_{\bar{k}} W^{\bar{k}} \cdot \det g_{\bar{q}p} + W^{\bar{k}} \partial_{\bar{k}} \det g_{\bar{q}p} \\ &= \partial_{\bar{k}} W^{\bar{k}} \cdot \det g_{\bar{q}p} + \det g_{\bar{q}p} \cdot W^{\bar{k}} \partial_{\bar{k}} \log \det g_{\bar{q}p} \\ &= (\det g_{\bar{q}p}) (\partial_{\bar{k}} W^{\bar{k}} + W^{\bar{k}} g^{p\bar{q}} \partial_{\bar{k}} g_{\bar{q}p}) \\ &= (\det g_{\bar{q}p}) (\partial_{\bar{k}} W^{\bar{k}} + g^{q\bar{p}} \partial_{\bar{k}} g_{\bar{p}q} \cdot W^{\bar{k}}) \end{aligned}$$

where in the 3<sup>rd</sup> step, we used that  $(\log \det A)' = \text{tr}(A')$ ,

Recall also that  $\Gamma_{q\bar{k}}^{\bar{l}} = g^{l\bar{p}} \partial_q g_{\bar{p}k} \Rightarrow g^{q\bar{p}} \partial_{\bar{k}} g_{\bar{p}q} = \Gamma_{k\bar{q}}^q$ . Thus the above equation becomes:

$$\begin{aligned} \partial_{\bar{k}} (W^{\bar{k}} \det g_{\bar{q}p}) &= \det g_{\bar{q}p} \overline{(\partial_{\bar{k}} W^{\bar{k}} + \Gamma_{k\bar{q}}^q W^{\bar{k}})} \\ &= \det g_{\bar{q}p} \overline{(\partial_{\bar{k}} W^{\bar{k}} + \Gamma_{q\bar{k}}^q W^{\bar{k}})} \quad \text{cby the Kähler condition)} \\ &= \det g_{\bar{q}p} \overline{\nabla_{\bar{k}} W^{\bar{k}}} \\ &= \det g_{\bar{q}p} \nabla_{\bar{k}} W^{\bar{k}} \end{aligned}$$

Summing up, we obtain:

$$\begin{aligned}
& -\frac{1}{p!q!} \int_X (g^{k\bar{j}} \varphi_{\bar{j}I}^\alpha) (\partial_{\bar{k}} W_{k\alpha}^{I\bar{k}} \cdot \frac{\omega^n}{n!}) \\
&= -\frac{1}{p!q!} \int_X g^{k\bar{j}} \varphi_{\bar{j}I}^\alpha (\nabla_{\bar{k}} W_{k\alpha}^{I\bar{k}}) \frac{\omega^n}{n!} \\
&= \frac{1}{p!q!} \int_X g^{k\bar{j}} \varphi_{\bar{j}I}^\alpha (-g^{k\bar{\ell}} \nabla_k \psi_{\bar{k}M\bar{\ell}}^\beta) H_{\bar{\beta}\alpha} g^{I\bar{M}} \cdot \frac{\omega^n}{n!}
\end{aligned}$$

Hence by definition:

$$(\bar{\partial}^+ \psi)_{\bar{k}M}^\beta = -g^{k\bar{\ell}} \nabla_k \psi_{\bar{k}M\bar{\ell}}^\beta.$$

Now we can derive the formula for the Laplacian  $\square$ :  $\forall \varphi \in \Gamma(X, E \otimes \Lambda^{p,q})$

$$\begin{aligned}
(\bar{\partial}^+ \bar{\partial} \varphi)_{\bar{j}I} &= -g^{\ell\bar{m}} \nabla_{\bar{\ell}} (\bar{\partial} \varphi)_{\bar{j}I\bar{m}}^\alpha \\
&= -g^{\ell\bar{m}} \nabla_{\bar{\ell}} (\nabla_{\bar{m}} \varphi_{\bar{j}I}^\alpha - \nabla_{\bar{j}_1} \varphi_{\bar{j}_2 \dots \bar{j}_2 \bar{m}I}^\alpha - \dots - \nabla_{\bar{j}_q} \varphi_{\bar{m} \bar{j}_q \dots \bar{j}_1 I}^\alpha) \\
&= -g^{\ell\bar{m}} \nabla_{\bar{\ell}} \nabla_{\bar{m}} \varphi_{\bar{j}I}^\alpha + g^{\ell\bar{m}} (\nabla_{\bar{\ell}} \nabla_{\bar{j}_1} \varphi_{\bar{j}_2 \dots \bar{j}_2 \bar{m}I}^\alpha + \dots + \nabla_{\bar{\ell}} \nabla_{\bar{j}_q} \varphi_{\bar{m} \bar{j}_q \dots \bar{j}_1 I}^\alpha)
\end{aligned}$$

$$\begin{aligned}
(\bar{\partial} \bar{\partial}^+ \varphi)_{\bar{j}I} &= (\bar{\partial} \bar{\partial}^+ \varphi)_{\bar{j}_2 \dots \bar{j}_1 I}^\alpha \\
&= (-1)^p (\bar{\partial} \bar{\partial}^+ \varphi)_{\bar{j}_2 \dots \bar{j}_2 I \bar{j}_1}^\alpha \\
&= (-1)^p (\nabla_{\bar{j}_1} (\bar{\partial}^+ \varphi)_{\bar{j}_2 \dots \bar{j}_2 I}^\alpha - \nabla_{\bar{j}_2} (\bar{\partial}^+ \varphi)_{\bar{j}_2 \dots \bar{j}_2 \bar{j}_1 I}^\alpha - \dots - \nabla_{\bar{j}_q} (\bar{\partial}^+ \varphi)_{\bar{j}_1 \bar{j}_q \dots \bar{j}_2 I}^\alpha) \\
&= (-1)^p (\nabla_{\bar{j}_1} (-g^{\ell\bar{m}} \nabla_{\bar{\ell}} \varphi_{\bar{j}_2 \dots \bar{j}_2 I \bar{m}}^\alpha) + \nabla_{\bar{j}_2} (g^{\ell\bar{m}} \nabla_{\bar{\ell}} \varphi_{\bar{j}_2 \dots \bar{j}_2 \bar{j}_1 I \bar{m}}^\alpha) + \dots + \nabla_{\bar{j}_q} (g^{\ell\bar{m}} \nabla_{\bar{\ell}} \varphi_{\bar{j}_1 \bar{j}_q \dots \bar{j}_2 I \bar{m}}^\alpha)) \\
&= -g^{\ell\bar{m}} (\nabla_{\bar{j}_1} \nabla_{\bar{\ell}} \varphi_{\bar{j}_2 \dots \bar{j}_2 I \bar{m}}^\alpha - \nabla_{\bar{j}_2} \nabla_{\bar{\ell}} \varphi_{\bar{j}_2 \dots \bar{j}_2 \bar{j}_1 I \bar{m}}^\alpha - \dots - \nabla_{\bar{j}_q} \nabla_{\bar{\ell}} \varphi_{\bar{j}_1 \bar{j}_q \dots \bar{j}_2 I \bar{m}}^\alpha)
\end{aligned}$$

$$\begin{aligned}
\Rightarrow (\square \varphi)_{\bar{j}I}^\alpha &= \{ \bar{\partial}^+ (\bar{\partial} \varphi) \}_{\bar{j}I}^\alpha + \{ \bar{\partial} (\bar{\partial}^+ \varphi) \}_{\bar{j}I}^\alpha \\
&= -g^{\ell\bar{m}} \nabla_{\bar{\ell}} \nabla_{\bar{m}} \varphi_{\bar{j}I}^\alpha + g^{\ell\bar{m}} \{ [\nabla_{\bar{\ell}}, \nabla_{\bar{j}_1}] \varphi_{\bar{j}_2 \dots \bar{j}_2 \bar{m}I}^\alpha + \dots + [\nabla_{\bar{\ell}}, \nabla_{\bar{j}_q}] \varphi_{\bar{m} \bar{j}_q \dots \bar{j}_1 I}^\alpha \}
\end{aligned}$$

Note that the commutators can be replaced by curvature terms, for instance:

$$\begin{aligned}
[\nabla_{\bar{\ell}}, \nabla_{\bar{j}_1}] \varphi_{\bar{j}_2 \dots \bar{j}_2 \bar{m}I}^\alpha &= F_{\bar{j}_1 \bar{\ell}}^\alpha{}_\beta \varphi_{\bar{j}_2 \dots \bar{j}_2 \bar{m}I}^\beta \\
&+ R_{\bar{j}_1 \bar{\ell} \bar{j}_2}^{\bar{k}} \varphi_{\bar{k} \bar{j}_2 \dots \bar{j}_2 \bar{m}I}^\alpha + \dots + R_{\bar{j}_1 \bar{\ell} \bar{j}_2}^{\bar{k}} \varphi_{\bar{j}_2 \dots \bar{j}_2 \bar{k} \bar{m}I}^\alpha + R_{\bar{j}_1 \bar{\ell} \bar{m}}^{\bar{k}} \varphi_{\bar{j}_2 \dots \bar{j}_2 \bar{k} I}^\alpha \\
&- R_{\bar{j}_1 \bar{\ell}}^{\bar{k}}{}_\rho \varphi_{\bar{j}_2 \dots \bar{j}_2 \bar{m} \bar{k} \rho \dots \bar{i}_1}^\alpha - \dots - R_{\bar{j}_1 \bar{\ell}}^{\bar{k}}{}_{i_1} \varphi_{\bar{j}_2 \dots \bar{j}_2 \bar{m} I \rho \dots \bar{i}_2 \bar{k}}^\alpha.
\end{aligned}$$

Thm. (Kodaira Vanishing Theorem, Version 1)

Let  $E$  be a holomorphic line bundle over a compact Kähler manifold  $X$ . Let

$h$  be a metric on  $E$ ,  $F_{\bar{k}j}$  its curvature. Assume:

$$F_{\bar{k}j} + R_{\bar{k}j} \geq \varepsilon \cdot g_{\bar{k}j} \quad (*)$$

for some constant  $\varepsilon > 0$ . Then  $\ker \square|_{E \otimes \Lambda^{0,1}} = 0$ .

Pf: Apply Bochner-Kodaira formula to this case:

$$\begin{aligned} (\square \varphi)_j &= -g^{\ell\bar{m}} \nabla_{\bar{\ell}} \nabla_{\bar{m}} \varphi_j + g^{\ell\bar{m}} [\nabla_{\bar{\ell}}, \nabla_{\bar{m}}] \varphi_{\bar{m}} \\ &= -g^{\ell\bar{m}} \nabla_{\bar{\ell}} \nabla_{\bar{m}} \varphi_j + g^{\ell\bar{m}} \{ F_{j\bar{\ell}} \varphi_{\bar{m}} + R_{j\bar{\ell}\bar{m}}^{\bar{k}} \varphi_{\bar{k}} \} \end{aligned}$$

Under the Kähler condition:  $R_{j\bar{\ell}\bar{m}}^{\bar{k}} = R_{\bar{m}\bar{\ell}j}^{\bar{k}}$

$$\begin{aligned} (\square \varphi)_j &= -g^{\ell\bar{m}} \nabla_{\bar{\ell}} \nabla_{\bar{m}} \varphi_j + g^{\ell\bar{m}} F_{j\bar{\ell}} \varphi_{\bar{m}} + g^{\ell\bar{m}} R_{\bar{m}\bar{\ell}j}^{\bar{k}} \varphi_{\bar{k}} \\ &= -g^{\ell\bar{m}} \nabla_{\bar{\ell}} \nabla_{\bar{m}} \varphi_j + F_{j\bar{\ell}}^{\bar{m}} \varphi_{\bar{m}} + R_{j\bar{\ell}}^{\bar{m}} \varphi_{\bar{m}} \end{aligned}$$

Pair this with  $\varphi$ :

$$\begin{aligned} \langle \varphi, \square \varphi \rangle &= \int_X (\square \varphi)_j \bar{\varphi}_{\bar{k}} h g^{k\bar{j}} \frac{\omega^n}{n!} \\ &= - \int_X g^{\ell\bar{m}} \nabla_{\bar{\ell}} \nabla_{\bar{m}} \varphi_j \bar{\varphi}_{\bar{k}} g^{k\bar{j}} h \frac{\omega^n}{n!} + \int_X (F_{j\bar{\ell}}^{\bar{m}} \varphi_{\bar{m}} + R_{j\bar{\ell}}^{\bar{m}} \varphi_{\bar{m}}) \bar{\varphi}_{\bar{k}} h g^{k\bar{j}} \frac{\omega^n}{n!} \end{aligned}$$

$$\begin{aligned} \text{Now } & - \int_X g^{\ell\bar{m}} \nabla_{\bar{\ell}} \nabla_{\bar{m}} \varphi_j \bar{\varphi}_{\bar{k}} g^{k\bar{j}} h \frac{\omega^n}{n!} \\ &= - \int_X \nabla_{\bar{\ell}} (g^{\ell\bar{m}} \nabla_{\bar{m}} \varphi_j) \bar{\varphi}_{\bar{k}} g^{k\bar{j}} h \frac{\omega^n}{n!} \\ &= \int_X \nabla_{\bar{m}} \varphi_j \cdot \nabla_{\bar{\ell}} \bar{\varphi}_{\bar{k}} g^{\ell\bar{m}} g^{k\bar{j}} h \frac{\omega^n}{n!} \\ &= \|\nabla_{\bar{m}} \varphi_j\|^2 \end{aligned}$$

$$\begin{aligned} \text{Moreover, } & \int_X (F_{j\bar{\ell}}^{\bar{m}} \varphi_{\bar{m}} + R_{j\bar{\ell}}^{\bar{m}} \varphi_{\bar{m}}) \bar{\varphi}_{\bar{k}} h g^{k\bar{j}} \frac{\omega^n}{n!} \\ &= \int_X (F_{j\bar{\ell}} + R_{j\bar{\ell}}) g^{\ell\bar{m}} \varphi_{\bar{m}} \bar{\varphi}_{\bar{k}} g^{k\bar{j}} h \frac{\omega^n}{n!} \\ &\geq \int_X \varepsilon \cdot g_{j\bar{\ell}} g^{\ell\bar{m}} \varphi_{\bar{m}} \bar{\varphi}_{\bar{k}} g^{k\bar{j}} h \frac{\omega^n}{n!} \\ &= \varepsilon \int_X \varphi_j \bar{\varphi}_{\bar{k}} g^{k\bar{j}} \frac{\omega^n}{n!} \\ &= \varepsilon \|\varphi\|^2 \end{aligned}$$

$$\Rightarrow \langle \square \varphi, \varphi \rangle = \|\nabla_{\bar{m}} \varphi_j\|^2 + \varepsilon \|\varphi\|^2 \geq \varepsilon \|\varphi\|^2.$$

If  $\square \varphi = 0$ , then  $\varepsilon \|\varphi\|^2 = 0 \Rightarrow \varphi = 0$ . □

Rmk: The above estimate, together with Hodge decomposition thm (to be proven in what follows), gives:

(a). (Hodge)  $\Rightarrow H_{\bar{3}}^1(X, E) = 0$ , i.e.  $\forall f \in C^\infty(X, E \otimes \Lambda^{0,1})$ ,  $\bar{\partial} f = 0 \Rightarrow f = \bar{\partial} u$  for some  $u \in C^\infty(X, E)$ .

(b). Furthermore, we have  $\|u\|^2 \leq \frac{1}{\varepsilon} \|f\|^2$ .

• The bounds are independent of  $h$  as long as (\*) condition is satisfied. More precisely, fix a metric  $\hat{h}$  on  $\varphi$ , with curvature form  $\hat{F}_{\bar{k}\bar{j}}$ . Then:

Thm.  $\forall \varphi \in C^\infty(M, \mathbb{R}), h = e^{-\varphi} \hat{h}$ . Assume:

$$\partial_j \partial_{\bar{k}} \varphi + F_{\bar{k}j} + R_{\bar{k}j} \geq \epsilon g_{\bar{k}j}$$

Then  $\exists u$ , with  $\bar{\partial}u = f$ , with:

$$\int_X |u|^2 e^{-\varphi} \leq \frac{1}{\epsilon} \int_X |f|^2 e^{-\varphi}$$

$$\text{or } \|u\|^2 \leq \frac{1}{\epsilon} \|f\|^2 \quad \square$$

This is a very important theme in current research.

Thm. (Kodaira Vanishing Theorem, Version 2)

Let  $E \rightarrow X$  be a positive holomorphic line bundle over  $(X, g_{\bar{k}j})$ , compact Kähler manifold, i.e.  $\exists h$  a metric on  $E$  with  $F_{\bar{k}j} = -\partial_j \partial_{\bar{k}} \log h > 0$ . Then  $\exists m_0 > 0$ , s.t.  $\forall m \geq m_0$ , we have:

$$\text{Ker } \square |E^m \otimes \Lambda^{0,1} = 0$$

Pf: Bochner-Kodaira formula for  $E^m \otimes \Lambda^{0,1}$  is:

$$\square \varphi_j = -g^{\ell\bar{p}} \nabla_{\bar{\ell}} \nabla_{\bar{p}} \varphi_j + (m F_{j\bar{\ell}} + R_{j\bar{\ell}}) \varphi^{\ell}$$

Thus when  $m \geq m_0 \gg 0$ , we have:

$$\langle \square \varphi, \varphi \rangle \geq \epsilon \|\varphi\|^2$$

and the thm. follows. □

Kodaira-Akizuki-Nakano Formulas.

Previously,  $\square$  is compared with the metric Laplacian. However, on Kähler manifolds we have another Laplacian of  $\partial$ :  $\bar{\square} \triangleq \partial \partial^{\dagger} + \partial^{\dagger} \partial$ , so we would like to compare  $\square$  and  $\bar{\square}$ . To do this, we shall use the following operator  $\Lambda$ :

$$\Lambda: \Gamma(X, E \otimes \Lambda^{p+1, q+1}) \ni \Phi \longmapsto (\Lambda \Phi)_{\bar{j}I} = g^{\ell\bar{m}} \Phi_{\bar{m}\ell\bar{j}I} \in \Gamma(X, E \otimes \Lambda^{p, q}).$$

Then we have the identities:

$$\begin{cases} [\partial, \Lambda] = \bar{\partial}^{\dagger} \\ [\bar{\partial}, \Lambda] = -\partial^{\dagger} \end{cases}$$

Having this, we can prove:

Thm (KAN)

$$\square = \bar{\square} + [F, \Lambda]$$

Pf:  $\square - \bar{\square} = \bar{\partial}\bar{\partial}^\dagger + \bar{\partial}^\dagger\bar{\partial} - \partial\partial^\dagger - \partial^\dagger\partial$

$$\begin{aligned}
&= \bar{\partial}[\partial, \Lambda] + [\partial, \Lambda]\bar{\partial} + \partial[\bar{\partial}, \Lambda] + [\bar{\partial}, \Lambda]\partial \\
&= \bar{\partial}\partial\Lambda - \bar{\partial}\Lambda\bar{\partial} + \partial\Lambda\bar{\partial} - \Lambda\partial\bar{\partial} + \partial\bar{\partial}\Lambda - \partial\Lambda\bar{\partial} + \bar{\partial}\Lambda\bar{\partial} - \Lambda\bar{\partial}\partial \\
&= (\bar{\partial}\partial + \partial\bar{\partial})\Lambda - \Lambda(\partial\bar{\partial} + \bar{\partial}\partial) \\
&= [(\bar{\partial}\partial + \partial\bar{\partial}), \Lambda]
\end{aligned}$$

Recall that, for any  $\varphi \in \Gamma(X, E \otimes \Lambda^{p,q})$

$$d_A^2 \varphi = F \wedge \varphi$$

and

$$d_A^2 \varphi = (\partial + \bar{\partial})^2 \varphi = (\partial\bar{\partial} + \bar{\partial}\partial)\varphi$$

$\Rightarrow$

$$\square - \bar{\square} = [F, \Lambda]. \quad \square$$

Proof of the identities:

- $[\partial, \Lambda] = \bar{\partial}^\dagger$

$$[\partial, \Lambda]\varphi = \partial\Lambda\varphi - \Lambda\partial\varphi.$$

$$\begin{aligned}
\{\partial\Lambda\varphi\}_{\bar{L}Mj} &= \nabla_j(\Lambda\varphi)_{\bar{L}m_p \dots m_1} - \nabla_{m_1}(\Lambda\varphi)_{\bar{L}m_p \dots m_2 j} - \dots - \nabla_{m_p}(\Lambda\varphi)_{\bar{L}j m_{p-1} \dots m_1} \\
&= \nabla_j(g^{a\bar{b}}\varphi_{\bar{b}a})_{\bar{L}m_p \dots m_1} - \nabla_{m_1}(g^{a\bar{b}}\varphi_{\bar{b}a})_{\bar{L}m_p \dots m_2 j} - \dots - \nabla_{m_p}(g^{a\bar{b}}\varphi_{\bar{b}a})_{\bar{L}j m_{p-1} \dots m_1}.
\end{aligned}$$

$$\begin{aligned}
\{\Lambda\partial\varphi\}_{\bar{L}Mj} &= g^{a\bar{b}}(\partial\varphi)_{\bar{b}a}\bar{L}Mj \\
&= g^{a\bar{b}}(\nabla_j\varphi_{\bar{b}a})_{\bar{L}M} - \nabla_{m_1}\varphi_{\bar{b}a}\bar{L}m_p \dots m_2 j - \dots - \nabla_{m_p}\varphi_{\bar{b}a}\bar{L}j m_{p-1} \dots m_1 - \nabla_a\varphi_{\bar{b}j}\bar{L}M
\end{aligned}$$

Subtracting, using Ricci's lemma, we obtain:

$$\begin{aligned}
\{[\partial, \Lambda]\varphi\}_{\bar{L}Mj} &= g^{a\bar{b}}\nabla_a\varphi_{\bar{b}j}\bar{L}M \\
&= g^{a\bar{b}}\nabla_a\varphi_{\bar{L}M}\bar{\delta}j \\
&= -g^{a\bar{b}}\nabla_a\varphi_{\bar{L}Mj}\bar{\delta} \\
&= (\bar{\partial}^\dagger\varphi)_{\bar{L}Mj}
\end{aligned}$$

(Here recall that  $(\bar{\partial}^\dagger\psi)_{\bar{r}kM}^\beta = -g^{k\bar{\ell}}\nabla_k\psi_{\bar{r}M\bar{\ell}}^\beta$ .)

To show the second identity, we first compute  $\partial^\dagger$ . Given  $\varphi \in \Gamma(X, E \otimes \Lambda^{p,q})$

$\psi \in \Gamma(X, E \otimes \Lambda^{p+1,q})$ , by definition, we have  $\langle \partial\varphi, \psi \rangle = \langle \varphi, \partial^\dagger\psi \rangle$ , i.e.

$$\begin{aligned}
\langle \varphi, \partial^\dagger\psi \rangle &= \frac{1}{(p+1)!q!} \int_X (\partial\varphi)_{\bar{L}Mj} \overline{\psi_{\bar{p}a\bar{r}}^\beta} g^{p\bar{L}} g^{M\bar{a}} g^{j\bar{r}} H_{\alpha\bar{\beta}} \frac{\omega^n}{n!} \\
&= \frac{1}{p!q!} \int_X \nabla_j\varphi_{\bar{L}M}^\alpha \overline{\psi_{\bar{p}a\bar{r}}^\beta} g^{p\bar{L}} g^{M\bar{a}} g^{j\bar{r}} H_{\alpha\bar{\beta}} \frac{\omega^n}{n!}
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{p!q!} \int_X \nabla_j \varphi_{\bar{L}M}^\alpha \overline{\psi_{\bar{P}QR}^\beta g^{r\bar{j}}} g^{p\bar{L}} g^{m\bar{a}} H_{\alpha\bar{\beta}} \frac{\omega^n}{n!} \\
&= -\frac{1}{p!q!} \int_X \varphi_{\bar{L}M}^\alpha \overline{\nabla_j (g^{r\bar{j}} \psi_{\bar{P}QR}^\beta)} g^{p\bar{L}} g^{m\bar{a}} H_{\alpha\bar{\beta}} \frac{\omega^n}{n!}
\end{aligned}$$

$$\Rightarrow (\partial^\dagger \psi)_{\bar{P}Q}^\beta = -\nabla_j g^{r\bar{j}} \psi_{\bar{P}QR}^\beta$$

$$\bullet [\bar{\partial}, \wedge] = -\partial^\dagger$$

Take  $\varphi \in \Gamma(X, \wedge^{p,q} \otimes E)$ , then  $\wedge \varphi \in \Gamma(X, \wedge^{p-1, q+1} \otimes E)$ ,  $\bar{\partial} \wedge \varphi \in \Gamma(X, \wedge^{p-1, q} \otimes E)$

and  $\bar{\partial} \varphi \in \Gamma(X, \wedge^{p, q+1} \otimes E)$ ,  $\wedge \bar{\partial} \varphi \in \Gamma(X, \wedge^{p-1, q} \otimes E)$

$$\begin{aligned}
\{ \bar{\partial} \wedge \varphi \}_{\bar{j}_1 \bar{j}} &= \nabla_{\bar{j}} (\wedge \varphi)_{\bar{j}_1} - \nabla_{\bar{j}_1} (\wedge \varphi)_{\bar{j}_2 \dots \bar{j}_2 \bar{j}_1} - \dots - \nabla_{\bar{j}_{q-1}} (\wedge \varphi)_{\bar{j}_1 \bar{j}_2 \dots \bar{j}_1} \\
&= \nabla_{\bar{j}} g^{a\bar{b}} \varphi_{\bar{b} a \bar{j}_1} - \nabla_{\bar{j}_1} g^{a\bar{b}} \varphi_{\bar{b} a \bar{j}_2 \dots \bar{j}_2 \bar{j}_1} - \dots - \nabla_{\bar{j}_{q-1}} g^{a\bar{b}} \varphi_{\bar{b} a \bar{j}_1 \bar{j}_2 \dots \bar{j}_1}
\end{aligned}$$

$$\begin{aligned}
\{ \wedge \bar{\partial} \varphi \}_{\bar{j}_1 \bar{j}} &= g^{a\bar{b}} (\bar{\partial} \varphi)_{\bar{b} a \bar{j}_1} \\
&= g^{a\bar{b}} (\nabla_{\bar{j}} \varphi_{\bar{b} a \bar{j}_1} - \nabla_{\bar{b}} \varphi_{\bar{j} a \bar{j}_1} - \nabla_{\bar{j}_1} \varphi_{\bar{b} a \bar{j}_2 \dots \bar{j}_2 \bar{j}_1} - \dots - \nabla_{\bar{j}_{q-1}} \varphi_{\bar{b} a \bar{j}_1 \bar{j}_2 \dots \bar{j}_1})
\end{aligned}$$

Subtracting gives:

$$\begin{aligned}
([\bar{\partial}, \wedge] \varphi)_{\bar{j}_1 \bar{j}} &= g^{a\bar{b}} \nabla_{\bar{b}} \varphi_{\bar{j} a \bar{j}_1} \\
&= g^{a\bar{b}} \nabla_{\bar{b}} \varphi_{\bar{j}_1 \bar{j} a} \\
&= (-\partial^\dagger \varphi)_{\bar{j}_1 \bar{j}}
\end{aligned}$$

□

Rmk: The signs of these identities can be easily checked as follows.

For  $\varphi \in \Gamma(X, \wedge^{0,1})$ ,  $\varphi = \varphi_{\bar{j}} d\bar{z}^{\bar{j}}$ , then

$$[\partial, \wedge] \varphi = (\partial \wedge - \wedge \partial) \varphi = -\wedge \partial \varphi$$

Now,  $\partial \varphi = \nabla_{\bar{k}} \varphi_{\bar{j}} dz^{\bar{k}} \wedge d\bar{z}^{\bar{j}} \Rightarrow (\partial \varphi)_{\bar{j}\bar{k}} = \nabla_{\bar{k}} \varphi_{\bar{j}}$ . Thus

$$[\partial, \wedge] \varphi = -g^{\bar{k}\bar{j}} (\partial \varphi)_{\bar{j}\bar{k}} = -g^{\bar{k}\bar{j}} (\nabla_{\bar{k}} \varphi_{\bar{j}}) = \bar{\partial}^\dagger \varphi$$

Similarly if we take  $\psi \in \Gamma(X, \wedge^{1,0})$ ,  $\psi = \psi_{\bar{k}} dz^{\bar{k}}$ , then

$$[\bar{\partial}, \wedge] \psi = (\bar{\partial} \wedge - \wedge \bar{\partial}) \psi = -\wedge \bar{\partial} \psi$$

where  $\bar{\partial} \psi = \nabla_{\bar{j}} \psi_{\bar{k}} d\bar{z}^{\bar{j}} \wedge dz^{\bar{k}}$ , or  $(\bar{\partial} \psi)_{\bar{j}\bar{k}} = -\nabla_{\bar{j}} \psi_{\bar{k}}$ . Hence

$$[\bar{\partial}, \wedge] \psi = -\wedge \bar{\partial} \psi = -g^{\bar{k}\bar{j}} (\bar{\partial} \psi)_{\bar{j}\bar{k}} = g^{\bar{k}\bar{j}} \nabla_{\bar{j}} \psi_{\bar{k}} = -\partial^\dagger \psi_{\bar{k}}$$

Lemma. Let  $E \rightarrow X$  be a line bundle with metric  $h$  and curvature  $F_{\bar{k}\bar{j}}$ .

At each  $z \in X$ , let  $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$  be the eigenvalues of  $F_{\bar{k}\bar{j}}$  w.r.t.  $g_{\bar{k}\bar{j}}$ .

Then :



$$\langle [F, \Lambda] \psi, \psi \rangle_{\mathbb{Z}} = \sum_{\bar{k} \bar{j}} (\sum_{a \in \mathbb{J}} \lambda_a + \sum_{b \in \bar{\mathbb{K}}} \lambda_b - \sum_{c=1}^n \lambda_c) |\psi_{\bar{k} \bar{j}}|^2$$

where we take a coordinate system s.t.  $dz^1, \dots, dz^n$  are orthonormal at  $\mathbb{Z}$ .

Rmk: Recall that since  $F_{\bar{k} \bar{j}}$  and  $g_{\bar{k} \bar{j}}$  are both Hermitian,  $T^{\bar{l}}_{\bar{j}} \triangleq g^{\bar{l} \bar{k}} F_{\bar{k} \bar{j}}$  is then a self-adjoint endomorphism of the holomorphic tangent space at  $\mathbb{Z}$ :

$$\langle T a, b \rangle_g = \langle a, T b \rangle_g$$

$$\text{Indeed, } g_{\bar{k} \bar{j}} T^{\bar{l}}_{\bar{j}} a^{\bar{l}} \bar{b}^{\bar{k}} = g_{\bar{k} \bar{j}} g^{i \bar{s}} F_{\bar{s} \bar{l}} a^{\bar{l}} \bar{b}^{\bar{k}} = F_{\bar{k} \bar{j}} a^{\bar{l}} \bar{b}^{\bar{k}},$$

$$\text{and } g_{\bar{s} \bar{j}} a^{\bar{j}} T^{\bar{s}}_{\bar{k}} \bar{b}^{\bar{k}} = g_{\bar{s} \bar{j}} a^{\bar{j}} g^{\bar{s} \bar{r}} F_{\bar{r} \bar{k}} \bar{b}^{\bar{k}} = g_{\bar{s} \bar{j}} g^{\bar{r} \bar{s}} F_{\bar{k} \bar{r}} a^{\bar{j}} \bar{b}^{\bar{k}} = F_{\bar{k} \bar{j}} a^{\bar{j}} \bar{b}^{\bar{k}}.$$

Thus we may talk about the eigenvalues of  $T$ :  $\lambda_1 \leq \lambda_2 \leq \dots \leq \lambda_n$ .

Pf: We compute:

$$\begin{aligned} (\Lambda F \psi)_{\bar{k} \bar{j} \bar{r} \bar{s}} &= g^{n \bar{m}} (F \psi)_{\bar{m} \bar{n} \bar{k} \bar{j} \bar{r} \bar{s}} \\ &= g^{n \bar{m}} (F \psi)_{\bar{k} \bar{j} \bar{m} \bar{n} \bar{r} \bar{s}} \\ &= g^{n \bar{m}} (F_{\bar{r} \bar{s}} \psi_{\bar{k} \bar{j} \bar{m} \bar{n}} + \text{anti-symmetric terms of } (\bar{r}, \bar{k}, \bar{m}), (s, \bar{j}, n)) \\ &= g^{n \bar{m}} (F_{\bar{r} \bar{s}} \psi_{\bar{k} \bar{j} \bar{m} \bar{n}} \\ &\quad \textcircled{1} \\ &\quad - \underbrace{\sum_a F_{\bar{k} a s} \psi_{\bar{k} \bar{q} \dots \bar{k}_{a+1} \bar{r} \bar{k}_{a-1} \dots \bar{k}_i \bar{j} \bar{m} \bar{n}}}_{\textcircled{2}} - F_{\bar{m} \bar{s}} \psi_{\bar{k} \bar{j} \bar{r} \bar{n}} \\ &\quad \textcircled{3} \\ &\quad - \sum_b F_{\bar{r} \bar{j} b} \psi_{\bar{k} \bar{j} p \dots j b s j b-1 \dots j_i \bar{m} \bar{n}} - F_{\bar{r} \bar{n}} \psi_{\bar{k} \bar{j} \bar{m} \bar{s}} \\ &\quad \textcircled{4} \\ &\quad + \underbrace{\sum_a \sum_b F_{\bar{k} a j b} \psi_{\bar{k} \bar{q} \dots \bar{r} \dots \bar{k}_i j p \dots s \dots j_i \bar{m} \bar{n}}}_{\textcircled{4}} + \sum_a F_{\bar{k} a n} \psi_{\bar{k} \bar{q} \dots \bar{r} \dots \bar{k}_i \bar{j} \bar{m} \bar{s}} \\ &\quad + \sum_b F_{\bar{m} j b} \psi_{\bar{k} \bar{j} p \dots s \dots j_i \bar{r} \bar{n}} + F_{\bar{m} \bar{n}} \psi_{\bar{k} \bar{j} \bar{r} \bar{s}}) \end{aligned}$$

Next.

$$\begin{aligned} (F \Lambda \psi)_{\bar{k} \bar{j} \bar{r} \bar{s}} &= F_{\bar{r} \bar{s}} g^{n \bar{m}} \psi_{\bar{m} \bar{n} \bar{k} \bar{j}} + \text{anti-symmetric terms in } (\bar{r}, \bar{k}), (s, \bar{j}) \\ &= F_{\bar{r} \bar{s}} g^{n \bar{m}} \psi_{\bar{m} \bar{n} \bar{k} \bar{j}} \\ &\quad \textcircled{1} \\ &\quad - \underbrace{\sum_a F_{\bar{k} a s} g^{n \bar{m}} \psi_{\bar{m} \bar{n} \bar{k} \bar{q} \dots \bar{r} \dots \bar{k}_i \bar{j}}}_{\textcircled{2}} - \underbrace{\sum_b F_{\bar{r} \bar{j} b} g^{n \bar{m}} \psi_{\bar{m} \bar{n} \bar{k} \bar{j} p \dots s \dots j_i}}_{\textcircled{3}} \\ &\quad \textcircled{4} \\ &\quad + \underbrace{\sum_a \sum_b F_{\bar{k} a j b} g^{n \bar{m}} \psi_{\bar{m} \bar{n} \bar{k} \bar{q} \dots \bar{r} \dots \bar{k}_i j p \dots s \dots j_i}}_{\textcircled{4}} \end{aligned}$$

Subtracting gives (notice the underlined cancelation relation)

$$([\Lambda, F]\psi)_{\bar{k}j\bar{r}s} = -g^{n\bar{m}} F_{\bar{m}s} \psi_{\bar{k}j\bar{r}n} - g^{n\bar{m}} F_{\bar{r}n} \psi_{\bar{k}j\bar{m}s} \\ + \sum_a g^{n\bar{m}} F_{\bar{k}an} \psi_{\bar{k}a\bar{r}\dots\bar{r}\dots\bar{k}j\bar{m}s} + \sum_b g^{n\bar{m}} F_{\bar{m}jb} \psi_{\bar{k}j\bar{r}\dots s\dots j\bar{r}n} + g^{n\bar{m}} F_{\bar{m}n} \psi_{\bar{k}j\bar{r}s}$$

Now in a local coordinate system where  $g_{\bar{m}n} = \delta_{mn}$ ,  $F_{\bar{k}j} = \lambda_j \delta_{kj}$ , we have:

$$([\Lambda, F]\psi)_{\bar{k}j\bar{r}s} = -\lambda_s \psi_{\bar{k}j\bar{r}s} - \lambda_r \psi_{\bar{k}j\bar{r}s} + \sum_a \lambda_{ka} \psi_{\bar{k}a\bar{r}\dots\bar{r}\dots\bar{k}j\bar{r}as} \\ + \sum_b \lambda_{jb} \psi_{\bar{k}j\bar{r}\dots s\dots j\bar{r}jb} + \sum_{c=1}^n \lambda_c \psi_{\bar{k}j\bar{r}s} \\ = -\lambda_s \psi_{\bar{k}j\bar{r}s} - \lambda_r \psi_{\bar{k}j\bar{r}s} - \sum_a \lambda_{ka} \psi_{\bar{k}j\bar{r}s} - \sum_b \lambda_{jb} \psi_{\bar{k}j\bar{r}s} + \sum_{c=1}^n \lambda_c \psi_{\bar{k}j\bar{r}s}. \\ = (-\sum_{a \in \mathbb{R}, \bar{r}} \lambda_a - \sum_{b \in \mathbb{J}, s} \lambda_b + \sum_{c=1}^n \lambda_c) \psi_{\bar{k}j\bar{r}s}. \quad \square$$

As an immediate corollary, we obtain:

Thm (KAN)

Let  $E \rightarrow X$  be a positive line bundle, with metric  $h$  and curvature

$$F = -\partial_j \bar{\partial}_{\bar{k}} \log h > 0$$

Choose the curvature form of  $E$  as metric on  $X$ , i.e.  $g_{\bar{k}j} = F_{\bar{k}j}$  (which implies all  $\lambda_a = 1$ ,  $a = 1, \dots, n$ ). Then:

$$\langle [F, \Lambda]\psi, \psi \rangle = (p+q-n) \|\psi\|^2$$

Consequently,

$$\text{Ker } \square|_{E \otimes \Lambda^{p,q}} = 0 \text{ for } p+q > n.$$

In particular,

$$\text{Ker } \square|_{E \otimes \Lambda^{n,q}} = 0 \text{ for any } q > 1. \quad \square$$

Observations:

1. Together with Hodge thm, this again gives solvability with  $L^2$ -bounds  $\bar{\partial}u = f$ .
2. There are versions of all these where  $X$  is not compact but complete and pseudo-convex. (Demailly)
3. There are versions for  $X$  bounded pseudo-convex domains in  $\mathbb{C}^n$  (not complete) (Hörmander,  $L^2$ -estimate)

## §7. Hodge Decomposition Theorem

Let  $E \rightarrow X$  be a holomorphic vector bundle,  $H_{\bar{\alpha}\beta}$  a metric on  $E$ ,  $g_{\bar{k}j}$  a metric on  $X$  (not necessarily Kähler). Define  $\square = \bar{\partial}\bar{\partial}^\dagger + \bar{\partial}^\dagger\bar{\partial}$  on  $C^\infty(X, E \otimes \Lambda^{p,q})$ . The main result of this chapter is the following:

Thm. (Hodge, Kodaira)

Let  $L^2(X, E \otimes \Lambda^{p,q})$  denote the space of  $L^2$   $(p,q)$ -forms valued in  $E$ :

(a).  $\exists$  an orthonormal basis  $\{\psi_\ell\}$  of  $L^2(X, E \otimes \Lambda^{p,q})$ ,  $\psi_\ell \in C^\infty(X, E \otimes \Lambda^{p,q})$  and

$$\square \psi_\ell = \lambda_\ell \psi_\ell$$

for each  $\ell \in \mathbb{N}$ .

(b)  $\exists$  an operator  $G: L^2 \rightarrow L^2$ , bounded, self-adjoint operator so that

$$G\bar{\partial} = \bar{\partial}G$$

$$G\bar{\partial}^\dagger = \bar{\partial}^\dagger G$$

$$\square G = G\square = \text{Id} - \pi$$

where  $\pi: L^2 \rightarrow L^2$  is the orthogonal projection of  $L^2$  inner product space onto  $\text{Ker } \square = \{\varphi \in C^\infty(X, E \otimes \Lambda^{p,q}) \mid \square\varphi = 0\}$ .

(c). For each  $\lambda$ , the eigenspace  $\{\varphi \in C^\infty(X, E \otimes \Lambda^{p,q}) \mid \square\varphi = \lambda\varphi\}$  is finite dimensional and  $\lambda_m \geq C \cdot m^\delta$  for some  $\delta > 0$ .

Cor. 1.  $\forall \varphi \in C^\infty(X, E \otimes \Lambda^{p,q})$ ,  $\varphi$  can be written as  $(\text{Id} = \square G + \pi)$ :

$$\varphi = \pi\varphi + \square G\varphi = \pi\varphi + \bar{\partial}(\bar{\partial}^\dagger G\varphi) + \bar{\partial}^\dagger(\bar{\partial}G\varphi)$$

i.e.  $C^\infty(X, E \otimes \Lambda^{p,q}) = \text{Ker } \square \oplus \text{Range } \bar{\partial} \oplus \text{Range } \bar{\partial}^\dagger$ , where  $\oplus$  means that the summands are mutually orthogonal to each other. □

Def. (Dolbeault cohomology). For the complex:

$$\dots \xrightarrow{\bar{\partial}} \Gamma(X, E \otimes \Lambda^{p,q-1}) \xrightarrow{\bar{\partial}} \Gamma(X, E \otimes \Lambda^{p,q}) \xrightarrow{\bar{\partial}} \Gamma(X, E \otimes \Lambda^{p,q+1}) \xrightarrow{\bar{\partial}} \dots$$

it's Dolbeault cohomology is defined as:

$$H_{\bar{\partial}}^{p,q}(X, E) \hat{=} (\text{Ker } \bar{\partial}|_{E \otimes \Lambda^{p,q}}) / (\text{Im } \bar{\partial}|_{E \otimes \Lambda^{p,q-1}})$$

i.e.  $\forall \varphi, \varphi' \in \text{Ker } \bar{\partial}|_{E \otimes \Lambda^{p,q}}$ ,  $\varphi \sim \varphi' \Leftrightarrow \varphi - \varphi' = \bar{\partial}\psi$  for some  $\psi \in \Gamma(X, E \otimes \Lambda^{p,q-1})$ .

Cor 2. If  $\varphi \in C^\infty(X, E \otimes \Lambda^{p,q})$  and  $\bar{\partial}\varphi = 0$ , then

$$\varphi = \pi\varphi + \bar{\partial}\bar{\partial}^\dagger G\varphi$$

i.e. any Dolbeault cohomology class  $[\varphi]$  can be represented by a harmonic representative  $\pi\varphi \in \text{Ker } \square|_{E \otimes \Lambda^{p,q}}$ .  $\square$

Rmk:  $\text{Ker } \square|_{E \otimes \Lambda^{p,q}}$  depends apparently on metrics while  $H_{\bar{\partial}}^{p,q}$  is independent of metric. Thus with the vanishing thms proven previously, we can obtain information (vanishing) of  $H_{\bar{\partial}}^{p,q}$ .

Sobolev Spaces  $H_{(s)}(X, E \otimes \Lambda^{p,q})$

$X$ : compact. We define the Sobolev norm  $\|\cdot\|_{(s)}$  on  $C^\infty(X, E \otimes \Lambda^{p,q})$ .

$$\|\varphi\|_{(s)}^2 \triangleq \sum_{k \leq s} \int_X \nabla_{a_1 \dots a_k} \varphi^\alpha \overline{\nabla_{b_1 \dots b_k} \varphi^\beta} g^{A\bar{B}} H_{\bar{\beta}\alpha} \frac{\omega^n}{n!},$$

where  $a_i, b_j$  range in  $\{1, \dots, n, \bar{1}, \dots, \bar{n}\}$ .  $H_{\bar{\beta}\alpha}$  is the metric on  $E \otimes \Lambda^{p,q}$ .

For e.g.

$$\|\varphi\|_{(0)}^2 = \int_X \varphi^\alpha \bar{\varphi}^\beta H_{\bar{\beta}\alpha} \frac{\omega^n}{n!} = \|\varphi\|_{L^2}^2$$

$$\|\varphi\|_{(1)}^2 = \|\varphi\|_{(0)}^2 + \int_X \nabla_j \varphi^\alpha \overline{\nabla_{\bar{k}} \varphi^\beta} g^{j\bar{k}} H_{\bar{\beta}\alpha} \frac{\omega^n}{n!} + \int_X \nabla_{\bar{j}} \varphi^\alpha \overline{\nabla_k \varphi^\beta} g^{k\bar{j}} H_{\bar{\beta}\alpha} \frac{\omega^n}{n!}$$

$$\begin{aligned} \|\varphi\|_{(2)}^2 = & \|\varphi\|_{(1)}^2 + \int_X \nabla_j \nabla_{\bar{k}} \varphi^\alpha \overline{\nabla_{\bar{\ell}} \nabla_m \varphi^\beta} g^{j\bar{\ell}} g^{k\bar{m}} H_{\bar{\beta}\alpha} \frac{\omega^n}{n!} \\ & + \int_X \nabla_{\bar{j}} \nabla_k \varphi^\alpha \overline{\nabla_{\bar{\ell}} \nabla_m \varphi^\beta} g^{\ell\bar{j}} g^{m\bar{k}} H_{\bar{\beta}\alpha} \frac{\omega^n}{n!} \\ & + \int_X \nabla_j \nabla_{\bar{k}} \varphi^\alpha \overline{\nabla_{\bar{\ell}} \nabla_m \varphi^\beta} g^{j\bar{\ell}} g^{m\bar{k}} H_{\bar{\beta}\alpha} \frac{\omega^n}{n!} \\ & + \int_X \nabla_{\bar{j}} \nabla_k \varphi^\alpha \overline{\nabla_{\bar{\ell}} \nabla_m \varphi^\beta} g^{\ell\bar{j}} g^{k\bar{m}} H_{\bar{\beta}\alpha} \frac{\omega^n}{n!} \quad \text{etc.} \end{aligned}$$

Def.  $H_{(s)}(X, E \otimes \Lambda^{p,q}) = \overline{C^\infty(X, E \otimes \Lambda^{p,q})}$  w.r. t.  $\|\cdot\|_{(s)}$ .

( = Space of Cauchy sequences in  $C^\infty(X, E \otimes \Lambda^{p,q})$ ,  
Cauchy w.r. t.  $\|\cdot\|_{(s)}$  ).

i.e.  $\varphi \in H_{(s)}$ ,  $\varphi = \{\varphi_\ell\}_{\ell \in \mathbb{N}}$ .  $\|\varphi_\ell - \varphi_m\|_{(s)} \rightarrow 0$ , as  $\ell, m \rightarrow \infty$ .

By def. of  $\|\cdot\|_{(s)}$ ,  $\varphi_\ell \rightarrow \varphi$  in  $H_{(s)}$  for some  $\varphi \in L^2$ , thus we may think of  $H_{(s)}$  as a subspace of  $L^2$ .

Observation:  $\|\varphi_\ell - \varphi_m\|_{C^s} \rightarrow 0 \Leftrightarrow \|\nabla^k \varphi_\ell - \nabla^k \varphi_m\|_{L^2} \rightarrow 0$  for all  $k \leq s$ , i.e.

$\nabla^k \varphi_\ell \rightarrow \psi_k$  in  $L^2$ , for some  $\psi_k \in L^2$ ,  $\forall k \leq s$ . Thus  $\varphi \in H_{loc}^s$  gives rise to a sequence  $\{\varphi_\ell: \varphi_\ell \in C^\infty(X, E \otimes \Lambda^{p,q})\}$ ,  $\varphi_\ell \rightarrow \varphi$  in  $L^2$ , and  $\nabla^k \varphi_\ell$  converges in  $L^2$  (to some  $\psi_k$ ) for all  $k \leq s$ .

Def. The  $k$ -th derivative of  $\varphi$  (in the sense of distributions) is defined by

$$\nabla^k \varphi \triangleq \lim_\ell \nabla^k \varphi_\ell = \psi_k \in L^2$$

for all  $k \leq s$ .

Formal adjointness

We know that,  $\forall \Phi, \Psi \in C^\infty$ ,

$$\langle \square \Phi, \Psi \rangle = \langle \Phi, \square \Psi \rangle,$$

since  $\square = \bar{\partial} \bar{\partial}^* + \bar{\partial}^* \bar{\partial}$  and

$$\begin{aligned} \langle \square \Phi, \Psi \rangle &= \langle \bar{\partial} \bar{\partial}^* \Phi, \Psi \rangle + \langle \bar{\partial}^* \bar{\partial} \Phi, \Psi \rangle \\ &= \langle \bar{\partial}^* \Phi, \bar{\partial} \Psi \rangle + \langle \bar{\partial} \Phi, \bar{\partial}^* \Psi \rangle \end{aligned}$$

This remains true if  $\varphi, \psi \in H_{loc}^2$

$$\langle \square \varphi, \psi \rangle = \langle \varphi, \square \psi \rangle$$

where the inner product taken in  $L^2$ .

pf: Take  $\varphi_\ell \rightarrow \varphi$ ,  $\psi_\ell \rightarrow \psi$ . Then  $\square \varphi_\ell$  and  $\square \psi_\ell$  are smooth

$$\langle \square \varphi, \psi \rangle = \lim_{\ell \rightarrow \infty} \langle \square \varphi_\ell, \psi_\ell \rangle = \lim_{\ell \rightarrow \infty} \langle \varphi_\ell, \square \psi_\ell \rangle = \langle \varphi, \psi \rangle.$$

Basic facts about Sobolev spaces

(a). The operator  $\square$  satisfies,  $\forall \Phi \in C^\infty(X, E \otimes \Lambda^{p,q})$

$$\|\square \Phi\|_{C^s} \leq C_s \|\Phi\|_{C^{s+2}}$$

Thus  $\square$  extends uniquely to a linear bounded operator  $\square: H_{loc}^{s+2} \rightarrow H_{loc}^s$ .

Indeed,  $\varphi \in H_{loc}^{s+2}$ , write  $\varphi = \lim_\ell \varphi_\ell$  w/r. t.  $\|\cdot\|_{C^{s+2}}$ , then

$$\|\square \varphi_\ell - \square \varphi_m\|_{C^s} \leq C_s \|\varphi_\ell - \varphi_m\|_{C^{s+2}} \rightarrow 0 \text{ as } \ell, m \rightarrow \infty.$$

Hence  $\{\square \varphi_\ell\}$  is Cauchy w/r. t.  $\|\cdot\|_{C^s}$ , thus defines an element of  $H_{loc}^s$ .

(b). Sobolev lemma.

Let  $n_{\mathbb{R}}$  be the real dimension of  $X$ . If  $s > \frac{n_{\mathbb{R}}}{2}$ , then  $\exists C_s$ , s.t.  $\forall \Phi \in C^\infty$ ,

$$\|\Phi\|_{C^0} \leq C_s \|\Phi\|_{H^s}$$

where  $\|\Phi\|_{C^0}$  is the sup norm:

$$\|\Phi\|_{C^0} = \sup_x (\Phi_{x_1}^\alpha \Phi_{x_2}^\beta \dots \Phi_{x_n}^\gamma g^{j_1 \bar{j}_1} g^{l_1 \bar{l}_1})$$

Two corollaries are immediate:

Cor 1. If  $\varphi \in H^s$  for  $s > \frac{n_{\mathbb{R}}}{2}$ , then  $\varphi$  is continuous.

Pf: Let  $\varphi_\ell \rightarrow \varphi$  in  $L^2$ ,  $\varphi_\ell \in C^\infty$ ,  $\|\varphi_\ell - \varphi_m\|_{H^s} \rightarrow 0$  as  $\ell, m \rightarrow \infty$ . By Sobolev lemma,  $\|\varphi_\ell - \varphi_m\|_{C^0} \leq C_s \|\varphi_\ell - \varphi_m\|_{H^s} \rightarrow 0$  as  $\ell, m \rightarrow \infty$ . i.e.  $\{\varphi_\ell\}$  is uniformly Cauchy and thus converges to a continuous function  $\tilde{\varphi}$ . But  $\varphi_\ell \rightarrow \tilde{\varphi}$  in  $L^2$  for some  $\tilde{\varphi}$ . By uniqueness of limit,  $\varphi = \tilde{\varphi}$ , and thus  $\varphi$  is continuous.  $\square$

Cor 2. If  $s > \frac{n_{\mathbb{R}}}{2} + m$ , then  $\varphi \in H^s \Rightarrow \varphi \in C^m$ . Thus  $\bigcap_{s>0} H^s = C^\infty$   $\square$

(c). Rellich's lemma.

$X$ : compact.  $s < t$ . Let  $\{\varphi_j \in H^t\}$  be a sequence s.t.  $\|\varphi_j\|_{H^t} \leq 1$ . Then  $\exists$  a subsequence  $\{\varphi_{j_k}\}$ , and  $\varphi \in H^s$ , and  $\varphi_{j_k} \rightarrow \varphi$  in  $H^s$ .

Linear elliptic PDE

(a). Let  $\square$  be the Laplacian, then  $\forall s, \exists C_s$  s.t.  $\forall \varphi \in C^\infty$ ,

$$\|\varphi\|_{H^{s+2}} \leq C_s (\|\square\varphi\|_{H^s} + \|\varphi\|_{H^{s+1}})$$

This depends crucially on the ellipticity of Laplacian, and is called "A Priori estimate". Similar as above, this translates directly to  $H^{s+2}$ .

(b). Regularity Thm.

Let  $\Omega \subseteq X$  be an open set and  $u, f \in L^2$  s.t.  $\square u = f$  on  $\Omega$  in the sense of distributions, (i.e. by def.  $\langle u, \square\varphi \rangle = \langle f, \varphi \rangle$ ,  $\forall \varphi \in C_c^\infty(\Omega)$ ) and  $\square u = f$  in the standard sense.

## Construction of Green's operator $G$

(a). Let  $\text{Ker } \square = \{ \varphi \in H_{(2)} ; \square \varphi = 0 \}$ . Then  $\text{Ker } \square$  is finite dimensional

Pf: If  $\{ \varphi_j \}$  were an infinite orthonormal basis of  $\text{Ker } \square$  in  $H_{(2)}$ . By a priori estimate, we have, for  $\varphi_j - \varphi_\ell$  ( $j \neq \ell$ )

$$\begin{aligned} \|\varphi_j - \varphi_\ell\|_{(2)} &\leq C \cdot (\|\square(\varphi_j - \varphi_\ell)\|_{(0)} + \|\varphi_j - \varphi_\ell\|_{(1)}) \\ &= C \cdot \|\varphi_j - \varphi_\ell\|_{(1)} \end{aligned}$$

Since  $\varphi_j, \varphi_\ell \in \text{Ker } \square$ . But  $\|\varphi_j\|_{(2)} = 1$ , and thus by Rellich's lemma, we may assume  $\|\varphi_j - \varphi_\ell\|_{(1)} \rightarrow 0$ , by passing to a subsequence. This tells us that

$$\sqrt{2} = \|\varphi_j - \varphi_\ell\|_{(2)} \rightarrow 0 \text{ as } j, \ell \rightarrow \infty.$$

Contradiction. □

Similarly, we can show that each eigenspace  $\text{Ker}(\square - \lambda)$  is finite dimensional, since we have

$$\begin{aligned} \|\varphi\|_{(s+2)} &\leq C \cdot (\|\square \varphi\|_{(s)} + \|\varphi\|_{(s+1)}) \\ &\leq C \cdot (\|\square \varphi - \lambda \varphi\|_{(s)} + |\lambda| \|\varphi\|_{(s)} + \|\varphi\|_{(s+1)}) \end{aligned}$$

and the above proof applies.

Note that by regularity thm,  $\text{Ker } \square$  actually consists of smooth functions.

(b). Define the range of  $\square$  by:

$$R(\square) \triangleq \{ \varphi \in H_{(0)}, \varphi = \square \psi \text{ for some } \psi \in H_{(2)} \}$$

We have the key fact that:

- $R(\square)$  is closed

Observations: In general,  $W \subseteq H$ ,  $H$ : Hilbert space and  $W$  a subspace, we have

$$H = \overline{W} \oplus W^\perp$$

Then since the range is closed, we can write

$$L^2 = R(\square) \oplus R(\square)^\perp$$

i.e.  $\forall \varphi \in L^2$ ,  $\varphi = \square \psi + \varphi_0$ , with  $\varphi_0 \in R(\square)^\perp$ . We may further choose  $\psi \perp \text{Ker } \square$  in the  $L^2$ -sense. (One may worry that throwing away from  $\psi$  its component in  $\text{Ker } \square$  may not be well-defined. But there is no trouble since  $\text{Ker } \square$  is finite dimensional thus closed, and in fact consists of smooth functions).

We have shown that  $R(\square)$  is closed in §4. We reproduce the proof here for the sake of completeness:

Claim: a priori estimate  $\Rightarrow \exists C > 0$ , s.t.  $\forall \varphi \in C^\infty$ ,  $\varphi \perp \text{Ker } \square$ , we have

$$\|\varphi\|_{L^2} \leq C \cdot \|\square\varphi\|_{L^2} \quad (\text{AP}')$$

Then,  $\forall \Phi \in R(\square)$ , and  $\Phi_n = \square\varphi_n$ ,  $\varphi_n \perp \text{Ker } \square$ , and  $\Phi_n \rightarrow \Phi$  in  $H_{(2)}$ . Now, by (AP')  $\|\varphi_n - \varphi_m\|_{L^2} \leq C \cdot \|\square\varphi_n - \square\varphi_m\|_{L^2} = C \|\Phi_n - \Phi_m\|_{L^2} \rightarrow 0 \Rightarrow \{\varphi_n\}$  converges in  $H_2$ , say,  $\varphi_n \rightarrow \varphi$ . Then  $\square\varphi = \lim_n \square\varphi_n = \Phi$ .

Pf of claim:

Otherwise,  $\forall n \in \mathbb{N}$ ,  $\exists \varphi_n \in \Gamma(X, L)$ ,  $\varphi_n \perp \text{Ker } \square$  with  $\|\varphi_n\|_{L^2} \geq n \|\square\varphi_n\|_{L^2}$ . Define:

$$\psi_n \triangleq \varphi_n / \|\varphi_n\|_{L^2}. \quad \text{Then } \|\square\psi_n\|_{L^2} \leq \frac{1}{n}, \quad \|\psi_n\|_{L^2} = 1.$$

By a priori estimate,  $\|\psi_n - \psi_m\|_{L^2} \leq D \cdot (\|\square\psi_n - \square\psi_m\|_{L^2} + \|\psi_n - \psi_m\|_{L^1})$  for some  $D > 0$ . Rellich's lemma  $\Rightarrow \exists$  subsequence, which we may assume to be  $\{\psi_n\}$  to start with, s.t.  $\|\psi_n - \psi_m\|_{L^1} \rightarrow 0$  as  $n, m \rightarrow \infty \Rightarrow \psi_n \rightarrow \psi$  in  $H_{(2)}$ , with  $\|\psi\|_{L^2} = \lim \|\psi_n\|_{L^2} = 1$ , and  $\square\psi = \lim_n \square\psi_n = 0 \Rightarrow \psi \in \text{Ker } \square \subseteq H_{(2)}$ . On the other hand,  $\psi_n \in (\text{Ker } \square)^\perp \Rightarrow \psi \in (\text{Ker } \square)^\perp \subseteq H_{(2)}$ . It follows that  $\psi = 0$ , contradiction to  $\|\psi\|_{L^2} = 1$ .

Define Green's operator  $G: L^2 \rightarrow H_{(2)}$ ,  $\varphi \mapsto \psi$ .

Then by def., we have

$$\square G\varphi = \square\psi = \varphi - \varphi_0 = (\text{Id} - \Pi_{R(\square)^\perp})\varphi$$

(c). We can identify  $R(\square)^\perp = \text{Ker}(\square)$  (in the standard sense,  $\text{Ker } \square$  on  $C^\infty$ )  
Indeed,  $\forall f \in R(\square)^\perp$ , i.e.  $\langle f, \square\psi \rangle = 0$ ,  $\forall \psi \in H_{(2)}$ . In particular, take  $\psi \in C^\infty$ ,  $\Rightarrow \square f = 0$  in the sense of distributions. By elliptic regularity thm,  $f$  is smooth and  $\square f = 0$  in the usual sense. Thus  $f \in \text{Ker } \square$ .

Conversely, suppose  $f \in \text{Ker } \square$ , consider  $\psi \in H_{(2)}$ . Now  $f \in C^\infty \subseteq H_{(2)}$

$$\Rightarrow \langle f, \square\psi \rangle_{L^2} = \langle \square f, \psi \rangle_{L^2} = 0$$

$$\Rightarrow f \in R(\square)^\perp.$$

In summary,  $\forall \varphi \in L^2$ ,  $\exists! \psi \in H_{(2)}$ ,  $\varphi_0 \in \text{Ker } \square$  s.t.  $\varphi = \square\psi + \varphi_0$ , and we defined



$G\varphi = \psi$ ,  $\psi \perp \text{Ker } \square$ . Then:

$$\square G\varphi = \square \psi = \varphi - \varphi_0 = \varphi - \pi\varphi = (\text{Id} - \pi)\varphi.$$

where  $\pi: L^2 \rightarrow \text{Ker } \square$  is the orthogonal projection. By composing  $H_{(2)} \hookrightarrow L^2$ , we obtain  $G: L^2 \rightarrow L^2$ .

Green's operator

Observation: The a priori estimate implies the following estimate, (actually AP'):

$$\|G\varphi\|_{(2)} \leq C \cdot \|\varphi\|_{(0)} \quad (\text{AP}'')$$

Indeed, in the above notation,  $G\varphi = \psi$ , and  $\square\psi = \varphi - \varphi_0$ ,  $\psi \perp \text{Ker } \square$ . By AP'

$$\|G\varphi\|_{(2)} = \|\psi\|_{(2)} \leq C \|\square\psi\|_{(0)} = C \|\varphi - \varphi_0\|_{(0)} \leq C \|\varphi\|.$$

Def. An operator  $G: H \rightarrow H$ , where  $H$  is a Hilbert space, is said to be compact if:

(i).  $G$  is bounded:  $\|G\varphi\| \leq C \cdot \|\varphi\|$ ,  $\forall \varphi \in H$

(ii).  $\forall \{\varphi_j\}$  a bounded sequence in  $H$ ,  $\{G\varphi_j\}$  contains a convergent subsequence (pre-compactness).

Claim: The Green's operator  $G$  is compact.

Pf: Let  $\{\varphi_j\} \in L^2$ ,  $\|\varphi_j\|_{(0)} \leq 1$ . By the estimate (AP'')

$$\|G\varphi_j\|_{(2)} \leq C \cdot \|\varphi_j\|_{(0)} \leq C.$$

Rellich's lemma  $\Rightarrow \forall t < 2$ , say,  $t=0$ ,  $\{G\varphi_j\}$  contains a convergent subsequence w.r. t.  $\|\cdot\|_{(0)}$  norm.

Claim:  $G$  is self-adjoint and positive.

Pf:  $\forall \varphi, \tilde{\varphi} \in L^2$ , then  $\varphi = \square\psi + \varphi_0$ ,  $\tilde{\varphi} = \square\tilde{\psi} + \tilde{\varphi}_0$ , and  $\psi, \tilde{\psi} \in H_{(2)}$ ,  $\psi, \tilde{\psi} \perp \text{Ker } \square$  w.r. t. the  $L^2$ -norm. Now:

$$\langle G\varphi, \tilde{\varphi} \rangle_{L^2} = \langle \psi, \square\tilde{\psi} + \tilde{\varphi}_0 \rangle_{L^2} = \langle \psi, \square\tilde{\psi} \rangle_{L^2}$$

Similarly:  $\langle \varphi, G\tilde{\varphi} \rangle_{L^2} = \langle \square\psi + \varphi_0, \tilde{\psi} \rangle_{L^2} = \langle \square\psi, \tilde{\psi} \rangle_{L^2}$

Hence  $\langle G\varphi, \tilde{\varphi} \rangle_{L^2} = \langle \varphi, G\tilde{\varphi} \rangle_{L^2}$ . Taking  $\tilde{\varphi} = \varphi \Rightarrow \langle G\varphi, \varphi \rangle_{L^2} = \langle \psi, \square\psi \rangle_{L^2} \geq 0$ .

Claim:  $G\bar{\partial} = \bar{\partial}G$ ,  $G\bar{\partial}^\dagger = \bar{\partial}^\dagger G$

Pf: Since  $L^2 = \text{Ker } \square \oplus R(\square)$ , it suffices to check  $G\bar{\partial} = \bar{\partial}G$  on each factor.

Let  $\varphi \in \text{Ker } \square$ , i.e.  $(\bar{\partial}^\dagger \bar{\partial} + \bar{\partial} \bar{\partial}^\dagger)\varphi = 0 \iff \bar{\partial}^\dagger \varphi = 0, \bar{\partial} \varphi = 0$ . Then  $G\bar{\partial} \varphi = 0$ . On the other hand,  $\bar{\partial}G\varphi = 0$  since  $G\varphi = 0$ .

Next, let  $\varphi \in R(\square)^\perp$ ,  $\varphi = \square \psi$ . Then  $\bar{\partial}G\varphi = \bar{\partial}\psi$ . On the other hand,

$$\begin{aligned} G(\bar{\partial}\varphi) &= G(\bar{\partial}(\bar{\partial} \bar{\partial}^\dagger + \bar{\partial}^\dagger \bar{\partial})\psi) \\ &= G(\bar{\partial} \bar{\partial}^\dagger \bar{\partial} \psi) \\ &= G((\bar{\partial} \bar{\partial}^\dagger + \bar{\partial}^\dagger \bar{\partial}) \bar{\partial} \psi) \\ &= G \square \bar{\partial} \psi \\ &= (\text{Id} - \pi) \bar{\partial} \psi. \end{aligned}$$

Thus it suffices to show that  $\pi \bar{\partial} \psi = 0$ .  $\forall \eta \in C^\infty$ , we have

$$\langle \pi \bar{\partial} \psi, \eta \rangle = \langle \bar{\partial} \psi, \pi \eta \rangle = \langle \psi, \bar{\partial}^\dagger \pi \eta \rangle.$$

But since  $\pi \eta \in \text{Ker } \square \implies \bar{\partial}^\dagger \pi \eta = \bar{\partial} \pi \eta = 0 \implies \pi \bar{\partial} \psi = 0$ .

Taking adjoint of  $\bar{\partial}G = G\bar{\partial}$  gives  $G\bar{\partial}^\dagger = \bar{\partial}^\dagger G$ , since  $G$  is self-adjoint.

The above claims finish part b) of Hodge-Kobayashi thm.

A bit functional analysis: Spectrum of compact self-adjoint operators.

Lemma. Let  $G: H \rightarrow H$  be a compact, non-negative, self-adjoint operator on a Hilbert space  $H$ . Then:


(i). The upper bound  $\mu = \sup_{\|u\|=1} \langle Gu, u \rangle$  is an eigenvalue, i.e.  $\exists 0 \neq v \in H$  with  $Gv = \mu v$ . If  $\mu = 0$ ,  $G \equiv 0$ .

(ii). Each eigen-space  $\{v \mid Gu = \lambda v\}$  is finite dimensional.

(iii). The only accumulation points of the eigenvalues  $\lambda > 0$  of  $G$  is  $\lambda = 0$

(iv). The span of (i.e. the space of finite linear combinations of) eigenspaces with positive eigenvalues is dense in  $\text{Range}(G)$  (=  $R(G)$  for short).

(v).  $G$  admits an orthonormal basis of eigen-vectors.



Pf: Elementary observation: Let  $A$  be a self-adjoint, non-negative, bounded operator. Then

$$\|Au\| \leq \langle Au, u \rangle^{\frac{1}{2}} \|A\|^{\frac{1}{2}}$$

In fact, if we define an inner product  $[u, v] \triangleq \langle Au, v \rangle$ , which is non-negative, possibly degenerate inner product. Thus by Cauchy-Schwartz:

$$\begin{aligned} \|Au\|^2 &= \langle Au, Au \rangle = [u, Au] \\ &\leq [u, u]^{\frac{1}{2}} [Au, Au]^{\frac{1}{2}} \\ &= \langle Au, u \rangle^{\frac{1}{2}} \langle A(Au), Au \rangle^{\frac{1}{2}} \\ &\leq \langle Au, u \rangle^{\frac{1}{2}} \|AAu\|^{\frac{1}{2}} \|Au\|^{\frac{1}{2}} \\ &\leq \langle Au, u \rangle^{\frac{1}{2}} \|A\|^{\frac{1}{2}} \|Au\| \\ \Rightarrow \|Au\| &\leq \langle Au, u \rangle^{\frac{1}{2}} \|A\|^{\frac{1}{2}} \end{aligned}$$

(i). Let  $A \triangleq \mu \text{Id} - G$ , non-negative, self-adjoint and bounded. The above inequality:

$$\|(\mu \text{Id} - G)u\| \leq C \cdot \langle (\mu \text{Id} - G)u, u \rangle^{\frac{1}{2}}$$

Let  $\{u_j\}$  satisfy  $\langle Gu_j, u_j \rangle \rightarrow \mu$ , and  $\|u_j\| = 1$ . Thus

$$\|(\mu \text{Id} - G)u_j\| \rightarrow 0 \text{ when } j \rightarrow \infty.$$

$$\Rightarrow (\mu u_j - Gu_j) \rightarrow 0 \text{ as vectors in } H.$$

Since  $G$  is compact and  $\{u_j\}$  is bounded, by passing to a subsequence if necessary, we may assume that  $Gu_j \rightarrow v \in H$ .

If  $\mu \neq 0$ ,  $u_j = \frac{1}{\mu} Gu_j \rightarrow \frac{1}{\mu} v$  and  $Gu = \lim_j \mu Gu_j = \mu v$ .

If  $\mu = 0$ ,  $A \equiv 0 \Rightarrow G \equiv 0$  and the statement is trivially true.

(ii). Let  $\{u_j\}$  be an orthonormal basis of the  $\lambda$ -eigenspace. Since  $\lambda > 0$ ,  $u_j = \frac{1}{\lambda} Gu_j$ .

If the  $\lambda$ -eigenspace were infinite dimensional, since  $G$  is compact, by going to a subsequence if necessary, we may assume that  $Gu_j$  converges. This implies  $\lambda \sqrt{2} = \lambda \|u_j - u_k\| = \|Gu_j - Gu_k\| \rightarrow 0$  ( $j, k \rightarrow \infty$ ), contradiction.

(iii). Let  $\lambda_j > 0$  be eigenvalues and  $\lambda_j \rightarrow \lambda$ . If  $\lambda > 0$ , we may choose as above a sequence  $u_j$ ,  $Gu_j = \lambda_j u_j$  and  $\{Gu_j\}$  converges, i.e.  $\|Gu_j - Gu_{j+1}\| \rightarrow 0$  when

$j \rightarrow \infty$ . But

$$\|Gu_j - Gu_{j+1}\| = \|\lambda_j u_j - \lambda_{j+1} u_{j+1}\| \rightarrow \sqrt{2}\lambda, \quad j \rightarrow \infty$$

contradiction.

(iv). Define  $\lambda_1 = \sup_{\|u\|=1} \langle Gu, u \rangle$ , and assume that  $\lambda_1 > 0$  ( $G \neq 0$ , by (a)). Then  $\exists u_1$  eigenvector:  $Gu_1 = \lambda_1 u_1$ . Let  $H_1 = (\mathbb{C}\{u_1\})^\perp$ , then  $G|_{H_1}: H_1 \rightarrow H_1$ . Indeed,  $\forall v \in H_1, \langle Gu, u_1 \rangle = \langle v, Gu_1 \rangle = \langle v, \lambda_1 u_1 \rangle = 0$ . Define  $\lambda_2 = \sup_{\|u\|=1, u \in H_1} \langle Gu, u \rangle$ , and by a) again, we find  $u_2 \in H_1, Gu_2 = \lambda_2 u_2 \dots$

Inductively, at the  $k$ -th stage, we can find  $Gu_1 = \lambda_1 u_1, \dots, Gu_k = \lambda_k u_k$  and define  $H_k = \mathbb{C}\{u_1, \dots, u_k\}^\perp$ , then  $G|_{H_k}: H_k \rightarrow H_k$ .

Claim:  $\|G|_{H_k}\| \leq \lambda_{k+1}$ , where  $\lambda_{k+1} = \sup_{\|u\|=1, u \in H_k} \langle Gu, u \rangle$ .

Recall from elementary observation that:

$$\begin{aligned} \|G|_{H_k}\| &= \sup_{\|u\|=1, u \in H_k} \|Gu\| \leq \sup_{\|u\|=1, u \in H_k} \langle Gu, u \rangle^{\frac{1}{2}} \|G|_{H_k}\|^{\frac{1}{2}} \\ &= \lambda_{k+1}^{\frac{1}{2}} \|G|_{H_k}\|^{\frac{1}{2}} \\ \Rightarrow \|G|_{H_k}\| &\leq \lambda_{k+1} \end{aligned}$$

By (iii),  $\lambda_{k+1} \rightarrow 0$ . Let  $\pi_k$  be the orthogonal projection onto  $\mathbb{C}\{u_1, \dots, u_k\}$ . Then

$G\pi_k = \pi_k G: \forall u \in H$ .

$$\begin{aligned} G\pi_k u &= G\left(\sum_{j=1}^k \langle u_j, u \rangle u_j\right) \\ &= \sum_{j=1}^k \langle u_j, u \rangle \lambda_j u_j \\ &= \sum_{j=1}^k \langle Gu_j, u \rangle u_j \\ &= \sum_{j=1}^k \langle u_j, Gu \rangle u_j \\ &= \pi_k Gu. \end{aligned}$$

Now let  $u = Gu \in R(G)$ ,  $u - \pi_k u = Gu - \pi_k Gu = G(u - \pi_k u)$ . Note that

$$u - \pi_k u \in H_k \Rightarrow \|u - \pi_k u\| = \|G(u - \pi_k u)\| \leq \lambda_{k+1} \|u - \pi_k u\| \leq \lambda_{k+1} \|u\| \rightarrow 0 \text{ as } k \rightarrow \infty.$$

Thus  $\mathbb{C}\{u_1, \dots, u_k, \dots\}$  is dense in  $R(G)$ .

(v). Since  $\mathbb{C}\{u_1, \dots, u_k, \dots\}$  is dense in  $R(G) \Rightarrow$  the span is dense in  $\overline{R(G)}$ . Moreover, since  $G$  is bounded and self-adjoint,  $R(G)^\perp = \text{Ker } G$ . ( $\because \forall u, v \in H, \langle Gu, v \rangle = \langle u, Gv \rangle \Rightarrow Gv = 0$  iff  $v \perp$  all  $Gu$ 's). Now  $H = \overline{R(G)} \oplus \text{Ker } G$ . Take an

orthonormal basis for  $\text{Ker } G$  and adjoint it to the basis  $\{u_j\}$  obtained above, then we have an orthonormal basis of eigenvectors.  $\square$

Now it follows from our construction that  $L^2$  admits a basis of eigenvectors  $\{\varphi_\lambda\}$  of  $G$ ,  $G\varphi_\lambda = \lambda\varphi_\lambda$ . Next we shall prove the growth of these eigenvalues.

Growth of eigenvalues.

Let  $\{\varphi_\lambda(z)\}$  be an orthonormal basis of eigenfunctions of  $\square$ . Given  $\Lambda > 0$ , define the spectral function of  $\square$ :

$$e_\Lambda(\omega, z) \triangleq \sum_{\lambda \leq \Lambda} \varphi_\lambda(\omega) \overline{\varphi_\lambda(z)}.$$

More precisely,  $\varphi_\lambda(\omega) = \{\varphi_{\lambda \bar{j} i}^\alpha(\omega)\}$  and

$$e_\Lambda(\omega, z) = \frac{1}{p!q!} \sum_{\lambda \leq \Lambda} \varphi_{\lambda \bar{j} i}^\alpha(\omega) \overline{\varphi_{\lambda \bar{k} l}^\beta(z)} g^{i\bar{l}}(\omega) g^{k\bar{j}}(\omega) H_{\bar{\beta}\alpha}(\omega)$$

Note that  $\varphi_{\lambda \bar{k} l}^\beta(z) \in E_z \otimes \Lambda_z^{p,q}$ , and  $\varphi_{\lambda \bar{j} i}^\alpha g^{i\bar{l}} g^{k\bar{j}} H_{\bar{\beta}\alpha}(\omega) \in (E_\omega \otimes \Lambda_\omega^{p,q})^*$

$\Rightarrow e_\Lambda(\omega, z) \in (E_\omega \otimes \Lambda_\omega^{p,q})^* \otimes (E_z \otimes \Lambda_z^{p,q})$  and  $e_\Lambda(z, z)$  is a scalar.

We shall use the spectral function to study the growth of  $\lambda$ 's.

Observation 1:  $\int_X e_\Lambda(z, z) \frac{\omega^n}{n!} = \sum_{\lambda \leq \Lambda} \|\varphi_\lambda\|_{L^2}^2 = \#\{\lambda \leq \Lambda\}$

Observation 2: To show the growth  $\lambda_m \geq D \cdot m^S$ , it suffices to show that  $\exists C > 0$ , independent of  $z$ , so that:

$$|e_\Lambda(z, z)| \leq C \cdot \Lambda^p \quad (*).$$

Indeed, this would imply by integrating over  $M$ :

$$\#\{\lambda \leq \Lambda\} \leq D \cdot \Lambda^p \quad \text{for some } D.$$

Reorder  $\lambda$ 's so that  $\lambda_1 \leq \lambda_2 \leq \dots$ . Then the eigenvalues satisfy:

$$\lambda_{[D \cdot \Lambda^p + 1]} \geq \Lambda$$

Set  $N = [D \cdot \Lambda^p + 1] \leq D' \Lambda^p \Rightarrow \lambda_N \geq C \cdot N^{\frac{1}{p}}$ .

Proof of (\*).

Strategy: By the Sobolev lemma, the desired estimate should follow from estimates

for  $\|\cdot\|_{(s)}$ , where  $s > n$ ,  $n = \dim_{\mathbb{C}} X$ .

Recall the a priori estimate:

$$\|\varphi\|_{(s+2)} \leq C \cdot (\|\square\varphi\|_{(s)} + \|\varphi\|_{(s+1)})$$

which is equivalent to:

$$\|\varphi\|_{(s+2)}^2 \leq C' \cdot (\|\square\varphi\|_{(s)}^2 + \|\varphi\|_{(s+1)}^2)$$

We shall now derive a technical improvement of this estimate:

$$\|\varphi\|_{(s+2)}^2 \leq C \cdot (\|\square\varphi\|_{(s)}^2 + \|\varphi\|_{(s)}^2).$$

$C$  depending on  $s$  only. We do this, for instance, for  $s=0$ .

$$\text{AP: } \|\varphi\|_{(2)}^2 \leq C \cdot (\|\square\varphi\|_{(0)}^2 + \|\varphi\|_{(1)}^2)$$

It suffices to bound  $\|\varphi\|_{(1)}^2$  by  $\|\square\varphi\|_{(0)}^2$  and  $\|\varphi\|_{(0)}^2$ . But by def.

$$\|\varphi\|_{(1)}^2 = \|\varphi\|_{(0)}^2 + \int_X \nabla_j \varphi \overline{\nabla_{\bar{k}} \varphi} g^{j\bar{k}} \frac{\omega^n}{n!} + \dots$$

Using integration by parts, we have

$$\int_X \nabla_j \varphi \overline{\nabla_{\bar{k}} \varphi} g^{j\bar{k}} \frac{\omega^n}{n!} = - \int_X \varphi \overline{g^{\bar{k}j} \nabla_j \nabla_{\bar{k}} \varphi} \frac{\omega^n}{n!}$$

Note that by Bochner-Kodaira type formulas,  $-g^{\bar{k}j} \nabla_j \nabla_{\bar{k}}$  differs from  $\square$  only by tensor terms (with torsion occurring in non-Kähler cases), which are not differential operators (or diff. op. of order 0), and these terms can be bounded by  $C \cdot \|\varphi\|_{(0)}^2$ . Thus the result follows.

$$\text{Now from } \|\varphi\|_{(2)}^2 \leq C \cdot (\|\square\varphi\|_{(0)}^2 + \|\varphi\|_{(0)}^2)$$

$$\Rightarrow \|\varphi\|_{(4)}^2 \leq C \cdot (\|\square\varphi\|_{(2)}^2 + \|\varphi\|_{(2)}^2)$$

$$\leq C' \cdot (\|\square^2\varphi\|_{(0)}^2 + \|\square\varphi\|_{(0)}^2 + \|\varphi\|_{(0)}^2)$$

$\Rightarrow \dots$

$$\Rightarrow \|\varphi\|_{(2k)}^2 \leq C \left( \sum_{j=1}^k \|\square^j \varphi\|_{(0)}^2 + \|\varphi\|_{(0)}^2 \right) \text{ for any } \varphi \in C^\infty$$

Suppose  $\varphi \in H^\Lambda \cong \text{Span}\{u_1, \dots, u_\lambda\}$ ,  $\lambda \leq \Lambda$ . Then  $\|\square\varphi\| \leq \Lambda \|\varphi\|$ . Indeed, we have

$$\varphi = \sum_{\lambda \leq \Lambda} \langle u_\lambda, \varphi \rangle u_\lambda$$

$$\Rightarrow \square\varphi = \sum_{\lambda \leq \Lambda} \lambda \langle u_\lambda, \varphi \rangle u_\lambda$$

$$\Rightarrow \|\square\varphi\|^2 = \sum \lambda^2 |\langle u_\lambda, \varphi \rangle|^2 \leq \Lambda^2 \sum |\langle u_\lambda, \varphi \rangle|^2 = \Lambda^2 \|\varphi\|^2.$$

Now by induction, we have:

$$\|\square^l \varphi\|^2 \leq \Lambda^{2l} \|\varphi\|^2$$

Thus if  $\varphi \in H^\Lambda$ , then:

$$\|\varphi\|_{(2k)}^2 \leq C \cdot (\Lambda^{2k+1}) \|\varphi\|_{(0)}^2$$

or equivalently:

$$\|\varphi\|_{(2k)} \leq C_k (\Lambda^{k+1}) \|\varphi\|_{(0)}$$

Now by Sobolev's inequality, if  $2k > n$ , we have:

$$\|\varphi\|_{C^0} \leq \|\varphi\|_{(2k)} \leq C_k (\Lambda^{k+1}) \|\varphi\|_{(0)}$$

for any  $\varphi \in H^\Lambda$ . Now  $\forall f \in L^2$ , consider the orthogonal projection of  $f$  onto  $H^\Lambda$ :

$$\pi_\Lambda(f)(z) = \langle f(\omega), e_\Lambda(\omega, z) \rangle_{L^2},$$

integrating w.r. t.  $\omega$ . Since now  $\pi_\Lambda(f) \in H^\Lambda$ , this implies that:

$$\|\pi_\Lambda(f)\|_{C^0} \leq C_k (\Lambda^{k+1}) \|\pi_\Lambda(f)\|_{(0)} \leq C_k (\Lambda^{k+1}) \|f\|_{(0)}$$

In particular:

$$\begin{aligned} \|e_\Lambda(\cdot, z)\|_{L^2} &= \sup_{f \neq 0} \frac{|\langle f, e_\Lambda(\cdot, z) \rangle|}{\|f\|_{(0)}} \\ &\leq C_k (\Lambda^{k+1}) \quad (**) \end{aligned}$$

On the other hand, for a fixed  $z_0$ :

$$\|e_\Lambda(\cdot, z_0)\|_{L^2}^2 = \langle e_\Lambda(\cdot, z_0), e_\Lambda(\cdot, z_0) \rangle = e_\Lambda(z_0, z_0)$$

Thus (\*\*) implies that

$$|e_\Lambda(z_0, z_0)| \leq C_k^2 (\Lambda^{k+1})^2$$

for any  $z_0 \in X$ . Thus (\*) is proved.

Remaining steps

(1). A priori inequality:  $\forall \varphi \in C_0^\infty$ ,

$$\|\varphi\|_{(s+2)}^2 \leq C (\|\square \varphi\|_{(s)}^2 + \|\varphi\|_{(s+1)}^2)$$

(2). Prove the basic property of Sobolev spaces:

- Sobolev's lemma
- Rellich's lemma
- Equivalence between different definitions.

(3). Regularity thms.

Elliptic linear PDE's on  $\mathbb{R}^n$ .

Let  $\mathbb{R}^n = \{(x_1, \dots, x_n)\}$  and  $D^\alpha \triangleq \left(\frac{1}{i} \frac{\partial}{\partial x_1}\right)^{\alpha_1} \dots \left(\frac{1}{i} \frac{\partial}{\partial x_n}\right)^{\alpha_n}$ , where  $\alpha = (\alpha_1, \dots, \alpha_n) \in \mathbb{N}^n$ , and  $|\alpha| = \sum \alpha_i$ . In this notation, the formal Taylor series of a function  $u(x)$  is:

$$u(x) = \sum_{\alpha} \frac{1}{\alpha!} D^\alpha u(0) (ix)^\alpha$$

where  $\alpha! \triangleq \alpha_1! \dots \alpha_n!$  and  $x^\alpha = x_1^{\alpha_1} \dots x_n^{\alpha_n}$ .

Def: (1)  $L$  is a linear partial differential operator of order  $m$  if

$$(Lu)(x) = \sum_{|\alpha| \leq m} a_\alpha(x) D^\alpha u(x)$$

(2). The symbol  $\sigma_L(x, \xi)$  of  $L$  is the following function

$$\sigma_L(x, \xi) = \sum_{|\alpha|=m} a_\alpha(x) \xi^\alpha$$

for any  $x \in \mathbb{R}^n$ ,  $\xi \in \mathbb{R}^n$ .

(3).  $L$  is said to be elliptic at  $x_0$  if  $|\sigma_L(x_0, \xi)| \geq C \cdot |\xi|^m$ . It's elliptic if it's elliptic at all points.

Rmk:  $|\sigma_L(x_0, \xi)| \geq C \cdot |\xi|^m$  is equivalent to the requirement that  $\sigma_L(x_0, \xi) \neq 0$ ,  $\forall \xi \neq 0$ . Indeed " $\Rightarrow$ " is easy to see. Conversely,  $|\sigma_L(x, \cdot)|$  achieves minimal value  $C$  on the (compact) unit sphere. Thus  $\forall \xi$ .

$$\begin{aligned} |\sigma_L(x, \xi)| &= |\sigma_L(x, \frac{\xi}{|\xi|})| |\xi|^m && (\sigma_L(x, \xi) \text{ homogeneous of degree } m) \\ &= |\xi|^m |\sigma_L(x, \frac{\xi}{|\xi|})| \\ &\geq C \cdot |\xi|^m \end{aligned}$$

E.g.  $\square$  is elliptic of order 2. Recall from Bochner-Kodaira formula:

$$\begin{aligned} \square \varphi_{\bar{j}1}^\alpha &= -g^{j\bar{k}} \nabla_{\bar{j}} \nabla_{\bar{k}} \varphi_{\bar{j}1}^\alpha + \text{Curvature terms} \\ &= -g^{j\bar{k}} (\partial_{\bar{j}} + \Gamma_{\bar{j}}) (\partial_{\bar{k}} + \Gamma_{\bar{k}}) \varphi_{\bar{j}1}^\alpha \\ &= -g^{j\bar{k}} \partial_{\bar{j}} \partial_{\bar{k}} \varphi_{\bar{j}1}^\alpha + \text{lower degree terms.} \end{aligned}$$

Thus the symbol of  $\square$  is given by:

$$\sigma_\square(z, \xi) = -g^{j\bar{k}}(z) \xi_{\bar{j}} \xi_{\bar{k}} = -|\xi|^2$$

for any  $\xi \in \Lambda^{1,0}$ . Thus  $\square$  is elliptic of order 2. Actually the symbol of  $\square$  is a well-defined function on  $\Lambda^1 M$ .



Sobolev spaces in  $\mathbb{R}^n$ .

Def: (1),  $\forall \varphi \in C_0^\infty(\mathbb{R}^n)$ ,  $\|\varphi\|_{(k)}^2 \triangleq \sum_{|\alpha| \leq k} \|D^\alpha \varphi\|^2$

(2),  $H(k)(\mathbb{R}^n) \triangleq \overline{C_0^\infty(\mathbb{R}^n)}$  w.r.t. the norm  $\|\cdot\|_{(k)}$ .

Basic properties about Fourier transform.

Let  $\mathcal{S}$  be the space of Schwartz functions. i.e.  $\mathcal{S} \triangleq \{\varphi \in C^\infty(\mathbb{R}^n) \text{ s.t. } \forall \alpha, N, (1+|x|^2)^N |D^\alpha \varphi(x)| \leq C_{N,\alpha} \text{ for all } x \in \mathbb{R}^n\}$  (rapidly decaying condition.) The space contains  $C_0^\infty(\mathbb{R}^n)$  and functions like  $e^{-\frac{|x|^2}{2}}$  etc.

Define the Fourier transform:

$$\varphi(x) \longrightarrow \hat{\varphi}(\xi) \triangleq \frac{1}{(2\pi)^n} \int_{\mathbb{R}^n} e^{-i\langle x, \xi \rangle} \varphi(x) dx$$

Then:

(1),  $\forall \varphi \in \mathcal{S}$ ,  $\hat{\varphi} \in \mathcal{S}$ . The inverse is given by Fourier inversion formula.

$$\varphi(x) = \int_{\mathbb{R}^n} e^{i\langle x, \xi \rangle} \hat{\varphi}(\xi) d\xi.$$

(2), Plancherel formula:  $\|\varphi\|_{L^2} = \|\hat{\varphi}\|_{L^2}$ .

(3),  $(D^\alpha \varphi)^\wedge(\xi) = \xi^\alpha \hat{\varphi}(\xi)$ .

These properties imply that there is a simple characterization of Sobolev norms:

$\forall \varphi \in \mathcal{S}(\mathbb{R}^n)$ :

$$\begin{aligned} \|\varphi\|_{(k)}^2 &= \sum_{|\alpha| \leq k} \|D^\alpha \varphi\|_{L^2}^2 \\ &= \sum_{|\alpha| \leq k} \|(D^\alpha \varphi)^\wedge\|_{L^2}^2 \\ &= \sum_{|\alpha| \leq k} \|\xi^\alpha \hat{\varphi}(\xi)\|_{L^2}^2 \\ &= \sum_{|\alpha| \leq k} \int_{\mathbb{R}^n} |\xi^\alpha|^2 |\hat{\varphi}(\xi)|^2 d\xi. \end{aligned}$$

Observe that  $\exists C, C > 0$  constants s.t.

$$C(1+|\xi|^2)^k \leq \sum_{|\alpha| \leq k} |\xi^\alpha|^2 \leq C \cdot (1+|\xi|^2)^k.$$

Hence:

$$C \int_{\mathbb{R}^n} (1+|\xi|^2)^k |\hat{\varphi}(\xi)|^2 d\xi \leq \|\varphi\|_{(k)}^2 \leq C \cdot \int_{\mathbb{R}^n} (1+|\xi|^2)^k |\hat{\varphi}(\xi)|^2 d\xi$$

and the norm defined by  $\int_{\mathbb{R}^n} (1+|\xi|^2)^k |\hat{\varphi}(\xi)|^2 d\xi$  is equivalent to  $\|\cdot\|_{(k)}$ .

Proof of the a priori estimate:

Step 1. Let  $L_m = \sum_{|\alpha|=m} C_\alpha D^\alpha$  be an elliptic differential operator, homogeneous of degree  $m$  and with constant coefficients  $C_\alpha$ . Then  $\forall \varphi \in C_0^\infty(\mathbb{R}^n)$ :

$$\|\varphi\|_{C^{m+s}}^2 \leq C \cdot (\|L_m \varphi\|_{C^s}^2 + \|\varphi\|_{C^{m+s-1}}^2)$$

Pf: We just show  $s=0$ .  $m>0$  is similar.

$$\begin{aligned} \|L_m \varphi\|_{C^0}^2 &= \int_{\mathbb{R}^n} |(L_m \varphi)^\wedge(\xi)|^2 d\xi \\ &= \int_{\mathbb{R}^n} |\sum_{|\alpha| \leq m} C_\alpha \xi^\alpha \hat{\varphi}(\xi)|^2 d\xi \\ &= \int_{\mathbb{R}^n} |\sigma_{L_m}(\xi) \hat{\varphi}(\xi)|^2 d\xi \\ &= \int_{\mathbb{R}^n} |\sigma_{L_m}(\xi)|^2 |\hat{\varphi}(\xi)|^2 d\xi \end{aligned}$$

Now by def.,  $|\sigma_{L_m}(\xi)|^2 \geq C |\xi|^{2m}$ :

$$\begin{aligned} \Rightarrow \|L_m \varphi\|_{C^0}^2 &\geq C \cdot \int_{\mathbb{R}^n} |\xi|^{2m} |\hat{\varphi}(\xi)|^2 d\xi \\ \Rightarrow \|L_m \varphi\|_{C^0}^2 + \|\varphi\|_{C^0}^2 &\geq C \cdot \int_{\mathbb{R}^n} (|\xi|^{2m} + 1) |\hat{\varphi}(\xi)|^2 d\xi \\ &\geq C \cdot \|\varphi\|_{C^m}^2. \end{aligned}$$

Step 2. Let  $z_0$  be any given point in  $\mathbb{R}^n$ ,  $L$  any elliptic linear differential operator of order  $m$ . Then  $\exists V_{z_0} \ni z_0$ ,  $\bar{V}_{z_0}$  compact, s.t.

$$\|\varphi\|_{C^m} \leq C (\|L\varphi\|_{C^0} + \|\varphi\|_{C^{m-1}})$$

for any  $\varphi \in C_0^\infty(V_{z_0})$ .

Pf: Define  $L_m$  by  $L_m \varphi = \sum_{|\alpha|=m} a_\alpha(z_0) D^\alpha \varphi(z)$ . Then

$$\|L\varphi\|_{C^0} \geq \|L_m \varphi\|_{C^0} - \|(L - L_m)\varphi\|_{C^0}$$

$$\text{and } \|(L - L_m)\varphi\|_{C^0} = \sum_{|\alpha|=m} \|(a_\alpha(z) - a_\alpha(z_0)) D^\alpha \varphi\|_{C^0} + \sum_{|\alpha| \leq m-1} \|a_\alpha(z) D^\alpha \varphi\|_{C^0}$$

Claim:  $\forall \varepsilon > 0$ ,  $\exists V_{x_0} \ni x_0$  open nhd,  $\bar{V}_{x_0}$  compact so that  $\forall \varphi \in C_0^\infty(V_{x_0})$

$$\|(L - L_m)\varphi\|_{C^0} \leq \varepsilon \cdot \|\varphi\|_{C^m} + C' \|\varphi\|_{C^{m-1}}$$

Then assuming this claim,  $\forall \varphi \in C_0^\infty(V_{x_0})$ ,

$$\begin{aligned} \|\varphi\|_{C^m} &\leq C \cdot (\|L_m \varphi\|_{C^0} + \|\varphi\|_{C^{m-1}}) \\ &\leq C \cdot (\|(L - L_m)\varphi\|_{C^0} + \|L\varphi\|_{C^0} + \|\varphi\|_{C^{m-1}}) \\ &\leq C \cdot \varepsilon \|\varphi\|_{C^m} + C \cdot C' \|\varphi\|_{C^{m-1}} + C \|L\varphi\|_{C^0} + C \|\varphi\|_{C^{m-1}} \end{aligned}$$

Taking  $C \cdot \varepsilon = \frac{1}{2}$ , we obtain:

$$\begin{aligned} \|\varphi\|_{C^m} &\leq 2C \|L\varphi\|_{C^0} + 2C(C'+1) \|\varphi\|_{C^{m-1}} \\ &\leq C'' (\|L\varphi\|_{C^0} + \|\varphi\|_{C^{m-1}}). \end{aligned}$$

Now we check the claim:

$$(L - L_m)\varphi = \sum_{|\alpha| \leq m-1} a_\alpha(x) D^\alpha \varphi + \sum_{|\alpha|=m} (a_\alpha(x) - a_\alpha(x_0)) D^\alpha \varphi$$

For  $|\alpha| \leq m-1$  terms,

$$\|a_\alpha(x) D^\alpha \varphi\|_{(0)}^2 = \int_{\mathbb{R}^n} |a_\alpha(x) D^\alpha \varphi(x)|^2 dx$$

Thus on any open subset of  $\mathbb{R}^n$  with compact closure,  $|a_\alpha(x)|$  is bounded, and

$$\|a_\alpha(x) D^\alpha \varphi\|_{(0)}^2 \leq C \cdot \|\varphi\|_{(m-1)}^2$$

Now choose  $V_{x_0}$  so that  $\sup_{x \in V_{x_0}} |a_\alpha(x) - a_\alpha(x_0)| < \varepsilon$ . Then  $\forall \varphi \in C_0^\infty(V_{x_0})$ ,  $|\alpha|=m$  we have:

$$\begin{aligned} \|(a_\alpha(x) - a_\alpha(x_0)) D^\alpha \varphi\|_{(0)}^2 &= \int_{\mathbb{R}^n} |(a_\alpha(x) - a_\alpha(x_0)) D^\alpha \varphi|^2 dx \\ &= \int_{V_{x_0}} |(a_\alpha(x) - a_\alpha(x_0)) D^\alpha \varphi|^2 dx \\ &\leq \varepsilon \int_{V_{x_0}} |D^\alpha \varphi|^2 dx \\ &= \varepsilon \int_{\mathbb{R}^n} |D^\alpha \varphi|^2 dx \\ &\leq \varepsilon \|\varphi\|_{(m)}^2 \end{aligned}$$

Now, to prove  $\|\varphi\|_{(m+s)} \leq C(\|L\varphi\|_{(s)} + \|\varphi\|_{(s+m-1)})$ , we apply similar techniques as above. Observe that  $\forall s \in \mathbb{N}$ , we have:

$$\|(L - L_m)\varphi\|_{(s)} \leq \left\| \sum_{|\alpha| \leq m-1} a_\alpha(x) D^\alpha \varphi \right\|_{(s)} + \left\| \sum_{|\alpha|=m} (a_\alpha(x) - a_\alpha(x_0)) D^\alpha \varphi \right\|_{(s)}$$

For  $|\alpha| \leq m-1$  terms:

$$\|a_\alpha(x) D^\alpha \varphi\|_{(s)}^2 = \sum_{|\beta| \leq s} \|D^\beta (a_\alpha(x) D^\alpha \varphi)\|_{(0)}^2$$

and fix each  $\beta$ ,

$$D^\beta \left( \sum_{|\alpha| \leq m-1} a_\alpha(x) D^\alpha \varphi \right) = \sum_{|\gamma| \leq m+s-1} b_\gamma(x) D^\gamma \varphi$$

Thus

$$\left\| \sum_{|\alpha| \leq m-1} a_\alpha(x) D^\alpha \varphi \right\|_{(s)} \leq C \cdot \|\varphi\|_{(m+s-1)}$$

For  $|\alpha|=m$  terms,

$$\begin{aligned} \|(a_\alpha(x) - a_\alpha(x_0)) D^\alpha \varphi\|_{(s)}^2 &= \sum_{|\beta| \leq s} \|D^\beta (a_\alpha(x) - a_\alpha(x_0)) D^\alpha \varphi\|_{(0)}^2 \\ &= \sum_{|\beta| \leq s} \|(a_\alpha(x) - a_\alpha(x_0)) D^{\beta+\alpha} \varphi + \sum_{\gamma \leq \beta+|\alpha|-1} b_\gamma(x) D^\gamma \varphi\|_{(0)}^2 \\ &\leq \sum_{|\beta| \leq s} \|(a_\alpha(x) - a_\alpha(x_0)) D^{\beta+\alpha} \varphi\|_{(0)}^2 + \sum_{|\beta| \leq s} \left\| \sum_{\gamma \leq s+m-1} b_\gamma D^\gamma \varphi \right\|_{(0)}^2 \\ &\leq \varepsilon \cdot \|\varphi\|_{(m+s)}^2 + C \|\varphi\|_{(m+s-1)}^2. \end{aligned}$$

Rmk: Even for  $s$  non-integers or negative numbers, we may define the Sobolev norm by:

$$\|\varphi\|_{(s)}^2 \triangleq \int_{\mathbb{R}^n} (1+|\xi|^2)^s |\hat{\varphi}(\xi)|^2 d\xi$$

The same a priori estimate holds for all  $s \in \mathbb{R}$ . The proof requires some more difficult arguments.

Step 3. Fix a compact set  $K \subseteq \mathbb{R}^n$ ,  $K \neq \emptyset$ , we prove the AP estimate for any  $\varphi \in C_0^\infty(K)$ .

Clearly,  $K \subseteq \bigcup_{w \in K} V_w$ , where  $V_w$  is constructed as in step 2. Extract a finite subcover,  $K \subseteq \bigcup_{j=1}^N V_{x_j}$ , and take a partition of unity subordinate to  $V_{x_j}$ :

$$0 \leq \rho_j \in C_0^\infty(V_{x_j}), \quad \sum_{j=1}^N \rho_j \equiv 1.$$

Then:  $\forall \varphi \in C_0^\infty(K)$ ,  $\varphi = \sum_{j=1}^N \rho_j \varphi$  with  $\rho_j \varphi \in C_0^\infty(V_{x_j})$ ,

$$\begin{aligned} \|\varphi\|_{(m)} &\leq \sum_{j=1}^N \|\rho_j \varphi\|_{(m)} \\ &\leq C \cdot \sum_{j=1}^N (\|L(\rho_j \varphi)\|_{(0)} + \|\rho_j \varphi\|_{(m-1)}) \end{aligned}$$

Observation 1:  $\forall s \in \mathbb{N}$ ,  $\|\rho_j \varphi\|_{(s)} \leq C \cdot \|\varphi\|_{(s)}$ :

$$\begin{aligned} \|\rho \varphi\|_{(s)}^2 &= \sum_{|\alpha| \leq s} \int |D^\alpha(\rho \varphi)|^2 dx \\ &= \sum_{|\alpha| \leq s} \int |b_\alpha D^\alpha \varphi|^2 dx \\ &\leq \sum_{|\alpha| \leq s} C \cdot \int |D^\alpha \varphi|^2 dx \\ &\leq C \cdot \|\varphi\|_{(s)}^2 \end{aligned}$$

Observation 2:  $\|L(\rho_j \varphi) - \rho_j L\varphi\|_{(0)} \leq C \cdot \|\varphi\|_{(m-1)}$ :

$$\begin{aligned} \text{In fact } L(\rho \varphi) - \rho L\varphi &= \sum_\alpha a_\alpha D^\alpha(\rho \varphi) - \sum \rho a_\alpha D^\alpha \varphi \\ &= \sum_{\beta+\gamma=\alpha} a_\alpha D^\beta \rho D^\gamma \varphi - \sum \rho a_\alpha D^\alpha \varphi \\ &= \sum_{\beta+\gamma=\alpha, |\gamma| < |\alpha| \leq m} a_\alpha D^\beta \rho D^\gamma \varphi \\ \Rightarrow \|L(\rho \varphi) - \rho L\varphi\|_{(0)} &\leq C \cdot \|\varphi\|_{(m-1)} \end{aligned}$$

Hence

$$\begin{aligned} \|\varphi\|_{(m)} &\leq C \cdot \sum_{j=1}^N (\|L(\rho_j \varphi)\|_{(0)} + \|\rho_j \varphi\|_{(m-1)}) \\ &\leq C \cdot \sum_{j=1}^N (\|\rho_j L\varphi\|_{(0)} + \|L(\rho_j \varphi) - \rho_j L\varphi\|_{(0)} + C' \|\varphi\|_{(m-1)}) \\ &\leq C \cdot \sum_{j=1}^N (\rho_j \|L\varphi\|_{(0)} + C \cdot C'' \|\varphi\|_{(m-1)} + C \cdot C' \|\varphi\|_{(m-1)}) \\ &\leq C (\|L\varphi\|_{(0)} + \|\varphi\|_{(m-1)}) \end{aligned}$$

□

Rmk: This step is also true for all  $s \in \mathbb{R}$ , by using some integral inequality.

Proof of Sobolev lemma:

If  $s > \frac{n}{2}$ , then  $\forall \varphi \in C_0^\infty(\mathbb{R}^n)$

$$\|\varphi\|_{C^0} \leq C \cdot \|\varphi\|_{(s)}$$

Pf: By the Fourier inversion formula:

$$\varphi(x) = \int e^{i\langle x, \xi \rangle} \hat{\varphi}(\xi) d\xi$$

and thus

$$\begin{aligned} |\varphi(x)| &\leq \int |\hat{\varphi}(\xi)| d\xi \\ &\leq \left( \int (1+|\xi|^2)^s |\hat{\varphi}(\xi)|^2 d\xi \right)^{\frac{1}{2}} \cdot \left( \int (1+|\xi|^2)^{-s} d\xi \right)^{\frac{1}{2}} \quad (\text{Cauchy-Schwartz}) \end{aligned}$$

Now if  $s > \frac{n}{2}$ ,  $\int_{\mathbb{R}^n} (1+|\xi|^2)^{-s} d\xi < \infty$  and thus

$$|\varphi(x)| \leq C \cdot \|\varphi\|_{(s)}$$

$$\Rightarrow \|\varphi\|_{C^0} \leq C \cdot \|\varphi\|_{(s)}. \quad \square$$

Similarly, we can show that  $\forall \varphi \in C_0^\infty(\mathbb{R}^n)$ , if  $s > \frac{n}{2} + k$ .

$$\|\varphi\|_{C^k} \leq C \cdot \|\varphi\|_{(s)}$$

In fact,  $\forall k \in \mathbb{N}$ ,  $|\alpha| = k$ , and  $s > \frac{n}{2} + k$ :


$$\begin{aligned} |D^\alpha \varphi(x)| &\leq \int |\xi^\alpha| |\hat{\varphi}(\xi)| d\xi \\ &\leq \int (1+|\xi|^2)^{\frac{|\alpha|}{2}} |\hat{\varphi}(\xi)| d\xi \\ &\leq \left( \int (1+|\xi|^2)^s |\hat{\varphi}(\xi)|^2 d\xi \right)^{\frac{1}{2}} \left( \int (1+|\xi|^2)^{-s+|\alpha|} d\xi \right)^{\frac{1}{2}} \\ &\leq C \cdot \|\varphi\|_{(s)} \end{aligned}$$

and the result follows.

Proof of Rellich's lemma

Recall that if  $0 \leq s < t$ , and  $\{\varphi_j \in H^{(t)} \mid \|\varphi_j\|_{(t)} \leq 1, \text{supp } \varphi_j \subseteq K \subseteq \mathbb{R}^n\}$ . Then Rellich's lemma  $\Rightarrow \exists$  subsequence  $\{\varphi_{j_\epsilon}\}$  converging w.r.t.  $\|\cdot\|_{(s)}$ .

The lemma is clearly wrong without the assumption that  $K$  is a fixed compact subset of  $\mathbb{R}^n$ . For instance:



The translated bump functions  $\|\varphi_j\|_{L^1} \leq C$ , but has no convergent subsequence at all.

$$\begin{aligned}
 \text{Pf: } \|\varphi_j - \varphi_k\|_{L^1}^2 &= \int_{\mathbb{R}^n} (1+|\xi|^2)^s |\hat{\varphi}_j(\xi) - \hat{\varphi}_k(\xi)|^2 d\xi \\
 &= \int_{\mathbb{R}^n} (1+|\xi|^2)^{-t-s} (1+|\xi|^2)^t |\hat{\varphi}_j(\xi) - \hat{\varphi}_k(\xi)|^2 d\xi \\
 &= \int_{|\xi| \leq R} + \int_{|\xi| > R}
 \end{aligned}$$

To extract a convergent subsequence, we need the above two integrals to converge. Now  $\forall \xi \in \mathbb{R}^n$ ,

$$\hat{\varphi}(\xi) = \int e^{-i\langle x, \xi \rangle} \varphi(x) dx$$

We will show that  $\{\hat{\varphi}_j(\xi)\}$  is an equicontinuous family, and then the second integral will converge once  $R \gg 0$ , and the first integral will converge by passing to a uniformly convergent subsequence on the compact set  $\{|\xi| \leq R\}$ .

Step 1:  $\{\hat{\varphi}_j(\xi)\}$  is equi-continuous.

(a).  $\{\hat{\varphi}_j(\xi)\}$  is uniformly bounded.

$$\begin{aligned}
 |\hat{\varphi}_j(\xi)| &\leq \int_{\mathbb{R}^n} |\varphi_j(x)| dx = \int_K |\varphi_j(x)| dx \\
 &\leq \left( \int_K |\varphi_j(x)|^2 dx \right)^{\frac{1}{2}} \cdot \left( \int_K 1 \cdot dx \right)^{\frac{1}{2}} \\
 &\leq \text{Vol}(K)^{\frac{1}{2}} \cdot \|\varphi_j\|_{L^1} \\
 &\leq C
 \end{aligned}$$

(b).  $\left\{ \frac{\partial}{\partial \xi_j} \hat{\varphi}_j(\xi) \right\}$  is uniformly bounded.

$$\begin{aligned}
 \left| \frac{\partial}{\partial \xi_j} \hat{\varphi}_j(\xi) \right| &\leq \int_{\mathbb{R}^n} |ix_j e^{-i\langle x, \xi \rangle} \varphi_j(x)| dx \\
 &= \int_K |x_j| |\varphi_j(x)| dx \\
 &\leq \left( \int_K |\varphi_j(x)|^2 dx \right)^{\frac{1}{2}} \left( \int_K |x_j|^2 dx \right)^{\frac{1}{2}} \\
 &\leq C \cdot \|\varphi_j\|_{L^1} \\
 &\leq C
 \end{aligned}$$

This implies equi-continuity: By the mean value thm:

$$|\hat{\Phi}_j(\xi) - \hat{\Phi}_j(\eta)| \leq \max_{\xi, \eta} |\nabla \hat{\Phi}_j| \cdot |\xi - \eta| \\ \leq C \cdot |\xi - \eta|$$

Hence by Arzela-Ascoli thm,  $\forall$  compact subset of  $\mathbb{R}^n$ ,  $\exists$  subsequence  $\{\hat{\Phi}_{j_k}(\xi)\}$  converging uniformly on this compact set.

Step 2.  $\forall \varepsilon > 0$ ,  $\exists R \gg 0$  s.t.  $(1+|R|^2)^{-ct-s} \leq \frac{\varepsilon}{8}$ . By step 1, we have a subsequence  $\{\varphi_{j_k}\}$  s.t.  $\{\hat{\Phi}_{j_k}(\xi)\}$  converges uniformly on  $\{|\xi| \leq R\}$ . Now

$$\int_{|\xi| > R} (1+|\xi|^2)^{-ct-s} (1+|\xi|^2)^t |\hat{\Phi}_{j_k}(\xi) - \hat{\Phi}_{j_l}(\xi)|^2 d\xi \\ \leq 2 \int_{|\xi| > R} \frac{\varepsilon}{4} (1+|\xi|^2)^t (|\hat{\Phi}_{j_k}(\xi)|^2 + |\hat{\Phi}_{j_l}(\xi)|^2) d\xi \\ \leq \frac{\varepsilon}{4} \int_{\mathbb{R}^n} (1+|\xi|^2)^t (|\hat{\Phi}_{j_k}(\xi)|^2 + |\hat{\Phi}_{j_l}(\xi)|^2) d\xi \\ = \frac{\varepsilon}{4} (\|\varphi_{j_k}\|_{ct+s}^2 + \|\varphi_{j_l}\|_{ct+s}^2) \\ \leq \frac{\varepsilon}{2}$$

Next, by equi-continuity, we may choose  $N \in \mathbb{N} \gg 0$  so that whenever  $l, k \geq N$ ,

$$\sup_{|\xi| \leq R} |\hat{\Phi}_{j_k}(\xi) - \hat{\Phi}_{j_l}(\xi)|^2 \leq (1+R^2)^{-s} \cdot (\text{vol}(\{|\xi| \leq R\}))^{-1} \cdot \frac{\varepsilon}{2}$$

Then

$$\int_{|\xi| \leq R} (1+|\xi|^2)^s |\hat{\Phi}_{j_k}(\xi) - \hat{\Phi}_{j_l}(\xi)|^2 d\xi \leq \int_{|\xi| \leq R} (1+|\xi|^2)^s \sup_{|\xi| \leq R} |\hat{\Phi}_{j_k}(\xi) - \hat{\Phi}_{j_l}(\xi)|^2 d\xi \\ \leq \int_{|\xi| \leq R} (1+R^2)^s \cdot (1+R^2)^{-s} (\text{vol})^{-1} \frac{\varepsilon}{2} d\xi \\ \leq \frac{\varepsilon}{2}$$

Summing up, we obtain the desired:

$$\|\varphi_{j_k} - \varphi_{j_l}\|^2 \leq \varepsilon \quad (l, k \geq N).$$

□

Reducing compact manifolds to  $\mathbb{R}^n$ .


Recall that on a compact manifold  $X$ ,  $s \in \mathbb{N}$ ,

$$\|\varphi\|_{(s)} \triangleq \sum_{|\alpha| \leq s} \int |\nabla^\alpha \varphi|^2 \sqrt{g} dx$$

We may also introduce another norm as follows. Write  $X = \bigcup_{\mu=1}^N X_\mu$ , where  $X_\mu$  corresponds to the unit ball in  $\mathbb{R}^n$ , and  $\{\rho_\mu\}$  a partition of unity subordinate to this cover. Define:

$$\|\varphi\|_{(s)} \triangleq \sum_{\mu=1}^N \|\rho_\mu \varphi\|_{(s, \mathbb{R}^n)}^2$$

where the r.h.s. uses the usual Sobolev norm on  $\mathbb{R}^n$ .



Claim: These two norms are equivalent, i.e.  $\exists c, C > 0$  s.t.

$$c \cdot \|\varphi\|_{(s)} \leq \|\varphi\|_{(s)} \leq C \cdot \|\varphi\|_{(s)}$$

This follows from the next two simple observations:

(1). For each  $X_\mu$ , we have:  $\forall \varphi \in C_0^\infty(X_\mu)$

$$\|\varphi\|_{(s)} \sim \|\varphi\|_{(s, \mathbb{R}^n)}$$

(2). For either norm, we always have,

$$\forall a \in C^\infty(X) \quad \|\mathbf{a}\varphi\|_{(s)} \leq C_a \|\varphi\|_{(s)}$$

$$\forall b \in C_0^\infty(\mathbb{R}^n) \quad \|b\varphi\|_{(s, \mathbb{R}^n)} \leq C_b \|\varphi\|_{(s, \mathbb{R}^n)}$$

(2) is easy. We check (1):

$$(s=0): \quad \|\varphi\|_{(0)}^2 = \int_X |\varphi|^2 \sqrt{g} dx = \int_{\Phi_\mu^{-1}(X_\mu)} |\varphi \circ \Phi_\mu(y)|^2 \sqrt{g(y)} dy$$

$$\|\varphi\|_{(0, \mathbb{R}^n)}^2 = \int_{\Phi_\mu^{-1}(X_\mu)} |\varphi \circ \Phi_\mu(y)|^2 dy$$

Note that  $g = \det g_{ij} > 0$ , and on the compact set  $\Phi_\mu^{-1}(X_\mu) \subseteq \mathbb{R}^n$  is bounded from below and above. Hence  $\|\varphi\|_{(0)}^2 \sim \|\varphi\|_{(0, \mathbb{R}^n)}^2$

$$(s=1): \quad \|\varphi\|_{(1)}^2 = \|\varphi\|_{(0)}^2 + \int_X g^{ij} \nabla_i \varphi \nabla_j \varphi \sqrt{g} dx$$

$$\|\varphi\|_{(1, \mathbb{R}^n)}^2 = \|\varphi\|_{(0, \mathbb{R}^n)}^2 + \int_{\mathbb{R}^n} \sum_i |\partial_i (\varphi \circ \Phi_\mu(y))|^2 dy$$

From simple linear algebra, we know that if  $h_{ij}, \tilde{h}_{ij}$  are two positive definite bilinear forms on  $\mathbb{R}^n$ , then they are equivalent:  $\lambda h(u, u) \leq \tilde{h}(u, u) \leq \Lambda h(u, u), \forall u \in \mathbb{R}^n$ .

( $\lambda, \Lambda$  can be taken to be the minimal and maximal eigenvalues of the self-adjoint operator  $h^{ip} \tilde{h}_{pj}$ ). Now over a compact set  $\Phi_\mu^{-1}(X_\mu)$ , take  $h_{ij} = g^{ij}, \tilde{h}_{ij} = \delta^{ij}$ , then

$\exists c, C$  s.t.  $0 < c \leq \lambda(x) \leq \Lambda(x) \leq C, \forall x \in \Phi_\mu^{-1}(X_\mu)$ , hence

$$c \int_{\mathbb{R}^n} g^{ij} \nabla_i \varphi \nabla_j \varphi \sqrt{g} dx \leq \int_{\mathbb{R}^n} \sum_i |\nabla_i \varphi|^2 dx \leq C \cdot \int_{\mathbb{R}^n} g^{ij} \nabla_i \varphi \nabla_j \varphi \sqrt{g} dx$$

Now note that  $\nabla_i \varphi = \partial_i \varphi + A_i \varphi \Rightarrow |\nabla_i \varphi|^2 \leq |\partial_i \varphi|^2 + |A_i \varphi|^2$ , but  $|A_i|$  is

bounded on the compact set  $\Phi_\mu^{-1}(X_\mu)$ . Similarly  $|\partial_i \varphi|^2 \leq |\nabla_i \varphi|^2 + |A_i \varphi|^2$ . Thus



by adding  $\|\varphi\|_{L^2(\Omega)}$ ,  $\|\varphi\|_{L^2(\mathbb{R}^n)}$ , these norms are again equal.

The general case follows from similar discussions. Now we can prove the equivalence of  $\|\cdot\|_{L^2(\Omega)}$  and  $\|\cdot\|_{L^2(\mathbb{R}^n)}$ .

By triangle inequality,

$$\|\varphi\|_{L^2(\Omega)} \leq \sum_{\mu=1}^N \|\rho_\mu \varphi\|_{L^2(\mathbb{R}^n)} \leq C \cdot \sum_{\mu=1}^N \|\rho_\mu \varphi\|_{L^2(\mathbb{R}^n)}$$

On the other hand, each term  $\mu=1, \dots, N$

$$\|\rho_\mu \varphi \circ \Phi_\mu(y)\|_{L^2(\mathbb{R}^n)} \leq C' \|\rho_\mu \varphi\|_{L^2(\Omega)} \leq C'_\mu \|\varphi\|_{L^2(\Omega)}$$

Summing up, the claim follows.

Using this equivalence of norms, we may transform a priori estimate, Rellich's lemma, Sobolev lemma from  $\mathbb{R}^n$  to compact manifolds. For instance, we prove a priori estimate for compact manifolds.

$\forall \varphi \in C^\infty(X)$ , write  $\varphi = \sum_\mu \rho_\mu \varphi$ . Now

$$\begin{aligned} \|\varphi\|_{L^2(\Omega)} &\leq \sum_\mu \|\rho_\mu \varphi\|_{L^2(\mathbb{R}^n)} \\ &\leq C \cdot \sum_\mu \|(\rho_\mu \varphi) \circ \Phi_\mu\|_{L^2(\mathbb{R}^n)} \\ &\leq C' \sum_\mu (\|L((\rho_\mu \varphi) \circ \Phi_\mu)\|_{L^2(\mathbb{R}^n)} + \|(\rho_\mu \varphi) \circ \Phi_\mu\|_{L^2(\mathbb{R}^n)}) \end{aligned}$$

Observation:  $(\rho_\mu \varphi) \circ \Phi_\mu = (\rho_\mu \circ \Phi_\mu) \cdot (\varphi \circ \Phi_\mu)$  and  $\rho_\mu \circ \Phi_\mu \in C^\infty(\mathbb{R}^n)$ . Then:

$$\begin{aligned} \|L((\rho_\mu \varphi) \circ \Phi_\mu)\|_{L^2(\mathbb{R}^n)} &\leq \|(\rho_\mu \circ \Phi_\mu) \circ L(\varphi \circ \Phi_\mu)\|_{L^2(\mathbb{R}^n)} \\ &\quad + \|\sum_{|\nu| \leq m-1} (\hat{\rho}_\mu \circ \Phi_\mu) \cdot D^\nu \varphi\|_{L^2(\mathbb{R}^n)} \\ &\leq C \cdot (\|(\rho_\mu \circ \Phi_\mu) \circ L(\varphi \circ \Phi_\mu)\|_{L^2(\mathbb{R}^n)} \\ &\quad + \|(\rho_\mu \circ \Phi_\mu)\|_{L^2(\mathbb{R}^n)}). \end{aligned}$$

where  $\hat{\rho}_\mu$  are order  $\geq 1$  derivatives of  $\rho_\mu$ , coming from derivatives of  $L$  by Leibnitz rule. (we may also require  $|\partial^\nu \rho_\mu| \leq \rho_\mu$ ) Thus:

$$\begin{aligned} \|\varphi\|_{L^2(\Omega)} &\leq C' \sum_\mu (\|\rho_\mu \circ \Phi_\mu \cdot L(\varphi \circ \Phi_\mu)\|_{L^2(\mathbb{R}^n)} + \|(\rho_\mu \circ \Phi_\mu)\|_{L^2(\mathbb{R}^n)}) \\ &\leq C' (\|L\varphi\|_{L^2(\Omega)} + \|\varphi\|_{L^2(\Omega)}) \end{aligned}$$

The transformation of Rellich's lemma and Sobolev lemma from  $\mathbb{R}^n$  to  $X$  is similar. □

Elliptic regularity

Basic technique: Method of difference quotients.

Assume  $u \in H^{(s)}(\mathbb{R}^n)$ .  $\forall h \neq 0$ , let

$$\Delta_h^j u \triangleq \frac{1}{h} (u(x_1, \dots, x_i+h, \dots, x_n) - u(x_1, \dots, x_n))$$

Then :  $\sup_h (\sum_{i=1}^n \|\Delta_h^j u\|_{(s)}) < \infty \iff u \in H^{(s+1)}(\mathbb{R}^n)$

Pf: " $\Leftarrow$ "  $\|\Delta_h^j u\|_{(s)}^2 = \int (1+|\xi|^2)^s |(\Delta_h^j u)^\wedge(\xi)|^2 d\xi$

We compute:  $(\Delta_h^j u)^\wedge(\xi) = \int_{\mathbb{R}^n} e^{-i\langle x, \xi \rangle} \frac{1}{h} (u(x_1+h_1, \dots, x_n) - u(x_1, \dots, x_n)) dx$   
 $= \frac{1}{h} (e^{ih\xi_1} - 1) \hat{u}(\xi)$

Note that  $|\frac{1}{h}(e^{ih\xi_1} - 1)| = |\xi_1| \int_0^1 e^{it\xi_1} dt| \leq |\xi_1|$

$$\Rightarrow \|\Delta_h^j u\|_{(s)}^2 \leq \int (1+|\xi|^2)^s |\xi_1|^2 |\hat{u}(\xi)|^2 d\xi$$

$$\leq \int (1+|\xi|^2)^{s+1} |\hat{u}(\xi)|^2 d\xi$$

$$= \|u\|_{(s+1)}^2$$

" $\Rightarrow$ "  $\int (1+|\xi|^2)^s |\xi_1|^2 |\hat{u}(\xi)|^2 d\xi = \int \lim_{h \rightarrow 0} (1+|\xi|^2)^s |\frac{e^{ih\xi_1} - 1}{h}|^2 |\hat{u}(\xi)|^2 d\xi$

By Fatou's lemma ( $f_j \geq 0$ , then  $\int \liminf f_j d\mu \leq \liminf \int f_j d\mu$ )

$$\int \lim_{h \rightarrow 0} (1+|\xi|^2)^s |\frac{e^{ih\xi_1} - 1}{h}|^2 |\hat{u}(\xi)|^2 d\xi \leq \lim_{h \rightarrow 0} \|\Delta_h^j u\|_{(s)}^2 \leq \text{const. independent of } h. \quad \square$$

Manipulations with  $\Delta_h^j$ .

(1).  $\Delta_h^j (\frac{\partial u}{\partial x_k}) = \frac{\partial}{\partial x_k} (\Delta_h^j u)$

Indeed,  $\Delta_h (\frac{\partial u}{\partial x_k}) = \frac{1}{h} (\frac{\partial u}{\partial x_k}(x+h) - \frac{\partial u}{\partial x_k}(x)) = \frac{\partial}{\partial x_k} (\frac{1}{h} (u(x+h) - u(x)))$ .

(2). Let  $a \in C_0^\infty(\mathbb{R}^n)$ , then:

$$\|\Delta_h^j (au) - a \Delta_h^j u\|_{(s)} \leq C_a \|u\|_{(s)}$$

where  $C_a$  is a constant independent of  $u$  and  $h$ .

Note that, since  $\Delta_h$  behaves like a differential, we trivially have

$$\|\Delta_h (au)\|_{(s)} \leq C_a \|u\|_{(s+1)}.$$

Pf:  $\Delta_h (au)(x) = \frac{1}{h} (a(x+h)u(x+h) - a(x)u(x))$   
 $= \frac{1}{h} (a(x+h)u(x+h) - a(x)u(x+h) + a(x)u(x+h) - a(x)u(x))$   
 $= u(x+h) \Delta_h (a)(x) + a(x) \Delta_h (u)(x)$

$$\Rightarrow \Delta_h^j (au) - a \Delta_h^j u = \Delta_h (a) u(x+h).$$

Observation 1:  $\|u(x+h)\|_{(s)} = \|u(x)\|_{(s)}$

Observation 2:  $\Delta_h (a)$  is smooth, and is bounded independently of  $h$ .

In fact: 
$$\begin{aligned} a(x_1+h, x_2, \dots) - a(x) &= \int_0^1 \frac{d}{dt} a(x_1+th, x_2, \dots) dt \\ &= \int_0^1 \frac{\partial a}{\partial x_1}(x_1+th, x_2, \dots) h dt \\ &= h \int_0^1 \frac{\partial a}{\partial x_1}(x_1+th, x_2, \dots) dt \end{aligned}$$

$$\Rightarrow \Delta_h(a) = \int_0^1 \frac{\partial a}{\partial x_1}(x_1+th, x_2, \dots) dt$$

$$\Rightarrow \|\Delta_h(a)\|_{C^s} \leq \int_0^1 \left\| \frac{\partial a}{\partial x_1} \right\|_{C^s} dt = Ca$$

The result follows.

(3). More generally, if  $L = \sum a_\alpha(x) D^\alpha$ ,  $a_\alpha(x) \in C^\infty(\mathbb{R}^n)$ , similar proof shows that:

$$\|\Delta_h(Lu) - L(\Delta_h u)\|_{C^s} \leq C \cdot \|u\|_{C^{s+m}}$$

Note that our elliptic operator doesn't satisfy  $a_\alpha(x) \in C^\infty(\mathbb{R}^n)$ , but we will introduce a cut off trick to deal with this.

Proof of regularity thm.

Note that if  $u \in H^{s_1}(\mathbb{R}^n)$ , we have two ways of defining  $Lu$ . One way is to regard  $L: H^{s_1} \rightarrow H^{s_1-m}$ , and  $Lu$  is its image in  $H^{s_1-m}$ . i.e.  $\exists u_j \in C^\infty(\mathbb{R}^n)$ ,  $u_j \rightarrow u$  w.r.t.  $\|\cdot\|_{C^s}$ , and  $Lu = \lim Lu_j$  w.r.t.  $\|\cdot\|_{C^{s-m}}$ .

The second way is in the sense of distributions. We say that  $Lu = g$  for some function  $g$  if

$$\int_{\mathbb{R}^n} u \left( \sum_{|\alpha| \leq m} (-1)^{|\alpha|} D^\alpha \right) (a_\alpha \cdot \varphi) = \int_{\mathbb{R}^n} g \varphi,$$

$\forall \varphi \in C^\infty(\Omega)$ . In case  $u \in C^\infty(\Omega)$ , the l.h.s. becomes  $\int_{\mathbb{R}^n} \left( \sum_{|\alpha| \leq m} a_\alpha D^\alpha u \right) \varphi$ , by integration by parts.

Observation: If  $Lu = f$  in the first sense and  $Lu = g$  in the second sense, then  $f = g$ .

Indeed, take  $u_j \rightarrow u$  in  $H^{s_1}$ ,  $u_j \in C^\infty(\mathbb{R}^n)$ . Denote the operator  $\sum_{|\alpha| \leq m} (-1)^{|\alpha|} D^\alpha (a_\alpha \cdot)$  by  $L^t$ , we have,  $\forall \varphi \in C^\infty(\mathbb{R}^n)$ ,

$$\begin{aligned} \left| \int_{\mathbb{R}^n} (u_j - u_k) L^t \varphi dx \right| &= \left| \int_{\mathbb{R}^n} (u_j - u_k) \hat{\cdot} (L^t \varphi) \hat{\cdot} d\mathbb{B}_1 \right| \\ &\leq \left\| \int_{\mathbb{R}^n} |u_j - u_k| \hat{\cdot} (1 + |\mathbb{B}_1|^2)^s d\mathbb{B}_1 \right\| \left\| \int_{\mathbb{R}^n} |L^t \varphi| \hat{\cdot} (1 + |\mathbb{B}_1|^2)^{-s} d\mathbb{B}_1 \right\| \\ &= \|u_j - u_k\|_{C^s} \|L^t \varphi\|_{C^{-s}} \rightarrow 0 \quad (j, k \rightarrow \infty), \end{aligned}$$

since  $L^t \varphi$  is compactly supported and thus  $\|L^t \varphi\|_{C^{-s}} < \infty$ . Letting  $j \rightarrow \infty$ , and

integrating by parts for  $u_j$ , we have:

$$|\int_{\mathbb{R}^n} f\varphi - \int_{\mathbb{R}^n} u_k L^t \varphi| \rightarrow 0 \text{ as } k \rightarrow \infty,$$

which means  $\lim_{k \rightarrow \infty} u_k = f$  in the sense of distributions.

Next, we introduce local Sobolev space: Given  $\Omega \subseteq \mathbb{R}^n$ ,  $H_{(s)}^{loc}(\Omega) \triangleq \{\varphi \mid \forall \chi \in C_0^\infty(\Omega), \chi\varphi \in H_{(s)}(\mathbb{R}^n)\}$ .

Observation:  $\bigcap_s H_{(s)}^{loc}(\Omega) = C^\infty(\Omega)$ .

Claim:  $Lu = f$  in  $\Omega$ ,  $f \in C^\infty(\Omega)$ , and  $u \in H_{(s+m)}^{loc}(\Omega)$ , then  $u \in H_{(s+m+1)}^{loc}(\Omega)$ .


Then by the above observation,  $u \in C^\infty(\Omega)$ .

Recall the AP estimate,  $\forall K \subseteq \Omega$  compact subset ( $K \neq \emptyset$ ),  $\forall \varphi \in C_0^\infty(K)$

$$\|\varphi\|_{(s+m)} \leq C(\|L\varphi\|_{(s)} + \|\varphi\|_{(s+m-1)})$$

Let  $H_{(s+m)}^{comp}(K) \triangleq \overline{C_0^\infty(K)}$  w/r.t.  $\|\cdot\|_{(s+m)}$  norm, then  $H_{(s+m)}^{comp}(K) \subseteq H_{(s+m)}^{loc}(\Omega)$ .

Want:  $\forall \chi \in C_0^\infty(\Omega)$ ,  $\chi u \in H_{(s+1)}(\mathbb{R}^n)$ . Since  $\text{Supp } \chi \subseteq \Omega$ , we may pick inclusion of open neighborhoods:



We wish to show that  $\sup_h \|\Delta_h(\chi u)\|_{(s+m)} < \infty$  as in difference quotient.

Now we have:

$$\|\Delta_h(\chi u)\|_{(s+m)} \leq C(\|L(\Delta_h(\chi u))\|_{(s)} + \|\Delta_h(\chi u)\|_{(s+m-1)})$$

But


$$\|L(\Delta_h(\chi u)) - \Delta_h(L(\chi u))\|_{(s)} \leq C \cdot \|\chi u\|_{(s+m)} < \infty$$

and

$$\begin{aligned} \|\Delta_h(L\chi u) - \Delta_h(\chi Lu)\|_{(s)} &\leq \|L\chi u - \chi Lu\|_{(s+1)} \\ &\leq C \cdot \|u\|_{(s+m)} < \infty \end{aligned}$$

□

A small technical remark: In using difference quotient, we need  $L$  to have its coefficients compactly supported. This could not happen for elliptic operators! Thus, we need to pick another  $\hat{\chi} \in C_0^\infty(\Omega)$ ,  $\hat{\chi} \equiv 1$  on  $\text{supp } \chi$  to do a cut-off



and we may so pick  $\hat{X}$  that the support of  $\hat{X}L$  after small  $h$  shifts still remain inside  $K$ , and on  $\text{supp } X$  we can apply our AP estimate.

## §8. Geometry of Subbundles

Def: Let  $E \rightarrow X$  be a holomorphic vector bundle. A subbundle  $E' \rightarrow X$  of  $E$  is holomorphic vector bundle s.t.  $E'_z$  is a subspace of  $E_z$ .  $\forall z \in X$ .

In terms of transition functions,  $X = \cup X_\mu$ ,  $E \leftrightarrow \{t_{\mu\nu}(z)\}$ ,  $E' \leftrightarrow \{t'_{\mu\nu}(z)\}$ , then being a subbundle means:

$$t_{\mu\nu}(z) = \begin{pmatrix} t'_{\mu\nu}(z) & b_{\mu\nu} \\ 0 & t''_{\mu\nu} \end{pmatrix}$$

where  $t'_{\mu\nu}(z)$  is an  $r' \times r'$  matrix,  $t''_{\mu\nu}$  an  $(r-r') \times (r-r')$  matrix,  $b_{\mu\nu}$  an  $r' \times (r-r')$  matrix.

Given  $E \rightarrow X$  a holomorphic vector bundle,  $\{H_{\bar{\alpha}\beta}\}$  a metric on  $E$ , then we have a unitary Chern connection on  $E$ :

$$\begin{cases} \nabla_j \varphi^\alpha = \partial_j \varphi^\alpha \\ \nabla_{\bar{j}} \varphi^\alpha = H^{\alpha\bar{\gamma}} \partial_{\bar{j}} (H_{\bar{\gamma}\beta} \varphi^\beta) \end{cases}$$

Let  $E'$  be a subbundle (with  $\text{rank } E' = r' < r = \text{rank } E$ ), we have two ways of constructing a connection on  $E'$ : (In the following, let  $\mu \in \{1, \dots, n, \bar{1}, \dots, \bar{n}\}$ )

(1).  $\{H_{\bar{\alpha}\beta}\}$  restricts to a metric on  $E'$ , with the requirement to be a Chern connection, it defines a unique connection.

(2).  $\forall \varphi \in \Gamma(X, E') \subseteq \Gamma(X, E)$ , we can first take  $\nabla_\mu \varphi \in \Gamma(X, E)$ , and take the orthogonal projection from  $E$  to  $E'$ :

$$\nabla'_\mu \varphi = \pi'(\nabla_\mu \varphi)$$

Note that for any scalar function,

$$\begin{aligned} \nabla'_\mu (f\varphi) &= \pi'(\partial_\mu(f)\varphi) + f \nabla_\mu \varphi \\ &= \partial_\mu(f)\varphi + f \nabla'_\mu \varphi \end{aligned}$$

and thus defines a connection on  $E'$ . This agrees with the first construction since  $\forall \varphi, \varphi_1, \varphi_2 \in \Gamma(X, E')$ :

$$\begin{aligned} \nabla'_j \varphi &= \pi'(\partial_j \varphi) = \partial_j \varphi \quad (\text{since } E' \text{ is holomorphic}) \\ \partial_\mu \langle \varphi_1, \varphi_2 \rangle &= \langle \nabla_\mu \varphi_1, \varphi_2 \rangle + \langle \varphi_1, \nabla_{\bar{\mu}} \varphi_2 \rangle \\ &= \langle \pi' \nabla_\mu \varphi_1, \varphi_2 \rangle + \langle \varphi_1, \pi' \nabla_{\bar{\mu}} \varphi_2 \rangle \end{aligned}$$

$$= \langle \nabla'_\mu \varphi_1, \varphi_2 \rangle + \langle \varphi_1, \nabla'_\mu \varphi_2 \rangle$$

Def.  $\forall \varphi \in \Gamma(X, E')$ ,  $B_\mu \varphi \triangleq \nabla_\mu \varphi - \nabla'_\mu \varphi = \pi''(\nabla_\mu \varphi)$ , where  $\pi''$  is the projection onto the orthogonal complement of  $E''$  of  $E'$ . (As real bundles  $E \cong E' \oplus E''$ ).

Observation :

$$\begin{aligned} B_\mu(f\varphi) &= \nabla_\mu(f\varphi) - \nabla'_\mu(f\varphi) \\ &= \partial_\mu(f) \cdot \varphi + f \cdot \nabla_\mu \varphi - \partial_\mu(f) \cdot \varphi - f \cdot \nabla'_\mu \varphi \\ &= f(\nabla_\mu \varphi - \nabla'_\mu \varphi) \\ &= f \cdot B_\mu \varphi. \end{aligned}$$

i.e.  $B_\mu \in \Gamma(X, \text{Hom}(E', E''))$ . Equivalently, if  $B = B_\mu dz^\mu$ , then  $B \in \Gamma(X, \Lambda^1 \otimes \text{Hom}(E', E''))$ .  
Moreover,  $B_j \varphi = \partial_j \varphi - \partial_j \varphi = 0 \Rightarrow B \in \Gamma(X, \Lambda^{1,0} \otimes \text{Hom}(E', E''))$ .  $B$  is called the second fundamental form of  $E'$

Connection on  $E''$ .

$E''$  is a smooth subbundle of  $E$ , but it's not necessarily holomorphic. We can define a connection on  $E''$  by,  $\forall \psi \in \Gamma(X, E'')$ :

$$\nabla_\mu \psi = \pi''(\nabla_\mu \psi)$$

Again this is a unitary connection, but in general, it's not a Chern connection since  $E''$  is not necessarily holomorphic. Also set:

$$C_\mu \psi \triangleq \nabla_\mu \psi - \nabla'_\mu \psi$$

Claim:  $B_\mu^\dagger = -C_{\bar{\mu}}$ .

Pf:  $\forall \varphi \in \Gamma(X, E')$ ,  $\psi \in \Gamma(X, E'')$ ,  $\langle \varphi, \psi \rangle = 0$

$$\begin{aligned} \Rightarrow 0 &= \partial_\mu \langle \varphi, \psi \rangle = \langle \nabla_\mu \varphi, \psi \rangle + \langle \varphi, \nabla_\mu \psi \rangle \\ &= \langle \pi'' \nabla_\mu \varphi, \psi \rangle + \langle \varphi, \pi' \nabla_\mu \psi \rangle \\ &= \langle B_\mu \varphi, \psi \rangle + \langle \varphi, C_{\bar{\mu}} \psi \rangle \end{aligned}$$

Hence  $C \triangleq C_\mu dz^\mu \in \Gamma(X, \Lambda^{0,1} \otimes \text{Hom}(E'', E'))$ . It also follows that  $B \equiv 0 \Leftrightarrow C \equiv 0$ , which happens only when  $E''$  is holomorphic as well.

Curvatures of  $E, E', E''$ .

Let  $\Phi \in \Gamma(X, E)$ , write  $\Phi = \varphi + \psi$ , with  $\varphi \in \Gamma(X, E')$ ,  $\psi \in \Gamma(X, E'')$ . Then

$$\begin{aligned} \nabla_{\mu} \Phi &= \nabla_{\mu}(\varphi + \psi) \\ &= \nabla_{\mu}' \varphi + B_{\mu} \varphi + \nabla_{\mu}'' \psi + C_{\mu} \psi \\ &= \begin{pmatrix} \nabla_{\mu}' & C_{\mu} \\ B_{\mu} & \nabla_{\mu}'' \end{pmatrix} \begin{pmatrix} \varphi \\ \psi \end{pmatrix} \end{aligned}$$

Write

$$d_{\nabla} \Phi = \begin{pmatrix} d_{\nabla}' & C \\ B & d_{\nabla}'' \end{pmatrix} \begin{pmatrix} \varphi \\ \psi \end{pmatrix},$$

then the curvature:

$$\begin{aligned} d_{\nabla}^2 \Phi &= F \Phi \\ &= \begin{pmatrix} d_{\nabla}'^2 + C \wedge B & d_{\nabla}' C + C d_{\nabla}'' \\ B d_{\nabla}' + d_{\nabla}'' B & d_{\nabla}''^2 + B \wedge C \end{pmatrix} \begin{pmatrix} \varphi \\ \psi \end{pmatrix} \end{aligned}$$

Explicitly, recall that  $F = F_{\bar{k}j} dz^j \wedge d\bar{z}^k$ ,  $B = B_j dz^j$  and  $C = C_{\bar{k}} d\bar{z}^k$ . In our notation,  $(C \wedge B)_{\bar{k}j} = C_{\bar{k}} d\bar{z}^k \wedge B_j dz^j = -C_{\bar{k}} B_j dz^j \wedge d\bar{z}^k$ .

$$\Rightarrow F_{\bar{k}j} |_{\text{Hom}(E', E')} = F'_{\bar{k}j} - C_{\bar{k}} B_j = F'_{\bar{k}j} + B_{\bar{k}}^{\dagger} B_j.$$

Thus

$$\begin{aligned} \langle F_{\bar{k}j} \varphi, \varphi \rangle &= \langle F'_{\bar{k}j} \varphi, \varphi \rangle + \langle B_{\bar{k}}^{\dagger} B_j \varphi, \varphi \rangle \\ &= \langle F'_{\bar{k}j} \varphi, \varphi \rangle + \langle B_j \varphi, B_{\bar{k}} \varphi \rangle \end{aligned}$$

Since  $\langle B_j \varphi, B_{\bar{k}} \varphi \rangle$  is always positive definite,  $\{F_{\bar{k}j}\}$  is more "positive" than  $\{F'_{\bar{k}j}\}$ . The equation  $d_{\nabla}^2 \Phi$  can be cleaned up to be:

$$F = \begin{pmatrix} F' + B^{\dagger} \wedge B & -d_{\nabla} B^{\dagger} \\ d_{\nabla} B & F'' + B \wedge B^{\dagger} \end{pmatrix}$$

Necessary condition for Hermitian-Einstein metrics.

Let  $E \rightarrow X$  be a holomorphic vector bundle on  $(X, \omega)$  a compact Kähler manifold.  $\omega = \frac{i}{2} g_{\bar{k}j} dz^j \wedge d\bar{z}^k$  the Kähler form.

**Question:** When does there exist  $H_{\bar{\alpha}\beta}$  on  $E$ , so that  $\Lambda F = \mu \text{Id}$ , where



$\Lambda$  is the Hodge operator and  $\mu$  is a constant. i.e

$$(g^{j\bar{k}} F_{\bar{k}j})^\alpha{}_\beta = \mu \delta^\alpha{}_\beta \quad (H-E).$$

As  $F_{\bar{k}j} = -\partial_{\bar{k}}(H^{-1}\partial_j H)$ , this equation is non-linear in  $H$ .

Def: (Degree of  $E$ ). Given  $E \rightarrow (X, \omega)$ ,

$$\deg E \triangleq \frac{i}{2} \int_X \text{Tr}(F) \wedge \frac{\omega^{n-1}}{(n-1)!}$$

Note that if  $F = F_{\bar{k}j}^\alpha{}_\beta dz^j \wedge d\bar{z}^k$ , then  $\text{Tr}(F) = F_{\bar{k}j}^\alpha{}_\alpha dz^j \wedge d\bar{z}^k$ . It's a closed form whose cohomology class is  $C_1(E)$ . Thus the degree doesn't depend on the metric  $\{H_{\bar{k}j}\}$  or  $\{g_{\bar{k}j}\}$  but only  $E$  and  $[\omega]$ .

Def. (Slope of  $E$ ).  $\text{Slope}(E) \triangleq \deg E / \text{rank} E$

Key observation (Kobayashi, Lübke)

Thm. If  $E \rightarrow (X, \omega)$  admits a Hermitian-Einstein metric, then  $\forall E'$  which is a holomorphic subbundle of  $E$ ,

$$\text{slope}(E') \leq \text{slope}(E).$$

Equality holds iff  $E = E' \oplus E''$  with  $E''$  holomorphic.

Def. A holomorphic vector bundle  $E$  is said to be stable in the sense of Mumford-Takemoto if  $\forall E'$  holomorphic subbundle,  $\text{slope}(E') < \text{slope}(E)$ .

Pf of thm.

Let  $E' \subseteq E$  be a holomorphic subbundle. If a Hermitian-Einstein metric exists, we can use it to compute slope  $E$ .

$$\deg E = \frac{i}{2} \int_X (\text{Tr} F) \wedge \frac{\omega^{n-1}}{(n-1)!}$$

where  $F = F_{\bar{k}j}^\alpha{}_\beta dz^j \wedge d\bar{z}^k$ . To calculate  $\text{Tr} F \wedge \frac{\omega^{n-1}}{(n-1)!}$ , we may do it pointwise.

Write  $\omega = \frac{i}{2} (dz_1 \wedge d\bar{z}_1 + \dots + dz_n \wedge d\bar{z}_n)$ , then

$$\frac{\omega^{n-1}}{(n-1)!} = \left(\frac{i}{2}\right)^{n-1} \left( \sum_{j=1}^n dz_1 \wedge d\bar{z}_1 \wedge \dots \wedge \widehat{(dz_j \wedge d\bar{z}_j)} \wedge \dots \wedge dz_n \wedge d\bar{z}_n \right)$$

and thus:

$$\begin{aligned} F \wedge \frac{\omega^{n-1}}{(n-1)!} &= \sum_{j=1}^n F_{j\bar{j}} \alpha_{\beta} dz_j \wedge d\bar{z}_{\bar{j}} \wedge dz_1 \wedge d\bar{z}_1 \wedge \dots \wedge \widehat{(dz_j \wedge d\bar{z}_{\bar{j}})} \wedge \dots \wedge dz_n \wedge d\bar{z}_n \\ &= g^{j\bar{k}} F_{\bar{k}j} \alpha_{\beta} \frac{\omega^n}{n!} \end{aligned}$$

(Generally for any 2-form valued tensor  $T: T \wedge \frac{\omega^{n-1}}{(n-1)!} = g^{j\bar{k}} T_{\bar{k}j} \frac{\omega^n}{n!}$ ).

Hence:

$$\begin{aligned} \text{Tr}(F) \wedge \frac{\omega^{n-1}}{(n-1)!} &= \text{Tr}(\Lambda F) \frac{\omega^n}{n!} \\ &= \text{Tr}(\mu \text{Id}) \frac{\omega^n}{n!} \\ &= \mu \cdot \text{rank} E \frac{\omega^n}{n!} \end{aligned}$$

and

$$\int_X \text{Tr}(F) \wedge \frac{\omega^{n-1}}{(n-1)!} = \mu \cdot \text{rank} E \cdot \text{vol}(X)$$

On the other hand, recall that we have:

$$\begin{aligned} \Rightarrow \quad F_{\bar{k}j}|_{E'} &= F'_{\bar{k}j} + B_{\bar{k}}^{\dagger} B_j \\ \text{deg } E' &= \int_X \text{Tr}(F_{\bar{k}j}|_{E'} - B_{\bar{k}}^{\dagger} B_j) \frac{\omega^{n-1}}{(n-1)!} \\ &= \int_X \text{Tr}(g^{j\bar{k}} (F_{\bar{k}j}|_{E'} - B_{\bar{k}}^{\dagger} B_j)) \frac{\omega^n}{n!} \\ &= \int_X \text{Tr}(\Lambda F|_{E'}) \frac{\omega^n}{n!} - \int_X \text{Tr}(g^{j\bar{k}} \bar{B}_{\bar{k}} B_j) \frac{\omega^n}{n!} \\ &= \mu \cdot \text{rank } E' \cdot \text{vol}(X) - \|B\|_{L^2}^2 \\ &\leq \mu \cdot \text{rank } E' \cdot \text{vol}(X) \end{aligned}$$

$$\Rightarrow \quad \text{slope}(E') \leq \mu \cdot \text{vol}(X) = \text{slope}(E)$$

Note that "=" holds iff  $B \equiv 0$ , i.e. iff  $E$  splits as direct sum of holomorphic vector bundles:  $E \cong E' \oplus E''$ .  $\square$

Another example of a global aspect in canonical metrics: The Yang-Mills equation.

Recall that from electromagnetism, we are interested in:

- The functional:  $A = A_{\mu} dx^{\mu} \mapsto I(A) \triangleq \int_{\mathbb{R}^{3,1}} |F_{\mu\nu}|^2$ ,  $F \triangleq dA$ .
- The critical points of  $I$ , i.e. those connections  $A_{\mu} dx^{\mu}$  satisfying  $\frac{\delta I}{\delta A_{\mu}} = 0$

We are interested in the following generalizations.

$E \rightarrow (X, \omega)$ : holomorphic vector bundle over a compact Kähler manifold.

$A = A_j dz^j$ : a connection on  $E$ . Consider the Yang-Mills functional:

$$I(A) \triangleq \int_X g^{l\bar{k}} g^{j\bar{m}} F_{\bar{k}j}^\alpha \overline{F_{lm}^\gamma} H_{\bar{p}\alpha} H^{\beta\bar{s}} \frac{\omega^n}{n!}$$

Introduce the Hodge  $*$  operator,  $*$ :  $\Lambda^{p,q} \rightarrow \Lambda^{n-p, n-q}$ , with the following defining property:  $\varphi \wedge * \bar{\psi} = \langle \varphi, \bar{\psi} \rangle \frac{\omega^n}{n!}$ . In this notation, we have:

$$I(A) = \int_X \text{Tr}(F \wedge * F)$$

In case  $\dim X = 2$ ,  $*$ :  $\Lambda^{1,1} \rightarrow \Lambda^{1,1}$ , and  $*^2 = \text{id}_{\Lambda^{1,1}}$ . Consider the eigenspace orthogonal decomposition  $\Lambda^{1,1} \cong \Lambda_+^{1,1} \oplus \Lambda_-^{1,1}$ ,  $F = F_+ + F_-$  with  $*F_+ = F_+$  and  $*F_- = -F_-$ . Notice that:

$$\begin{aligned} I(A) &= \int_X \text{Tr}((F_+ + F_-) \wedge *(F_+ + F_-)) \\ &= \int_X \text{Tr}(F_+ \wedge *F_+ + F_+ \wedge *F_- + F_- \wedge *F_+ + F_- \wedge *F_-) \\ &= \|F_+\|_{L^2}^2 + \|F_-\|_{L^2}^2 \end{aligned}$$

On the other hand, observe that:

$$\begin{aligned} C_2(E) &= \int_X \text{Tr}(F \wedge F) \\ &= \int_X \text{Tr}((F_+ + F_-) \wedge (*F_+ - *F_-)) \\ &= \|F_+\|_{L^2}^2 - \|F_-\|_{L^2}^2 \end{aligned}$$

is topological.  $\Rightarrow I(A) \geq C_2(E) = \text{const}$ . Thus if we can find a connection with  $F_- = 0$ ,  $I(A)$  achieves minimum. The equation  $F_- = 0$  itself is easier than the non-linear YM equation.

## §9. Kähler Manifolds

Recall that, for a complex manifold  $X$  and  $E \rightarrow X$  holomorphic vector bundle we have:

$$\bar{\partial}: \Gamma(X, E \otimes \Lambda^{p,q}) \rightarrow \Gamma(X, E \otimes \Lambda^{p,q+1})$$

$$\phi \mapsto \bar{\partial}\phi$$

$\forall \phi, \psi \in \Gamma_c(X, E \otimes \Lambda^{p,q})$ , we have

$$\langle \phi, \psi \rangle = \frac{1}{p!q!} \int_X \phi_{\bar{j}_1 \dots \bar{j}_p}^{\alpha} \overline{\psi_{\bar{k}_1 \dots \bar{k}_q}^{\beta}} g^{k\bar{j}} g^{i\bar{l}} H_{\bar{\alpha}\beta} \cdot \text{vol.}$$

is an  $L^2$ -inner product. To define this we only needed:

- (1). A hermitian metric on  $E$  and on  $TX$ .
- (2). A volume element

Recall that if  $(X, g)$  is Kähler, then:

$$(\bar{\partial}^{\dagger} \psi)_{\bar{k}\bar{l}}^{\beta} = -g^{k\bar{j}} \nabla_{\bar{k}} \psi_{\bar{j}\bar{l}}^{\beta}$$

Rmk: Kähler condition enters as we used integration by parts. Furthermore,  $\nabla$  is the connection on  $E \otimes \Lambda^{p,q}$ , which is the Levi-Civita and Chern connection.

Kähler condition

Let  $(X, J)$ , complex manifold,  $J$ : almost complex structure,  $g$ : Riemannian metric, compatible with  $J$ :

$$g(Ju, Jv) = g(u, v).$$

i.e. it's a Hermitian metric. Let  $\omega(u, v) = g(Ju, v)$ ,  $\omega$  is alternating. Note that  $g$  induces the following structure:

- (1).  $\nabla_{LC}$ : Levi-Civita connection on  $T^{\mathbb{C}}X = TX \otimes_{\mathbb{R}} \mathbb{C} = T^{1,0}X \oplus T^{0,1}X$ .
- (2). A hermitian metric on  $T^{1,0}$  and thus the Chern connection  $\nabla_{Ch}$ .

Thm. The following are equivalent:

- (1).  $(X, J, g)$  is Kähler
- (2).  $\nabla_{LC} J = 0$
- (3).  $\nabla_{LC} \omega = 0$
- (4).  $\nabla_{LC} = \nabla_{Ch}$

(5).  $\forall p \in X, \exists U \ni p$  s.t.  $z(p) = 0$  and  $g_{\bar{k}j} = \delta_{kj} + O(|z|^2)$

Generalized Jacobi's identity.

Let  $A \in \Gamma(X, \Lambda^a \otimes \text{End}(E)), B \in \Gamma(X, \Lambda^b \otimes \text{End}(E))$ , we may define their generalized Lie bracket  $[A, B] \in \Gamma(X, \Lambda^{a+b} \otimes \text{End}(E))$ :

$$[A, B] \triangleq AB - (-1)^{ab} BA$$

We have Jacobi's identity:

$$ad_{[A, B]} = [ad_A, ad_B]$$

i.e.  $\forall C \in \Gamma(X, \Lambda^c \otimes \text{End}(E))$ :

$$[[A, B], C] = [A, [B, C]] - (-1)^{ab} [B, [A, C]].$$

Recall that we defined

$$L: \Gamma(X, E \otimes \Lambda^{p,q}) \rightarrow \Gamma(X, E \otimes \Lambda^{p+1, q}), \psi \mapsto \omega \wedge \psi$$

and its adjoint:

$$\Lambda: \Gamma(X, E \otimes \Lambda^{p,q}) \rightarrow \Gamma(X, E \otimes \Lambda^{p-1, q-1})$$

Thm. (Hodge identities)  $(X, J, g)$  Kähler and  $(E, H)/X$  a hermitian holomorphic vector bundle. Then:

$$(1). [\bar{\partial}^\dagger, L] = \sqrt{-1} \partial$$

$$(2). [\partial^\dagger, L] = -\sqrt{-1} \bar{\partial}$$

$$(3). [L, \bar{\partial}] = -\sqrt{-1} \partial^\dagger$$

$$(4). [L, \partial] = \sqrt{-1} \bar{\partial}^\dagger$$

where  $\partial = \nabla_{CH}^{1,0}$  on  $E$ .

Pf: We have shown this in local coordinates before. Here is the proof in the notation introduced above.

Note that (1) $^\dagger \Rightarrow$  (3) and (2) $^\dagger \Rightarrow$  (4), and (1)  $\Rightarrow$  (2). Thus it suffices to show (1).

Since  $\bar{\partial}\psi = d\bar{z}^\beta \wedge \nabla_{\bar{\beta}}\psi$ , we have  $\bar{\partial}^\dagger\psi = -\iota(dz^\alpha) \nabla_{\alpha}\psi$ , where  $\iota(dz^\alpha)$  acts on forms by:

(i). On 1-forms,  $\iota(dz^\alpha)(\theta) = \langle \theta, dz^\alpha \rangle_g$ , the Euclidean inner product. Thus

$$\iota(dz^\alpha)(dz^\beta) = 0, \quad \iota(dz^\alpha)(d\bar{z}^\beta) = g^{\alpha\bar{\beta}}.$$

(ii).  $(dz^\alpha)$  acts as a derivation: For  $\varphi \wedge \psi$ ,

$$L(dz^\alpha)(\varphi \wedge \psi) = (L(dz^\alpha)\varphi) \wedge \psi + (-1)^{|\varphi|} \varphi \wedge (L(dz^\alpha)\psi).$$

Then:

$$\begin{aligned} \bar{\partial}^\dagger L\psi &= \bar{\partial}^\dagger(\omega \wedge \psi) \\ &= -L(dz^\alpha) \nabla_{\bar{\alpha}}(\omega \wedge \psi) \\ &= -L(dz^\alpha) \omega \wedge \nabla_{\bar{\alpha}} \psi \\ &= -(L(dz^\alpha)\omega) \wedge \nabla_{\bar{\alpha}} \psi - \omega \wedge (L(dz^\alpha) \nabla_{\bar{\alpha}} \psi) \\ &= -(L(dz^\alpha)\omega) \wedge \nabla_{\bar{\alpha}} \psi + L \bar{\partial}^\dagger \psi \end{aligned}$$

But  $\omega = \sqrt{-1} g_{\bar{\beta}\gamma} dz^\gamma \wedge d\bar{z}^\beta$

$$\begin{aligned} \Rightarrow L(dz^\alpha)(\omega) &= \sqrt{-1} g_{\bar{\beta}\gamma} (L(dz^\alpha)(dz^\gamma) \wedge d\bar{z}^\beta - \sqrt{-1} g_{\bar{\beta}\gamma} dz^\gamma \wedge (L(dz^\alpha)(d\bar{z}^\beta)) \\ &= -\sqrt{-1} g_{\bar{\beta}\gamma} g^{\alpha\bar{\beta}} dz^\gamma \end{aligned}$$

$$\begin{aligned} \Rightarrow -(L(dz^\alpha)(\omega)) \wedge \nabla_{\bar{\alpha}} \psi &= \sqrt{-1} g_{\bar{\beta}\gamma} g^{\alpha\bar{\beta}} dz^\gamma \wedge \nabla_{\bar{\alpha}} \psi \\ &= \sqrt{-1} dz^\alpha \wedge \nabla_{\bar{\alpha}} \psi \\ &= \sqrt{-1} \partial \psi \end{aligned}$$

□

### Applications

(1). (KAN identity)  $\square_{\bar{\partial}} - \square_{\partial} = [L, F, \wedge]$ .

This is proven before. In particular, if  $E = \mathcal{O}_X$  is trivial,  $F = 0$ .

$$\square_{\partial} = \square_{\bar{\partial}}$$

Thm. For Kähler manifolds,  $\Delta_d = 2\square_{\partial} = 2\square_{\bar{\partial}}$ .

Pf: Recall that  $\Delta_d = dd^\dagger + d^\dagger d$

$$= [d, d^\dagger] \quad (d \text{ has degree } 1, d^\dagger \text{ degree } -1)$$

$$= [\partial + \bar{\partial}, \partial^\dagger + \bar{\partial}^\dagger]$$

$$= [\partial, \partial^\dagger] + [\bar{\partial}, \bar{\partial}^\dagger] + [\bar{\partial}, \partial^\dagger] + [\partial, \bar{\partial}^\dagger]$$

$$= \square_{\partial} + \square_{\bar{\partial}} + [\bar{\partial}, \partial^\dagger] + [\partial, \bar{\partial}^\dagger]$$

Thus it suffices to show that  $[\partial, \bar{\partial}^\dagger] = [\bar{\partial}, \partial^\dagger] = 0$ .

$$[\partial, \bar{\partial}^\dagger] = [-\sqrt{-1} [\bar{\partial}^\dagger, L], \bar{\partial}^\dagger]$$

$$= -\sqrt{-1} ([\bar{\partial}^\dagger, [L, \bar{\partial}^\dagger]] - (-1)^2 [L, [\bar{\partial}^\dagger, \bar{\partial}^\dagger]])$$

$$\begin{aligned}
&= -\sqrt{-1} ([ [L, \bar{\partial}^\dagger], \bar{\partial}^\dagger ] - [L, 2\bar{\partial}^\dagger \bar{\partial}^\dagger ]) \\
&= \sqrt{-1} ([ [\bar{\partial}^\dagger, L], \bar{\partial}^\dagger ]) \\
&= - [ \partial, \bar{\partial}^\dagger ].
\end{aligned}$$

$\Rightarrow [ \partial, \bar{\partial}^\dagger ] = 0$ . Similarly for  $[ \bar{\partial}, \partial^\dagger ]$ . □

Cor.  $\Delta d$  preserves  $(p, q)$ -types, i.e.  $\psi \in \Gamma(X, \Lambda^{p,q}) \Rightarrow \Delta d\psi \in \Gamma(X, \Lambda^{p,q})$ . □

Thm. (Hodge decomposition)  $X$ : compact Kähler manifold, then:

$$H^r(X) = \bigoplus_{p+q=r} H^{p,q}(X)$$

Pf: Take a Kähler metric on  $X$ . By Hodge's thm:  $H^r(X) \cong \mathcal{H}^r(X)$ , the space of harmonic  $d$ -forms.  $\forall \psi \in \mathcal{H}^r(X)$ , we may decompose it into its  $(p, q)$ -type:

$$\psi = \sum_{p+q=r} \psi^{p,q}$$

and

$$\begin{aligned}
0 &= \Delta d\psi = \sum_{p+q=r} (\Delta d\psi)^{p,q} \\
&= \sum_{p+q=r} \Delta d\psi^{p,q}
\end{aligned}$$

$\Rightarrow \Delta d\psi^{p,q} = 2\Box \bar{\partial}\psi^{p,q} = 0 \Rightarrow \psi^{p,q} \in \mathcal{H}^{p,q}(X)$ . □

Cor. If  $X$  is a compact Kähler manifold, then  $h^r(X)$  is even for  $r$  odd.

Pf: Indeed by the thm,  $H^r(X) \cong H^{r,0}(X) \oplus H^{r-1,1}(X) \oplus \dots \oplus H^{1,r-1}(X) \oplus H^{0,r}(X)$ .

Note that  $(\bar{\cdot}): H^{p,q}(X) \xrightarrow{\cong} H^{q,p}(X)$  is a complex conjugate linear isomorphism of vector spaces  $\Rightarrow h^r(X) = 2h^{r,0} + 2h^{r-1,1} + \dots$  is even. □

E.g. Hopf surface: Non-Kähler complex manifold.

$S^3 \times \mathbb{R} \cong \mathbb{R}^4 \setminus \{0\} \cong \mathbb{C}^2 \setminus \{0\}$  is a complex manifold. Define  $\mathbb{Z} \curvearrowright S^3 \times \mathbb{R}$  by

$$(\vec{v}, \lambda) \mapsto (e^\lambda \cdot \vec{v})$$

$$n \cdot (\vec{v}, \lambda) = (\vec{v}, \lambda + n)$$

(or on  $\mathbb{C}^2 \setminus \{0\}$ ,  $n$  acts by multiplication by  $e^n$ ).  $\mathbb{Z}$  preserves the complex structure on  $\mathbb{C}^2 \setminus \{0\}$ . Thus  $S^3 \times \mathbb{R} / \mathbb{Z} \cong S^3 \times S^1$  is a complex manifold.

But it's not Kähler since  $h^1(X) = 1$  is odd.

(ii) ( $\partial\bar{\partial}$ -lemma) If  $\varphi \in \Omega^{p,q}$  is  $d$ -exact, i.e.  $\varphi = d\psi$  for some  $\psi$ , then  $\exists \eta \in \Omega^{p-1,q-1}$  s.t.  $\varphi = \partial\bar{\partial}\eta$ .

Rmk: The converse is easy:  $\varphi = \partial\bar{\partial}\eta = (\partial + \bar{\partial})\bar{\partial}\eta = d\bar{\partial}\eta$ .

Pf of lemma.

$\varphi$  is  $d$ -exact  $\Rightarrow [\varphi] = 0$  in  $H^r(X)$ ,  $r = p+q$ . Let

$$\Gamma(X, \Lambda^{p,q}) = \mathcal{H}^{p,q}(X) \oplus \text{Im } \bar{\partial} \oplus \text{Im } \bar{\partial}^*$$

$$\Rightarrow \varphi = \square_{\bar{\partial}} G_{\bar{\partial}} \varphi = (\bar{\partial}\bar{\partial}^* + \bar{\partial}^*\bar{\partial}) G_{\bar{\partial}} \varphi.$$

Since  $[\bar{\partial}, G_{\bar{\partial}}] = 0$ ,

$$\begin{aligned} \Rightarrow \varphi &= \bar{\partial}\bar{\partial}^* G_{\bar{\partial}} \varphi + \bar{\partial}^* G_{\bar{\partial}} \bar{\partial} \varphi \\ &= \bar{\partial}\bar{\partial}^* G_{\bar{\partial}} \varphi \end{aligned}$$

Let  $\eta = \bar{\partial}^* G_{\bar{\partial}} \varphi \in \Gamma(X, \Lambda^{p,q-1})$ , then

$$\begin{aligned} \partial\eta &= \partial\bar{\partial}^* G_{\bar{\partial}} \varphi \\ &= \bar{\partial}^* \partial G_{\bar{\partial}} \varphi \quad ([\partial, \bar{\partial}^*] = 0) \\ &= \bar{\partial}^* \partial G_{\partial} \varphi \quad (G_{\bar{\partial}} = G_{\partial}) \\ &= \bar{\partial}^* G_{\partial} \partial \varphi \\ &= 0 \end{aligned}$$

$\Rightarrow \eta$  is  $\partial$ -closed  $\Rightarrow \eta = \pi_{\partial} \eta + \square_{\partial} G_{\partial} \eta$  ( $\pi_{\partial} \eta \in \mathcal{H}_{\partial}^{p,q} = \mathcal{H}_{\bar{\partial}}^{p,q}$ )

$$\begin{aligned} \Rightarrow \varphi &= \bar{\partial}(\pi_{\partial} \eta + (\partial\bar{\partial}^* + \bar{\partial}^*\partial) G_{\partial} \eta) \\ &= \bar{\partial}\partial\bar{\partial}^* G_{\partial} \eta. \end{aligned}$$

□



## §9. The Calabi Conjecture

$X$ : compact complex manifold.

**Question:** Let  $\omega_0 = \frac{i}{2} g_{\bar{k}j} dz^j \wedge d\bar{z}^k$  be a Kähler form on  $X$ . Let  $T_{\bar{k}j}$  be a (1,1) form in  $C_1(X)$ . Is there a Kähler form  $\omega$  in the same class as  $\omega_0$  satisfying

$$R_{\bar{k}j}(\omega) = T_{\bar{k}j}?$$

Here  $C_1(X) \cong C_1(T^{1,0}X)$ .

Calabi conjectured that the answer is yes, and such an  $\omega$  is unique within the class of  $\omega_0$ .

**Thm. (Yau)** The Calabi conjecture is true.

**Cor.** Let  $X$  be a compact complex manifold with  $C_1(X) = 0$ . Then if  $\omega_0$  is any Kähler class, there is a unique  $\omega \in [\omega_0]$  with

$$R_{\bar{k}j}(\omega) = 0$$

Reduction to a Monge-Ampère equation

$\omega \in [\omega_0]$  means that  $\omega - \omega_0 = d\Phi$  is exact. By the  $\partial\bar{\partial}$ -lemma, we have  $\omega - \omega_0 = \frac{i}{2} \partial\bar{\partial}\varphi$  where  $\varphi \in C^\infty(X, \mathbb{R})$ . Thus we are looking for with:

$$\omega \triangleq \omega_0 + \frac{i}{2} \partial\bar{\partial}\varphi > 0 \quad (\omega_0\text{-pluri-subharmonic condition})$$

and

$$R_{\bar{k}j}(\omega) = T_{\bar{k}j} \quad (*)$$

The equation (\*) may be rewritten as:

$$-\partial\bar{\partial} \log \omega^n = T_{\bar{k}j} \quad (*')$$

Observe that  $\omega^n$  is not a scalar function on  $X$ , but if we change variables, we get  $\omega^n(z) = \tilde{\omega}^n(w) \left| \frac{\partial w}{\partial z} \right|^2$ , and  $\log \omega^n = \log \tilde{\omega}^n + \log \left( \frac{\partial w}{\partial z} \right) + \log \left( \frac{\partial w}{\partial \bar{z}} \right)$ . Thus

$$\begin{aligned} -\partial\bar{\partial} \log \omega^n &= -\partial\bar{\partial} \log \tilde{\omega}^n - \partial\bar{\partial} \log \left( \frac{\partial w}{\partial z} \right) - \partial\bar{\partial} \log \left( \frac{\partial w}{\partial \bar{z}} \right) \\ &= -\partial\bar{\partial} \log \tilde{\omega}^n \end{aligned}$$

is a well-defined (1,1)-form.

To write the l.h.s. of  $(*)'$  in terms of scalars, note that

$$\begin{aligned} -\partial_j \partial_{\bar{k}} \log \omega^n &= -\partial_j \partial_{\bar{k}} \log \omega_0^n - \partial_j \partial_{\bar{k}} \log \frac{\omega^n}{\omega_0^n} \\ &= R_{\bar{k}j}(\omega_0) - \partial_j \partial_{\bar{k}} \log \frac{\omega^n}{\omega_0^n} \end{aligned}$$

where  $\frac{\omega^n}{\omega_0^n}$  is now a scalar function. Then  $(*)'$  is equivalent to

$$\begin{aligned} -\partial_j \partial_{\bar{k}} \log \left( \frac{\omega^n}{\omega_0^n} \right) &= T_{\bar{k}j} - R_{\bar{k}j}(\omega_0) \\ &= \partial_j \partial_{\bar{k}} F \end{aligned}$$

In other words,

$$-\partial_j \partial_{\bar{k}} \left( \log \frac{\omega^n}{\omega_0^n} - F \right) = 0$$

where  $F$  is a given smooth function on  $X$  (by  $\partial\bar{\partial}$ -lemma again).

Claim:  $\log \frac{\omega^n}{\omega_0^n} - F \equiv c$ .

$$\begin{aligned} \text{Pf: } \partial_j \partial_{\bar{k}} h = 0 &\Rightarrow \int_X \partial_j \partial_{\bar{k}} h \cdot \bar{h} g^{j\bar{k}} \frac{\omega^n}{n!} = 0 \\ &\Rightarrow 0 = \int_X \partial_{\bar{k}} h \overline{\partial_j h} g^{j\bar{k}} \frac{\omega^n}{n!} \\ &= \|\bar{\partial} h\|_{L^2}^2 \end{aligned}$$

Thus  $h$  is globally holomorphic and thus a constant □

Changing  $F$  into  $F+c$ , we are reduced to solve:

$$\omega^n = e^F \omega_0^n$$

More explicitly:

$$\det(g_{\bar{k}j}^0 + \partial_j \partial_{\bar{k}} \varphi) = e^F \det(g_{\bar{k}j}^0)$$

where  $F$  is a global smooth function determined up to a constant. This is the Monge-Ampère equation. Note that the constant can be fixed as follows:

$$\begin{aligned} \int_X \omega^n &= \int_X (\omega_0 + \frac{i}{2} \partial\bar{\partial}\varphi)^n \\ &= \int_X \omega_0^n + n \omega_0^{n-1} \wedge (\frac{i}{2} \partial\bar{\partial}\varphi) + \dots + (\frac{i}{2} \partial\bar{\partial}\varphi)^n \\ &= \int_X \omega_0^n \end{aligned}$$

since  $\partial\bar{\partial}\varphi = d(\bar{\partial}\varphi)$  is exact and  $\omega_0$  is closed. Thus  $\omega^n = e^F \omega_0^n$  gives

$$\int_X e^F \omega_0^n = \int_X \omega^n = \int_X \omega_0^n$$

and this condition uniquely determines  $F$  in the Monge-Ampère equation.

Method of Continuity.

Introduce the following family of equations, for the unknown  $\varphi_t$  ( $0 \leq t \leq 1$ ):

$$\det(g_{\bar{k}\bar{j}}^0 + \partial_{\bar{j}} \partial_{\bar{k}} \varphi_t) = e^{tF - C_t} \det(g_{\bar{k}\bar{j}}^0) \quad (*)_t$$

where  $C_t$  is so chosen that

$$\int_X e^{tF - C_t} \omega_0^n = \int_X \omega_0^n$$

i.e.  $C_t = \log \frac{\int_X e^{tF} \omega_0^n}{\int_X \omega_0^n}$

Define  $I = \{t \in [0, 1] \mid (*)_t \text{ admits a smooth solution satisfying the } \omega_0\text{-pluri-subharmonic condition } g_{\bar{k}\bar{j}}^0 + \partial_{\bar{j}} \partial_{\bar{k}} \varphi_t > 0\}$

We shall show that:

- (a).  $I \neq \emptyset$  (easy, since  $0 \in I$ )
- (b).  $I$  is open
- (c).  $I$  is closed

Then it will follow that  $I = [0, 1]$  and  $(*)$  admits a smooth solution.

Proof of (b). — Implicit function thm. for Banach spaces.

Rewrite  $(*)_t$  as:


$$\log \frac{\det(g_{\bar{k}\bar{j}}^0 + \partial_{\bar{j}} \partial_{\bar{k}} \varphi)}{\det(g_{\bar{k}\bar{j}}^0)} - (tF - C_t) = 0 \quad (*)_t$$

We want to show that if  $(*)_t$  admits a solution  $\varphi_{t_0}$  at  $t_0$ , then  $\exists \varepsilon > 0$  s.t.  $\forall t: |t - t_0| < \varepsilon$ ,  $(*)_t$  admits a  $C^\infty$ ,  $\omega_0$ -pluri-subharmonic solution  $\varphi_t$ .

Recall the standard implicit function thm on  $\mathbb{R}^2$ : let  $E(x, y)$  be a  $C^2$  function on  $\mathbb{R}^2$  and assume that:

$$\begin{cases} E(x_0, y_0) = 0 \\ \frac{\partial E}{\partial y}(x_0, y_0) \neq 0 \end{cases}$$

Then  $\exists \varepsilon > 0$  and a nhd  $V$  of  $y_0$  s.t.  $\forall x: |x - x_0| < \varepsilon$ ,  $\exists! y \in V$  s.t.  $E(x, y) = 0$ .



Thm. (Implicit function thm for Banach spaces)

Let  $B_i, i=1, 2, 3$  be Banach spaces, and  $E: B_1 \times B_2 \rightarrow B_3$  be a  $C^1$ -continuous function. Assume that  $E(x_0, y_0) = 0$  for some  $x_0 \in B_1$  and  $y_0 \in B_2$ .  $\frac{\partial E}{\partial y}(x_0, y_0)$  is an invertible operator from  $B_2$  to  $B_3$ , with bounded inverse. Then  $\exists$  nhd  $V$  of  $x_0$ ,  $\tilde{V}$  of  $y_0$  so that  $\forall x \in V, \exists! y \in \tilde{V}$  s.t.  $E(x, y) = 0$ .

We need to set up the proof of (b) so as to apply the implicit function thm. Let

$$E(t, \varphi) \triangleq \log \frac{\det(g_{ij}^0 + \partial_i \partial_{\bar{k}} \varphi)}{\det(g_{ij}^0)} - (tF - Ct)$$

where  $E: \mathbb{R} \times B_2 \rightarrow B_3$  for appropriate Banach spaces  $B_2, B_3$ .

Def. (Schauder spaces) On  $\mathbb{R}^n$ , fix  $0 < \alpha < 1$ .

$$\|\varphi\|_{C^{0,\alpha}} \triangleq \sup |\varphi| + \sup_{x \neq y} \frac{|\varphi(x) - \varphi(y)|}{|x - y|^\alpha}$$

$C^{0,\alpha}(\mathbb{R}^n) \triangleq \{\varphi \mid \|\varphi\|_{C^{0,\alpha}} < \infty\}$ . Observe that  $\varphi, \psi \in C^{0,\alpha} \Rightarrow \varphi \cdot \psi \in C^{0,\alpha}, e^\varphi \in C^{0,\alpha}$ .

More generally,  $C^{k,\alpha}(\mathbb{R}^n) \triangleq \{\varphi \mid \partial^{\nu} \varphi \in C^{0,\alpha}(\mathbb{R}^n), \forall |\nu| \leq k\}$ .

On a compact manifold, similar spaces  $C^{k,\alpha}(X)$  can be defined via taking a partition of unity.

Next, observe that the solution to  $(*)_t$  is not unique:  $\varphi_t$  satisfies  $(*)_t \Rightarrow$  so does  $\varphi_t + \text{const}$ . Introduce:

$$C_0^{k,\alpha}(X) \triangleq \{\varphi \in C^{k,\alpha}(X) \mid \int_X \varphi \omega_0^n = 0\}$$

Then we claim that  $E(t, \varphi): \mathbb{R} \times C_0^{2,\alpha} \rightarrow C_0^{0,\alpha}$ . Indeed:

$$\int_X \left( \frac{\omega^n}{\omega_0^n} - e^{tF - Ct} \right) \omega_0^n = \int_X \omega^n - \int_X e^{tF - Ct} \omega_0^n = 0$$

since  $C_t$  is so chosen.

Verifying the hypotheses of the IFT.

Given  $f: B_2 \rightarrow B_3$ , we say that  $f$  is differentiable at  $y_0 \in B_2$  if  $\exists$  a bounded linear map  $L: B_2 \rightarrow B_3$  (bounded:  $\exists C > 0$  s.t.  $\forall y \in B_2, \|Ly\|_{B_3} \leq C\|y\|_{B_2}$ ).

s.t. 
$$\lim_{y \rightarrow y_0} \frac{\|f(y) - f(y_0) - L_{y_0}(y - y_0)\|_{B_3}}{\|y - y_0\|_{B_2}} = 0$$

$\text{Hom}(B_2, B_3) = \{\text{bounded linear operators sending } B_2 \text{ into } B_3\}$  is a Banach space, with norm defined by:

$$\|T\| \triangleq \sup_{\substack{y \in B_2 \\ y \neq 0}} \frac{\|Ty\|_{B_3}}{\|y\|_{B_2}}$$

$E \in C^1(\Omega)$  if  $y \mapsto L_y$  is defined for all  $y \in \Omega$  and is continuous as a mapping  $\Omega \rightarrow \text{Hom}(B_2, B_3)$ .

Thus now in our case, we want to show that

- (1).  $E \in C^1$
- (2). Compute  $\frac{\delta E}{\delta \varphi} \in \text{Hom}(C^{2,\alpha}, C^{0,\alpha})$  (boundedness)
- (3).  $\frac{\delta E}{\delta \varphi}$  has a bounded inverse.

Computation of  $\frac{\delta E}{\delta \varphi}$ :

By variation: imagine that  $\varphi \mapsto \varphi + \delta\varphi$ .

$$\begin{aligned} \delta(E) &= \delta\left(\frac{\det(g_{\bar{k}j}^0 + \partial_j \partial_{\bar{k}} \varphi)}{\det g_{\bar{k}j}^0} - e^{tF - Ct}\right) \\ &= \frac{1}{\det g_{\bar{k}j}^0} \delta(\det(g_{\bar{k}j}^0 + \partial_j \partial_{\bar{k}} \varphi)) \end{aligned}$$

Observe that if  $G$  is an invertible matrix:

$$\begin{aligned} \delta(\det G) &= (\det G)^{-1} \delta(\det G) \cdot \det G \\ &= \delta(\ln \det G) \cdot \det G \\ &= \text{Tr}(G^{-1} \delta G) \det G \end{aligned}$$

Hence:

$$\begin{aligned} \delta E &= \frac{1}{\det(g_{\bar{k}j}^0)} \det(g_{\bar{k}j}^0) \cdot g^{j\bar{k}} \delta g_{\bar{k}j} \\ &= \frac{1}{\det(g_{\bar{k}j}^0)} \det(g_{\bar{k}j}^0) \cdot g^{j\bar{k}} \partial_j \partial_{\bar{k}} \delta\varphi \\ &= \frac{1}{\det(g_{\bar{k}j}^0)} \det g_{\bar{k}j}^0 \cdot \Delta_g \delta\varphi. \end{aligned}$$

and

$$L: \delta\varphi \mapsto \frac{\det g_{\bar{k}j}}{\det g_{\bar{k}j}^0} \Delta_g \delta\varphi.$$

Claim:  $L\varphi$  admits a bounded inverse, i.e.  $\forall \mu \in C_0^{0,\alpha}$ ,  $\exists! \nu \in C_0^{2,\alpha}$  s.t.

$$L\nu = \mu$$

Furthermore,

$$\|\nu\|_{C^{2,\alpha}} \leq C \|\mu\|_{C^{0,\alpha}}$$

for some constant  $C > 0$ . This follows from the following:

Basic fact from linear analysis:

Let  $(X, g_{ij})$  be any compact Riemannian manifold, then the equation

$$\Delta_g \nu = \mu$$

admits a solution iff

$$\int_X \mu \sqrt{g} = 0.$$

The solution  $\nu$  is unique if we require the normalization condition:

$$\int_X \nu \sqrt{g} = 0$$

Now in our case,  $L\varphi \nu = \mu \iff \Delta_g \nu = \frac{\omega_0^n}{\omega^n} \mu$ . This is solvable since  $\mu \in C_0^{0,\alpha}$ , and the solution is unique if we specify

$$\int_X \nu \omega_0^n = 0$$

The similar a priori estimates hold for Schauder spaces, which implies the boundedness of the inverse of  $L$  (the Green's operator)

Proof of (c):  $I$  is closed.

We want to show that  $t_j \in I$ ,  $t_j \rightarrow T \implies T \in I$ . i.e.  $\varphi_{t_j}$  admits a solution of  $(*)_{t_j}$ , and  $t_j \rightarrow T$ , then  $\exists \varphi$  a solution of  $(*)_T$  satisfying the Monge-Ampère equation and the pluri-subharmonic condition.

To start, we would wish to show that a subsequence, renamed still by  $\{\varphi_j\}$ , converges in  $C^{3,\alpha}$  (or all  $C^{m,\alpha}$ ). If so, and the limit  $\varphi$  satisfies the M-A equation, we see that the determinant of  $g_{\bar{k}j}^0 + \partial_j \partial_{\bar{k}} \varphi$  is fixed pointwise. If we can further show that the eigenvalues of  $g_{\bar{k}j}^0 + \partial_j \partial_{\bar{k}} \varphi$  is bounded from above, they will also be bounded from below, whence  $g_{\bar{k}j}^0 + \partial_j \partial_{\bar{k}} \varphi > 0$ .

To achieve this, it suffices to show that,  $\forall m$ ,

$$\|\varphi_{t_j}\|_{C^m} \leq Am. \quad (AP)_m,$$

since  $\|\varphi_{t_j}\|_{C^1} + \|\varphi_{t_j}\|_{C^0} \leq \text{const} \Rightarrow \{\varphi_{t_j}\}$  is equicontinuous. (The standard trick as used in Hodge theory:  $|\varphi_{t_j}(x) - \varphi_{t_j}(y)| \leq \text{Sup} \|\nabla \varphi\| \cdot |x - y|$ .) Then by Arzela-Ascoli thm, there would be a convergent subsequence.

### Schauder theory

$(AP)_m, 0 \leq m \leq 3 \Rightarrow (AP)_m$  for all  $m$ . (In fact,  $0 \leq m \leq 2 + \varepsilon$  will do. This is because a priori estimate is valid only for  $\varepsilon > 0 : \|\varphi\|_{C^{2+\varepsilon}}$ ). The 4 cases

$m=3$  (Calabi)

$m=2$  (Aubin-Yau, Pogorelov)

$m=1$

$m=0$  (This is the hardest part, done by Yau using Moser's iteration).

Furthermore,  $(m=2) + (m=0) \Rightarrow (m=1)$ .

10 years ago, Kolodziej proved a stronger estimate:

Thm. (Kolodziej).

$$\det(g_{\bar{i}\bar{j}} + \partial_j \partial_{\bar{k}} \varphi) = f \det g_{\bar{i}\bar{j}}$$

with  $f > 0$ . Then  $\forall p > 1, \exists \alpha > 0$ , so that whenever  $\|f\|_{L^p} < \infty$ , we have

$$\|\varphi\|_{C^\alpha} < \text{Const}$$

### Proof of the $C^0$ -estimate

Moser iteration in the simplest case.

Consider  $\Delta u = f$ , where  $\Delta$  is the Laplacian on a compact manifold  $(X, g_{ij})$ . Assume that  $u$  is normalized:  $\int_X u \sqrt{g} dx = 0$ . We want to estimate for  $\|u\|_{C^0}$ .

$$\begin{aligned} (a). \quad \int_X u \cdot f \sqrt{g} dx &= \int_X u \cdot \Delta u \sqrt{g} dx \\ &= \int_X u \cdot \frac{1}{\sqrt{g}} (\partial_j (\sqrt{g} g^{ij} \partial_i u)) \sqrt{g} dx \\ &= - \int_X (\partial_j u \cdot \partial_i u) g^{ij} \sqrt{g} dx \\ &= - \int_X \|\nabla u\|^2 \sqrt{g} dx \end{aligned}$$

$$\Rightarrow \|\nabla u\|_{L^2}^2 \leq \left| \int_X u \cdot f \sqrt{g} dx \right| \leq \|u\|_{L^2} \|f\|_{L^2}$$

Recall Poincaré's inequality:

$$\|u\|_{L^2}^2 \leq C \cdot (\|\nabla u\|_{L^2}^2 + \left(\int_X u \sqrt{g}\right)^2)$$

Combined with above it gives

$$\|u\|_{L^2}^2 \leq C \cdot \|u\|_{L^2} \|f\|_{L^2}$$

$$\Rightarrow \|u\|_{L^2} \leq C \cdot \|f\|_{L^2}$$

(b). Moser iteration: Let  $\beta \equiv \frac{n}{n-2} (> 1)$ ,  $n = \dim_{\mathbb{R}} X$ . Then  $\forall p \geq 2$ ,

$$\max(1, \|u\|_{L^{p\beta}}) \leq (C \cdot p)^{\frac{1}{\beta}} \max(1, \|u\|_{L^p})$$

where  $C$  is a constant independent of  $p$ .

Assuming this, we can bound  $\|u\|_{L^\infty}$  ( $= \|u\|_{C^0}$ ) in the following manner:

Start with  $p$  (thinking of  $p=2$  as the starting point from (a)).

$$\log \max(1, \|u\|_{L^{p\beta}}) \leq \frac{1}{\beta} \log(C \cdot p) + \log \max(1, \|u\|_{L^p})$$

Iterate with  $p$  replaced by  $p\beta$ :

$$\begin{aligned} \log \max(1, \|u\|_{L^{p\beta^2}}) &\leq \frac{1}{p\beta} \log(C \cdot p\beta) + \log \max(1, \|u\|_{L^{p\beta}}) \\ &\leq \frac{1}{p\beta} \log(C \cdot p\beta) + \frac{1}{p} \log(C \cdot p) + \log \max(1, \|u\|_{L^p}) \end{aligned}$$

After  $k$ -steps, we get:

$$\log \max(1, \|u\|_{L^{p\beta^k}}) \leq \underbrace{\sum_{l=1}^{k-1} \frac{1}{p\beta^l} \log(C \cdot p\beta^l)}_{\text{converges when } k \rightarrow \infty} + \log \max(1, \|u\|_{L^p})$$

Hence in the limit,

$$\log \max(1, \|u\|_{L^\infty}) \leq A + \log \max(1, \|u\|_{L^p})$$

(c). Proof of the Moser iteration step

Recall from calculus: What's the anti-derivative of  $|x|^\alpha$ ?

$$(x \cdot |x|^\alpha)' = (\alpha+1) |x|^\alpha$$

Then if  $\Delta u = f$ ,

$$u \cdot |u|^\alpha \Delta u = |u|^\alpha \cdot f$$

$\Rightarrow$

$$\int_X |u|^\alpha \Delta u \sqrt{g} dx = \int_X |u|^\alpha f \sqrt{g} dx$$

Integrating by parts, this gives:



$$\begin{aligned}
\int_X u |u|^{\alpha} f \sqrt{g} dx &= - \int_X \nabla(u |u|^{\alpha}) \cdot \nabla u \sqrt{g} dx \\
&= - \int_X (\alpha+1) |u|^{\alpha} \|\nabla u\|^2 \sqrt{g} dx \\
&= - (\alpha+1) \int_X |u|^{\frac{\alpha}{2}} \|\nabla u\|^2 \sqrt{g} dx \\
&= - \frac{(\alpha+1)}{(1+\frac{\alpha}{2})^2} \int_X \|\nabla(u \cdot |u|^{\frac{\alpha}{2}})\|^2 \sqrt{g} dx
\end{aligned}$$

$\Rightarrow$

$$- \frac{(\alpha+1)}{(1+\frac{\alpha}{2})^2} \int_X \|\nabla(u \cdot |u|^{\frac{\alpha}{2}})\|^2 \sqrt{g} dx = \int_X u |u|^{\alpha} f \sqrt{g} dx$$

Similar as using Poincaré's inequality above, we need the Sobolev inequality:

$$\|u\|_{L^{2\beta}}^2 \leq C \cdot (\|\nabla u\|_{L^2}^2 + \|u\|_{L^2}^2)$$

In particular, apply Sobolev's inequality to  $v = u \cdot |u|^{\frac{\alpha}{2}}$ , we have:

$$\|u \cdot |u|^{\frac{\alpha}{2}}\|_{L^{2\beta}}^2 \leq C \cdot (\|\nabla(u \cdot |u|^{\frac{\alpha}{2}})\|_{L^2}^2 + \|u \cdot |u|^{\frac{\alpha}{2}}\|_{L^2}^2)$$

i.e.

$$\left( \int_X |u|^{(1+\frac{\alpha}{2})2\beta} \sqrt{g} dx \right)^{\frac{1}{\beta}} \leq C \cdot \left( \frac{(1+\frac{\alpha}{2})^2}{(\alpha+1)} \int_X |u|^{\alpha+1} f \sqrt{g} dx + \int_X |u|^{\alpha+2} \sqrt{g} dx \right)$$

Set  $p = 2 + \alpha$ . Then

$$\left( \int_X |u|^{p\beta} \sqrt{g} dx \right)^{\frac{1}{\beta}} \leq C \cdot (p \cdot \left| \int_X |u|^{\alpha+1} f \sqrt{g} dx \right| + \left| \int_X |u|^p \sqrt{g} dx \right|)$$

Note that

$$\begin{aligned}
\left| \int_X |u|^{\alpha+1} f \sqrt{g} dx \right| &\leq \|f\|_{C^0} \int_X |u|^{\alpha+1} \sqrt{g} dx \\
&\leq \|f\|_{C^0} \left( \int_X |u|^{\alpha+2} \sqrt{g} dx \right)^{\frac{\alpha+1}{\alpha+2}} \left( \int_X 1 \cdot \sqrt{g} dx \right)^{\frac{1}{\alpha+2}} \quad (\text{Jordan-Hölder}) \\
&\leq C' \cdot (\|u\|_{L^p}^p)^{\frac{\alpha+1}{\alpha+2}} \quad \left( \int |fg| \leq \left( \int |f|^p \right)^{\frac{1}{p}} \left( \int |g|^q \right)^{\frac{1}{q}} \right)
\end{aligned}$$

Combined, since  $\frac{\alpha+1}{\alpha+2} < 1$ , there are two cases: if  $\|u\|_{L^p} \leq 1$ ,  $C' \cdot (\|u\|_{L^p}^p)^{\frac{\alpha+1}{\alpha+2}} \leq C' \cdot 1$ ; otherwise,  $C' \cdot (\|u\|_{L^p}^p)^{\frac{\alpha+1}{\alpha+2}} \leq C' \cdot \|u\|_{L^p}^p \Rightarrow$

$$\|u\|_{L^{p\beta}}^p \leq C' p \max(1, \|u\|_{L^p}^p).$$

Moser iteration for the Monge-Ampère equation.

Notation:  $\omega = \frac{i}{2} g_{\bar{j}k} dz^j \wedge d\bar{z}^k$ .  $\omega_{\phi} = \frac{i}{2} (g_{\bar{j}k} + \partial_{\bar{b}} \partial_j \phi) dz^j \wedge d\bar{z}^k$ . Then the Monge-Ampère equation becomes:

$$\omega_{\phi}^n = e^f \omega^n, \quad (f = (tF - Ct))$$

Similar as the simple case above, we want to control  $\|\nabla u\|_{L^2}^2$ . The trick is the following:

$$e^f \omega^n - \omega^n = \omega_{\phi}^n - \omega^n$$

$$\begin{aligned}
&= (\omega_\varphi - \omega)(\omega_{\bar{\varphi}}^{n-1} + \omega_{\bar{\varphi}}^{n-2}\omega + \dots + \omega^{n-1}) \\
&= \frac{i}{2} \partial \bar{\partial} \varphi (\omega_{\bar{\varphi}}^{n-1} + \dots + \omega^{n-1}).
\end{aligned}$$

As in the Laplacian case, we integrate by parts.

$$\begin{aligned}
\int_X \varphi |\varphi|^\alpha (e^f - 1) \omega^n &= \int_X \varphi |\varphi|^\alpha \left( \frac{i}{2} \partial \bar{\partial} \varphi \right) \wedge (\omega_{\bar{\varphi}}^{n-1} + \dots + \omega^{n-1}) \\
&= \int_X \varphi |\varphi|^\alpha \left( \frac{i}{2} d \bar{\partial} \varphi \right) \wedge (\omega_{\bar{\varphi}}^{n-1} + \dots + \omega^{n-1}) \\
&= - \int_X d(\varphi |\varphi|^\alpha) \left( \frac{i}{2} \bar{\partial} \varphi \right) \wedge (\omega_{\bar{\varphi}}^{n-1} + \dots + \omega^{n-1}) \\
&= - \int_X (\alpha+1) |\varphi|^\alpha \cdot \frac{i}{2} \partial \varphi \wedge \bar{\partial} \varphi \wedge (\omega_{\bar{\varphi}}^{n-1} + \dots + \omega^{n-1}) \\
&= - \frac{(\alpha+1)}{(1+\frac{\alpha}{2})^2} \int_X \partial(\varphi |\varphi|^{\frac{\alpha}{2}}) \wedge \bar{\partial}(\varphi |\varphi|^{\frac{\alpha}{2}}) \wedge (\omega_{\bar{\varphi}}^{n-1} + \dots + \omega^{n-1})
\end{aligned}$$

Now we make two observations:

$$(1). \int_X \partial \psi \wedge \bar{\partial} \psi \wedge \omega^{n-1} = \|\nabla \psi\|_{L^2}^2.$$

This follows, since for any  $(1,1)$ -form  $T_{\bar{j}j} dz^j \wedge d\bar{z}^{\bar{k}}$

$$\int_X T_{\bar{j}j} dz^j \wedge d\bar{z}^{\bar{k}} \wedge \omega^{n-1} = \int_X g^{j\bar{k}} T_{\bar{j}j} \omega^n$$

Thus,

$$\begin{aligned}
\int_X \partial \psi \wedge \bar{\partial} \psi \wedge \omega^{n-1} &= \int_X g^{j\bar{k}} \partial_{\bar{k}} \psi \partial_j \psi \cdot \omega^n \\
&= \|\nabla \psi\|_{L^2}^2.
\end{aligned}$$

In particular,

$$\frac{\alpha+1}{(1+\frac{\alpha}{2})^2} \int_X i(\partial(\varphi |\varphi|^{\frac{\alpha}{2}}) \wedge \bar{\partial}(\varphi |\varphi|^{\frac{\alpha}{2}}) \wedge \omega^{n-1}) = \frac{(\alpha+1)}{(1+\frac{\alpha}{2})^2} \int_X |\nabla(\varphi |\varphi|^\alpha)|^2 \omega^n$$

(2). The remaining terms  $\int i(\partial(\varphi |\varphi|^{\frac{\alpha}{2}}) \wedge \bar{\partial}(\varphi |\varphi|^{\frac{\alpha}{2}}) \wedge \omega_{\bar{\varphi}}^k \wedge \omega^{n-k-1})$  are all non-negative.

In fact, the integrand is non-negative pointwise. More precisely, we say a  $(k,k)$ -form is positive if w.r.t. some  $(1,0)$  frame  $\{e_i\}$ , it can be written as a positive linear combination of terms of the form:

$$(ie_{i_1} \wedge \bar{e}_{\bar{i}_1}) \wedge \dots \wedge (ie_{i_k} \wedge \bar{e}_{\bar{i}_k}).$$

Note that this agrees with the standard definition of positivity for  $(1,1)$  forms.

Moreover, if  $\psi, \psi'$  are positive  $\implies$  so is  $\psi \wedge \psi'$ . Hence

$$\underbrace{i(\partial \psi \wedge \bar{\partial} \bar{\psi})}_{\geq 0} \wedge \underbrace{\omega_{\bar{\varphi}}^k}_{\geq 0} \wedge \underbrace{\omega^{n-k-1}}_{\geq 0} \geq 0$$

In fact, as an  $(n,n)$ -form, it's a positive multiple of the volume form.

$$\psi \prod_{i=1}^n \left( \frac{i}{2} dz^i \wedge d\bar{z}^{\bar{i}} \right), \text{ with } \psi \geq 0.$$

Combining these, we obtain the desired bound for  $\|\nabla(|\varphi|^{2\frac{n}{n-1}})\|_{L^2}$ . Applying Moser's iteration as above and we get the  $C^0$  bound. This is the major contribution of Yau.

Proof of the  $C^2$ -estimate.

In fact we shall just control something weaker:

$$(\omega + \frac{i}{2} \partial \bar{\partial} \varphi)^n = e^f \omega^n \quad (**)$$

Then  $\exists C > 0$ , depending only on  $f$ ,  $\omega$  and  $\text{osc}(\varphi) (\triangleq \sup \varphi - \inf \varphi)$ , which can be treated now as a known quantity, by the  $C^0$ -estimate above), s.t.

$$\|\Delta \varphi\|_{C^0} \leq C$$

The  $C^2$ -estimate follows from this and a priori estimate (similar as done in Hodge theory).

Idea of proof (Aubin, Yau, Pogorelov)

Apply the maximum principle to the expression  $\log(n + \Delta \varphi) - A\varphi$ . More precisely, we claim that  $\exists A, C_1, C_2 > 0$  s.t.

$$(*) \implies \Delta'(\log(n + \Delta \varphi) - A\varphi) \geq C_1 \left( \sum_{j=1}^n \frac{1}{\lambda_j} \right) - C_2 \quad (**)$$

at the maximum point of  $\log(n + \Delta \varphi) - A\varphi$ . Here  $\lambda_i$ 's are the eigenvalues of  $\omega_\varphi$  w.r.t.  $\omega$ ,  $\Delta'$  is the Laplacian of  $g'_{\bar{k}j} = g_{\bar{k}j} + \partial_j \partial_{\bar{k}} \varphi$ .

Assuming this claim, then since  $p$  is a maximum,

$$0 \geq C_1 \left( \sum_{j=1}^n \frac{1}{\lambda_j} \right) - C_2$$

( $\Delta'(\log(n + \Delta \varphi) - A\varphi)$  is the trace of the Hessian of  $\log(n + \Delta \varphi) - A\varphi$ , which is a negative semi-definite matrix since  $p$  is a maximum  $\implies$  the trace  $\leq 0$ ).

$$\implies \frac{1}{\lambda_j} \leq C_3, \quad \forall j \quad (\text{or } \lambda_j \text{ bounded from below})$$

However, by (\*),  $\prod_{j=1}^n \lambda_j = e^f(p)$  is bounded, hence  $\lambda_j$ 's are bounded from below and above at  $p$ . i.e.  $\exists C_4 \geq C_5$  s.t.

$$C_5 \leq \lambda_j \leq C_4$$

Now consider a general point. Since  $p$  is a maximum,

$$\log(n + \Delta \varphi(z)) - A\varphi(z) \leq \log(n + \Delta \varphi(p)) - A\varphi(p)$$

$$\begin{aligned}
\Rightarrow \log(n + \Delta\varphi(z)) &\leq \log(n + \Delta\varphi(p)) + A(\varphi(z) - \varphi(p)) \\
&= \log(\text{Tr}\omega') + A(\varphi(z) - \varphi(p)) \\
&\leq C_6 + A \cdot \text{osc}(\varphi) \\
\Rightarrow (n + \Delta\varphi(z)) &\leq e^{C_6 + A \cdot \text{osc}(\varphi)} \leq C_7.
\end{aligned}$$

Proof of (\*\*):

The trick here is to use the endomorphism  $h^{\ell}_m = g^{\ell\bar{p}} g'_{\bar{p}m}$  to calculate  $\Delta h$ . We begin with:

$$\begin{aligned}
\text{Tr}h &= h^{\ell}_\ell \\
&= g^{\ell\bar{p}} (g'_{\bar{p}\ell} + \partial_\ell \partial_{\bar{p}} \varphi) \\
&= n + \Delta\varphi
\end{aligned}$$

and compute:

$$\begin{aligned}
\Delta'(\text{Tr}h) &= (g')^{j\bar{k}} \partial_j \partial_{\bar{k}} (\text{Tr}h) \\
&= (g')^{j\bar{k}} \partial_{\bar{k}} \partial_j (\text{Tr}h) \\
&= (g')^{j\bar{k}} \text{Tr}(\nabla_{\bar{k}}' ((\nabla_j' h) h^{-1} \cdot h)) \\
&= (g')^{j\bar{k}} \text{Tr}\{(\nabla_{\bar{k}}' (\nabla_j' h) h^{-1}) \cdot h + (\nabla_j' h) \cdot h^{-1} \cdot \nabla_{\bar{k}}' h\}
\end{aligned}$$

where  $\nabla'$  is the covariant derivative with respect to  $g'_{\bar{j}}$ . Now recall that  $R_{\bar{k}j} = -\partial_{\bar{k}}((g')^{-1} \partial_j g)$ : curvature form in matrix notation. Remember that  $h = g^{-1} g' \in \Gamma(X, \text{End}(T^{1,0}X))$  (a holomorphic bundle with induced metrics). Then:

$$\begin{aligned}
R_{\bar{k}j} &= -\partial_{\bar{k}}(g^{-1} \partial_j g) \\
&= -\partial_{\bar{k}}(h g'^{-1} \partial_j (g' h^{-1})) \\
&= -\partial_{\bar{k}}\{(h g'^{-1} \partial_j g' \cdot h^{-1} + h g'^{-1} g' \partial_j (h^{-1}))\} \\
&= -\partial_{\bar{k}}\{(h g'^{-1} \partial_j g' \cdot h^{-1} + h \partial_j (h^{-1}))\}
\end{aligned}$$

Recall that covariant differential of an endomorphism  $T$  takes the form:

$$\nabla_j T = \partial_j T + A_j T - T A_j$$

where  $A_j = g^{-1} \partial_j g$ . Thus  $\nabla_j'(h^{-1}) = \partial_j(h^{-1}) + g'^{-1} \partial_j g' \cdot h^{-1} - h^{-1} (g'^{-1} \partial_j g')$   $\Rightarrow$   
 $h \nabla_j'(h^{-1}) = h \partial_j(h^{-1}) + h g'^{-1} \partial_j g' \cdot h^{-1} - (g'^{-1} \partial_j g')$ . Plugging in we get:

$$\begin{aligned}
R_{\bar{k}j} &= -\partial_{\bar{k}}(h \nabla_j' h^{-1} + (g'^{-1} \partial_j g')) \\
&= -\partial_{\bar{k}}(g'^{-1} \partial_j g') + \partial_{\bar{k}}(\nabla_j' h \cdot h^{-1}) \\
&= R'_{\bar{k}j} + \partial_{\bar{k}}(\nabla_j' h \cdot h^{-1}) \\
&= R'_{\bar{k}j} + \nabla_{\bar{k}}'(\nabla_j' h \cdot h^{-1})
\end{aligned}$$

Hence we obtain:

$$\Delta'(\text{Tr}h) = g'^{j\bar{k}} \text{Tr}((R_{\bar{k}j} - R'_{\bar{k}j}) \cdot h) + g'^{j\bar{k}} \text{Tr}((\nabla_j' h \cdot h^{-1})(\nabla_{\bar{k}}' h))$$

Then:

$$\begin{aligned}
\Delta' \log(\text{Tr}h) &= g'^{j\bar{k}} \partial_j \partial_{\bar{k}} (\log \text{Tr}h) \\
&= g'^{j\bar{k}} \partial_j \frac{\partial_{\bar{k}}(\text{Tr}h)}{\text{Tr}h} \\
&= \frac{\Delta'(\text{Tr}h)}{\text{Tr}h} - g'^{j\bar{k}} \frac{(\partial_{\bar{k}} \text{Tr}h)(\partial_j \text{Tr}h)}{(\text{Tr}h)^2} \\
&= \frac{\Delta'(\text{Tr}h)}{\text{Tr}h} - \frac{\|\nabla' \text{Tr}h\|^2}{(\text{Tr}h)^2}
\end{aligned}$$

Plugging in  $\Delta'(\text{Tr}h)$ :

$$\begin{aligned}
\Delta' \log(\text{Tr}h) &= \frac{g'^{j\bar{k}} \text{Tr}((R_{\bar{k}j} - R'_{\bar{k}j}) \cdot h)}{\text{Tr}h} + \\
&\quad \left\{ \frac{g'^{j\bar{k}} \text{Tr}((\nabla_j' h \cdot h^{-1})(\nabla_{\bar{k}}' h))}{\text{Tr}h} - \frac{g'^{j\bar{k}} \partial_j \text{Tr}h \cdot \partial_{\bar{k}} \text{Tr}h}{(\text{Tr}h)^2} \right\}
\end{aligned}$$

Basic observations:

- 1). Yau's inequality: the term in the bracket  $\{ \cdot \} \geq 0$  (to be proved)
- 2). The  $(g')^{j\bar{k}} \text{Tr}(R'_{\bar{k}j} \cdot h)$  can be simplified as follows:

$$\begin{aligned}
(g')^{j\bar{k}} \text{Tr}(R'_{\bar{k}j} \cdot h) &= (g')^{j\bar{k}} R'_{\bar{k}j}{}^\ell{}_m h^m{}_\ell \\
&= (\text{Ric})^\ell{}_m h^m{}_\ell \\
&= (\text{Ric})^\ell{}_m g^{m\bar{p}} g'_{\bar{p}\ell} \\
&= (\text{Ric})_{\bar{p}m} g^{m\bar{p}} \quad (\bullet)
\end{aligned}$$

However, we know that  $(\text{Ric})_{\bar{k}j} = -\partial_j \partial_{\bar{k}} \log \det(g'_{\bar{q}p})$ . Thus

$$(\text{Ric})_{\bar{p}m} = -\partial_m \partial_{\bar{p}} \log \det(g'_{\bar{q}p})$$

and by the Monge - Ampère equation

$$\det(g'_{\bar{q}p}) = \det(g_{\bar{q}p}) \cdot e^{+F - Ct}$$

we have:

$$\begin{aligned} (\text{Ric}')_{\bar{p}m} &= -\partial_m \partial_{\bar{p}} \log \det(g_{\bar{p}p}) - \partial_m \partial_{\bar{p}} (tF - Ct) \\ &= \underbrace{(\text{Ric})_{\bar{p}m} - \partial_m \partial_{\bar{p}} tF}_{\text{(unlike Ric', this quantity is prescribed)}} \end{aligned}$$

Substitute this into (•), we get:

$$\begin{aligned} (g')^{j\bar{k}} \text{Tr}(R'_{\bar{k}j} \cdot h) &= \theta^{m\bar{p}} ((\text{Ric})_{\bar{p}m} - \partial_m \partial_{\bar{p}} (tF)) \\ &= R - \Delta(tF) \end{aligned}$$

where  $R$  denotes the scalar curvature of  $(g_{\bar{k}j})$ .

3). We estimate

$$\begin{aligned} (g')^{j\bar{k}} \text{Tr}(R_{\bar{k}j} \cdot h) &= (g')^{j\bar{k}} (R_{\bar{k}j}{}^{\ell m} h^m{}_{\ell}) \quad (\text{Recall that } h = g^{-1}g' \Rightarrow (g')^{-1} = h^{-1}g^{-1}) \\ &= (h^{-1})^{j_r} g^{r\bar{k}} R_{\bar{k}j}{}^{\ell m} h^m{}_{\ell} \\ &= (h^{-1})^{j_r} R^r{}_j{}^{\ell m} h^m{}_{\ell} \end{aligned}$$

Since  $R^r{}_j{}^{\ell m}$  is the curvature tensor of a known metric (a.k.a. bi-sectional curvature), it can be bounded from below by  $\min_{i,j} |R^i{}_j{}^i{}_j| \triangleq B$

$$(g')^{j\bar{k}} \text{Tr}(R_{\bar{k}j} \cdot h) \geq -B \text{Tr}(h) \text{Tr}(h^{-1})$$

(This can be proved, for instance, by diagonalizing  $h$ , then

$$\text{l.h.s.} = \lambda_j^{-1} R^j{}_j{}^{\ell \ell} \lambda_{\ell} \quad )$$

From these observations, we have

$$\begin{aligned} \Delta' \text{Tr}(\log h) &\geq -B(\text{Tr} h^{-1}) - \frac{1}{\text{Tr} h} (R - \Delta(tF)) \\ &\geq -B(\text{Tr} h^{-1}) - C_3 (\text{Tr} h)^{-1} \\ &= -C_4 \text{Tr} h^{-1} \end{aligned}$$

(the last inequality follows by diagonalizing  $h$  (positive endomorphism), and

$$(\text{Tr} h)^{-1} = (\lambda_1 + \dots + \lambda_n)^{-1} \leq \lambda_i^{-1} \leq \text{Tr}(h^{-1}) \quad ).$$

This is close to the desired inequality, but the constant has wrong sign. That's why we need to subtract  $A \Delta' \varphi$ :

$$\begin{aligned} \Delta' \varphi &= (g')^{j\bar{k}} \partial_j \partial_{\bar{k}} \varphi \\ &= (g')^{j\bar{k}} ((g')_{\bar{k}j} - g_{\bar{k}j}) \\ &= n - \text{Tr}(h^{-1}) \quad (h^{-1} = g^{-1} \cdot g) \end{aligned}$$

Altogether, choose  $A \gg 0$ , we have:

$$\begin{aligned} \Delta(\log \text{Tr} h - A\varphi) &\geq -C_4 \text{Tr} h^{-1} - A(n - \text{Tr} h^{-1}) \\ &= (A - C_4) \text{Tr} h^{-1} - A \cdot n \\ &\geq C_1 \text{Tr} h^{-1} - C_2, \end{aligned}$$

as claimed in (\*\*).

Proof of Yau's inequality.

This is a tensorial inequality, and thus it suffices to work in a coordinate system so that at  $z$ :

$$\begin{cases} g_{\bar{k}j}(z) = \delta_{\bar{k}j}, \\ g'_{\bar{k}j}(z) = \lambda_j \delta_{\bar{k}j} \quad (\lambda_j = 1 + \varphi_{j\bar{j}}), \\ \nabla'_p = \partial_p \text{ at } z. \end{cases}$$

Thus we want to show that:

$$(g'^{j\bar{k}})(\partial_{\bar{j}} \text{Tr} h)(\partial_{\bar{k}} \text{Tr} h) \leq \text{Tr} h \cdot \{ (g')^{p\bar{q}} \text{Tr}(\nabla'_p h \cdot h^{-1} \cdot \nabla'_{\bar{q}} h) \}$$

i.e.

$$\begin{aligned} \sum_{p,\bar{q}} \frac{\delta^{p\bar{q}}}{(1+\varphi_{p\bar{p}})} (\partial_p (\sum_i (1+\varphi_{i\bar{i}}))) \partial_{\bar{q}} (\sum_j (1+\varphi_{j\bar{j}})) &= \sum_p \frac{1}{(1+\varphi_{p\bar{p}})} \sum_{i,j} (\partial_p \varphi_{i\bar{i}}) (\partial_{\bar{p}} \varphi_{j\bar{j}}) \\ &\leq \{ \sum_i (1+\varphi_{i\bar{i}}) \} \cdot \left\{ \sum_{p,\bar{q}} \frac{\delta^{p\bar{q}}}{(1+\varphi_{p\bar{p}})} \sum_j (\partial_p (1+\varphi_{j\bar{j}})) \frac{1}{1+\varphi_{j\bar{j}}} \partial_{\bar{q}} (1+\varphi_{j\bar{j}}) \right\} \\ &= \left\{ \sum_i (1+\varphi_{i\bar{i}}) \right\} \cdot \left\{ \sum_p \frac{1}{(1+\varphi_{p\bar{p}})} \cdot \sum_j \frac{\partial_p \varphi_{j\bar{j}} \cdot \partial_{\bar{p}} \varphi_{j\bar{j}}}{(1+\varphi_{j\bar{j}})} \right\} \end{aligned}$$

This is just the Cauchy-Schwartz inequality. Indeed, write:

$$\begin{aligned} \sum_p \frac{1}{(1+\varphi_{p\bar{p}})} \sum_{i,j} (\partial_p \varphi_{i\bar{i}}) (\partial_{\bar{p}} \varphi_{j\bar{j}}) &= \sum_{i,j} \left\{ \sum_p \left( \frac{\partial_p \varphi_{i\bar{i}}}{(1+\varphi_{p\bar{p}})^{\frac{1}{2}}} \right) \left( \frac{\partial_{\bar{p}} \varphi_{j\bar{j}}}{(1+\varphi_{p\bar{p}})^{\frac{1}{2}}} \right) \right\} \\ &\leq \sum_{i,j} \left( \sum_p \frac{|\partial_p \varphi_{i\bar{i}}|^2}{(1+\varphi_{p\bar{p}})} \right)^{\frac{1}{2}} \left( \sum_p \frac{|\partial_{\bar{p}} \varphi_{j\bar{j}}|^2}{(1+\varphi_{p\bar{p}})} \right)^{\frac{1}{2}} \quad (\text{Cauchy-Schwartz in } p, \text{ for fixed } i,j) \\ &= \left( \sum_i \left( \sum_p \frac{|\partial_p \varphi_{i\bar{i}}|^2}{(1+\varphi_{p\bar{p}})} \right)^{\frac{1}{2}} \right) \cdot \left( \sum_j \left( \sum_p \frac{|\partial_{\bar{p}} \varphi_{j\bar{j}}|^2}{(1+\varphi_{p\bar{p}})} \right)^{\frac{1}{2}} \right) \\ &= \left( \sum_i \left( \sum_p \frac{|\partial_p \varphi_{i\bar{i}}|^2}{(1+\varphi_{p\bar{p}})} \right)^{\frac{1}{2}} \right)^2 \\ &= \left( \sum_i (1+\varphi_{i\bar{i}}) \right)^{\frac{1}{2}} \cdot \left( \sum_p \frac{|\partial_p \varphi_{i\bar{i}}|^2}{(1+\varphi_{i\bar{i}})(1+\varphi_{p\bar{p}})} \right)^{\frac{1}{2}} \\ &\leq \left\{ \sum_i (1+\varphi_{i\bar{i}}) \right\} \cdot \left\{ \sum_{i,p} \frac{|\partial_p \varphi_{i\bar{i}}|^2}{(1+\varphi_{i\bar{i}})(1+\varphi_{p\bar{p}})} \right\} \quad (\text{Cauchy-Schwartz in } i) \end{aligned}$$

Proof of the  $C^3$ -estimate

We shall prove the bound:  $S \leq \text{Const}$ , where  $S = |\nabla' \bar{\nabla}' \nabla' \varphi|^2$ . More precisely,  $S = (g')^{j\bar{m}} (g')^{p\bar{k}} \varphi_{j\bar{k}\bar{\ell}} \overline{\varphi_{m\bar{p}q}} (g')^{\ell\bar{q}}$ , with  $\varphi_{j\bar{k}\bar{\ell}} = \nabla_{\bar{\ell}}' \varphi_{\bar{k}j}$  w/r.t. the metric  $g'_{\bar{k}j} = g_{\bar{k}j} + \partial_j \bar{\partial}_{\bar{k}} \varphi$ . The proof is due to Cartan, Nirenberg and Yau.

Claim:  $\exists$  constants  $A_1, C_5, C_6 > 0$  s.t.

$$\Delta'(S - A_1 \Delta \varphi) \geq C_5 S - C_6 \quad (***)$$

Assuming this claim, we may apply the maximum principal again. At a point  $p$ , where  $S - A_1 \Delta \varphi$  attains its maximum, similar as before, we have

$$\Delta'(S - A_1 \Delta \varphi)(p) \leq 0$$

$$\Rightarrow S(p) \leq C_7$$

Then for a general  $z \in M$ ,

$$S(z) - A_1 \Delta \varphi(z) \leq S(p) - A_1 \Delta \varphi(p)$$

$$\begin{aligned} \Rightarrow S(z) &\leq S(p) + A_1 (\Delta \varphi(z) - \Delta \varphi(p)) \\ &\leq C_7 + A_1 \cdot 2 \cdot \|\Delta \varphi\|_{C^0} \\ &\leq C_8 \end{aligned}$$

Proof of (\*\*\*)

For this, observe that:  $S = |\nabla' h \cdot h^{-1}|_{g'}^2$ , where  $h^j_{\bar{k}} = g^{j\bar{p}} g'_{\bar{p}k}$ . Indeed

$$\begin{aligned} |\nabla' h \cdot h^{-1}|_{g'}^2 &= \nabla'_j h^a_{\bar{b}} (h^{-1})^b_{\bar{c}} \overline{\nabla'_k h^d_{\bar{e}} (h^{-1})^e_{\bar{f}}} (g')^{j\bar{k}} (g')^{\bar{a}d} (g')^{c\bar{f}} \\ &= (\nabla'_j g^{a\bar{u}}) g'_{\bar{u}b} g'^{b\bar{v}} g'_{\bar{v}c} \overline{(\nabla'_k g^{d\bar{s}}) g'_{\bar{s}e} g'^{e\bar{r}} g'_{\bar{r}f}} \cdot (g')^{j\bar{k}} (g')^{\bar{a}d} (g')^{c\bar{f}} \\ &= (\nabla'_j g^{a\bar{u}} \cdot g'_{\bar{u}c}) \cdot \overline{(\nabla'_k g^{d\bar{r}}) g'_{\bar{r}f}} \cdot (g')^{j\bar{k}} (g')^{\bar{a}d} (g')^{c\bar{f}} \\ &= (-g^{a\bar{m}} \nabla'_j (g'_{\bar{m}n}) g'^{n\bar{u}} g'_{\bar{u}c}) \cdot \overline{(-g^{d\bar{s}} \nabla'_k (g'_{\bar{s}t}) g'^{t\bar{r}} \cdot g'_{\bar{r}f})} \cdot (g')^{j\bar{k}} \cdot (g')^{\bar{a}d} \cdot (g')^{c\bar{f}} \\ &= (g^{a\bar{m}} \nabla'_j (\partial_{\bar{m}} \partial_c \varphi) \cdot \overline{g^{d\bar{s}} \nabla'_k (\partial_{\bar{s}} \partial_f \varphi)}) \cdot (g')^{j\bar{k}} \cdot (g')^{\bar{a}d} \cdot (g')^{c\bar{f}} \\ &= \end{aligned}$$



Now we compute  $\Delta'S$ :

$$\begin{aligned}\Delta'S &= \Delta' |\nabla'h \cdot h^{-1}|_{g'}^2 \\ &= \Delta' \langle \nabla'h \cdot h^{-1}, \nabla'h \cdot h^{-1} \rangle\end{aligned}$$

where  $\Delta' = (g')^{j\bar{k}} \nabla_j \nabla_{\bar{k}}$ . Then:

$$\begin{aligned}\Delta'S &= (g')^{j\bar{k}} (\nabla_j \langle \nabla_{\bar{k}} \nabla'h \cdot h^{-1}, \nabla'h \cdot h^{-1} \rangle + \nabla_j \langle \nabla'h \cdot h^{-1}, \nabla_{\bar{k}} \nabla'h \cdot h^{-1} \rangle) \\ &= (g')^{j\bar{k}} (\langle \nabla_j \nabla_{\bar{k}} \nabla'h \cdot h^{-1}, \nabla'h \cdot h^{-1} \rangle + \langle \nabla_{\bar{k}} \nabla'h \cdot h^{-1}, \nabla_j \nabla'h \cdot h^{-1} \rangle \\ &\quad + \langle \nabla_j \nabla'h \cdot h^{-1}, \nabla_{\bar{k}} \nabla'h \cdot h^{-1} \rangle + \langle \nabla'h \cdot h^{-1}, \nabla_j \nabla_{\bar{k}} \nabla'h \cdot h^{-1} \rangle) \\ &= \langle \Delta' \nabla'h \cdot h^{-1}, \nabla'h \cdot h^{-1} \rangle + \langle \nabla'h \cdot h^{-1}, \bar{\Delta}'(\nabla'h \cdot h^{-1}) \rangle \\ &\quad + |\bar{\nabla}'(\nabla'h \cdot h^{-1})|^2 + |\nabla'(\nabla'h \cdot h^{-1})|^2\end{aligned}$$

Next, note that

$$\begin{aligned}\Delta'(\nabla_j'h \cdot h^{-1}) &= g'^{l\bar{k}} \nabla_l' \nabla_{\bar{k}}'(\nabla_j'h \cdot h^{-1}) \\ &= g'^{l\bar{k}} \nabla_l'(R_{\bar{k}j} - R'_{\bar{k}j}) \quad (\text{by an earlier computation in } C^2\text{-estimate}) \\ &= g'^{l\bar{k}} (\nabla_l'R_{\bar{k}j} - \nabla_j'R'_{\bar{k}l}) \quad (\text{by the 2}^{nd}\text{- Bianchi's identity: } \nabla_j'R_{\bar{k}l} = \nabla_l'R_{\bar{k}j}) \\ &= g'^{l\bar{k}} \nabla_l'R_{\bar{k}j} - \nabla_j'(Ric')\end{aligned}$$

Up to now, we haven't used the condition that  $(g'_{\bar{k}j})$  solves the MA equation, which says that  $Ric'$  is known:

$$\begin{aligned}Ric'_{\bar{a}b} &= -\partial_b \partial_{\bar{a}} \log \det(g'_{\bar{p}q}) \\ &= Ric_{\bar{a}b} - \partial_b \partial_{\bar{a}} F \cdot t\end{aligned}$$

Thus  $\nabla_j'(Ric')$  is bounded and so is  $g'^{l\bar{k}} \nabla_l'R_{\bar{k}j}$ . It follows that:

$$\|\langle \Delta' \nabla'h \cdot h^{-1}, \nabla'h \cdot h^{-1} \rangle\|_{g'} \leq C_9 S + C_{10}$$

(There is some subtle details here: the norm here is taken w.r.t.  $g'_{\bar{k}j}$ , which is a metric to be solved. However, by the  $C^2$ -estimate,  $\{g'_{\bar{k}j}\}$  and  $\{g_{\bar{k}j}\}$  are uniformly equivalent:

$$\left. \begin{aligned}Tr h &= g^{j\bar{k}} g'_{\bar{k}j} = n + \Delta\varphi \\ \Delta\varphi &\text{ bounded}\end{aligned} \right\} \Rightarrow \text{the sum of eigenvalues of } h \text{ is bounded:}$$

Moreover,  $\prod \lambda_i$  is fixed  $\Rightarrow$  Each  $\lambda_i$  is bounded. Thus  $h$  is a uniformly bounded endomorphism  $\Rightarrow g' = gh$  is uniformly bounded.)

Hence:

$$\begin{aligned} \Delta' S &\geq \langle \Delta' \nabla' h \cdot h^{-1}, \nabla' h \cdot h^{-1} \rangle + \langle \nabla' h \cdot h^{-1}, \bar{\Delta}'(\nabla' h \cdot h^{-1}) \rangle \\ &\geq -C_{11} S - C_{12}. \end{aligned}$$

where we leave out the positive terms  $|\bar{\nabla}'(\nabla' h \cdot h^{-1})|^2 + |\nabla'(\nabla' h \cdot h^{-1})|^2$ , and  $\bar{\Delta}'(\nabla' h \cdot h^{-1}) = \Delta' + \text{Curvature terms of } (g_{\bar{a}\bar{b}})$ . The curvature terms are bounded again by the  $C^2$ -estimate.

Next, we compute  $\Delta' \Delta \varphi$ : (c.f.  $C^2$ -estimate)

$$\begin{aligned} \Delta'(\Delta \varphi) &= \Delta'(Tr h) \\ &= g'^{p\bar{q}} (Tr(\nabla_{\bar{q}}(\nabla_p h \cdot h^{-1} \cdot h))) \\ &= g'^{p\bar{q}} (Tr(R_{\bar{q}p} \cdot h - R'_{\bar{q}p} \cdot h) + Tr(\nabla_p h \cdot h^{-1} \cdot \nabla_{\bar{q}} h)) \\ &= g'^{p\bar{q}} (\underbrace{Tr(R_{\bar{q}p} \cdot h)}_{\text{known}}) - (R - \Delta F_t) + g'^{p\bar{q}} Tr(\underbrace{\nabla_p h \cdot h^{-1} \cdot \nabla_{\bar{q}} h}_{\text{of order } S}). \end{aligned}$$

By our  $C^2$ -estimate,  $\{g_{p\bar{q}}\}$  and  $\{g'_{p\bar{q}}\}$  are equivalent metrics, and the last term is of order  $S$ , we have:

$$\Delta'(\Delta \varphi) \geq -(C_{13} S + C_{14})$$

Combining these two estimates as we did for  $C^2$ -estimate, we obtain the claimed inequality (\*\*\*) .

A final remark. Why these estimates imply the desired convergence.

Differentiating the MA equation w.r. t.  $\nabla_{\bar{e}}$  gives

$$(g')^{j\bar{k}} \nabla_{\bar{e}} \nabla_{\bar{j}} \nabla_{\bar{k}} \varphi = \text{known quantity}$$

$\Rightarrow \nabla_{\bar{e}} \varphi$  satisfies the Laplacian equation with  $C^0$  r.h.s. By the classical Schauder elliptic regularity, which is part of classical linear PDE theory, one gets that  $\nabla_{\bar{e}} \varphi$  is of class  $C^{1,\nu}$ ,  $\forall \nu \in (0,1)$ . This in turn implies that  $\varphi$  is of class  $C^{2,\nu}$ , ... (Black box). This gives the desired convergence and we may apply the Ascoli-Arzelà thm.